

DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY
GEORGE OTIS SMITH, DIRECTOR

BULLETIN 408

A RECONNAISSANCE
OF
SOME MINING CAMPS IN ELKO, LANDER,
AND EUREKA COUNTIES, NEVADA

BY

WILLIAM H. EMMONS



WASHINGTON
GOVERNMENT PRINTING OFFICE
1910



CONTENTS.

	Page.
Introduction.....	9
Scope of report.....	9
Field work and acknowledgments.....	9
Bibliography.....	11
Topography, climate, and vegetation.....	12
History.....	14
Production.....	14
General geology.....	15
Sedimentary rocks.....	15
Introduction.....	15
Cambrian system.....	17
Prospect Mountain quartzite.....	17
Eldorado limestone.....	17
Secret Canyon shale.....	18
Hamburg limestone.....	18
Dunderberg shale.....	18
Ordovician.....	18
Pogonip limestone.....	18
Eureka quartzite.....	18
Lone Mountain limestone.....	18
Devonian system.....	19
Carboniferous system.....	20
White Pine shale.....	20
Diamond Peak quartzite.....	20
Lower coal measures.....	20
Weber conglomerate.....	20
Upper coal measures.....	21
Mesozoic rocks.....	21
Tertiary rocks.....	21
Eocene.....	21
Miocene.....	22
Pliocene.....	22
Quaternary deposits.....	23
Glacial deposits.....	23
Alluvial deposits.....	23
Igneous rocks.....	24
Intrusive granitic rocks.....	24
Distribution and composition.....	24
Contact relations and age.....	27
Porphyries associated with intrusive granitic rocks.....	29
Tertiary eruptive rocks.....	30
Rhyolite.....	30
Basalt.....	31
Andesite.....	32
Dacite.....	33
Succession of the Tertiary eruptive rocks.....	33
Age of the Tertiary eruptive rocks.....	34

	Page.
General geology—Continued.	
Deformation of the rocks.....	35
Résumé of geologic history.....	37
Ore deposits.....	39
General statement.....	39
Earlier deposits (Cretaceous?).....	40
General statement.....	40
Contact-metamorphic deposits.....	40
Irregular replacement deposits in limestone.....	41
Replacement veins and sheeted zones in limestone and in shale.....	42
Fissure veins in quartzite.....	42
Fissure veins in the older intrusive rocks.....	43
Later deposits (Miocene).....	43
Fissure veins and sheeted zones in andesite.....	44
Fissure veins and fracture zones in rhyolite.....	44
Placer deposits.....	45
Prospecting.....	46
Mountain ranges and mining districts.....	47
Owyhee Bluffs.....	47
General features.....	47
Geologic features.....	47
Midas or Gold Circle district.....	48
Location and history.....	48
Geology.....	48
General outline.....	48
Hydrothermal metamorphism.....	49
Fissuring and faulting.....	50
Ore deposits.....	50
General features.....	50
Prospecting the lodes.....	52
Ore shoots.....	52
Secondary enrichment.....	53
Résumé of geology.....	53
Mine descriptions.....	54
Rex mine.....	54
Gold Crown lode.....	54
St. Paul-Banner lode.....	56
Golden Chariot claim.....	56
Esmeralda mine.....	56
Water Witch mine.....	56
Elko Prince lode.....	57
Midas mine.....	57
Gold Circle claim.....	57
Belvoir claim.....	57
Eastern Star mine.....	57
Independence Range.....	57
General features.....	57
Tuscarora.....	58
History.....	58
Ore deposits.....	60
Silver lodes.....	60
Gold deposits.....	61
Falcon mine.....	62
Cornucopia.....	63
Good Hope district.....	65

Mountain ranges and mining districts—Continued.	Page.
Burner Hills.....	66
General features.....	66
Mint mine.....	67
Centennial Range.....	67
General features.....	67
Ore deposits.....	69
Lime Mountain.....	70
Columbia and Aura.....	71
General statement.....	71
Mine descriptions.....	71
Infidel mine.....	71
Big Four mine.....	72
Columbia Queen mine.....	72
Blue Jacket mine.....	72
Jack Pot mine.....	73
California mine.....	74
Tiger lode.....	74
Humboldt mine.....	74
Polaris mine.....	74
Aura King mine.....	74
Edgemont.....	75
Lucky Girl group.....	75
Bull Run mine.....	79
Mountain City.....	80
Location and history.....	80
Geology.....	80
Ore deposits.....	81
Mine descriptions.....	82
Protection mine.....	82
Resurrection mine.....	82
Nelson mine.....	83
Mountain City mine.....	83
Van Duzer Creek placers.....	84
Lone Mountain.....	84
Cortez Range between the Carlin and Dalton peaks.....	86
General features.....	86
Mine descriptions.....	87
Nevada Star mine.....	87
Copper King claim.....	87
Richmond district.....	87
Lynn district.....	87
Pinyon Range.....	88
General features.....	88
Bullion.....	89
Geology and ore deposits.....	89
Mine descriptions.....	93
Standing Elk mine.....	93
Tripoli mine.....	93
Red Bird mine.....	94
Copper Belle mine.....	94
Delmas mine.....	94
Other claims.....	95

Mountain ranges and mining districts—Continued.

	Page.
Pinyon Range—Continued.	
Mineral Hill.....	95
Location and history.....	95
Geology.....	96
Ore deposits.....	97
Alpha.....	99
Cortez Range south of Humboldt River.....	99
General features.....	99
Geologic features.....	100
Cortez and Mill Canyon district.....	100
Location and history.....	100
Geology.....	101
Cortez mines.....	103
Garrison mine.....	103
Valley View mine.....	106
White Horse turquoise mine.....	106
Mill Creek mines.....	106
General features.....	106
Bullion Hill mines.....	107
Empire State mine.....	108
Hidden Treasure mine.....	108
Lewis Canyon claims.....	108
Falconer and Berlin mines.....	109
Caledonia mine.....	110
Safford district.....	110
General features.....	110
Mine descriptions.....	111
Zenoli mine.....	111
Onondaga mine.....	111
Ruby claim.....	112
Humboldt mine.....	112
Bonanza mine.....	112
Pittsburg and Palisade tunnel.....	112
West iron mine.....	112
Shoshone Range.....	113
General features.....	113
Geologic features.....	114
Tenabo.....	114
General statement.....	114
Geology.....	115
Ore deposits.....	116
Mine descriptions.....	116
Little Gem mine.....	116
Phoenix mine.....	117
Gold Quartz mine.....	117
Violet claim.....	118
Two Widows claim.....	118
Lander.....	118
General features.....	118
Mine descriptions.....	118
Bonnie Jean mine.....	118
Silver Prize vein.....	119
Silver Side mine.....	119

Mountain ranges and mining districts—Continued.

	Page.
Shoshone Range—Continued.	
Mud Springs.....	119
General features.....	119
Mine descriptions.....	119
Triumph mine.....	119
Big Bug claim.....	120
Bridal Wreath claim.....	120
Uncle Sam claim.....	120
Grey Eagle mine.....	120
Hilltop.....	120
Maysville mine.....	121
Lewis and Dean.....	122
Location and history.....	122
Geology and ore deposits.....	122
Mine descriptions.....	123
Morning Star and Pittsburg mines.....	123
Starr Grove mine.....	126
Betty O'Neil mine.....	126
Index.....	127

ILLUSTRATIONS.

	Page.
PLATE I. Topographic map of southern part of area.....	60
II. Topographic map of south-central part of area.....	64
III. Topographic map of north-central part of area.....	68
IV. Topographic map of northern part of area.....	72
V. Geologic map of an area in northeastern Nevada, between meridians 116° and 117° and parallels 40° and 41° 28'.....	In pocket.
FIGURE 1. Index map showing location of area described.....	10
2. Outline map showing areas covered by Plates I, II, III, and IV...	13
3. Geologic sketch map of the Gold Circle mining district.....	51
4. Plan of Rex mine, 65-foot level, Midas (Gold Circle) district.....	55
5. Geologic sketch showing Panther vein, Cornucopia district.....	63
6. Plan of adit level of Buckeye and Ohio mine, Good Hope district..	66
7. Cross section of Big Four lode, Columbia.....	72
8. Plan of tunnel level, Columbia Queen mine, Columbia.....	73
9. Sketch of wall, level 8, 1,200 feet from portal, Lucky Boy mine, Edgemont.....	76
10. Section S: 40° E. through Lucky Girl vein, Edgemont.....	77
11. Sketch of the northwest wall, level 8, Lucky Boy mine, Edgemont.	77
12. Cross section of Lucky Boy vein, Edgemont.....	78
13. Generalized section through Bull Run mine, S. 10° E. from tun- nel 1 to tunnel 4.....	79
14. Cross section through Ravens Nest, Pinyon Range.....	89
15. Sketch showing the relation of the ore zone to granodiorite and lime- stone on the Sweepstakes claim of the Delmas group, Bullion district.....	92
16. Sketch illustrating the structure at Mineral Hill.....	96
17. Cross section of Star Chamber stope, Mineral Hill.....	97
18. Section eastward through Tenabo Peak, showing underground workings at Garrison mine.....	102
19. Section through Garrison mine, Cortez.....	105
20. Cross section of Arctic mine, Cortez district. Near top of incline..	106
21. Cross section of ore body, Falconer mine, Mill Canyon.....	109
22. Cross section along Mayo incline, Morning Star lode, Dean.....	124

A RECONNAISSANCE OF SOME MINING CAMPS IN ELKO, LANDER, AND EUREKA COUNTIES, NEVADA.

By WILLIAM H. EMMONS.

INTRODUCTION.

SCOPE OF REPORT.

The following report gives the results of a reconnaissance of the mining districts in northeastern Nevada which are included between the one hundred and sixteenth and one hundred and seventeenth meridians and the fortieth and forty-second parallels. The eastern boundary of the area crosses the Bullion and Lone Mountain mining districts, and is about 12 miles west of Elko and 6 miles east of Carlin; the northern boundary is the Idaho line; and the western boundary passes about 12 miles west of the Gold Circle district and about 3 miles west of Battle Mountain. The general outline of the area and its position with respect to neighboring places are shown on the index map (fig. 1). As thus defined it includes about 7,000 square miles, embracing the western part of Elko County and the northern parts of Lander and Eureka counties. The report includes brief descriptive notes on the geology and ore deposits of the following camps: Gold Circle (Midas), Tuscarora, Lime Mountain, Edgemont, Aura, Columbia, Mountain City, Van Duzer Creek, Cornucopia, Good Hope, Burner, Falcon, Lone Mountain, Palisade (Safford district), Bullion (Railroad district), Mineral Hill, Alpha, Lynn, Cortez, Mill Canyon, Dean, Lewis, Maysville, Pittsburg, Hilltop, Tenabo, Lander, Mud Springs, and Grey Eagle.

FIELD WORK AND ACKNOWLEDGMENTS.

The work in the field was done between July 15 and October 20, 1908. During this time all the important mining districts were visited, but no geologic mapping was undertaken except that incidental to examination of the ore deposits. The reports of the United States Geological Exploration of the Fortieth Parallel, which covered about two-thirds of the area described herein, have been of great service as an aid in the study of the ore deposits. Plate V is a generalized reproduction, with some minor changes, of a part of Plate IV from the atlas of the Fortieth Parallel Survey.




FIGURE 1.—Index map showing location of area described.

A great many of the mines were inaccessible when the camps were visited. Some of them had not been worked since the eighties, when silver mining was at high tide. The notes on such mines are of the most general character and treat mainly the surface geology and the general relations of the ore deposits.

The writer is indebted to Mr. Waldemar Lindgren, in charge of the section of metalliferous ore deposits of the United States Geological Survey, for criticism and advice, and to the mine owners, operators, and prospectors of the area, who without exception have offered every facility to aid the work.

BIBLIOGRAPHY.

The following publications include the most important literature relating to the area described and adjacent territory in Nevada. Many notes of great historical interest may be found in the early reports to the United States Government by R. W. Raymond and J. Ross Brown, and in later reports of the Bureau of the Mint. A pamphlet treating of the mineral wealth of Elko County, published in 1907 and distributed gratis by the Chamber of Commerce of Elko, Nev., contains interesting details regarding some of the mines.

BANCROFT, HUBERT HOWE. History of Nevada, p. 322. A reliable and entertaining account of the history of Nevada from the earliest settlements to 1888. Includes a narrative of the development of the Comstock lode and other important districts, and some valuable data concerning the early history of Elko, Lander, and Eureka counties.

CURTIS, J. S. Silver-lead deposits of Eureka, Nev. Mon. U. S. Geol. Survey, vol. 7, 1884: A study of the geology and ore deposits of the Eureka district, with detailed descriptions of the mines.

GILBERT, G. K. Lake Bonneville. Mon. U. S. Geol. Survey, vol. 1, 1890; also Rept. Geog. and Geol. Surveys W. 100th Mer., vol. 3, 1875. Contain some valuable generalizations respecting geology of Nevada. The area described in the present reconnaissance was not covered by Lake Bonneville.

HAGUE, ARNOLD. Geology of the Eureka district, Nevada, with an atlas. Mon. U. S. Geol. Survey, vol. 20, 1892. Describes the geology and ore deposits of the Eureka district and gives the best section of Paleozoic sedimentary rocks available for northeast Nevada.

KING, CLARENCE. Geological exploration of the fortieth parallel. This great work, including six volumes and an atlas, is the result of the joint labors of Clarence King, S. F. Emmons, Arnold Hague, James Hague, and others. The area described is a belt nearly 2° wide along the Southern Pacific and Union Pacific railroads from a point east of Cheyenne, Wyo., to a point west of Pyramid Lake, near the east front of the Sierra Nevada. It includes the central and southern portions of the area covered in the present reconnaissance, and the reports are the source of many of the geologic data given herein. References to special parts of this work are made in appropriate places.

LINDGREN, WALDEMAR. The geological features of the gold production in North America. Trans. Am. Inst. Min. Eng., vol. 33, 1902, p. 790. Discusses the precious-metal production of Nevada and suggests that Tuscarora affords a connecting link between the Tertiary veins of Nevada and those of Idaho.

- LOUDERBACK, GEORGE DAVIS. Basin range structure of the Humboldt region. *Bull. Geol. Soc. America*, vol. 15, 1904, pp. 289-346. Concludes that the present topography of certain of the basin ranges is due to faulting, and modified to only a slight extent by erosion.
- McCLELLAN, E. C. Map of Elko County, Nev. A reliable hachure map of Elko County, a part of which is reproduced herein. Published by the author, Elko, Nev.
- RUSSELL, I. C. Geological history of Lake Lahontan, a Quaternary lake of northwestern Nevada. *Mon. U. S. Geol. Survey*, vol. 11, 1885. Contains some generalizations respecting the geology of Nevada.
- SPURR, J. E. Origin and structure of the basin ranges. *Bull. Geol. Soc. America*, vol. 12, 1901, pp. 217-270. Includes a comprehensive review of the literature relating to the structure of the basin ranges and concludes that their present topographic expression is due mainly to erosion.
- SPURR, J. E. The succession and relation of lavas of the Great Basin region. *Jour. Geology*, vol. 8, 1900, pp. 621-646. A summary of the age relations of the Tertiary rocks of Nevada, with some generalizations bearing on the problems of magmatic differentiation.
- SPURR, J. E. Descriptive geology of Nevada south of the fortieth parallel and adjoining portions of California. *Bull. U. S. Geol. Survey No. 208*, 1903. An account of a reconnaissance of the area indicated by the title, which includes a review of previous work done in the area, together with a geologic map. The area borders on the south that described in the present paper.

TOPOGRAPHY, CLIMATE, AND VEGETATION.

The area here described (see Pls. I, II, III, and IV; fig. 2) is a part of the Great Basin and has the characteristic basin topography. The country is a low plateau traversed by long, narrow mountain ranges, which trend northward and are approximately parallel. Between the mountains are wide valleys parallel to the ranges. The lowest parts of the plateau are a little less than 5,000 feet above sea level and the summits of the ranges are in the main between 8,000 and 10,000 feet. The drainage of the country has a gridiron arrangement, the chief stream being Humboldt River, which flows westward across the mountain ranges to the Carson Sink, receiving tributaries at right angles from the valleys between the north-south ranges on both sides. The north third of the area is drained by Owyhee River, a tributary of Snake River, which empties into the Columbia.

The summers are long and hot, the winters are rather severe for the latitude, and the air is dry. In the summer there are frequent showers of short duration and occasional cloud-bursts. In the winter the precipitation is mainly snow, which in the highest mountains accumulates in sufficient quantity to last well into the middle of the summer. The lower country is covered by sagebrush, which grows to great size, especially in the lower parts of the valleys. High on the mountain slopes there are several varieties of pine trees, junipers, and south of Humboldt River some mountain mahogany. Farming is carried on and wild grasses are cut along the Humboldt and its tributaries, and wherever there is sufficient water for irrigation




FIGURE 2.—Outline map showing areas covered by Plates I, II, III, and IV.

the valley flats produce a great variety of products, which find a good market at the ranches and mines near by. The country, especially the northern part, is an excellent range for cattle and sheep and contains some large and successful ranches.

HISTORY.

As early as 1825 the region of Humboldt River was known to British and American trappers, but settlement was slow until the discoveries of gold in California in 1849. Many of the gold seekers of that time traveled by way of the Humboldt Valley, which, during dry seasons, furnished a good roadbed, and consequently this country was well known before the more remote territory was explored. Many of the early settlers were Mormons from Utah, but after the discovery of the mines these were greatly outnumbered. The presence of silver on the Comstock lode in paying quantities became generally known in 1859, and discoveries of lode deposits were made about the same time in the Reese River country and elsewhere in Nevada. As a result of successful operations on the Comstock lode in 1860, prospecting became the principal business of a large part of the rapidly increasing population, and in the following ten years discoveries were made at nearly all the mining camps in the central and northern parts of the State.

Silver mining reached high tide in the seventies and eighties and suffered a rapid decline in the nineties. The more recent discoveries of gold and silver in the southern part of the State gave a new zest to prospecting, and in 1906 and 1907 nearly every one of the old camps was overhauled, the mines and dumps were sampled, and in some of them new ore bodies were discovered. Several new camps were established, among them Gold Circle, Lynn, Tenabo, Hilltop, and Aura. As a result of the business depression of 1907, there was a sudden cessation of prospecting and many of the mines were closed. A number of unworthy flotations were stopped and some of the more promising projects were hindered by the withdrawal of capital from the purchase of mining shares. Since money has become more plentiful, some of the prospecting operations have been taken up again. Historical data respecting the development of the various camps are given where these camps are described.

PRODUCTION.

The total production of the mining camps of this area can not be closely approximated. Official reports relating to the periods of greatest production for most of the camps were not accessible to the writer, and the government reports which are at hand do not give sufficient detail to permit more than the rudest kind of an estimate. The production of Tuscarora, the largest camp, will probably fall

between \$25,000,000 and \$40,000,000. From the remaining camps, among which are Cortez, Mineral Hill, Bullion, Dean, Cornucopia, Edgemont, Mountain City, and others, the total is about \$25,000,000 more. A conservative estimate thus gives a total production of about \$50,000,000 for the entire area. On the other hand, if maximum figures are used, based largely on hearsay, the estimate of production would be considerably greater, reaching about \$70,000,000. Perhaps one-quarter of the total production is gold, a large part of which is from the placers at Tuscarora and the Dexter mine and from the silver bullion from various silver-mining camps, nearly all of which produce some gold. Except for relatively small amounts of lead, copper, and zinc which have been shipped to smelters in recent years, the remaining three-fourths of the production is silver.

In 1907 the production of the area was approximately \$350,000. Of this amount Edgemont, Aura, and Mountain City produced \$161,834, of which \$145,525 was gold, \$16,004 silver, and the remainder lead and zinc. The larger portion of this production came from the Lucky Girl mine, at Edgemont. The mines at Bullion in the Railroad district, the Zenoli mine in the Safford district, and the Grey Eagle mine produced altogether \$106,252, of which \$2,857 was gold, \$77,800 silver, \$15,454 copper, \$7,926 lead, and \$2,195 zinc. The remainder of the production came from Mill Canyon, Cortez, Tenabo, and Lander.

GENERAL GEOLOGY.

SEDIMENTARY ROCKS.

INTRODUCTION.

The country covered by this reconnaissance is an area of Paleozoic sedimentary rocks cut by granodiorite and other intrusive rocks and overlain by Tertiary lake beds and lava flows. In this part of Nevada none of the sedimentary or igneous rocks have suffered deep-seated dynamic metamorphism such as results in the development of gneissoid or schistose structure and none of them have been deformed in the zone of flow. Around the intrusive granular rocks there is noticeable contact metamorphism, with garnet zones, especially in limestone and in shales. Because of the abundance of faults in this region it would be necessary to work with great detail to gain an adequate knowledge of the sedimentary column. Such work has been done in the Eureka district, where the Paleozoic column is best known, but as that work was undertaken after the Fortieth Parallel report had been completed, the sedimentary rocks in the country covered by the King report have not been studied in the light of the information gained at Eureka. Consequently, only broad statements can be made respecting the age of the sedimentary

rocks, and on account of the small number of described exposures containing fossils much confidence should not be placed in the correlations of the rocks which have been made in a very hasty

Geologic section in Eureka region, Nevada.

System.	Formation.	Thickness (feet).	Character.
Carboniferous.	Upper coal measures.....	500	Light-colored blue and drab limestones.
	Weber conglomerate.....	2,000	Coarse and fine conglomerates, with angular fragments of chert; layers of reddish-yellow sandstone.
	Lower coal measures.....	3,800	Heavy-bedded dark-blue and gray limestone, with intercalated bands of chert; argillaceous beds near the base.
	Diamond Peak quartzite....	3,000	Massive gray and brown quartzite, with brown and green shales at the summit.
	White Pine shale.....	2,000	Black argillaceous shales, more or less arenaceous, with intercalations of red and reddish-brown friable sandstone, changing rapidly from one locality to another; plant impressions.
Devonian.	Nevada limestone.....	6,000	Lower strata indistinctly bedded, saccharoidal texture, gray color, passing up into strata distinctly bedded, brown, reddish brown, and gray in color, in places finely striped, producing a variegated appearance; upper rocks massive, well bedded, bluish-black in color; highly fossiliferous.
Ordovician.	Lone Mountain limestone....	1,800	Black, gritty beds at the base, passing into a light-gray siliceous rock, with all traces of bedding obliterated; Trenton fossils at the base; <i>Halysites</i> in the upper portion.
	Unconformity.		
	Eureka quartzite.....	500	Compact vitreous quartzite, white, blue, passing into reddish tints near the base; indistinct bedding.
	Pogonip limestone.....	2,700	Interstratified limestone, argillites, and arenaceous beds at the base, passing into purer, fine-grained limestone of a bluish-gray color, distinctly bedded; highly fossiliferous.
Cambrian.	Dunderberg shale ^a	350	Yellow argillaceous shale; layers of chert nodules throughout the bed, but more abundant near the top.
	Hamburg limestone.....	1,200	Dark-gray and granular limestone; surface weathering rough and ragged; only slight traces of bedding.
	Secret Canyon shale.....	1,600	Yellow and gray argillaceous shales, passing into shaly limestone; near the top interstratified layers of shale and thinly bedded limestones.
	Eldorado limestone ^b	3,050	Gray, compact limestone; lighter in color than the Hamburg limestone, traversed with thin seams of calcite; bedding planes very imperfect.
	Prospect Mountain quartzite.	1,500	Bedded brownish-white quartzites, weathering dark brown; ferruginous near the base; intercalated thin layers of arenaceous shales; beds whiter near the summit.

^a This name replaces "Hamburg" shale. See Walcott, C. D., Smithsonian Misc. Coll., vol. 53, No. 1812, 1908, p. 184.

^b This name replaces "Prospect Mountain" limestone. Idem.

reconnaissance. Devonian fossils have been described^a from several places in the Pinyon Range, among them Chimney station, Hot Spring Creek, and Pinyon Pass. A fauna closely related to the Helderberg fauna has been found at White's ranch, about 20 miles north of Beowawe, and was described by Hall and Whitfield.^b Carboniferous fossils^c have been found at Moleen Peak, at Lone Mountain, and at Railroad Canyon, on the north end of the Diamond Range, which is about 15 miles east of Chimney station.^a

Plate V shows the general distribution of the sedimentary rocks for part of the area. The descriptions of the sedimentary rocks in the table on page 16 are after Hague and Walcott,^d based on the Eureka section, with modifications by E. O. Ulrich as to the age of the Pogonip and Lone Mountain limestones and the Eureka quartzite, by G. H. Girty as to the age of the White Pine shale, and by C. D. Walcott as to the "Hamburg" shale and "Prospect Mountain" limestone.

CAMBRIAN SYSTEM.

Prospect Mountain quartzite.—This formation is the earliest of the Paleozoic rocks and consists of 1,500 feet of bedded brownish-white quartzite weathering dark brown, but white near the top. The quartzites are ferruginous near the base and contain intercalated layers of arenaceous and micaceous shale only a few feet thick. The Prospect Mountain differs from the Eureka quartzite, the next overlying siliceous formation, in being more ferruginous and in general less uniform in texture, carrying throughout more or less clayey material, whereas the Eureka is a nearly pure, highly altered sandstone. No fossils were found in the Prospect Mountain quartzite.

Eldorado limestone.—The Eldorado ("Prospect Mountain") limestone overlies the Prospect Mountain quartzite and consists of 3,050 feet of gray compact limestone imperfectly bedded and of light color, which at some places is traversed by thin seams of calcite. It is difficult to define sharply the characteristic features of this formation, as changes were frequent in the deposition of the sediments, not only in vertical but also in lateral extension. In general, however, the formation has a light bluish-gray tint when observed over large areas, although nearly all colors from white to black are found in the limestone, which at the same time is characterized throughout its thickness by seams of calcite varying from one-half inch to 6 inches in width and locally forming a network of white bands. The limestone is crystalline and granular over wide areas and at many places stratification is not well shown. The *Olenellus* fauna is found in shaly beds at the base of the limestone.

^a U. S. Geol. Expl. 40th Par., vol. 3, 1870, p. 557.

^b Idem, p. 609.

^c Idem, vol. 1, 1878, p. 248.

^d Hague, Arnold, Mon. U. S. Geol. Survey, vol. 20, 1892, p. 13.

Secret Canyon shale.—The Eldorado limestone grades into the yellow and gray argillaceous Secret Canyon shale, which is 1,600 feet thick. Near the top are interstratified layers of shale and thinly bedded limestones.

Hamburg limestone.—The Hamburg limestone, which overlies the Secret Canyon shale, consists of 1,200 feet of dark-gray granular limestone that shows only traces of bedding. Layers of fine sandstone and hard cherty bands occur at irregular intervals.

Dunderberg shale.—The Dunderberg ("Hamburg") shale overlies the Hamburg limestone and consists of yellow argillaceous shale with layers of chert nodules throughout the bed but more abundant near the top. Across its broadest development it measures 350 feet, but it rarely maintains a uniform thickness for long distances. The Dunderberg shale carries a well-developed upper Cambrian fauna.

ORDOVICIAN SYSTEM.

Pogonip limestone.—The Dunderberg shale passes gradually into the Pogonip limestone, which is of Ordovician age. This formation is 2,700 feet thick and consists of argillites and arenaceous beds at the base, passing into purer fine-grained limestone of a bluish-gray color, distinctly bedded and highly fossiliferous. The limestone is rich in Ordovician fossils.

Eureka quartzite.—The Eureka quartzite, of Ordovician age, consists of 500 feet of compact vitreous quartzite, white or blue, passing into reddish tints near the base, with indistinct bedding. The formation is massive and very thickly bedded and has a tendency to form cliffs or escarpments. After the Eureka quartzite was deposited it was raised above sea level, but it was not greatly eroded, for in the sections studied it is everywhere present. During the subsequent subsidence it was not everywhere submerged, and so different formations will be found to overlie it at different places.

Lone Mountain limestone.—The change is everywhere abrupt between the Eureka quartzite and the Lone Mountain limestone, and the latter is not everywhere present above the Eureka and at some places is very thin. At Eureka the Lone Mountain limestone is 1,800 feet thick. The black gritty beds at the base pass into a light-gray siliceous rock without traces of bedding. Trenton fossils are found at the base and *Halysites* in the upper portion.

C. D. Walcott^a made the following section across Lone Mountain:

	Feet.
Dark-gray limestone, with brown and variegated layers interbedded; typical Devonian fauna (Nevada limestone).....	1,500
Siliceous bluish-gray limestone, breaking up into shaly bands carrying abundant fossils of the Lower Devonian (Nevada limestone).	200

^a Hague, Arnold, Geology of the Eureka district, Nevada: Mon. U. S. Geol. Survey, vol. 20, 1892, pp. 61-62.

	Feet.
Siliceous limestone, light brown, gray, and buff in color, with <i>Halysites catenulatus</i> near the base; passing up into beds almost white, with blue and gray tints, followed by alternating dark and light beds (Lone Mountain limestone).....	2,000
White quartzite (Eureka quartzite).....	200
Dark-gray limestone, massive bedding, with intercalated shaly layers carrying a typical Silurian [Ordovician] fauna (Pogonip limestone).....	300
Siliceous cherty limestone.....	75
	4,275

DEVONIAN SYSTEM.

The Lone Mountain limestone passes by transition upward into the Devonian rocks. Of these the Nevada limestone is 6,000 feet thick at Eureka. The lower strata are indistinctly bedded, have a saccharoidal texture and a gray color, and pass upward into shales which are distinctly bedded, brown, reddish brown, and gray in color, and in places finely striped or variegated. The upper strata are massive, well bedded, bluish black, and highly fossiliferous. The intercalated bands of argillaceous shale and quartzite vary greatly in width but do not mark any especial part of the limestone, except that they are more abundant in the middle portion than elsewhere. The limestones are everywhere more or less magnesian and nearly pure dolomites occur at many places in narrow layers. The Modoc section ^a of the Nevada limestone is given below:

Section of Nevada limestone across ridge between Signal and Modoc peaks, Nevada.

	Feet.
Dark-gray to bluish-black massive limestone poor in fossils; quite well bedded; weathering partly smooth and dark colored; partly rough and pitted and of lighter color; mostly compact and massive, also of uneven texture; with numerous calcite seams.....	1,200
Light and dark colored limestone with <i>Stromatopora</i> and <i>Chaetetes</i> ; contains two layers thinly bedded (fissile).....	150
Compact light-yellow sandstone.....	60
Light and dark colored limestone in layers 10 to 20 feet thick, with <i>Stromatopora</i> and <i>Chaetetes</i>	240
Dark-colored limestone with <i>Stromatopora</i> and <i>Chaetetes</i>	180
Alternating layers (about 10 feet thick) of dark and light gray limestone, finely banded and lined; weathering brownish gray; in places bearing <i>Chaetetes</i>	900
Compact yellow sandstone.....	50
Dark and light gray limestone; indistinct bedding.....	150
Compact yellow sandstone.....	50
Dark and light colored limestone interbedded in layers from 4 to 10 feet thick.....	250
Light-gray siliceous limestone; very siliceous near base.....	270
Alternating beds of dark and light gray limestone; at base 30 feet [of] very siliceous limestone; with cross-bedding on weathered surface.....	180

^a Hague, Arnold, op. cit., p. 66.

	Feet.
Compact yellow sandstone.....	30
Dark and light gray limestone in thick belts of dark, lighter, and gray colors.....	575
Dark dense limestone, well bedded; bearing fossils.....	225
Shaly limestone, rich in fossils.....	200
Light-gray siliceous limestone, with fine lines of bedding; in upper portion weathering in almost rectangular fragments; growing less siliceous toward the bottom.....	550
Light-gray, highly crystalline, saccharoid dolomite; not siliceous..	140

The Nevada limestone, like the Lone Mountain, was deposited in a sinking sea bottom, and consequently at some places it rests upon the Eureka quartzite.

CARBONIFEROUS SYSTEM.

White Pine shale.—This formation, included in the Devonian in the early reports, has for several years been regarded as early Carboniferous by G. H. Girty, who correlates it with the Caney shale of Oklahoma in Bulletin 377 of the United States Geological Survey. It consists of 2,000 feet of black, argillaceous, somewhat sandy shales, with intercalations of red and reddish-brown friable sandstones. It rests conformably upon the Nevada limestone and occupies a clearly defined stratigraphic position, with a marked change in the character of sedimentation and a fauna distinct from those of both the underlying and overlying formations. Fossil plant remains are common and all evidences indicate that the formation is a shallow-water deposit.

Diamond Peak quartzite.—This formation consists of 3,000 feet of massive gray and brown quartzite with brown and green shales at the top. At the base fine conglomerates lie next to the argillaceous White Pine shale, but a short distance up these beds pass into a more massive, usually vitreous quartzite with a characteristic gray-brown color, which breaks irregularly with a fluty fracture. A narrow belt of blue limestone occurring about 200 feet above the base of the formation carries Carboniferous fossils. This formation seems to be absent in sections described in the report of the Fortieth Parallel Survey.

Lower coal measures.—The lower coal measures consist of 3,800 feet of heavy-bedded dark-blue and gray limestones with intercalated bands of chert and with argillaceous beds near the base. The beds rest conformably upon the Diamond Peak quartzite and are very extensively developed in Utah and northern Nevada. They contain an abundant fauna representing a commingling of species from the upper and lower Carboniferous.

Weber conglomerate.—The Weber is 2,000 feet thick at Eureka and consists of coarse and fine conglomerates with angular fragments of chert and layers of reddish-yellow sandstone. At Eureka the material

gives abundant evidence of shallow-water deposition and shows the existence of a land mass not very far removed at the time of deposition. The fragments are rounded pebbles of limestone, quartzite, flint, and jasperoid, evidently derived from early Paleozoic rocks. In the Shoshone Range there is a great thickness of fine-grained quartzites or siliceous shales, with conglomerates very subordinate in quantity, which are said to belong to the Weber. According to Clarence King^a the Weber is 6,000 feet thick in the Wasatch Mountains and 8,000 feet thick in the Oquirrh Mountains.

Upper coal measures.—These beds at Eureka are 500 feet thick and consist of blue and drab limestones carrying fossils of the upper Carboniferous period. Clarence King describes the upper Carboniferous limestone as a body 2,000 feet thick, which over the Great Basin country is prevailingly made up of light-gray or drab limestone and is as a rule thinly bedded. Some carbonaceous shales were found within these measures, but no workable coal has been discovered in the Great Basin. Near the base of the upper coal measures in the Eureka district there is an intraformational conglomerate in the pebbles of which there are organic remains such as are common in rocks below the Weber conglomerate. No Permian beds are known in this region.

MESOZOIC ROCKS.

No Mesozoic sedimentary rocks are represented in the area here discussed. Throughout the Triassic and Jurassic periods either this area was a land mass in which no beds were laid down or else such beds, if they were deposited, were subsequently eroded. Triassic beds are extensively developed in the Humboldt, Pahute, and Havallah ranges, north and east of Carson Lake, and Jurassic beds occur in the Montezuma Range and in the Eugene Mountains.

TERTIARY ROCKS.

Eocene.—No exposures of Eocene rocks are known to be present in the area here described, but a few miles east of the eastern border Eocene beds with coal seams are found at several places. The western border of the Eocene was presumably east of the one hundred and sixteenth meridian. On the northwest slope of the Elko Range, about 3 or 4 miles east of Elko, the Eocene beds are exposed, dipping 35° E.^b These beds consist of very thin shales, calcareous at some places and siliceous at others, and they contain seams of impure coal. They are overlain unconformably by the Humboldt formation (Pliocene), which here consists of white, porous, volcanic ash. In the Dixie Hills region, near the head of Dixie Valley, there are finely bedded calcareous shales and marls containing carbonaceous seams

^a U. S. Geol. Expl. 40th Par., vol. 1, 1878, p. 240.

^b Idem, vol. 2, 1877, p. 602.

and coal. The beds dip 30° E. and resemble those at Elko in composition. These exposures are farther west than any others of Eocene rocks. On the east front of the River Range, about 14 miles north of Osino, Eocene beds, including coal seams, dip 45° S. and are overlain by beds of volcanic ash belonging to the Humboldt formation (Pliocene).

Miocene.—No sedimentary rocks of Miocene age are known to have been deposited in the area studied, but west of it Miocene beds are extensively developed. Fossiliferous Miocene beds, which King has called the Truckee group, appear in the Kawsoh Mountains and along the south end of the Montezuma Range, and beds lithologically similar are present in the Reese River canyon. The Miocene beds contain much volcanic débris of a rhyolitic character and nearly everywhere are overlain by rhyolite.

Pliocene.—In Pliocene time a great lake occupied almost the whole territory between the Wasatch Range on the east and the Sierra Nevada on the west, extending northward far into Idaho and southward to an unknown distance. This lake Clarence King ^a has named Shoshone Lake, and the beds laid down in it are called the Humboldt formation. These beds are composed mainly of friable gray, white, and drab sandstone and marly limestones and at many places contain abundant volcanic material, chiefly tuff of a rhyolitic character. Some of the siliceous beds are made up largely of diatomaceous earth. In Bone Valley, just west of the Mallard Hills, which are some 30 miles north of Halleck, Pliocene fossils have been found.

The thickness of the Humboldt formation is at most places unknown, for complete sections are very rare. In the Huntington Valley, according to King, it can not be less than 600 or 800 feet, and occurrences at other places give the impression that the thickness is greater.

The Humboldt formation covers large areas in this part of Nevada. It is found on the east and west slopes of the Pinyon Range, on the east slope of the Cortez Range, and in large areas along Maggie Creek, in Rock Creek and Squaw valleys, and in the Owyhee Desert. At several places it lies approximately flat and rests unconformably on the upturned edges of the steeply tilted Eocene beds. In the eastern foothills of the Cortez Range, east of Cortez Peak,^b the Humboldt beds overlies an extensive basalt flow. Here the beds are arenaceous and near the top pass into fine clays; some of them contain a large proportion of lime carbonate and sodium carbonate. The formation seems to extend over the whole of Pine and Garden valleys, but only on the outer margins of these valleys are the Quaternary gravels cut through so as to expose the Humboldt. About 5 miles north of Mineral Hill the strata, which are here highly calcareous, form ver-

^a U. S. Geol. Expl. 40th Par., vol. 1, 1878, p. 454.

^b Idem, vol. 2, 1877, p. 583.

tical bluffs 50 feet high. At this place the Pliocene was clearly eroded before the Quaternary beds were deposited.

In the Squaw Valley region there are extensive beds of the Humboldt composed largely of volcanic ash. Farther north, in the Owyhee Desert, there are very large areas of the same rocks, interbedded with basalt flows. In that region the Humboldt beds are at some places tilted, and east of the Burner Hills they form a monocline, dipping steeply away from that uplift. As a rule, however, the Humboldt formation is not so steeply tilted as earlier formations, and at several places the beds are flat-lying, resting upon the more highly tilted Eocene beds. In the Centennial Range east of the Bull Run mine, and high up on the slope south of Bull Run Creek, there are Tertiary beds which contain volcanic material of considerable size. These beds carry some thin seams of impure coal and are tilted and faulted against the Paleozoic sedimentary rocks.

QUATERNARY DEPOSITS.

The Quaternary formations comprise the alluvial deposits which nearly everywhere cover the broad valleys between the mountains and the glacial deposits which are found on the slopes of the most lofty of the mountain ranges.

Glacial deposits.—Many of the mountains show no evidence of glaciation. Their summits are rounded domes and the streams which drain them flow through sharp, narrow canyons that are barren of glacial drift, but in some of the higher ranges the evidence of glaciation is clear and unmistakable. In the higher part of the Shoshone Range to the south of Battle Mountain the amphitheater at the head of the streams could have been formed only by the action of ice. The glacial features of the range were described by Hague,^a who records the presence of glacial débris and striæ at the foot of Shoshone Peak. The central portion of the Centennial Range is glaciated also, as is shown by the morainal material in the small canyons which trend from the summits of the range toward Aura. The Jack Creek Range (or north end of the Seetoya Range) was not visited by the writer; the amphitheaters at the heads of the canyons which drain this range indicate that it was subject to erosion by ice. All these glaciated groups rise more than 9,000 feet above sea level, but most of the mountains of this region do not reach that elevation and were not glaciated.

Alluvial deposits.—Nearly everywhere the valleys between the mountain ranges are covered by great accumulations of Quaternary gravels. These alluvial deposits were probably formed before, during, and after the glacial period, and it is not likely that they were affected by the local glaciation to any important extent.

^a U. S. Geol. Expl. 40th Par., vol. 2, 1877, p. 620.

In Quaternary time—according to King, in the glacial period—two great lakes formed between the Wasatch Range on the east and the Sierra Nevada on the west. Lake Bonneville, the remnants of which are to-day represented by Great Salt Lake and Utah Lake, extended from the foot of the Wasatch Range to the Gosiute Range. Lake Lahontan, which was nearly as large, extended eastward from the region of Pyramid Lake nearly to Battle Mountain. The country herein described, except a small area along the valley of Humboldt River near Battle Mountain, which is represented on the King map ^a as occupying a long embayment of Lake Lahontan, was situated in the higher country between these two lakes. There were, however, two smaller lakes east of the Shoshone Range. One of these was in the Crescent Valley west of the Cortez Range, and another was west of Carico Peak, its remnant being now represented by Carico Lake.

The Quaternary deposits are made up of fragmental material of various sizes. At some places the fragments are well rounded and at others they are angular. All the sedimentary and igneous rocks are represented in these fragments, the proportions depending on the nearness and size of their outcrops. Some of the alluvial deposits are doubtless subaerial accumulations formed largely by freshets resulting from the frequent cloud-bursts, but others were formed under water, as is shown by their stratification and by the fact that they are rich in saline matter, which has resulted from the drying up of the inclosed basin. In the Crescent Valley, between the Cortez and Shoshone ranges, ^b the Quaternary beds include deposits of relatively pure sodium chloride which has been used for metallurgical purposes.

IGNEOUS ROCKS.

INTRUSIVE GRANITIC ROCKS.

DISTRIBUTION AND COMPOSITION.

The oldest igneous rocks in the area are stocks and dikes of granular rocks which intrude the Paleozoic sedimentary rocks at many places. These intrusive rocks, which are commonly called granite, are as a rule medium grained and of light-gray color and are composed of quartz, feldspar, and biotite, with a variable amount of hornblende. None of the granitic rocks are sheared or contorted in the manner which results when deeply buried rocks are metamorphosed. Under the microscope they show a considerable range in composition. Some of them may properly be called quartz diorites or quartz monzonites, but most of them could be classified as granodiorites, if this term is used with its broadest application. Granodiorite is, according to Lindgren, ^c

^a U. S. Geol. Expl. 40th Par., vol. 1, 1878, Pl. VI.

^b *Idem*, p. 503.

^c Lindgren, Waldemar, *Am. Jour. Sci.*, 3d ser., vol. 46, 1893, p. 203.

a light-gray granitic rock consisting in typical development of feldspar, quartz, biotite, and hornblende, with a medium-grained texture. The soda-lime feldspars are usually considerable and to a variable extent in excess of the alkali feldspars. The silica varies between 60 and 73 per cent; the amount of lime is variable, but it rarely exceeds and usually falls somewhat short of the alkalis, while in some varieties, which can not be distinguished from the others in the field, there is more potash than soda; a frequently occurring relation is 2 per cent K_2O to 4 per cent Na_2O . It will be seen that the rock very closely approaches some quartz-mica diorites and often might be indicated by that name.

Specimens from four of the older intrusive masses are low in potash and relatively high in soda and lime. One of these intrusive bodies is in the Railroad mining district near Bullion; another forms the summit of Lone Mountain; a third is a large body forming the mountain on which the Grey Eagle mine is located; and a fourth is near the head of Willow Creek, about 8 miles west of Tuscarora. These stocks could properly be called quartz diorites and are higher in lime and in soda than any of the other stocks. They are composed characteristically of feldspars, quartz, biotite, and a little hornblende, with magnetite and apatite as accessory minerals. The feldspars are mainly plagioclase and for the most part andesine. Only a little orthoclase is present.

Intrusive rocks of this group which approach more nearly the mean composition of granodiorite are a number of broad dikes and stocks in the Centennial Range near Edgemont and Aura and several large masses in the Shoshone Range near Dean and Tenabo. The constituent minerals are the same as in the rocks described above, but there is a little more orthoclase and quartz and plagioclase is not quite so abundant.

An analysis ^a of the rock which constitutes the summit of Shoshone Peak was made by R. W. Woodward. This analysis gives all the iron as ferrous iron and the norm can not be calculated, but it is sufficient to confirm the classification of the rock as granodiorite. Its composition is very near that of a granodiorite from Pyramid Peak, Eldorado County, Cal., described by Lindgren,^b and that of other typical granodiorites in California.

Analysis of granodiorite from Shoshone Peak.

[By R. W. Woodward.]

SiO ₂	70.17
Al ₂ O ₃	14.53
FeO.....	4.03
CaO.....	2.29
MgO.....	.93
Na ₂ O.....	3.25
K ₂ O.....	3.35
H ₂ O.....	1.53
	100.08

^a U. S. Geol. Expl. 40th Par., vol. 2, 1877, p. 621.

^b Lindgren, Waldemar, Am. Jour. Sci., 4th ser., vol. 3, 1897, p. 306.

Still another group of the older intrusive rocks includes the stocks at Cortez and Mill Canyon, at Lone Mountain, and at Mountain City. The rocks of this group contain considerable orthoclase. Some of the specimens studied are composed of feldspar, quartz, mica, and hornblende. Orthoclase and quartz are more abundant than in the granodiorite intrusives above referred to. Plagioclase is less abundant and not so high in lime and is in the main oligoclase and andesine. A single stock may show notable differences in composition. The more acidic phases of these rocks are quartz monzonites and the more basic phases granodiorites.

An analysis made by Thomas M. Drown ^a of a specimen taken from Lone Mountain or Nannies Peak by S. F. Emmons is given below. This rock is a quartz monzonite and in composition approaches closely the quartz monzonite of the Idaho Hailey mine, Hailey, Idaho, described by Waldemar Lindgrén,^b and the quartz monzonite near San Miguel Peak, Telluride, Colo., described by Whitman Cross.^c

Analysis of quartz monzonite of Lone Mountain (Nannies Peak).

[By Thomas M. Drown.]

SiO ₂	70.77
Al ₂ O ₃	15.22
FeO.....	2.65
MnO ₂11
CaO.....	2.33
MgO.....	.71
Na ₂ O.....	3.75
K ₂ O.....	4.58
H ₂ O.....	.52
	100.64

The granitic mass in the Agate Pass region was not visited by the writer. An analysis of a specimen collected by S. F. Emmons is given below. It represents a typical soda-rich granite and is considerably higher in alkali than any of the rocks collected from the area studied.

Analysis of granite from Agate Pass.

[By Thomas M. Drown.]

SiO ₂	72.01
Al ₂ O ₃	14.75
FeO.....	2.35
MnO ₂17
CaO.....	.79
MgO.....	.65
Na ₂ O.....	4.21
K ₂ O.....	4.49
H ₂ O.....	.61
	100.03

^a U. S. Geol. Expl. 40th Par., vol. 2, 1877, p. 603.

^b Twentieth Ann. Rept. U. S. Geol. Survey, pt. 2, 1900, p. 81.

^c Telluride folio (No. 57), Geol. Atlas U. S., U. S. Geol. Survey, 1899, p. 6.

To the southwest of this body of soda-rich granite there is an area of quartz diorite. The mass that outcrops in Agate Pass forms the southwest half of the long area of granitic rock which is shown on the geologic map (Pl. V) on the northwest slope of Cortez Peak. As indicated in the geologic atlas of the Fortieth Parallel Survey, this quartz diorite mass is about 4 miles long and half as wide. It is composed of plagioclase, quartz, hornblende, and biotite, and the subjoined chemical analysis indicates that it is the most basic of all the granitic rocks of this group. As no ore deposits were reported in this area, it was not visited by the writer, but to judge from the description by Zirkel ^a it is highly sericitized along Agate Canyon and, as is well known, ore deposition very commonly accompanies sericitization. In composition the rock closely resembles a quartz diorite from Spanish Peak, Plumas County, Cal., described by H. W. Turner, ^b and a similar rock from Yaqui Creek, Mariposa County, Cal., described by the same writer. ^c

Analysis of quartz diorite from Agate Pass.

[By R. W. Woodward.]

SiO ₂	58.54
Al ₂ O ₃	16.68
FeO.....	5.62
CaO.....	6.00
MgO.....	5.22
Na ₂ O.....	2.76
K ₂ O.....	2.50
H ₂ O.....	2.15
	99.47

CONTACT RELATIONS AND AGE.

In the work of the Fortieth Parallel Survey some of the stocks were supposed to be of Archean age. The significance of contact phenomena was, however, fully recognized, for King, ^d speaking of their relations to the sedimentary rocks, says:

The configuration of the granite topography of the Archean surface prior to the deposition of the Paleozoic series was that of an area of mountain ranges, possessing some very abrupt precipitous walls, sharp lofty peaks, and broad low domes. Where these came to be uplifted together with superjacent strata, and afterward exhumed by erosion, which brought to light granite peaks piercing through highly inclined beds, it often becomes absolutely impossible to determine the relation of the two. In the absence of any granitic dikes penetrating the stratified series, or of peculiar local metamorphism, or of general evidence of intrusion, the bodies are usually referred to the old Archean topography. Only in cases where the granite is actually seen to penetrate either fissures or warped openings in the strata is it safe to refer it to a later origin than the sedimentary series.

^a U. S. Geol. Expl. 40th Par., vol. 2, 1877, p. 576.

^b Seventeenth Ann. Rept. U. S. Geol. Survey, pt. 1, 1896, p. 72.

^c Bull. U. S. Geol. Survey No. 150, 1898, p. 342.

^d King, Clarence, U. S. Geol. Expl. 40th Par., vol. 1, 1878, pp. 77, 100.

This statement of the criteria of intrusion is brief and comprehensive. It remains, therefore, to discuss the evidence of intrusion in some detail for each granitic mass. The more important intrusive bodies studied are located near Bullion, in the Railroad district, in the Cortez and Mill Canyon mining district, at several places in the Shoshone Range, at Lone Mountain, in the Centennial Range, and at Mountain City.

The stock near Bullion shows a very irregular contact and dikes of the granodiorite cut the limestone. The contact between the granodiorite and limestone is marked by a garnet zone several hundred feet wide.

At Cortez the relations between granodiorite and limestone are clearly crosscutting. At Mill Canyon the granodiorite stock sends out small apophyses into limestones, and near the contact the sedimentary rocks carry contact-metamorphic silicates.

In the Shoshone Range granitic rocks or their porphyries were noted at Tenabo, Mud Springs, Hilltop, Grey Eagle, and Dean. At Tenabo some veinlets of actinolite cut the quartzite near the contact with granodiorite porphyry, but no garnet zones were observed in this range. The contacts of the igneous and sedimentary rocks are distinctly crosscutting at Mud Springs, Hilltop, and Dean.

At Lone Mountain zones of contact-metamorphic silicates are developed near the contact of granodiorite and the sedimentary rocks.

In the Centennial Range the Paleozoic sedimentary rocks are cut by dikes and other intrusives of granodiorite which are clearly of later age.

At Mountain City the granitic rocks are clearly later than the sedimentary rocks. At California Hill garnet and other contact-metamorphic silicates are developed in limestone along the contact.

The contact relations of the granodiorite about 8 miles west of Tuscarora, near the head of Williams Creek, were not studied, but the rock has the same general composition as that of the stocks mentioned, and, like them, it does not show deep-seated metamorphism. In view of the evidence shown by the contacts of practically all the masses, it may be assumed that they are, without exception, intruded into the Paleozoic rocks and that they are therefore of post-Carboniferous age. They are known also to be older than the rhyolites and basalts and associated andesites, for they do not intrude these rocks. From evidence within the area studied, these stocks are therefore known to be younger than the Carboniferous and older than the Miocene, but on the assumption that the intrusives of this character throughout the West generally were formed at about the same time, the ages of these rocks may be estimated more closely. In the Humboldt Range F. L. Ransome found granodiorite stocks cutting

the Triassic sedimentary rocks. In California similar stocks cut the Jurassic. At Bisbee, Ariz., as shown by Ransome, granitic stocks cut the Carboniferous and are eroded and covered by sediments of Comanche (Lower Cretaceous) age—the “Bisbee group” of Ransome’s report.^a In the Clifton-Morenci district, according to Lindgren,^b the granite porphyry and related deep-seated porphyries cut Cretaceous beds which are equivalent to the Benton, and they are therefore probably of late Cretaceous or the very earliest Tertiary age. At Ely, Nev., at Bingham, Utah, and at other Utah camps deep-seated granular rocks intrude sedimentary rocks, but at those places no sedimentary rocks later than the Carboniferous are represented and the age of the intrusives can not be closely approximated. Near Philipsburg, Mont., large stocks of granites and nearly related rocks cut the Paleozoic and Mesozoic sediments, including beds as young as the Cretaceous. From the relations thus shown by the post-Paleozoic granular or porphyritic intrusive rocks elsewhere in the Western States it may with some reason be assumed that these intrusives are of about the same age, and that they were formed either at the beginning of the Cretaceous, during the Cretaceous, at the very end of Cretaceous time, or perhaps at the very beginning of the Tertiary.

PORPHYRIES ASSOCIATED WITH INTRUSIVE GRANITIC ROCKS.

Porphyritic dikes and other small intrusive masses are associated with some of the granitic stocks. Most of these are believed to be either the more rapidly cooled portions or the products of differentiation from the granodiorite magmas. In the main they were formed at about the same time that the stocks were intruded, but some of them may be younger and may belong to the late Tertiary period of volcanism.

An intruding quartz porphyry in the Standing Elk mine, near Bullion, contains abundant rounded phenocrysts of quartz and a smaller number of feldspar phenocrysts in a light-colored, fine-grained groundmass. Muscovite and biotite are present, but the dark-colored silicates are very sparingly developed and are much less abundant than in the granodiorite with which the porphyry is associated and from which it is probably derived.

At Lone Mountain granodiorite porphyry is associated with granodiorite and appears to grade into it. The granodiorite porphyry has a fine crystalline groundmass composed of quartz and orthoclase, in which there are phenocrysts of acidic plagioclase, resorbed quartz, biotite, and hornblende. Orthoclase and quartz are more abundant in the porphyry than in the granodiorite.

^a Ransome, F. L., Prof. Paper U. S. Geol. Survey No. 21, 1904, pp. 56-73.

^b Lindgren, Waldemar, Prof. Paper U. S. Geol. Survey No. 43, 1905, p. 85.

The granodiorite of the Phoenix mine, at Tenabo, is cut by a quartz porphyry, the microcrystalline groundmass of which contains abundant phenocrysts of orthoclase, quartz, and biotite, with little or no plagioclase. A similar porphyry cuts the sedimentary rocks in the Gold Quartz mine, near by. At the Two Widows mine, in the same camp, a small dike of quartz diorite porphyry cuts the siliceous sedimentary rocks. In the Maysville district and at Dean, on the north slope of the Shoshone Range, granodiorite porphyries cut the sedimentary rocks. These have a coarsely microcrystalline groundmass and grade into granodiorite.

The granodiorite at Mountain City is cut by small, light-colored dikes of fine-grained granite, some of which are typical aplite. They are probably differentiation products of the granodiorite magma.

At Cortez and at Mineral Hill several dikes cut the limestone. These are composed mainly of quartz, sericite, calcite, and iron oxide and are too much decomposed for determination. They are probably derived from the deep-seated intrusives.

A felsitic dike of quartz porphyry on the west slope of Lone Mountain and an intrusive of somewhat similar character below the Pittsburg mine, at Dean, are believed to belong to a period of volcanism later than that during which the granodiorites and related porphyries were formed. The groundmass of these quartz porphyries is much the same as that of rhyolite, indicating that they cooled near the surface. Quartz porphyries with rhyolites and andesites are discussed on another page.

TERTIARY ERUPTIVE ROCKS.

RHYOLITE.

Rhyolites are glassy igneous rocks which have about the same chemical composition as granites. In color they are white, pink, purple, or dark brown. The dense pasty groundmass may contain phenocrysts of quartz and orthoclase, with small amounts of biotite, augite, and hornblende. A little soda-rich plagioclase may also be present among the phenocrysts. They form surface flows and many of them show streaking due to flowage. Some of them are flow breccias formed of angular fragments of rhyolite in a matrix of the same. Such rocks result when a crust forms over the flow and is broken up by the movement of the still liquid portion, which, solidifying, forms the matrix for the fragments. Some of the rhyolites in the Gold Circle district are thin, fissile bands which resemble shales. The shaly appearance is probably due to banding that developed as the rhyolite flowed and was emphasized by subsequent weathering along the parting planes. Some of the rhyolites are vesicular. The small blebby holes in these rocks represent the places where imprisoned gases expanded when the pressure was

removed from the magma at the time of eruption. In some of the rhyolites in the Gold Circle district the vesicles are filled with amygdules of amethystine quartz, deposited by water after the rock had solidified.

The term rhyolite in this reconnaissance is used in a broad sense. Some of the rocks so called are probably latites. Some of the brown glasses which have been called obsidians contain, besides quartz phenocrysts, acidic plagioclase, augite, and some magnetite and in composition approach quartz-bearing andesites.

The rhyolites are the most abundant and most widely distributed of the Tertiary eruptive rocks; there is not an important mountain range which does not contain large bodies of them, but they are most numerous in the northern and western parts of the area. They are the oldest Tertiary eruptives and at many places rest directly on the Paleozoic sedimentary rocks. They are cut by andesites and overlain by flows of basalt. Their thickness has not been measured but must reach a maximum of over 2,000 feet. From many places they have probably been eroded. Their distribution in the various mountain ranges is discussed in greater detail where the rocks of the various ranges are described.

BASALT.

Basalts are dark basic igneous rocks with a dense compact ground-mass which may be finely microcrystalline, but which nearly everywhere contains dark-colored glass. Basic plagioclase, olivine, pyroxene, and magnetite are generally present, some of them as crystals visible to the eye. The darker color and the presence of the small crystals of greenish-yellow, glasslike olivine will usually serve to distinguish the basalt from andesites. Some of the basalts are dense and glassy and others are highly vesicular. In this area the basalts are not so widely distributed as the rhyolites, but there are several bodies of considerable size. One of these caps the great flat-topped Shoshone Mesa northeast of Battle Mountain and another covers Whirlwind Mesa west of Beowawe. A broad belt of basalt flanks the east slope of the Cortez Range for a distance of about 30 miles.

The basalts occur as flows which overlie the rhyolites and the older rocks, and as dikes which cut them. At some places they are overlain by Pliocene lake beds and at other places they rest upon these beds. The thickness of the basalt flows has not been determined at many places. The mass which caps Shoshone Mesa is, according to King, more than 1,000 feet thick. They are not known to have been intruded by the andesites nor have any ore deposits been found in them. They were presumably extravasated after the period of andesitic eruption and late Tertiary ore deposition.

ANDESITE.

Andesites are brown, gray, or greenish-gray rocks which are intermediate in composition between rhyolites and basalts. The groundmass is very finely crystalline or glassy. Andesites may be aphanitic, or without visible crystals, but as a rule they have phenocrysts of feldspar with a variable amount of augite, hornblende, and biotite. If the phenocrysts are conspicuous the rock may be termed andesite porphyry. Magnetite crystals and dots of magnetite are usually numerous in the groundmass and they occur as inclusions in the phenocrysts. In the andesites of this area nearly all the feldspar phenocrysts are plagioclase, but a little orthoclase is present in some. Andesine is the most abundant feldspar of the plagioclase group; oligoclase is present in some specimens and labradorite in the darker, more basic types. Some andesites carry small rounded phenocrysts of quartz, but where quartz is present in considerable quantity the rocks are called dacite. Although some of the andesites of this area show vesicular phases, this character is not so frequently developed as in the rhyolites and basalts. At Gold Circle, Tuscarora, Cornucopia, and Good Hope the andesites are associated with rhyolite. At Tenabo, Lander, and Lime Mountain they cut through the Paleozoic sedimentary rocks or form flows above them.

The andesite at Tuscarora has a dense greenish groundmass containing highly altered phenocrysts of andesine, hornblende, biotite, and magnetite, with some orthoclase. In an open cut at the Dexter mine the relations of the andesite and rhyolite are clearly crosscutting, and as the rhyolite is a flow, the andesite must have been intruded into it. At Cornucopia the andesite intrudes rhyolites and is a dark porphyry composed of a glassy groundmass which contains many phenocrysts of basic andesine and pyroxene, with large chloritic patches that seem to have resulted from the decomposition of hornblende.

At Gold Circle the andesite is relatively rich in augite and magnetite. Andesine predominates, but some of the plagioclase is labradorite. The andesite is mainly younger than the rhyolite, for dikes of andesite cut the rhyolite and flows of andesite which are presumably connected with the dikes overlie the rhyolite flows. At one place in this district a rhyolite flow caps the andesite and an intrusive mass of quartz porphyry which has the same composition as rhyolite was noted cutting andesite. It appears that the eruptions of the rhyolite took place in the main before the andesite was formed, but that some rhyolite eruptions followed those of andesite.

The contact relations between the andesites and the basalts have not been determined. No andesitic intrusions have been noted in the Pliocene lake beds (Humboldt formation) and the andesitic eruptions are therefore probably older than these beds. As the basalt

effusive rocks are closely associated with these beds, it is highly probable that the andesitic eruptions ceased before the basalts were extravasated.

Andesites cut the sedimentary rocks in the Shoshone Range and are exposed in the Gem mine, at Tenabo, and in the Lovie mine, at Lander. They have a dense glassy groundmass and at some places are vesicular, and so it is assumed that they represent flows or intrusives formed near the surface. They are dark dense rocks with an altered glassy groundmass containing phenocrysts of acidic plagioclase and some orthoclase.

An analysis of a specimen of somewhat altered andesite, collected by S. F. Emmons between Wagon Canyon and Palisade, in the Cortez Range, is given below.^a

Analysis of andesite near summit of Cortez Range, between Wagon Canyon and Palisade.

[By R. W. Woodward.]

SiO ₂	61.64
Al ₂ O ₃	17.44
Fe ₂ O ₃82
FeO.....	3.99
CaO.....	5.86
MgO.....	3.05
Na ₂ O.....	3.45
K ₂ O.....	1.15
H ₂ O.....	2.64
	100.04

DACITE.

When andesite contains a few phenocrysts of quartz it is termed quartz-bearing andesite, but if there is a considerable proportion of quartz—as much as 3 or 4 per cent of the volume—then the rock is dacite. The groundmass of dacite is glassy or fine grained, and the phenocrysts commonly present are feldspar, quartz, hornblende, biotite, and augite. The feldspars are mainly plagioclase and may be a little richer in soda than the plagioclases of andesite. The andesite at the Zenoli mine in the Safford district, near Palisade, is richer in quartz than the other andesites which form the country rock for the ore deposits. Northeast of this region there is a body of typical dacite containing a noticeable proportion of quartz. Another body of dacite is in the Cortez Range and extends southward from Palisade a distance of several miles. The dacites are closely related to andesites in composition and occurrence and are believed to have been erupted at about the same time.

SUCCESSION OF THE TERTIARY ERUPTIVE ROCKS.

The order of eruption or the succession of Tertiary igneous rocks in the Great Basin, as given by Baron von Richthofen,^b is (1) pro-

^a U. S. Geol. Expl. 40th Par., vol. 2, 1877, p. 587.

^b Richthofen, F. von, Principles of the natural system of volcanic rocks: Mem. California Acad. Sci., vol. 1, 1868, p. 36; also Jahrb. K. k. geol. Reichsanstalt, vol. 11.

pylite, (2) andesite, (3) trachyte, (4) rhyolite, (5) basalt. This sequencé in an amplified form was accepted by the geologists of the Fortieth Parallel Survey, but for rhyolite and basalt the term neolite was substituted. Subsequently, it was proved by G. F. Becker ^a that propylite is in the Washoe district an altered phase of andesite. Later still this form of alteration was shown to be common and so the term propylite has ceased to be used for the primary rock, but "propylitization" has been accepted for a certain kind of hydrothermal metamorphism. The trachytes, as determined by Zirkel for the Fortieth Parallel Survey, have been shown by Hague and Iddings ^b to be in the main andesites and dacites. Spurr, ^c who has studied the Tertiary lavas throughout a large part of the Great Basin, gives this general succession: (1) Rhyolite, (2) andesite, (3) rhyolite, (4) andesite, and (5) basalt.

In the area here considered the rhyolites were the first rocks erupted. They were the most extensive, for they cover a large part of the surface and their thickness is very great. Subsequently the andesites were erupted, but in much smaller amounts. They took the form of dikes and large intrusive masses as well as flows, and in that they differed from the rhyolites, which are flows. The eruption of the basalts followed that of the andesites and seems to have closed the period of volcanism in northern Nevada. The Pliocene lake beds (Humboldt formation), whose deposition followed the bulk of the rhyolite eruptions, were formed during the eruptions of the basalts, for basalt flows occur under them and interbedded with them. Some of the rhyolite appears also to have been erupted during the deposition of the Humboldt beds, for rhyolite pumice is interbedded with them, and this does not seem to be a sand washed from older rhyolites, but volcanic dust which settled directly in water. The periods of the extravasation of the various lavas are thus shown to have been to some extent overlapping, but the general succession is (1) rhyolite, (2) andesite, (3) basalt.

AGE OF THE TERTIARY ERUPTIVE ROCKS.

In Eocene time the area covered by this reconnaissance was, as already stated, dry land, but just east of it there was a great lake in which the Eocene beds were deposited. In the Dixie Hills, east of the Pinyon Range, some 25 miles southwest of Elko, these beds, according to S. F. Emmons, ^d are for the most part finely bedded calcareous shales containing carbonaceous members which carry seams of impure coal. Similar beds are found in Penn Canyon, ^e 14 miles north of

^a Geology of the Comstock lode and the Washoe district: Mon. U. S. Geol. Survey, vol. 3, 1882, p. 88.

^b Am. Jour. Sci., 3d ser., vol. 27, 1884, p. 453.

^c Jour. Geology, vol. 8, 1900, p. 621.

^d U. S. Geol. Expl. 40th Par., vol. 2, 1877, p. 562.

^e Idem, p. 595.

Osino, and on the northwest slope of the Elko Range,^a 4 miles east of Elko. There is no record of volcanic ejectamenta interstratified with these beds, and it is inferred that the country to the west of this lake was free from volcanism throughout the Eocene.

The area here discussed was dry land also during Miocene time, when there was a large lake to the west of it in which the Truckee formation was deposited. As already stated, typical exposures of these beds with fresh-water mollusks are found in the Kawsob Mountains and along the south end of the Montezuma Range, and beds of similar lithologic composition occur in the Reese River canyon about 12 miles southwest of the southwest corner of the area.^b These beds contain much volcanic material of rhyolitic character and nearly everywhere are overlain by rhyolite. Wherever observed in connection with basaltic eruptions they are cut through or overlain by the basalt.

In Pliocene time a lake occupied almost the whole territory between the Wasatch Range and the Sierra. In this lake were deposited the Humboldt beds, which contain abundant volcanic material, mainly of a rhyolitic character. These beds rest above basalts and flows of basalt are included in them.

The general succession of the eruptive rocks was, as already stated, (1) rhyolite, (2) andesite, (3) basalt. The rhyolite eruptions probably began in early Miocene time soon after the deposition of the Eocene lake beds, and continued through the Pliocene, for there are rhyolitic tuffs and lapilli in these lake beds. The andesites were intruded after there had been very extensive rhyolite eruptions, for great thicknesses of rhyolite are cut by andesite. As few if any andesites intrude the Pliocene lake beds, it is believed that the andesites were erupted mainly, if not altogether, late in Miocene time, probably at its close. The basalts were erupted during Pliocene time, for they are interbedded with the Pliocene lake beds.

DEFORMATION OF THE ROCKS.

The sedimentary rocks and lava flows are in few places found in the horizontal position, but nearly everywhere are tilted and faulted. In all the ranges where they are exposed the Paleozoic rocks are thrown into broad open folds and have dips which for the most part range from 15° to 45°. As shown by the atlas of the Fortieth Parallel Survey, the Cortez and Pinyon ranges are in the main anticlinal, the Seetoya or Jack Creek Range is synclinal, and the Shoshone Range is an eastward-dipping monocline modified by profound faulting. The Centennial Range is a northward-dipping faulted monocline modified by subordinate anticlines and synclines.

^a U. S. Geol. Expl. 40th Par., vol. 2, 1877, p. 601.

^b Idem, vol. 1, 1878, p. 412.

The Tertiary lava flows are at no place closely folded. The contorted bands of some of the shale-like rhyolites suggest that these rocks have been deformed by compression, but a close inspection shows that their convolutions are the result of movements which took place before the rocks had completely cooled, and are not caused by subsequent deformation. The lava beds are nearly everywhere in a tilted position, however, and at many places they are faulted. The tilting movements were evidently accomplished after the Paleozoic sedimentary rocks had been folded, and the lavas do not show so large an element of horizontal compression as the sediments.

The present attitude of the Paleozoic sedimentary rocks is doubtless due to various processes which operated at different times. As already stated, there are in the area no bedded rocks which were laid down between the close of the Carboniferous and the beginning of the Miocene, and consequently there is no evidence within the area itself which will show the time at which the deforming movements took place. Information respecting what happened during this time must therefore be gained outside of the area, at places where rocks of suitable age are known.

In California there is a general unconformity between the folded Jurassic beds and the Cretaceous beds, which are not so greatly folded. The time of the folding is therefore known to be at the end of the Jurassic. Folded Jurassic beds are found in northern Nevada as far east as the Pahute Range, and, although these beds are not covered by the Cretaceous, they were probably deformed at the time of the extensive mountain-making movements in California. It is not known whether this movement extended eastward across the Great Basin as far as the Wasatch Mountains, but at any rate the Wasatch and Uinta mountains and the country extending far to the east of them were strongly uplifted at the close of Cretaceous time. The effect of the movement at this time was probably felt also in northern Nevada.

A few miles east of Elko^a the Eocene beds are highly tilted and overlain by volcanic materials of Pliocene age. In the Dixie Hills, southwest of Elko, and in Penn Canyon, 14 miles north of Osino, beds of the same age are highly tilted. The Miocene lavas are not so greatly deformed, and so it is supposed that the movement took place before Miocene time, or near the close of the Eocene. The mountain-making movements which are recorded in the folded Paleozoic sedimentary rocks seem therefore to have taken place at the end of the Jurassic, at the end of the Cretaceous, and at the end of the Eocene, but the extent and relative importance of each can not be shown.

^a U. S. Geol. Expl. 40th Par., vol. 2, 1877, p. 595.

A period of deformation followed the eruption of the Miocene lavas, for these rocks at many places are highly tilted and faulted. This movement seems to have taken place without the strong compressional stresses which produced the pre-Miocene folds, and practically all the faults are of the normal type, in which the hanging wall appears to have dropped. Some of the mountain ranges were doubtless elevated at this time. Blocks of the Paleozoic sedimentary rocks which had been deformed by folding in pre-Miocene time were left in relatively exalted positions when other blocks sank away from them and formed the valleys. Since that time profound erosion has obliterated the fault scarps, but has not everywhere obliterated the entire effects of faulting, for the elevated block is still at a higher altitude than the depressed block.

A period of less intense crustal deformation followed the deposition of the Pliocene lake beds, which are in places gently warped and faulted.

RÉSUMÉ OF GEOLOGIC HISTORY.

In the area studied there is no record of pre-Cambrian events. The oldest rocks exposed are the quartzites and grits which outcrop on the crest of the Pinyon Range. Throughout Cambrian and early and middle Ordovician time this part of Nevada was the floor of a sea upon which quartzites, limestone, and shales were deposited, the whole series of pre-Silurian rocks having a thickness of more than 10,000 feet. Near the middle of the Ordovician period the sea became shallower and the Eureka quartzite was deposited. Subsequently the sea bottom was lifted above the water level without tilting the beds. The land mass did not remain long above the water, but sank slowly and gradually through late Ordovician time,^a and was again elevated at the close of the Ordovician, and throughout the Silurian remained above sea level. In Devonian time approximately 6,000 feet of limestone, sandstone, and shale were deposited. Sedimentation was uninterrupted between the Devonian and the Carboniferous, but in the early Carboniferous there was a shallowing of the waters, and at some places as much as 3,000 feet of sandstone was deposited. This was followed by the deposition of 3,800 feet of limestone and shales, above which was laid down the Weber conglomerate, having a thickness of 2,000 to 6,000 feet. This was followed by the deposition of the upper Carboniferous limestone, up to 2,000 feet thick.

During the Paleozoic era there were thus deposited between 30,000 and 40,000 feet of sedimentary rocks. These beds carried a considerable proportion of conglomerates, sandstones, and shales and must have been deposited, in part at least, in relatively shallow water and not far from the shore. According to Clarence King,^b the Paleozoic

^a Hague, Arnold, Mon. U. S. Geol. Survey, vol. 20, 1892, p. 57.

^b U. S. Geol. Expl. 40th Par., vol. 1, 1878, p. 247.

sea was east of a land mass whose shore was in northern Nevada near longitude $117^{\circ} 30'$. West of that meridian and north of the fortieth parallel there are no Paleozoic rocks in northern Nevada, but pre-Cambrian or post-Paleozoic rocks instead. As already stated, the sea bottom did not remain stationary, but oscillated from time to time and permitted a very great thickness of beds to be deposited. At the close of the Ordovician period parts of it rose and were not submerged again until the beginning of the Devonian. The close of Paleozoic time was marked by profound but relatively gentle continental movements. In the vicinity of the Havallah Range, about $117^{\circ} 30'$ west longitude, there was, according to King,^a a pivotal line. The area to the east of this line remained submerged in Paleozoic time, and that to the west was above sea level. In Mesozoic time the conditions were reversed—the country to the east became dry land and that to the west sank below the level of the sea and received thick contributions of Triassic and Jurassic sediments. According to King,^b immediately after the deposition of the Jurassic sediments they were folded with much horizontal compression, producing great north-south mountain ranges. The westernmost of these ranges was the Sierra Nevada, and in the Great Basin ranges were formed probably at the same time. The reason for this conclusion is that many of the mountain ranges parallel to the mountains in the area studied, but west of it, are composed of folded Jurassic rocks. The folding at this time was most intense in the Sierra Nevada and decreased eastward toward the Wasatch Range. At about this time, or else near the close of the Cretaceous, the granitic rocks, in the main granodiorites, were intruded in the Paleozoic beds. Many of the ore deposits were formed at the time of this intrusion.

With the advent of Tertiary time began a period marked by extensive inland lakes. The earliest of these in this region was the Gosiute Lake, in which were laid down the Eocene beds. Its western shore extended nearly to the eastern border of the area studied, this area constituting a land mass from which the sediments were derived. In this lake were deposited up to 2,000 feet of shales, clays, and limestones, with some beds of impure coal. The deposition of these beds was followed by mountain-making movements, in which they were tilted at some places as much as 45° . In Miocene time a great lake was formed west of the area. King has called this the Pahute Lake and the beds are called the Truckee formation. They contain much volcanic débris and record a time of great volcanic activity. This period of igneous activity differed from that of the post-Jurassic, as the rocks recording it are not granular rocks but are mainly lava flows and andesites, formed relatively near the surface. The rhyolites were first extravasated, and later these were intruded by andesites.

^a U. S. Geol. Expl. 40th Par., vol. 1, 1878, pp. 731, 759.

^b Idem, p. 733.

Many of the ore deposits were formed in connection with the andesitic intrusions.

In Pliocene time, when volcanism was still at its height, an extensive lake was formed which covered nearly the whole of the Great Basin. King ^a has called this Shoshone Lake and the beds that were laid down in it the Humboldt formation. In the lake, which may have contained lofty islands, a great thickness of sandstones, clays, and calcareous and diatomaceous shales was deposited, with a mass of volcanic material, principally of a rhyolitic character, which was blown out of volcanic vents and settled in the water of the lake. Extensive flows of basalt occurred at about this time.

The late Tertiary was marked by very extensive normal faulting, a large part of which took place after the lava flows were extravasated and after the second period of ore deposition. Many of the old mountain ranges were probably raised, and new ones may have been formed. In the Quaternary, mountain glaciers formed in the higher ranges during the glacial period. Small lakes occupied some of the depressions between the mountains, and extensive accumulations of débris, eroded from the mountains, filled the valleys.

ORE DEPOSITS.

GENERAL STATEMENT.

It has been shown that the igneous rocks of the area studied belong to two distinct periods of volcanism. The first of these was probably in Cretaceous time, and the rocks formed during that period are intrusive granular rocks and deep-seated porphyries, in the main granodiorites and granodiorite porphyries. All these rocks were formed at considerable depths, and since they were intruded this country has been greatly eroded and the capping which must have covered them when they solidified has been removed. These early intrusive rocks do not cut the Tertiary lake beds or lavas, but are confined to the Paleozoic sedimentary formations.

A large number of the ore deposits are in or near the early intrusives. These include deposits at Bullion (Railroad district), Lone Mountain, Edgemont, Columbia, Aura, Mountain City, Cortez, Mill Canyon, Grey Eagle, Dean, and Lewis, and some of the deposits at Tenabo. The deposits at Mineral Hill may also belong to this group.

The Tertiary lavas and associated rocks are in the main rhyolites, andesites, and basalts. These rocks are younger than the granodiorites and associated rocks and, for reasons given on pages 34-35, are thought to be of Miocene and Pliocene age. The ore bodies associated with the later eruptives include the deposits at Tuscarora, Cornucopia, Good Hope, Burner, Falcon, Stafford, Lynn, and Gold Circle, and probably some of those at Tenabo and Lander.

^a U. S. Geol. Expl. 40th Par., vol. 1, 1878, p. 456.

With respect to metal content, the deposits of both groups carry silver, gold, copper, and lead. Silver and gold are the most important metals of both groups, but the proportion of gold to silver is greater in the earlier deposits than in those of the later group. The deposits at Edgemont and at Dean belong to the earlier group, and at these places the only metal won in important quantity is gold. The copper and lead deposits are associated mainly with the older intrusive rocks. These metals occur also with the silver and gold ores of the later group, but in smaller proportion.

EARLIER DEPOSITS (CRETACEOUS?).

GENERAL STATEMENT.

The deposits associated with the older intrusive rocks are conservatively estimated to have produced \$22,000,000. The minerals of these deposits are given in the list below:

Actinolite	Chalcopyrite.	Hornblende.	Pyrrhotite.
Apatite.	Chlorite.	Kaolin.	Quartz.
Argentite.	Chrysocolla.	Limonite.	Sericite.
Arsenopyrite.	Copper.	Magnetite.	Silver.
Azurite.	Cuprite.	Malachite.	Specularite.
Barite.	Diopside.	Manganite.	Stephanite.
Biotite.	Enargite.	Molybdenite.	Stibnite.
Bismuthinite?	Epidote.	Muscovite.	Stromeyerite.
Bornite.	Freibergite.	Polybasite.	Tetrahedrite.
Bromyrite?	Fluorite.	Proustite.	Tremolite.
Calcite.	Galena.	Pyrrargyrite.	Zinc blende.
Cerussite.	Garnet.	Pyrite.	Zoisite.
Cerargyrite.	Gold.	Pyrolusite.	
Chalcanthite.	Gypsum.	Pyroxene.	
Chalcocite.	Hematite.	Pyromorphite.	

With respect to the intrusive rocks, the Cretaceous (?) ore deposits show various relations. The gold-bearing fissure veins at Edgemont and Bull Run are more than a mile away from the nearest granodiorite intrusions, but most of the deposits, especially those in limestone, are but a few rods away from the deep-seated intrusive rocks, or else they are associated with dikes which are probably connected with the larger intrusive bodies. The deposits of this group are (1) contact-metamorphic deposits, (2) irregular replacement deposits or chambers in limestone, (3) replacement veins and sheeted zones in limestones and shales, (4) fissure veins in quartzites, and (5) fissure veins in igneous rocks.

CONTACT-METAMORPHIC DEPOSITS.

The contact-metamorphic deposits are represented at Bullion (Railroad district), Lone Mountain, Lime Mountain, and Cortez. In point of production these deposits are not so important as other

deposits which are usually associated with them, but with respect to genesis they form a distinct and interesting type. At Bullion the contact-metamorphic ore consists of garnet, calcite, actinolite, tremolite, epidote, quartz, pyroxene, pyrite, chalcopyrite, bornite, galena, and zinc blende. The metals are copper, silver, and lead. All the minerals are intergrown and were formed at the same time. They are in limestone not more than a few rods from the contact with the igneous rock and were formed by gaseous solutions from that rock at the time of the intrusion. At Lone Mountain contact-metamorphic deposits occur near granodiorite but are not extensively developed. The ore consists of calcite, garnet, actinolite, magnetite, pyrite, chalcopyrite, and other minerals and carries values in copper and silver. At Lime Mountain some ore composed of calcite, white and black mica, tremolite, and copper-bearing sulphides is presumably of contact-metamorphic origin. The deposits of the Garrison mine, at Cortez, are in the main chambers of siliceous ore in limestone, but a small amount of contact-metamorphic ore is found near a dike of decomposed porphyry. This ore consists of calcite, tremolite, actinolite, quartz, and sericite, intergrown with which is a small amount of pyrite.

The contact-metamorphic silicates are developed at Mill Canyon, in the Cortez Range, and also at Mountain City, but no contact-metamorphic ore has been found at these places. So far as known there are no contact-metamorphic zones in the Shoshone Range along the border of the intrusive granodiorite, which nearly everywhere breaks through quartzite, a rock that is not favorable for contact metamorphism. Why the granodiorites of the central part of the Centennial Range did not cause contact metamorphism is not easily understood, for all the conditions so far as known seem to be similar to those prevailing where contact metamorphism has taken place.

IRREGULAR REPLACEMENT DEPOSITS IN LIMESTONE.

The irregular replacement deposits or chambers in limestone are among the most important ore bodies in the area studied. They include the silver-lead deposits at Bullion, the silver deposits at Cortez and Mineral Hill, some of those at Mill Canyon, and probably some of the inaccessible deposits at Lewis and Lone Mountain. These deposits are found in areas which are intruded by the granitic rocks or the related porphyries, and for the most part they have the form of chimneys or ribbons that are related to the intersections of fissures rather than to bedding planes. In all these deposits silver is the most important metal, but there are important amounts of lead and copper in the deposits at Bullion. Gold is usually present, but always in subordinate quantity. As a rule the ore is highly siliceous. The gangue minerals are quartz, barite, and calcite.

The principal sulphides are pyrite, galena, zinc blende, argentite, and chalcopyrite. Stibnite, stromeyerite, gray copper, polybasite, stephanite, and other minerals containing arsenic or antimony are present in some of the ore. Contact-metamorphic silicates are wanting. The ore bodies of this group are without exception near the deep-seated intrusives and are believed to have been deposited by solutions which were given off from these rocks as they cooled.

REPLACEMENT VEINS AND SHEETED ZONES IN LIMESTONE AND IN SHALE.

The replacement veins and sheeted zones in limestone and in shale are in many respects similar to the deposits just described, but instead of chimneys and irregular masses they are thin tabular bodies. The silver deposits at Columbia and Aura, in the Centennial Range, and some of the deposits at Lewis, in the Shoshone Range, belong to this class. The deposits occur in areas of calcareous sedimentary rocks intruded by granodiorites and related igneous rocks. Some of the veins are parallel to the bedding of the country rock, but most of them cut across it. They are clearly related to fissures and zones of movement and are apt to be wider in limestone than in shale, because the limestone is more favorable for openings and more readily replaced by the vein-forming solutions. The sheeted zones are in most respects similar to the replacement veins, but were deposited in two or more narrow, closely spaced, approximately parallel openings, and they usually include more or less of the brecciated country rock. As a rule the ore is highly siliceous. The gangue minerals are quartz, barite, and calcite; the sulphides are pyrite, galena, zinc blende, chalcopyrite, stibnite, and gray copper. Ruby silver and argentite are present in some of the deposits. At Mountain City and Mill Canyon there are some undeveloped prospects of ferruginous oxidized gold ore which probably belong to this group.

FISSURE VEINS IN QUARTZITE.

The fissure veins in quartzite are of considerable economic importance, for they include the gold deposits at Edgemont and at Bull Run and some of those at Dean. These deposits occur in areas of intrusive granodiorite or granodiorite porphyry, but those at Edgemont are more than a mile from known outcrops of igneous rocks. The ore is simple in composition, the gangue is quartz, and the sulphides are present only in small amounts. The primary sulphides are pyrite, galena, and arsenopyrite, with a small amount of chalcopyrite. Silver is present in subordinate quantity, but there is not enough copper to interfere with cyanide extraction. The gold is in the quartz and sulphides and a large proportion of it is free milling.

FISSURE VEINS IN THE OLDER INTRUSIVE ROCKS.

The group of fissure veins in the older intrusive rocks includes the principal deposits at Mountain City, Dean, Grey Eagle, and Mill Canyon. All of these deposits are in clear-cut fissures in granular rocks or in their porphyries, and were formed presumably soon after the intrusives had solidified. The veins were deposited by hot waters which probably rose from the deeper portions of the cooling intrusive mass. Along all these deposits the wall rock shows the effects of hydrothermal metamorphism: Pyrite, sericite, and quartz are everywhere developed in the wall rock near the veins and most extensively near the larger deposits. Calcite is usually associated with the sericite and was presumably formed at the same time. Adularia is not present. In the deposits at Mountain City, Grey Eagle, and Mill Canyon silver is the principal metal, but the ore carries also gold and lead. The sulphide ore is composed of quartz, pyrite, galena, zinc blende, gray copper, argentite, gold, arsenopyrite, and a little chalcopyrite. The deposits in granite at Mill Canyon are of the same general composition as those at Mountain City. At Dean the gold veins in granodiorite carry a very small amount of gray copper.

LATER DEPOSITS (MIOCENE).

After the early igneous activity had subsided there was a period of relative quiet, during which the country was eroded and supplied sediments for the lake lying just east of this area. In this lake the Eocene beds were deposited. In Miocene time volcanism on a grand scale was repeated. The first eruptions were rhyolites and later these were intruded by andesites and at some places covered with basalts. So far as the ore deposits are concerned the rhyolite and andesite only need be considered. Both of these form the country rock for the deposits of this group, but the andesites seem in all places to be the agents of mineralization. All the ore bodies of this group are in the andesites or in rhyolite near intrusive andesite. As the andesites were intruded into rhyolites, which are of Miocene age, and preceded basalts, which are Pliocene, they must have been intruded at or near the close of the Miocene. The ore deposits are therefore believed to be of late Miocene age. Conservatively estimated, they have yielded about \$28,000,000. A list of the minerals of these deposits is given below:

Adularia.	Chalcanthite.	Kaolin.	Pyromorphite.
Argentite.	Chalcocite.	Limonite.	Quartz.
Arsenopyrite.	Chalcopyrite.	Malachite.	Sericite.
Azurite.	Chlorite.	Manganite.	Sphalerite.
Barite.	Chrysocolla.	Muscovite.	Stephanite.
Bornite.	Enargite.	Orthoclase.	Stibnite.
Calcite.	Freibergite.	Pyrargyrite.	Tetrahedrite.
Cerussite.	Galena.	Proustite.	Turquoise.
Cerargyrite.	Gold.	Pyrite.	Zinc blende.

The principal deposits of this age may be divided into the following groups: (1) Fissure veins and sheeted zones in andesite; (2) fissure veins and fracture zones in rhyolite.

FISSURE VEINS AND SHEETED ZONES IN ANDESITE.

The deposits in andesite include the silver veins of Tuscarora and the principal deposits at Cornucopia, Burner, Falcon, and Stafford. They are banded siliceous veins which carry a variable amount of sulphides. Some pyrite, galena, and zinc blende are present, and in the richer deposits gray copper, ruby silver, stephanite, and other antimony and arsenic sulphides occur in appreciable quantities. Silver is the principal metal won, but the ore usually contains an important amount of gold.

Hydrothermal metamorphism of the propylitic type is everywhere pronounced in the andesite near these deposits. It is more conspicuous because it produces striking color changes in the rock. Sericite and pyrite are very extensively developed, and at Tuscarora adularia forms in considerable quantities. Between the most-altered phases of the andesite and the fresh, unaltered rock, chlorite, resulting from the decomposition of biotite and hornblende, forms in great abundance and some carbonates are deposited. At such places the rock is dark green and grades into the brownish-gray andesite on one side and into white decomposed sericitic andesite toward the ore deposits. The hydrothermal metamorphism of the andesite is confined to the areas of the ore deposits and was accomplished by the same solutions that deposited the ore in the fissures. The amount of change is directly proportional to the mineralization and the extent of the area affected depends on the distribution of the fissures. The bulk of the deposition was made in the open spaces, but where the walls were strongly fractured they were replaced by the ore minerals. Compared with the alteration of the walls caused by the vein-forming solutions along the veins in granodiorite, the changes along the younger veins in andesite are more extensive and the area affected is greater. There has been less erosion since the ores in andesite were deposited and consequently the ore bodies exposed were formed under thinner cover than the deposits associated with the older intrusive rocks. Lindgren has explained the extensive alteration which is almost invariably shown near the deposits in the younger eruptive rocks by the fact that the light covering at the time of deposition favored porosity and open spaces, which permitted the solutions to penetrate the rocks with greater freedom.

FISSURE VEINS AND FRACTURE ZONES IN RHYOLITE.

The ore bodies in rhyolite occur near intrusive andesite. The group includes the deposits of the Dexter mine, at Tuscarora, the

lodes at Gold Circle, and some prospects which have lately been discovered in the Lynn district. Gold is the principal metal, but a small amount of silver is present. Hydrothermal metamorphism is noticeable, but as a rule it is less intense and much less conspicuous than in the andesite. It consists in the main in the development of sericite, pyrite, and secondary quartz, and at the Dexter mine considerable adularia has been deposited. Chlorite is much less abundant than in the altered andesite and carbonates are wanting. The ore is very simple in composition. Most of it consists of quartz and pyrite. The deposition has usually taken place in many small openings in a broad zone of fracturing rather than in wide open spaces.

PLACER DEPOSITS.

Placer deposits are relatively unimportant in this area and have not been discovered at most of the camps. They have been found at Tuscarora, Aura, Van Duzer Creek, and Lynn. At Van Duzer Creek and Aura they are derived from the disintegration of the lode deposits of the earlier group; at Tuscarora and Lynn from the younger lodes in the Tertiary eruptive rocks.

The gold placers at Tuscarora were worked extensively in the seventies and are said to have yielded about \$7,000,000. At present no work is being done except that carried on by a few Chinese. The gold occurs as dust and as nuggets of considerable size. Its source is presumably some gold lodes which occur in rhyolite and andesite to the north and west of the diggings. A large acreage of ground west of Tuscarora has been located and sampled with drills. It is said that much of this ground will pay to work with dredges.

At Van Duzer Creek, which is between Aura and Mountain City, placer mining was carried on during the summer months for a number of years. Two small reservoirs have been built in this creek and steel pipes have been laid from these to supply water for monitors. There are few large boulders in this gulch, and when the water supply is sufficient operations may be carried on with success. The source of the gold consists of some undeveloped veins at the head of the stream. The mines were not worked in 1908.

In the Lynn district some of the gulches which head in an area of mineralized rhyolite carry placer gold and a few hundred dollars was recovered in pans and rockers during the season of 1908.

An attempt has been made to work the gravels of Bull Run Basin, near Aura, and an extensive system of ditches and flumes has been installed. After a few hundred yards had been washed out the project was abandoned. It is said that some gold was recovered, but on account of the size of the boulders in the gravels their exploitation was unprofitable. Some notes on the placers are given where the camps are described.

PROSPECTING.

From what has been said it is clear that the deposits of the older group are confined to the granular rocks and deep-seated porphyries and to the sedimentary formations in areas which have been intruded by those rocks. It would be worth while to scrutinize closely the country along the margins of the granular intrusive rocks and especially the limestones and quartzites some distance away. These rocks may be but little altered in the immediate vicinity of the deposits, and except where garnet zones are developed there may be in the country rock itself no conspicuous changes indicating mineralization. Silicified outcrops, iron-stained rocks, or gossans may mark the presence of the older deposits in sedimentary rocks. The ore-bearing solutions seem to have been capable of traveling farther with their burden when they were moving in fissures formed in quartzite than in the limestone. Quartzite is not readily replaced, especially by solutions rich in silica, and consequently the deposition does not occur until the solutions are cold enough to deposit by simple precipitation rather than by interchange of molecules with the calcareous wall rock. Deposits in quartzite are therefore likely to be farther from the intrusive granular rocks than the deposits in limestone. At Edgemont the lodes fill open spaces in quartzite and are more than a mile away from the nearest known intrusives. Along the lodes in the granodiorites the country rock is usually altered and leached white or pale green, but such alteration may not extend far from the lodes.

As already stated, a large part of the area is covered by late Tertiary lava flows, chiefly glassy rhyolites and vesicular basalts. The basalts are not known to contain any ore deposits, and the rhyolites are probably barren except where cut by later intrusives. The surface flows do not seem to have been agents of mineralization. If the magmas which formed them carried ore-bearing solutions these must have escaped during eruption and were lost. The deposits associated with the eruptive rocks of the younger period are in the intrusive andesites or in rhyolites cut by such intrusives. In prospecting for these deposits search should be made for the andesite, a dark-gray or brown rock easily recognized in the rhyolite areas. It is not so dark as the basalt and does not contain the green olivine crystals which are characteristic of that rock.

The later Tertiary deposits are limited also to areas marked by strong hot-water action. The prospector who is familiar with the deposits in the later eruptive rocks is well aware that they are confined to the leached areas. The "kindly look of the rock," or the dull white appearance resulting from devitrification, sericitization, and kindred processes, is quickly recognized. At many places these light-colored chalky areas may be distinguished far away, especially

where they are located on bare slopes and are surrounded by the darker andesite or by the glassy pinkish rhyolites or dark obsidians. Such areas of mineralization may yet remain to be discovered, but those which are now unknown are probably in obscure places partly covered by vegetation, soil, or rocky débris.

MOUNTAIN RANGES AND MINING DISTRICTS.

OWYHEE BLUFFS.

GENERAL FEATURES.

The Owyhee Bluffs, which are in the northwestern part of the area covered by this reconnaissance, form a lofty ridge trending north-eastward toward the Independence Mountains. Rose Mountain, one of the highest summits of this ridge, reaches an elevation of 7,949 feet, or about 2,000 feet above the level of Squaw Valley, which lies to the southeast of the bluffs. This valley is a broad expanse of agricultural land almost completely shut in by hills and low mountain ranges. Northward from the Owyhee Bluffs the country slopes very gently to the Owyhee Desert, a great expanse of level country which extends northward far into Idaho. Through this great plain the small tributaries to Little Humboldt River and to the forks of Owyhee River have sunk their channels, and here and there small rounded hills relieve the monotony of the otherwise featureless landscape. Owyhee Desert is better watered than the name implies and during a part of the year affords subsistence for live stock.

GEOLOGIC FEATURES.

The Owyhee Bluffs^a are made up almost exclusively of eruptive rocks of Tertiary age. The bedded rhyolites have by far the greatest distribution, forming the larger part of the crest of the ridge and extending downward to the base of the slopes. The rhyolites belong to the extensive series of flows which cover the larger part of the area of this reconnaissance north of Humboldt River. About 5 miles northeast of Midas, along the floor of a small canyon, there is a small area of shaly limestones surrounded by rhyolite, and so it is inferred that these flows rest upon the eroded surface of sedimentary rocks. The rhyolites are cut by andesite dikes and here and there are overlain by andesite flows, one of which is capped by rhyolite.

Squaw Valley borders the Owyhee Bluffs on the southeast and, like many of the smaller areas of low land between the mountain ranges, is covered for the most part with a deposit of Quaternary gravels, but in the lower end of this valley there are some stratified deposits of volcanic ash^a which on lithologic grounds have been referred to the Humboldt formation, of Pliocene age.

^a Emmons, S. F., U. S. Geol. Expl. 40th Par., vol. 3, 1870, p. 612.

MIDAS OR GOLD CIRCLE DISTRICT.

LOCATION AND HISTORY.

The Gold Circle district is situated in the hilly country along the southeastern slope of the Owyhee Bluffs, near the edge of Squaw Valley. It is about 45 miles north of Battle Mountain and approximately the same distance from Golconda, and is connected with both of these stations by stages which make round trips three times a week. In the summer of 1907 gold was discovered on several claims, and in March, 1908, as a result of a number of rich strikes, the district experienced one of the rushes which is characteristic of the method of settlement of mining camps in Nevada. A town site was laid out at Midas, and within a few weeks some 1,500 persons were established in this town. After the first excitement had passed away a majority of the newcomers left, and in September, 1908, the population of the camp had decreased to about 250 persons. Several of the claims were under development, and a number of lodes were being prospected with more or less success. A few tons of rich ore have been shipped to smelters, but the bulk of the ore that has been developed is not of a grade to pay the shipment charges, which are necessarily high, as they include a long wagon haul. The deepest shaft is sunk 200 feet; several other shafts are down 100 feet; and three or four tunnels have been driven to depths approximately 100 feet below the surface. Ground was broken in September, 1908, for a 10-stamp custom mill, which it was planned to erect at once, and two mining companies were contemplating the erection of mills in the near future.

GEOLOGY.

General outline.—The rocks of the Gold Circle district are rhyolite flows and flow breccias which are cut by dikes of andesite and overlain here and there by andesite flows. The rhyolites, which are the oldest rocks exposed in the district, cover the greater portion of the area. They occur in considerable variety, but the most common is a light-colored, dense, streaked rock composed in the main of a glassy or devitrified groundmass which contains scattered phenocrysts of feldspar and quartz. Other phases of the rhyolite are perlitic and some are vesicular. In the vicinity of Queen Canyon, east of the Esmeralda claims, the vesicles of rhyolite are filled with amygdules of beautiful amethystine quartz.

In Queen Canyon and at several other places in the Gold Circle district the rhyolite is highly fissile and thinly bedded, presenting the appearance of a silicified shale. The shaly appearance is probably due to banding that was developed as the rhyolite flowed and is emphasized by subsequent weathering and the deposition of iron oxide along the parting planes. The rhyolites include also flow

breccias which contain many angular fragments of streaked rhyolite cemented by a matrix of glassy rhyolite. This brecciation is due to movement of the flow while it was still more or less fluid, but after a crust had formed on the surface. The brecciated rhyolite, after weathering or alteration by hot waters, may closely resemble a friction breccia along a vein, and it may easily be mistaken for ore, unless the fact that the cementing material is glass and not quartz is noted.

The andesite which outcrops at many places in the district is a dark, fine-grained porphyritic rock, cutting through the rhyolite or forming flows interbedded with it. Under the microscope it is seen to be composed of a brown glassy groundmass containing crystals of andesine and labradorite feldspar, a considerable amount of augite, some magnetite, and a little quartz. Augite and the brown groundmass are partly altered to serpentine, calcite, and sericite. The feldspars contain too much soda for basalts, and no olivine was found in any of the thin sections. Dikes of andesite in rhyolite are well exposed on the Dixie claim, about half a mile northwest of Midas. The andesite caps the rhyolite, at some places forming only a thin veneer above it, as is shown on the Iron Mask claim, 1 mile east of Midas, and on the hillside to the south of this claim. Some of the andesite is highly vesicular, and the larger portion of it undoubtedly forms surface flows, the dikes representing the vents through which the flows rose to the surface. About $1\frac{1}{2}$ miles N. 15° E. of Midas, a few rods north of the Elko Prince Annex claim, there is a hill which is composed almost entirely of andesite. The rock forming the lower portion of the hill is solid porphyritic andesite; the upper 50 feet is highly vesicular and probably represents the upper portion of the same flow. Above the vesicular portion of the andesite is a bed of rhyolite, which represents a flow that was poured out subsequent to the extravasation of the andesite. In the country to the north of the Gold Circle district another bed of rhyolite was noted above andesite. Although the main mass of the rhyolite was erupted before the andesite, it is very clear that some of it is later than the andesite. This sequence is suggested also by a dike of felsitic quartz porphyry similar to rhyolite in composition which cuts andesite about one-fourth mile northwest of the Midas mine.

Hydrothermal metamorphism.—In a view of the rugged southeastern slope of the Owyhee Bluffs from the south, the Gold Circle district, including an area some 3 miles square, stands out in sharp contrast with the surrounding country. The rocks in this area are leached to a chalky white, stained to a light brown here and there by iron oxides. In the fresh glassy rhyolites which surround this area shades of pink and greenish gray predominate, the coloring matter being due to a very small amount of iron present in the glass. In the region of the ore deposits, which is in a broad way coincident with

the leached area, hot solutions have soaked into the country rock, causing devitrification of the glass and other mineralogical changes. Near the lodes, where the action was more intense, pyrite, quartz, chlorite, and sericite have been formed in the rhyolite. The few slides studied indicate that these minerals are rather closely restricted to the country rock within a few feet of the lodes and that devitrification has taken place farther away. The changes in andesite are less intense than in rhyolite; some feldspars are slightly sericitized and augite is partly altered to calcite and chlorite. Calcite, which is present in considerable quantity in the altered andesite, seems not to have formed in rhyolite, and that in the andesite may have been formed altogether by surface waters subsequent to the deposition of the ores.

Fissuring and faulting.—There has been considerable fissuring and faulting since the eruption of the rhyolite and andesite. All the ore deposits are related to planes of movement. At the Rex mine, on the Gold Circle claim, in the Sleeping Beauty tunnel and elsewhere, andesite and rhyolite are in faulted contact. The geologic sketch map (fig. 3) indicates the approximate distribution of the rhyolite and andesite. In work of a more detailed character the separate rhyolite flows could probably be distinguished and the details of the faulted structure could be worked out, but this was not done in the few days which were given to the study of the district. The lodes are plotted with a greater degree of accuracy than the boundaries of the geologic formations, which, at many places, were not traversed.

ORE DEPOSITS.

General features.—The deposits in the main are replacement veins and sheeted zones in rhyolite, which are located along prominent slickensided planes of movement. All the fissures strike northward and are with a few exceptions approximately parallel. In general they dip from 65° to 85° NE. In the commonest type a few inches of high-grade iron-stained siliceous ore occurs here and there along the slip planes, and in places the surrounding country rock for a distance of several feet is shattered and seamed with veinlets of quartz carrying gold. Thin drusy cavities with well-formed crystals of colorless quartz are found locally in these veinlets, and at some places this quartz is banded with a black silver-rich mineral, probably argentite. The rhyolite near the vein is devitrified, silicified, stained with iron oxide, and at many localities replaced by ore. In the St. Paul mine, where dark silver-bearing sulphides and pyrite occur in banded ribbons alternating with quartz and parallel to the walls of the vein, the deposit is a simple fissure filling, but in most of the lodes where the sulphides are shown the original openings were small and the deposition was mainly through replacement or impregnation




FIGURE 3.—Geologic sketch map of Gold Circle mining district,

of the rhyolite. The primary ore minerals are pyrite, quartz, gold, and probably argentite. The secondary minerals are quartz, iron oxide, manganese oxide, and horn silver. The gold is almost without exception associated with pyrite, with iron oxide, or with quartz highly stained with iron or manganese. At some places the sulphides begin to appear within a foot or two of the surface. The depth of the partly oxidized zone is from 100 to 150 feet below the surface.

Prospecting the lodes.—Well-defined fissures are very conspicuous in many of the replacement deposits. Some of these were formed before the ore was deposited, as is shown by the thin tabular bodies of oxidized ore which occur along some of the fissure planes and by the surfaces of the planes, which are corroded so that they do not show the polished striated surfaces that movement planes commonly exhibit. Along some of the lodes there are, however, slickensided planes which are clearly later than the ore. These are usually polished smooth and striated, and the ore along them is more or less brecciated. In some of the lodes this ore is ground to a gouge containing well-rounded fragments of quartz.

In developing a lode it is important to follow the fissures which were formed before the ore was deposited rather than those which are of later origin. In the oxidized zone it is difficult to distinguish the two classes of fissures, especially where there has been movement before and after deposition approximately along the same plane. The corrosion of the surface of the fissure and the presence of a thin tabular body of uncrushed quartz may distinguish these fissures from the slickensided planes of postmineral movement, which, in places, may carry crushed ore. Probably most of the faulting that brings the andesite into steep contact with the rhyolite took place after the deposition of the lodes, for, as a rule, such fault contacts show slickensiding or striæ, in contrast with the etched or roughened surfaces of fissures along which the uncrushed ore occurs. Where the lodes are walled on one side by the andesite they are, as a rule, highly crushed, but the andesite is relatively fresh, or at least is not nearly so much altered as the rhyolite. A faulted contact between these two rocks should be investigated with a view to finding a vein, for many of the planes of later faulting followed the zone of earlier fissuring, but it is not good prospecting to drift for great distances along faults which do not show mineralization or crushed ore, or which do not pan gold in the soil along the surface. Developments thus far have not shown any faults which cut across the lodes and displace the ore.

Ore shoots.—Notwithstanding the continued activity of lessees since the early discoveries, the search for shipping ore has thus far proved disappointing. Rich pannings and small bunches of high-grade ore are found near the surface in many places, but the present

state of the mines indicates that the production will have to come in the main from lower-grade deposits of milling ore. Some of the fissures along which the ore is found are regular and persistent, and have been opened here and there for half a mile or more along the strike, but these, where developed, do not carry ore of milling grade throughout their length, and development is not sufficient to show how extensive the ore shoots are. In the Rex mine a body of good milling ore 180 feet long has been developed on three sides to a depth of 65 feet, and there is also a considerable tonnage, partly in the sulphides, in the workings along the Gold Crown lode. A number of leases are in ore of milling grade, but these ore bodies have not been sufficiently developed to be regarded as ore in sight. On several of the undeveloped claims the surface showings seem sufficient to warrant further prospecting for milling ore.

Secondary enrichment.—The primary ore is auriferous pyrite and quartz, with which are associated a silver-bearing sulphide and other minerals. As the surface is worn away such ore is oxidized and the sulphur, together with most of the silica and iron and some of the gold, is carried away, but a larger proportion of the gold remains. As a result of this process there is likely to be a concentration of the gold in the upper part of the deposits, but to what extent such concentration has taken place in this district is not known. Some of the little seams of rich ore are solid and appear to have suffered slight change except oxidation. Some very good values have been found in the sulphide ore about 200 feet below the surface of the Gold Crown shaft, but there has been so little exploration in the primary sulphide ores that it is not possible to compare its value with that of the oxide ore.

RÉSUMÉ OF GEOLOGY.

A study of the Gold Circle district shows the following geologic history. In Tertiary time, probably in the Miocene, extensive flows of rhyolite were poured out upon a surface of Paleozoic sedimentary rocks. Subsequently the rhyolites were fissured and through these fissures andesite flows rose to the surface and covered the rhyolite. A portion of the magma remained in the fissures, forming dikes. The andesite was in turn cut by fissures which were filled with an acidic magma, of which one portion flowed out upon the surface and formed rhyolite and another portion, solidifying in the fissures, formed dikes of quartz porphyry. After the eruption of the andesite the country rock was strongly fissured, most of the planes of movement striking northwestward and dipping steeply to the northeast. Along some of these fissures auriferous pyrite and quartz with silver-bearing sulphides were deposited, the solutions dissolving portions of the country rock and replacing it where conditions were favorable with ore and other minerals. From the fissures the solutions spread

out into the country rock, causing devitrification and other changes. A second fissuring with some displacement occurred after the ores were deposited. These movements were mainly along the lodes and brecciated the quartz and sulphides. As a result, the ore shoots, which were already irregularly spaced along the earlier fissures, were strung out along a plane of later movement. As the rocks were eroded the ores were oxidized, pyrite changing to limonite, sericite and feldspar to kaolin. Hydrous silica was deposited in crevices, where it is associated with free gold and manganese oxides.

MINE DESCRIPTIONS.

Rex mine.—The Rex mine is on the eastern slope of a low ridge about a mile east of Midas. A shaft driven at an inclination of 64° is sunk to a depth of 65 feet. From the bottom of the shaft a drift is run 30 feet to the north and 120 feet to the south, with short crosscuts here and there. The lode is along a fault between rhyolite and andesite. It strikes about N. 15° W. and dips 66° W. It is a zone of crushed, silicified, iron-stained rhyolite from 5 to 16 feet wide and carries, according to C. G. Rothschild, from \$5 to \$28 a ton. The rhyolite is a dense, light-colored rock with a few phenocrysts of feldspar and quartz. In the lode it is silicified, iron stained, and cut by veinlets of quartz. The ore developed is highly oxidized, but a little pyrite is present in the bottom of the mine. There is a well-defined slickensided plane of movement along the foot wall between the andesite and the ore, and other fissures approximately parallel to this one cross the ore zone. The ore is restricted to the altered rhyolite, and the andesite, even where greatly crushed, is said to be barren. For this reason it seems probable that the displacement which brought the rhyolite and andesite into contact occurred after the deposition of the ore. Figure 4 is a plan of the Rex mine on the 65-foot level.

Gold Crown lode.—The Gold Crown lode between the Rex mine and Midas has been developed in a number of shafts, pits, and tunnels for a distance of nearly 3,000 feet along the strike. It is a zone of shattered rhyolite which strikes N. 67° W. and dips about 65° N. Wherever the lode has been developed there is a well-defined fissure which at some places is slickensided and carries gold values in crushed quartz. At the Snowstorm lease, at the west end of the lode, half a mile east of Midas, a vertical shaft is 84 feet deep and short levels are turned 70 feet and 84 feet below the surface. The shattered rhyolite for a width of 25 feet is said to carry milling ore. In the Climo lease, farther east on the lode, several pits and short tunnels expose a regular fissure along which some gold values have been obtained. Still farther east the lode is developed in the lower tunnel of the Gold Circle Crown Mining Company, where it is a wide zone of shattered rhyolite, through which considerable pyrite is disseminated. East of this tunnel, along the strike of the lode and higher on the hill, is a

second tunnel driven by the same company, and in this the lode is most extensively developed. The upper tunnel is a crosscut for a distance of 55 feet to a point where it encounters along the foot wall of the lode a smooth fissure that strikes S. 60° E. and dips 65° NE. This fissure is followed in the tunnel for a distance of 400 feet, and a shaft 200 feet deep is sunk in the hanging wall of the same fissure and intersects the tunnel at a depth of 80 feet. Small bodies of rich ore, from 1 to 10 inches wide, are found here and there along the fissure and the hanging-wall rhyolite is shattered, crushed, and cemented with veinlets of quartz and pyrite. A zone of the crushed rhyolite about 3 feet wide is said to be good milling ore. Some crushing has taken place since the ore was deposited, for fragments of quartz rounded by attrition are found here and there along the fissure in a mass of crushed leached rhyolite. The oxidized ore extends below the surface to a depth of about 110 feet, where the sulphides are encountered. The values in the oxidized and in the sulphide ore, so far as developed, are said to be approximately the same. The Emancipation lease is about 1,500 feet east of the Crown shaft and on the strike of the Gold Crown lode. Ninety feet below the collar of an inclined shaft driven in rhyolite there is a fissure which strikes south-eastward and dips 58° NE. and carries pockets of good milling ore.




FIGURE 4.—Plan of Rex mine, 65-foot level, Midas (Gold Circle) district.

The Emancipation lease is about 1,500 feet east of the Crown shaft and on the strike of the Gold Crown lode. Ninety feet below the collar of an inclined shaft driven in rhyolite there is a fissure which strikes south-eastward and dips 58° NE. and carries pockets of good milling ore.

St. Paul-Banner lode.—The St. Paul mine is about a mile southeast of Midas. A shaft inclined 76° E. is sunk on the lode to a depth of 132 feet and levels are turned 60 and 120 feet below the surface. The country rock is rhyolite and rhyolite flow breccia. The rhyolite near the lode is altered to kaolin and sericite and contains small crystals of pyrite, but these are said to be barren. The lode is a simple fissure filling which has a maximum width of about 12 inches and carries high values in gold and silver. It strikes northwestward and dips steeply to the northeast. The ore is oxidized to a depth of 70 feet. Below this depth it consists of banded quartz and dark argentiferous sulphides with faint red bands that are probably ruby silver. This deposit at the St. Paul mine differs from the other ore bodies of the Gold Circle district, the ore being a banded fissure filling rather than a replacement vein. A few tons of ore carrying \$100 to the ton have been shipped to smelters.

To the northwest, in the line of the strike of the St. Paul lode, exposures show values in several open cuts toward the Banner ground, and in one of these a little horn silver was found. On the Banner claim a fissure which is presumably the same as that of the St. Paul is exposed in three shafts. In the north shaft fine-grained andesite occurs on the west side of the fissure, which just north of this point passes through a flow of andesite that caps the rhyolite. On the Ripsaw claim, still farther north and in the strike of the Banner lode, a fissure with approximately the same strike is exposed in a long trench, where it dips 60° E. The Reco lode (No. 5) is 1,300 feet southwest of the Banner lode and approximately parallel to it. A shaft is sunk on a fissure in rhyolite which dips 86° E.

Golden Chariot claim.—On the Golden Chariot claim, 1 mile southeast of the St. Paul mine, three pits are sunk on the Gibson lode, which strikes a few degrees west of north and dips steeply eastward. Along the lode the rhyolite is highly shattered and stained with iron oxide. At the bottom of one of the pits there is a streak of rich ore composed of soft, rotten rhyolite, black manganese oxide, and hydrous silica, with numerous small flakes of free gold. On the Chariot vein near by a 50-foot shaft is sunk in shattered decomposed rhyolite, which carries pockets of rich ore.

Esmeralda mine.—The Esmeralda claim is about 3,500 feet south of the Golden Chariot. On the Charters-O'Byrne lease on this claim a 45-foot shaft is sunk on the promising outcrop of a lode which dips 76° SW. The strongly shattered rhyolite for a width of 3 feet along the lode carries low-grade gold ore. On the Riddle lease on the same lode, about 300 feet to the southeast, a shaft is sunk 50 feet in shattered iron-stained rhyolite, which is strongly mineralized and carries pockets of high-grade gold ore.

Water Witch mine.—The Water Witch claims are about half a mile north of Midas. In the Benan lease, on the Water Witch fraction, a

shaft is sunk in rhyolite on a sheeted zone which strikes a few degrees west of north. Along a prominent plane of movement there is from 1 to 12 inches of rich gold ore. A 3-foot zone of shattered rhyolite is said to carry \$20 to the ton. The screenings of the dump are said to carry shipping values.

Elko Prince lode.—The Elko Prince lode, about 1 mile northeast of Midas, is in altered rhyolite, strikes northwestward, and dips steeply to the northeast. It is developed in a shaft about 100 feet deep and in several surface pits near by. In places on the surface the lode is a sheeted zone of iron-stained rhyolite, but in the shaft it is a banded siliceous filling of an open space about a foot wide. The ore is said to carry \$20 a ton in gold and silver.

Midas mine.—The Midas claim is about 1½ miles north of the town of Midas. A shaft 100 feet deep is sunk in rhyolite on a sheeted zone which is from 2 to 5 feet wide and is said to carry \$8 to the ton in gold. The rusty ore from a small pay streak shows numerous specks of free gold.

Gold Circle claim.—The Gold Circle claim is about 1,200 feet northwest of the Midas. A zone of crushed silicified and highly iron-stained rhyolite along a fault between andesite and rhyolite pans free gold liberally. This deposit was discovered only a few days before the camp was visited by the writer and not more than 10 feet of work had been done.

Belvoir claim.—On the Belvoir claim, one-half mile south of the Midas, a sheeted zone of rhyolite is exposed in small pits and trenches. Some of the small fissures are filled with crushed rhyolite, cemented with iron-stained quartz which carries gold and silver.

Eastern Star mine.—The Eastern Star mine is on Frazier Creek about 4½ miles east of Midas, in an area of white devitrified rhyolite which is similar in appearance to that of the Gold Circle district. The two bodies of leached rhyolite are not directly connected, however, for a large area of fresh vitreous rhyolite lies between. A tunnel is driven northward for 160 feet along the strike of a zone of silicified iron-stained rhyolite which outcrops boldly on the summit of a low ridge. It is cut by three or four parallel veinlets up to 10 inches wide, which carry ribbons of quartz and argentite and in places show a liberal amount of free gold. Between the veinlets and for some distance on either side the rhyolite is impregnated with finely divided pyrite and other opaque minerals, so that a considerable mass of it is as dark as andesite.

INDEPENDENCE RANGE.

GENERAL FEATURES.

The Independence Range is a compact group of lofty mountains east of the Owyhee Bluffs and separated from them by a low pass

which is crossed by the road from Gold Creek to Burner. The range may be regarded as a northward extension of the Cortez Range, but the two are separated by a relatively low saddle between the headwaters of Soldier Creek and Independence Valley. Several of the highest mountains reach elevations from 9,000 to 10,000 feet above the sea, or about 4,000 feet above the level of Independence Valley.

As shown in the atlas of the Fortieth Parallel Survey, the range is made up mainly of thick rhyolite flows which here and there are eroded away to expose large outcrops of sedimentary rocks, chiefly the Weber quartzite. The rhyolite extends westward a great distance from Tuscarora and covers a large area surrounding Squaw Valley, including Gold Circle and other camps near by. At Tuscarora, Cornucopia, Good Hope, and Falcon the rhyolite is cut by large bodies of intrusive andesite. A body of granodiorite occurs near the headwaters of Willow Creek.

The mining camps situated in the Independence Range are Tuscarora, Cornucopia, Good Hope, and Falcon. The ore deposits are silver-bearing fissure veins of the Tertiary group and are in andesite or in rhyolite near intruding andesite. No deposits have yet been discovered in the sedimentary rocks or associated with the granodiorite intrusive rock.

TUSCARORA.

HISTORY.

Tuscarora is situated on the southeastern slope of Mount Blitzen, at an elevation of about 6,200 feet above sea level. It is near the west margin of Independence Valley, a broad area of flat hay land drained by Owyhee River, and is about 50 miles northwest of Elko, with which it is connected by a daily stage that also connects at Tuscarora with stages for Edgemont, Aura, Mountain City, and other northern points.

Placer deposits were found at Tuscarora in 1867, and several years later rich silver veins were discovered. In the seventies and eighties a number of silver mines were opened and a large production was sustained for a number of years. Most of the ore was milled at Tuscarora; only the very high grade ore was shipped to smelters. The six silver mills which were in operation employed the Reese River process, by which the ore was dried, stamped, roasted with salt, and amalgamated in silver pans. The Grand Prize mill employed a combination process, the ore being concentrated over vanners and the concentrates roasted and subsequently amalgamated in pans with the raw tailings. The silver mills had an aggregate of 80 stamps and are said to have given a satisfactory extraction.

The Dexter mine, which is a large deposit of low-grade gold ore, was discovered after the silver mines had been producing for several years. This mine was worked until 1898, when operations were discontinued

on account of the great volume of water which was encountered. The Dexter ore was treated in the Dexter mill, originally a 40-stamp amalgamating mill operated by electric power. Subsequently four Ellis tables and fourteen cyanide tanks with a capacity of about 450 tons were added to treat the tailings. In 1908 a second cyanide plant was installed.

Since 1898 there has been some leasing on the upper levels of the old silver mine, but extensive mining operations have been discontinued. In 1907 Arthur A. Brownlee obtained options on nearly all the mines and organized the Tuscarora Nevada Mines Company. This company spent considerable money sampling the mines and dumps, and is said to have found a large tonnage of low-grade ore in the old workings of the Dexter mine. The company plans to unwater the mine, to build a large cyanide plant, and to undertake many other improvements.

Accurate figures for the production of Tuscarora are not at hand. Various estimates range from \$25,000,000 to \$40,000,000. Most of this was obtained between 1872 and 1886 and the larger portion is silver. The gold placers are reported to have yielded \$7,000,000 and the Dexter mine \$5,000,000 in gold, the various silver mines being credited with the remainder.

When the camp was visited in 1908, practically all the mines were inaccessible. At the Dexter, which is situated at the lower edge of the mineralized area, the water table was within 20 feet of the surface and only some shallow workings and open pits could be entered. In the silver mines, which are at slightly higher elevations, a few workings above the 100-foot level were accessible.

The country rock of the Tuscarora mines is rhyolite and andesite porphyry, which at many places are covered by a thin layer of Quaternary gravel. Most of the deposits are in the porphyry, which in the vicinity of the mines is highly altered. It is the propylite of the Fortieth Parallel Survey. High on the southeastern slope of Mount Blitzen relatively fresh andesite outcrops at several places. The freshest andesite is composed of a dense greenish groundmass containing phenocrysts of andesine, orthoclase, hornblende, and biotite. Toward the mineralized area the andesite is greatly altered. At some distance from the veins chlorite is formed in great abundance and the rock is dark green, but in the highly mineralized area and within a few yards of the veins sericite and iron pyrite have been extensively deposited by replacement. On oxidation this rock alters to a brown iron-stained porphyry, which constitutes a large part of the various mine dumps. Some specimens of what seemed to be the least-altered phases of the porphyry in the mineralized area proved, on examination, to be rich in quartz and orthoclase. It will probably be found that the porphyritic rock commonly regarded as andesite

consists of several related kinds of rock, the separation of which will require great patience even under favorable conditions.

The rhyolite is when fresh a white or greenish dense or almost glassy rock; at many places it is a flow breccia. In the vicinity of the ore bodies it is extensively altered, and even the freshest specimens show, under the hand lens, a considerable amount of secondary pyrite.


In the open cut at the Dexter mine the relations of the andesite and the rhyolite are distinctly crosscutting, one rock intruding the other in a very irregular manner. Owing to hydrothermal action and subsequent oxidation it is not possible to ascertain which rock intrudes the other, but if the rhyolite is a flow, as is indicated by the banded shaly phases, then the andesite is the later rock and is intruded into the rhyolite.

ORE DEPOSITS.

The ore deposits, so far as could be ascertained from the limited observations which were possible when they were visited, are silver lodes in andesite, fractured zones or stockworks of gold ore in rhyolite, and gold placers.


Silver lodes.—The silver veins occur in an area of highly altered andesite, which contains small masses of rhyolite. Most of them strike a few degrees west of north and dip toward the west. The gangue is mainly quartz. The ore minerals are ruby silver, enargite, and other silver sulpharsenic and sulphantimony minerals, silver glance, galena, pyrite, and arsenopyrite. A little chalcopyrite and bornite were noted, but these are not abundant. Here and there is a little malachite, but copper is present only in small traces—as a rule, much less than 1 per cent. Shoots of gold ore are found in some of the silver lodes and most of the silver ore contains gold. Near the surface there was much horn silver and native silver. A single block of horn silver from the Commonwealth mine is said to have sold for \$30,000. The veins are fissure fillings between the walls of porphyry, and in some of them the ore surrounds numerous fragments of brecciated country rock. Locally the wall rock is replaced by workable ore, and at many places in the vicinity of the veins the country rock is said to carry low values in silver and gold.

The Navajo lode, which is about one-fourth mile west of Tuscarora, was the most productive system of veins. It strikes about N. 80° W. in the Navajo ground and it has been followed northwestward for about a mile. The deposits of the Navajo, Belle Isle, North Belle Isle, Nevada Queen, Commonwealth, and North Commonwealth mines are on this lode. At the North Commonwealth it bends and strikes about N. 60° W. The total production of these mines is said to have been about \$15,000,000. East of this lode, and situated in the main on lodes which are approximately parallel to it, are the



Topography from U. S. Geological
 Exploration of the Fortieth Parallel
 Clarence King, Geologist in charge

TOPOGRAPHIC MAP OF SOUTHERN QUARTER OF AREA SHOWN IN FIGURE 2



deposits of the Independence, P. & P., Eira, Silver Prize, Buckeye, De Frieze, Grand Prize, and other mines. According to report, most of the larger ore shoots were at the junction of fissures and pitched toward the northwest.

In the North Belle Isle, on the 70-foot level, several narrow fissure veins strike north and dip from 35° to 80° W. These carry values in silver and gold up to several hundred dollars a ton, and where they join, about 20 feet below the 70-foot level, they make a shoot of ore several feet wide from which \$1,000,000 is said to have been taken.

Gold deposits.—The most important gold deposit at Tuscarora is that of the Dexter mine. This is situated near the contact of rhyolite and andesite and dips northward at a low angle. A zone, mainly in the rhyolite, is strongly fractured for a distance of 1,400 feet east and west and about 200 feet north and south. Pockets of rich ore occur here and there in this zone, and it is said that the whole mass of this rock could be worked profitably by cyaniding on a large scale. The deposit is crossed by numerous closely spaced veinlets of quartz, which strike in all directions. In many of them the fissures are not completely filled, and the centers of the veinlets contain long, narrow druses lined on both sides with well-formed crystals of quartz half an inch long. Locally there is considerable adularia in crystals up to one-fourth inch long, deposited in drusy veinlets with quartz. These veinlets are said to carry high values in gold. The rhyolite of the fissured zone is strongly impregnated with pyrite, even the freshest specimens showing much pyrite under the hand lens. The gold seems to have been deposited through replacement of the rhyolite and also in numerous small open spaces. The deposit is crossed by several faults which strike northeastward and dip toward the northwest. The throw of all these faults is said to be toward the south; the hanging wall on the west side of the fault has moved downward, causing an offset of the ore zone toward the south. The faulting is therefore normal.

Some gold placers are located from 1 to 3 miles west of Tuscarora. They were extensively worked in the seventies and are said to have produced \$7,000,000, but little work has been done of late years except that carried on by a few Chinese. The deposits were worked mainly by ground sluicing. The gold occurs as dust and as nuggets of considerable size. One of these, about one-half gold and one-half quartz, weighed 9 ounces, and many nuggets have been found which weighed more than an ounce. The source of the gold is presumably some gold lodes which occur to the north and west of the diggings.

At the Rose mine, about $1\frac{1}{2}$ miles west of Tuscarora, several low-grade quartz veins have been found, and the country rock, which is altered porphyry, is said to carry appreciable values in gold.

On Beard Hill, 2 miles southwest of Tuscarora, the Surprise group is situated just above some old placer workings. The country rock

is shattered rhyolite, which is cut by many veinlets of iron-stained quartz that pans gold freely. Some of the placers were probably derived from these deposits.

A large acreage of ground west of Tuscarora has been located and sampled with drills. It is said that much of this ground will pay to work with dredges, and two companies are planning such operations. A large number of samples are reported to have given an average of about 14 cents per cubic yard.

FALCON MINE.

West of Tuscarora there is a great area of mountainous country, the higher peaks reaching elevations of 8,000 to 9,000 feet. The rocks are in the main Carboniferous quartzites capped by rhyolite and intruded by andesite and related rocks. A large mass of granodiorite, probably older than the rhyolite and porphyry but intrusive in the sedimentary rocks, is exposed at the headwaters of Willow Creek and Rock Creek.

The Falcon mine, at the head of a small tributary of Rock Creek, is about 12 miles by wagon road west of Tuscarora. The mine was worked from 1879 to 1881 and the ore was hauled to Tuscarora. In 1884 a four-pan silver mill was built, but this was not operated and is now in ruins. The deposit is a fissure vein from 2 to 5 feet wide and is approximately vertical. Two deep shafts are sunk on it and shallow pits are dug at several places. The country rock is andesite, which near the vein is altered to a light-gray rock composed largely of white mica, but the fresh dark andesite is exposed at several places within 300 or 400 feet of the vein. The ore is highly siliceous and contains a small proportion of finely divided pyrite and other dark sulphides, which are banded with the quartz and show comb and ribbon structure. The values are said to have been in ruby silver.

CORNUCOPIA.

The mines at Cornucopia, about 8 miles southwest of the stage station on Deep Creek, were operated actively in the seventies, when they produced, it is said, over a million dollars in silver. The ore was treated by pan amalgamation in a 20-stamp mill at Mill City, 2 miles below the town. The principal mines are the Leopard and the Panther, which were operated through shafts. The Leopard shaft is said to be 800 feet deep. When the camp was visited in 1908 all the deep workings were caved and only some shallow pits and surface stopes were accessible.

The country surrounding the Cornucopia district is a large area of low hills, which in the main are capped with rhyolite and obsidian. Under the microscope the denser rhyolite is seen to be composed of a glassy microlitic groundmass which contains phenocrysts of quartz,





FIGURE 5.—Geologic sketch map showing Panther vein, Cornucopia district.

oligoclase, pyroxene, magnetite, and a little hornblende. It is more basic than most rhyolites of this area and approaches andesite in composition. Some phases of rhyolite are vesicular and some are pumiceous. The rhyolite is cut by intrusive andesite, to which the accessible ore deposits are restricted. An exposure of the andesite in a relatively fresh condition may be observed near the ruins of a stone house at Cornucopia. The rock is a dark porphyry, composed of a glassy groundmass which contains many phenocrysts of basic andesine and pyroxene, with large chloritic patches that seem to have resulted from the decomposition of hornblende. At the exposure named it may be noted grading into a highly decomposed sericitic phase of the andesite, which is white or stained with yellowish-brown iron oxide. Everywhere in the vicinity of the ore deposits the andesite is similarly decomposed. Masses of quartz porphyry occur in the area of the andesite and are probably intruded into it. The quartz porphyry is light colored and is composed of a groundmass, presumably microcrystalline, which contains white mica, feldspars, and small phenocrysts of resorbed quartz. The rock outcrops in the tunnel of the Leopard mine, but none of the deposits, so far as known, is inclosed in it.

The ore deposits are sheeted zones in decomposed andesite. The ore is white quartz, which carries a very small proportion of dark sulphides, forming narrow ribbons in the quartz. Pyrite, argentite, and gray copper are present, and ruby silver is said to have been an important ore mineral. On the surface the ore minerals are mainly horn silver and a yellow mineral which is probably pyromorphite. The proportion of the sulphides present is very small; but they must have been rich, for the ore is said to have carried 400 ounces of silver to the ton for mill runs. In some of the ore the minerals are arranged symmetrically with respect to the walls, the quartz crystals pointing to the center of a druse, showing that the ore was deposited in open spaces. The country rock along the veins is, however, silicified and otherwise altered by the vein-forming solutions, and at some places carried workable values.

The Panther vein in the principal workings southwest of the silicified outcrop shown in figure 5 strikes S. 78° W. and dips 83° N. to 90°. Here underhand stoping has been carried down for a distance of 60 feet along the strike. The country rock is altered andesite, a soft kaolinized mass cut through by veinlets of white quartz. At the west end of the stope a smooth slickensided fault strikes northwestward, cutting off the vein. The surface of the fault shows striæ inclined northwestward to a line along the direction of steepest dip and making an angle of 15° with it. The country to the north has been prospected for the vein, but it has not been discovered in that direction on the southwest side of the fault. If the fault is



Topography from U.S. Geological
 Exploration of the Fortieth Parallel
 Clarence King, Geologist in charge

TOPOGRAPHIC MAP OF SOUTH-CENTRAL QUARTER OF AREA SHOWN IN FIGURE 2

10 5 0 10 Miles

1909

ENGRAVED AND PRINTED BY THE U.S. GEOLOGICAL SURVEY

normal, the vein to the west of it should be found south of the present workings, where possibly it is represented by some poorly defined masses of quartz which outcrop at that place.

A few rods northeast of the surface stope is a prominent ledge which strikes a few degrees north of east. This ledge, which is several hundred feet long, is the altered andesite somewhat silicified, fractured, and seamed with quartz veinlets. To the east of the ledge a vein in altered andesite has been developed for a few feet and some ore has been stoped. This vein is probably the faulted continuation of the silicified outcrop of quartz which lies between the two groups of workings.

GOOD HOPE DISTRICT.

In the Good Hope district, which is about 12 miles southwest of Cornucopia, mining was carried on in the early eighties, when the camp is said to have produced over \$100,000 in silver. The principal mines are the Buckeye and Ohio, the Snyder, and the Page & Kelley. All these mines were inaccessible in 1908 except the Buckeye and Ohio, which was under water above the adit level. The deposits are in the main sheeted zones in rhyolite flow breccia, but altered andesite is exposed at several places and it is probably the country rock for some of the ore deposits. The leaching of the country rock is extensive near some of the deposits, but is not so general as at Cornucopia. Pyritization is, however, more pronounced in the wall rock and the ore at Good Hope contains a much greater amount of the sulphides. The Buckeye and Ohio mine is on Fourmile Creek near its junction with Atlantic Cable Gulch. The mine was operated from 1882 to 1884, and the ore was run through a 5-stamp mill $1\frac{1}{2}$ miles below the mine. This mill was equipped with roaster, pans, and settlers and employed the Reese River process. A small concentrator was built in 1903, but the treatment employed was presumably unsuccessful, as only a small amount of ore was put through it.

On the surface above the adit level obsidian and other varieties of glassy lavas outcrop at several places. In the mine the rhyolite is a light-colored flow breccia, which is locally altered to a white claylike mass. Along a zone of movement it has been converted to gouge.

The lode is composed of veinlets of quartz and sulphides, which include masses of the country rock highly altered and partly silicified. A tunnel driven on the vein for 300 feet southwestward gives a depth below the surface of about 65 feet. At some places stopes are carried to the grass roots and several winzes are sunk upon the vein. The ore is composed of quartz, pyrite, arsenopyrite, freibergite, stibnite, and dark ruby silver.

The vein, as shown in figure 6, is followed by the adit for a distance of 300 feet. At the breast, where it dips steeply westward, it is cut




FIGURE 6.—Plan of adit level of Buckeye and Ohio mine, Good Hope district.

off by a fault which strikes 37° E. This fault follows a vein also and carries much crushed ore and some banded ribbons of unbroken ore. A stope from this vein is carried to the surface. Near the junction of the two veins a second fault, which deviates from the principal fault about 5° , cuts off the west vein. The ore therefore forms a letter V, which points to the south. If the faults are normal the continuation of the principal vein will be found south of the present workings, on the west side of the fault.

Near the head of Atlantic Cable Gulch, about 1 mile above the Buckeye and Ohio mine, several prospects show brecciated rhyolite silicified and cemented by quartz and dark sulphides. One of the lodes outcropping boldly on either side of the gulch strikes $N. 10^{\circ} E.$ and dips $35^{\circ} W.$, and its outcrop forms a V pointing up the gulch. In places it carries a considerable quantity of dark pyritic ore which has been crushed and recemented by white and

barren-looking quartz. At one place an abandoned incline is driven on the lode to a depth of about 30 feet, exposing a considerable mass of quartz and sulphides.

BURNER HILLS.

GENERAL FEATURES.

The Burner Hills, which are some 10 miles west of Good Hope, rise about 800 feet above the broad undulating plain of the Owyhee Desert, which lies to the east and north. From Good Hope to the

Burner Hills this plain is covered by beds of rhyolite and rhyolite-pumice. Nearly everywhere these beds are flat or dip at low angles in various directions, but as they approach the hills the marginal fringe of rhyolitic pumice becomes steeply upturned and dips away on the east side at angles up to 35°. Still higher on the hills siliceous shales with beds of intercalated limestone outcrop at many places. These beds are highly tilted and show a considerable variety of attitudes, but the prevailing dip appears to be away from the central axis of the hills, the summit of which is a fresh andesite showing massive columnar jointing. The andesite intrudes the sedimentary rocks and probably rhyolites also, although the contact at this place was not seen. A number of claims are located on these hills, but the Mint mine is the only one on which any considerable amount of work has been done. This mine was operated in the early eighties and shipped about \$30,000 worth of lead-silver ore to smelters. Active operations were suspended in 1893, and since that time but little work has been done.

MINT MINE.

At the Mint mine a tunnel is driven southwestward for 175 feet to the lode, which it follows for 300 feet. The lode strikes S. 25° W. and is approximately vertical. Here and there stopes have been carried upward and a winze is driven on the lode below the adit level. The ore consists of galena, sphalerite, pyrite, arsenopyrite, and chalcopyrite, in a gangue of quartz and calcite. Near the surface lead carbonate and iron oxide are present. The sulphides and quartz occur as ribbons parallel to the walls or as masses impregnating the andesite, which is somewhat altered by the vein-forming solutions. The high-grade ore is said to be irregular and buncy in the vein, but a zone up to 4 feet wide is regarded by the owners as available for concentrating, as a considerable proportion of the silver values is in galena.

South of the Mint mine are several small veins of iron-stained siliceous silver ore. Some of these cut across the sedimentary rocks; others occur as stringers parallel to the bedding.

CENTENNIAL RANGE.


GENERAL FEATURES.

The Centennial Range, which lies between Deep Creek on the south and East Fork of the Owyhee on the northeast, is about 20 miles long and from 5 to 10 miles wide, its principal axis trending northeastward from Lime Mountain to Montana City. The higher summits rise from 3,000 to 4,000 feet above the Chellis Valley, which topographically is a part of the Owyhee Desert, a great rolling plain of sagebrush that extends northwestward far into southern Idaho.

The range is separated from the Jack Creek Mountains, to the southeast, by Bull Run Basin, which is drained by Bull Run Creek. This stream flows westward through a deep V-shaped canyon that separates the central mountain mass of the Centennial Range from a narrow, lofty ridge which extends southward toward Deep Creek. The highest part of the range is a compact group of mountains that lie between Bull Run Creek and Blue Jacket Canyon and cluster about Porter Peak, the loftiest summit. North of Blue Jacket Canyon the hilly country extends to Mountain City and beyond that northward into Idaho. The topographic expression of the range is due to faulting modified by erosion and to a trivial extent by glaciation.


The rocks are mainly Paleozoic quartzites, limestone, and shales. On the slopes of Porter Peak the prevailing dip is northward, almost at right angles to the principal axis of the mountains. The great faults on which the chief structural features depend have so broken the geologic column that in the absence of fossils it was not possible to determine the age of the rocks satisfactorily in the limited time devoted to the work, but from the lithologic descriptions published by the King Survey for the country to the south it seems highly probable that the Carboniferous formations have the widest distribution. The great quartzite beds which form the southern portion of Porter Peak and which include the ore deposits at Edgemont and at Bull Run are regarded as Carboniferous. This is a medium-grained quartzite, which through great thicknesses shows comparatively slight variations. At the Bull Run mine there are some thin-bedded siliceous, shaly layers, and on the north slope of the hill south of Edgemont some fine conglomerates were noted, but the great mass of the formation is a dull gray or pink quartzite, massive, thick bedded, and strongly jointed, at many places showing too little evidence of stratification to define its attitude. On the ridge north of the stream which flows westward from Porter Peak through Edgemont the quartzite is overlain by a great series of limestones. This series is several thousand feet thick and on the north slope of Porter Peak grades into black shales which still farther north are overlain by a thick series of limestones and shales. The great expanse of hilly country between White Rock and Mountain City was not traversed in this reconnaissance, but at a distance it appears to be composed in the main of sedimentary rocks capped with rhyolite.

At a number of places, including Blue Jacket Canyon on the southeast side of the range and the ridge south of White Rock Canyon on the northwest side, the limestones are intruded by medium-grained granitic rocks. None of the outcrops of these intrusive masses are very extensive, but at the north end of the range they are numerous, and possibly they are connected in depth. A characteristic specimen from one of the larger masses on the ridge south



Topography from U.S. Geological
 Exploration of the Fortieth Parallel
 Clarence King, Geologist in charge

TOPOGRAPHIC MAP OF NORTH-CENTRAL QUARTER OF AREA SHOWN IN FIGURE 2



1909

of White Rock Canyon is a granodiorite composed of oligoclase, quartz, orthoclase, biotite, and hornblende. No garnetization of limestone was noted near these intrusions. At Mountain City the sedimentary rocks are intruded by a granodiorite, which is more coarsely crystalline and carries more quartz and orthoclase than the average granodiorite. North of Aura and also on California Hill, 2 miles above Mountain City, garnet, epidote, tremolite, and actinolite have formed extensively near the contact of granodiorite with limestone.

Here and there in the Bull Run Basin, protruding through the cover of granite, are outcrops of rhyolite and basalt, and in the country to the east, extending to a great but unknown distance, are thick beds of rhyolite with a subordinate amount of rhyolite flow breccia. Very extensive beds of rhyolite occur also in the lower country to the west and south of the Centennial Range. This rhyolite is much younger than the sedimentary rocks composing the main mass of the Centennial Range, and if it were in its original position it would now be above the sedimentary rocks instead of forming the floors of the lowest depressions, such as Bull Run Basin. The present structure was brought about by faulting and tilting. North of Bull Run Creek, on the south slope of the high mountains which form the central portion of the range and near the trail from Aura to the Bull Run mine, there are bluffs of volcanic agglomerates and tuffs composed of rhyolite fragments with a large proportion of diorite porphyry. At the base of the exposed portions of the beds there are some layers of shaly coal. The agglomerate beds are of Tertiary age, and as they dip toward the Paleozoic rocks which form the central and most elevated portion of the range, there must be a fault of several thousand feet between the two systems of rocks.

ORE DEPOSITS.

In 1869 a party composed of Cope, Dixon, and others, going from Silver City, Idaho, to White Pine, Nev., made discoveries and located claims near Columbia and in Blue Jacket Canyon near by. These discoveries attracted considerable attention to the district, and in the early seventies silver mills were built in Blue Jacket Canyon and at Mountain City. In 1875 Edward Stokes built a mill at Columbia to treat the ore from the Revenue, California, and Infidel mines. All these mills employed the Washoe or the Reese River process, and the conditions for operation were most trying, as it was necessary to haul the bulky supplies required for silver milling 75 miles or more by wagon road. Although a considerable tonnage of chloride ore, taken from the upper parts of the deposits, was put through these mills, it is a question whether much profit was made from the operations. Owing to the insuperable difficulties in

the way of getting cheap transportation, together with the faulted condition of the lodes, the operators lost heart and turned their attention to other fields. In the nineties the mining industry was revived by the discoveries of gold ores at Edgemont, and in 1906 the country again felt the stimulus of enthusiasm for prospecting which swept over Nevada from the southwest. In that year several gold veins were discovered near Aura and elaborate preparations were made to reopen some of the silver mines at Mountain City, but these operations were stopped in 1907 when it became difficult to obtain money for prospecting or development work.

The ore bodies are fissure veins which cut across the bedding of sedimentary rocks, bedding-plane deposits which follow the stratification, and fissure veins in granite. The sulphide ores fall into two general classes—(a) gold deposits of highly siliceous ore carrying a small percentage of pyrite and galena and (b) silver deposits carrying these minerals in greater abundance, together with a small proportion of arsenic and antimony minerals. At Edgemont and Aura the deposits are in the sedimentary rocks, but at Mountain City they are mainly in the granodiorite. They were formed before the faulting took place. In every mine where any considerable amount of development work has been done faults have been encountered. These are nearly everywhere of the normal type, which implies a downward movement of the hanging wall.

The rhyolites that flank the mountains are probably younger than the deposits which have been developed in the Centennial Range. At Gold Circle, Lynn, and elsewhere this rhyolite carries gold deposits that are of a different type from those of this range, and there is no reason why it should be avoided by the prospector, especially where it is intruded by dark rocks (andesite) and in areas where it is leached white by hot waters.

LIME MOUNTAIN.

Lime Mountain forms the southern extremity of an elevated ridge about 6 miles long, which lies between Bull Run Basin and Chellis Valley. This ridge, which may be regarded as a southward extension of the Centennial Range, is separated from it by the very steep canyon occupied by Bull Run Creek. The rocks at Lime Mountain are in the main dark-gray limestones, which at the summit of the mountain dip from 20° to 40° W. The limestone is cut by dikes and other intrusions of igneous rocks, which include quartz porphyry, andesite, and diabase. Locally the limestone is metamorphosed to a coarse-grained marble, but the metamorphic action is not intense and very little garnet or hornblende has been developed. The principal workings are at the Eldorado mine, where a 200-foot tunnel driven westward into the mountain is connected with a shaft 115

feet deep. At their intersection and upward to the surface much of the workings is in ore that is said to average several per cent of copper, with values in gold and silver. The ore consists of pyrite, chalcopyrite, and bornite, which are intergrown with white and black mica, calcite, and quartz. The deposit, which is probably of contact metamorphic origin, has been fissured since deposition, and there has been some secondary chalcocite enrichment of the copper-iron sulphides. The rock that caused the metamorphism could not be determined. About 1,000 feet northwest of the mine is a considerable area of quartz porphyry, and a diabase dike, outcropping on the crest of the hill, 500 feet above the mine, strikes toward it. On the crest of the ridge, at places which are much nearer to those rocks than the Eldorado, the limestone is not greatly metamorphosed, except in some localities where it is a very coarse marble.

COLUMBIA AND AURA.

GENERAL STATEMENT.

Columbia, which is situated at the north end of Bull Run Basin, was in the seventies the headquarters for prospecting and mining in the central part of the Centennial Mountains. In the boom times of 1906 Aura was founded a mile or two below Columbia, and it is now the post-office and supply point for the east side of the range.

Several mines are located near Columbia, but all of them except the Big Four have long been idle. This mine, which was discovered in recent years, made several shipments of ore in 1908 and is commonly regarded as a property of some promise. The silver mines which were abandoned years ago were relocated recently, but little work has been done except enough to hold the ground.

In the canyon of Blue Jacket Creek, which joins Columbia Creek in the north end of Bull Run Basin, there are several mines and prospects, but none of them have been extensively exploited except the Blue Jacket, which was worked in the seventies. Two 5-stamp amalgamation mills were built in Blue Jacket Canyon to work the gold ores, utilizing the water power of Blue Jacket Creek. Both were idle when the camp was visited in 1908. Some years ago a company was formed to mine the gravels of Bull Run Basin, which are said to carry gold, and considerable money was spent in ditches, flumes, and pipe. The effort was presumably not successful, and the project was abandoned after a few hundred yards of material had been put through the sluice boxes.

MINE DESCRIPTIONS.

Infidel mine.—The Infidel mine, which is located on the hill above Columbia, was worked from three tunnels driven northward into the hill. The mine has been idle for many years and the middle tunnel

only was accessible in August, 1908. The rocks near by are mainly limestones and shales, but between the Infidel and Big Four mines is a body of granodiorite of considerable size. The lode strikes 30° W. and lies approximately parallel to the bedding of the country rock, a dark shaly limestone. It is a sheeted zone composed of three or four veinlets of white quartz closely spaced and nearly parallel. The ore is white quartz, containing a very small proportion of galena, pyrite, and other dark sulphides, said to carry high values in silver.

Big Four mine.—The Big Four mine is on the east bank of Columbia Creek, about one-fourth of a mile above Columbia. The deposit is a flat vein which follows the bedding of a shaly limestone dipping from 10° to 15° S. Three tunnels, each about 100 feet long, are driven on the vein, exposing an ore body that has a maximum width of about 5 feet. The lode is a sheeted zone in the shaly limestone, and the ore is composed of quartz, calcite, pyrite, galena, zinc blende, and chalcopyrite, with iron oxide and some green copper carbonate. It is said to carry high values in both silver and gold, the cobbled ore





FIGURE 7.—Cross section of Big Four lode, Columbia.

being rich enough to pay for shipment. Near the lower tunnel the vein and the inclosing limestone are in faulted contact with granodiorite, as shown by figure 7.


Columbia Queen mine.—The Columbia Queen mine, formerly the Bonanza, is on the west side of Columbia Creek, about 300 yards west of the Big Four mine. Like the Infidel and Revenue, it was the property of the Stokes Company, which operated it in 1875. The deposit, which resembles that of the Infidel, is a sheeted zone in limestone. The lode is 3 feet wide and carries galena, zinc blende, pyrite, and gray copper, with high values in silver and some gold. As shown by figure 8, the deposit dips toward the south and is displaced by three northward-striking faults, each of which offsets the lode toward the north.

Blue Jacket mine.—The Blue Jacket mine is located at the head of Blue Jacket Creek near the divide between this stream and Silver Creek. The deposit was worked through two tunnels and a deep shaft, all of which were inaccessible when the mine was visited. The



Topography based on map of Elko County by E.C. McClellan

TOPOGRAPHIC MAP OF NORTHERN QUARTER OF AREA SHOWN IN FIGURE 2



country rock, is limestone and, to judge from the dump, the ore is similar to other silver deposits near by. It is composed of white quartz carrying a small percentage of galena, pyrite, and zinc blende, with copper carbonate, iron oxide, and other minerals. The ore was carried over a wire tramway to a 20-stamp pan-amalgamation mill, the ruins of which may be seen below the mine at a bend of Blue Jacket Canyon.

Jack Pot mine.—The Jack Pot mine is on the south side of Blue Jacket Canyon, about 300 yards below the lower tunnel of the Blue Jacket mine. It was discovered in 1906 by Blewett Brothers, who have done about 800 feet of development work. The country rock is




FIGURE 8.—Plan of tunnel level, Columbia Queen mine, Columbia.

limestone, which, some 200 yards south of the mine, is intruded by a broad dike of granodiorite. The lode, which is developed by four tunnels driven one above another, strikes northwestward and dips 35° SW. and carries shoots of ore which have a maximum width of about 5 feet. The ore is in the main decomposed, silicified limestone, stained with iron oxide. Galena, pyrite, chalcopyrite, and zinc blende are found in the lower workings. About 60 tons of the ore has been put through the Walker mill, but the saving by amalgamation was not satisfactory. A considerable tonnage on the dump and in the mine is said to be rich enough in gold to yield a profit on cyaniding.

California mine.—The California mine is located on the east slope of Porter Peak, near the head of a small tributary to Blue Jacket Creek. The lode is a siliceous replacement vein, which cuts across the bedding planes of limestone and dips southwestward at a high angle. The ore consists of white quartz, galena, and iron oxides, with some green copper carbonate. Where exposed in the principal working pit the lode is 4 or 5 feet wide.

Tiger lode.—The Tiger lode is on the north slope of Porter Peak, at the head of Silver Gulch, which drains westward from the central axis of the range. The lode dips 60° W. and is opened along the surface by five short tunnels driven one above another. Some ore from this mine is said to have been packed by mules to Mountain City in the seventies. The lode is a 3-foot vein of ore, very high in silica, and contains galena, pyrite, and a small amount of zinc blende. The surface ore carries copper carbonate, pyromorphite, and iron oxides and is said to be rich in silver. North of the Tiger lode, at the head of Silver Creek, there are a number of abandoned shafts that were driven on lodes of siliceous ores in limestone. Northwest of these shafts, on the ridge between White Rock and Silver Creek, dark limestones dip from 45° N. to 90° . Three or four silver veins striking northeastward cut across the limestone, and on the crest of the ridge these veins outcrop boldly at several places. With better transportation facilities these deposits should be regarded as worthy of more careful prospecting.

Humboldt mine.—The Humboldt mine is at the head of Polaris Gulch, about half a mile southeast of the Blue Jacket mine. The country rock is contorted shaly limestone, and the vein, which is $1\frac{1}{2}$ to 3 feet wide, is composed of white quartz, galena, pyrite, chalcocopyrite, and gray copper. An incline is driven southeastward at a low dip, but when the mine was visited most of it was under water. About 100 yards east of the Humboldt shaft a surface pit in granodiorite shows a strongly sheeted and leached zone, which is said to carry up to \$3 a ton in gold.

Polaris mine.—The Polaris mine is in Polaris Gulch, about one-fourth mile southeast of the Humboldt. An incline is sunk about 200 feet deep and from it two levels are turned. A fissure dipping 32° SE. cuts across the bedding of the limestone. Here and there along the fissure are masses of quartz and silver-bearing sulphides. This mine supplied some ore to the Columbia mill in the seventies.

Aura King mine.—The Aura King mine is in Blue Jacket Gulch, between the Walker and the Aura King mills. The vein, which is in limestone, dips 30° SW. and has been followed on the strike for 200 feet. It is from 3 inches to 1 foot wide and is said to carry high values in gold.

EDGEMONT.

Lucky Girl group.—Edgemont is an active little mining camp located on a branch of White Rock Creek, on the west slope of the Centennial Range. It is supported in the main by the mining and milling operations of the Montana Gold Mining Company, owning the Lucky Girl group of claims, which includes all of the deposits which have been extensively developed on the west side of the range. These claims were purchased in 1898 by Alex. Burrell, and a 20-stamp mill was built in 1902 and has been in continuous successful operation ever since, treating about 60 tons of ore a day. Electric power is transmitted from a plant installed on a tributary of White Rock Creek, 4 miles north of Edgemont. The mines of this company comprise about 5 miles of underground workings and extensive ore bodies have been developed. The deposits include the Lucky Boy, Lucky Girl, and Big Bob veins. The Lucky Boy vein is the most extensively developed and has supplied most of the ore to the mill. This group of mines and the Bull Run mine on the south slope of the mountain have yielded since they were opened about \$1,000,000, chiefly gold.

As now arranged, the ore is dropped to the fifth-level adit and is drawn by mules to the portal, where it is fed automatically to a bucket tramway, 3,600 feet long, which carries it to the mill 800 feet below. At the mill the ore falls on grizzlies and the large rock goes to a 9 by 11 Gates crusher. There are four batteries of five stamps each. The fine rock goes to the outside stamps, which weigh 750 pounds, and the coarse rock to the inside stamps, which weigh 1,000 pounds. The stamps drop 8 inches eighty-five times a minute. About 50 per cent of the values are caught on amalgamation plates, from which the pulp passes to three Wilfley tables and one Pinder concentrator, where the galena and pyrite are removed. To utilize a water supply from a gulch north of the mill, the tailings are passed through a flume to a dam about 2,000 feet below. Here the slimes are drained from the sands and the sands are loaded into nine 50-ton cyanide tanks. The slimes are collected at a second dam below and are loaded dry into a mixer, where they are broken and mixed with water to the desired consistency. From this they pass to the agitators and thence to decantation tanks. The total extraction is from 90 to 95 per cent, the ore carrying from \$5 to \$10 in gold and 1 ounce of silver to the ton. The ore yields about 0.6 per cent in concentrates that carry about \$115 in gold, silver, and lead.

The deposits are fissure veins in siliceous sedimentary rocks which are steeply tilted, folded, and faulted. The country rock is a brown or gray quartzite of rather uniform grain, with here and there thin beds of siliceous shale, which at some places is dark gray or nearly

black. Near the veins a little sericite or white mica has been developed in the quartzite, together with a small amount of pyrite, but, compared with deposits in the porphyries, the wall rock is but slightly changed in composition. Between the shaft of the Lucky Girl and the portal of the low-level tunnel, conglomerate beds are interstratified with the quartzite. These beds are only a few feet thick and consist in the main of well-rounded pebbles of uniform size, about one-half inch in diameter. Most of them are white quartz, with here and there a pebble of red jasper. In the Lucky Boy mine small folds may be observed at several places on the surface and underground, and the difficulties of interpreting the structure are increased also by pronounced sheeting across the bedding in the same general direction as the lodes and by jointing in other directions. The prevailing dip of the sedimentary rock is 30° or more northwestward, but at some places the dip is about 40° NE. Some of the quartz was deposited before the deformation of the rocks was completed, as is shown by the




FIGURE 9.—Sketch of wall on level 8, 1,200 feet from portal, Lucky Boy mine, Edgemont.

fractured ribbons of quartz indicated in figure 9. The principal deposits, however, are little affected by folding and their deposition must have followed the greater deforming movements.

The veins, which at most places cut across the bedding of the quartzite, outcrop plainly on the surface, where they are composed of white quartz slightly stained with iron oxide. The three lodes that are most developed strike northeastward and dip from 15° to 45° SE. The ore is highly siliceous and carries more than 90 per cent of quartz. The sulphides are pyrite, galena, and arsenopyrite. The gold is associated mainly with the sulphides or with their oxidation products. The oxidized and sulphide ores carry approximately the same values, from \$5 to \$10 a ton, and there is no evidence of secondary enrichment other than that due to the removal of soluble constituents; this, owing to the comparatively firm condition of the ore, is small. Copper carbonates are rare or altogether lacking, and no zinc minerals were noted. Considerable fissuring with slickensiding must have taken place before deposition, as is indicated by the section shown in figure 10, where

slickensided fissures stop at the vein, which they do not disturb or cross. Movement subsequent to deposition has produced much gouge along the walls, but this is barren except where it contains crushed quartz. The sharp contact of country rock and vein quartz




FIGURE 10.—Section S. 40° E. through Lucky Girl vein, Edgemont.

and the dependence of vein fillings on previous openings are illustrated by figure 11.

The Lucky Boy vein has been developed along the strike for a distance of about 3,200 feet and through a vertical depth of 400 feet. On the fifth level a fault which strikes approximately with the vein




FIGURE 11.—Sketch of the northwest wall on level 8, Lucky Boy mine, Edgemont.

and dips in the opposite direction, from 40° to 80° NW., may be located for a distance of 2,000 feet, being exposed at five places in various drifts and raises. The throw of the fault is about 30 feet, measured vertically, as shown in the section given in figure 12. It

results in a duplication of the vein on level 5. At some places the wide zone of faulting carries much gouge, and it is difficult to locate the main fault plane precisely or to determine the direction of dip, but in the main the slicken planes dip to the depressed side, indicating that the fault is normal. In a block of ground about 1,000 feet northeast of the portal of level 5 and extending for 500 feet toward the breast the vein has not been discovered, and on levels 6 and 8 in this block of ground its position is unknown. Presumably it has been shifted by faults which cross the great strike fault above mentioned and which may displace it also. To the northeast of this block of ground the vein continues with its usual dip and strike for 1,500 feet.




FIGURE 12.—Cross section of Lucky Boy vein, Edgemont, looking southwest, 775 feet southwest of intersection of vein with Gossip adit. Line of section trends N. 35° W.

The Lucky Girl vein, which lies several hundred feet northwest of the Lucky Boy, strikes N. 50° E. and dips from 35° to 45° SE. On the surface its outcrop is clearly exposed below the company bunk house. It cuts across massive quartzite which on the foot wall dips 22° NE. An incline is driven on the vein from which two levels are turned and a considerable portion of the lode above the lowest level has been removed. The vein is from 2 to 7 feet wide and resembles the oxidized portion of the Lucky Boy vein, being composed of banded, sheeted quartz, stained yellow with iron oxide and carrying several dollars to the ton in gold with about 1 ounce of silver. A section of the vein is shown in figure 10.

The Big Bob vein, which is similar to the Lucky Boy and Lucky Girl veins, outcrops a few yards above the mill, where it dips about 15° SE. It is opened on five levels, the lowest of which is driven about 130 feet vertically below the outcrop. The larger part of the workings on this vein were under water in 1908.

Bull Run mine.—The Bull Run mine is located high on the south slope of Bull Run Mountain, which lies just north of the great canyon through which Bull Run Creek flows to Owyhee River. In 1902 a 10-stamp amalgamating mill and cyanide plant were built near the portal of the lowest adit, but according to report the successful operation of this mill was impossible owing to the prohibitive expense of freighting fuel and supplies up the steep hill to the mine. About \$120,000 was recovered in 1902 and 1903, before the mill was shut down. The developments comprised about a mile of underground workings on the lode. The country rock is quartzite which carries thin layers of slightly micaceous shales. The quartzite, which resembles that of the Lucky Boy, is strongly sheeted and locally the




FIGURE 13.—Generalized section through Bull Run mine, S. 10° E. from tunnel 1 to tunnel 4.

shales have been contorted by compressive movements that took place before the deposition of the ore. The deposit is a well-defined fissure vein which has a maximum width of about 6 feet. It strikes northeast and dips from 22° to 38° SE.

The quartzite is sheeted parallel to the vein and in places its apparent stratification is parallel to the vein, but on the surface, where the true dip may more easily be made out, it strikes nearly eastward. The ore is composed of white quartz, stained here and there with iron oxides. In the lower levels pyrite and galena are present in small quantities, these sulphides having been leached out in the upper workings. The ore is said to carry several dollars to the ton in gold, a large proportion of which is free. The silver present is practically negligible. The composition of the ore, the country rock, and the structural features of the vein resemble very closely those of the deposits of the Lucky Girl group at Edgemont, which is not more than 2 miles to the north. The lode outcrops plainly in the steep cliffs of the mountain and may be followed through a vertical range of about 400 feet. As shown in figure 13, the lode is displaced by

two faults, both of which cross tunnels 2 and 3. The faults strike about N. 15° E. and displace the lode from 30 to 60 feet. The first one encountered in tunnel 3 dips 60° W. and the second is a wide crushed zone which shows slickensided planes dipping toward both walls with numerous fragments of quartz and much clay gouge. In both faults the west wall has dropped.

Between the Lucky Girl group of veins and the Bull Run mine there is a steep canyon which is occupied by a small stream that flows westward to Bull Run Creek. A good deal of this country is covered and so the conditions for prospecting are not favorable, but its position in the line of strike of the Bull Run and Edgemont veins would seem to warrant a closer scrutiny than it has received.

MOUNTAIN CITY.

LOCATION AND HISTORY.

Mountain City is situated in the northeastern part of the Centennial Range on the north fork of the Owyhee, about 1½ miles east of the boundary of the Duck Valley Indian Reservation, some 40 miles by stage from Tuscarora. The first discoveries were made in 1869 by Jesse Cope and others who were on their way from Silver City, Idaho, to the White Pine district, Nevada, and from this circumstance the Mountain City region is called the Cope mining district. In the seventies there was considerable activity in mining and three silver mills were in operation. These were small amalgamation mills of the Washoe pattern, equipped with stamps, pans, and settlers. It is said that over \$1,000,000 in silver was recovered prior to 1881, mainly from surface and shallow workings. Since 1881 considerable prospecting has been done, but the production of ore has been small. In late years three gold mills have been built and are still in good condition, but they were not running in the summer of 1908, when the camp was visited.

GEOLOGY.

The rocks at Mountain City are limestones and shales intruded by granodiorite and overlain by rhyolite and basalt. On the summit and south slope of California Hill, 2 miles south of Mountain City, and extending northwestward from that point, is a thick bed of light-buff marbleized limestone. On the north slope of this hill, near intruding granodiorite, tremolite, actinolite, epidote, white mica, garnet, and other silicates of contact-metamorphic origin are developed in the limestone. The granodiorite is composed of plagioclase, orthoclase, quartz, mica, and hornblende. In the coarser varieties, as at the Protection mine, some of the crystals of feldspar are 1 inch long. The granodiorite is cut by aplitic dikes composed of quartz and orthoclase. Around the border of the mineralized areas there are extensive flows of rhyolite, the commonest variety of which is a purplish-pink rhyolite with phenocrysts of quartz, feldspar, and a

little black mica. Other varieties of the rhyolite are flow breccias, brown glass, and black obsidian. Vesicular basalt, rich in olivine and pyroxene, was noted on the surface east of the Resurrection claims. The granodiorite is younger than the limestone, which it intrudes, causing contact metamorphism. Where the rhyolite flows were noted above the granodiorite the contact relations could not be made out, but from consideration of areas elsewhere the rhyolite and basalt are regarded as of later origin than the granodiorite, and at Mountain City they are probably later than the deposition of the silver ores.

ORE DEPOSITS.

The ore deposits are fissure veins in granite and in metamorphosed limestones. They outcrop plainly at the surface, where some of them carry good values in silver. The veins do not fall into well-defined parallel systems but strike in various directions, the prevailing dip being toward the south.

Some of the veins, as shown on California Hill, are later than the aplitic phase of the granite which cuts the normal coarse-grained granite. None of the developed deposits are in rhyolite or basalt, although some gold-bearing veins in rhyolite are said to occur in the country east of Mountain City. In the Nelson mine the lodes pass from granite to limestone without much change in width or value. There is little replacement of the limestone, for the walls are clear cut and angular fragments of the country rock are included in the veins. Where the wall rock is granite the dark silicates have been leached out and sericite and pyrite have been developed in the granite by secondary processes.

The unoxidized ore is composed of quartz, pyrite, galena, zinc blende, gray copper, argentite, gold, and arsenopyrite, with a little chalcopyrite. All of the ore is highly siliceous, quartz constituting as much as 90 per cent of the rock. The oxidized ore is composed of quartz, chalcedony, horn silver, pyromorphite, iron oxides, native gold and silver, lead carbonate, copper carbonate, and copper silicate. Brittle silver and dark ruby silver are said to be present also. The oxidation of the deposits is erratic, the sulphides occurring at some places within a few feet of the surface, while some of the minerals of oxidation are to be found as deep as the lodes have been explored, or about 250 feet below the surface. The greater proportion of the silver values are in decomposed chloride and lead carbonate ore. Specimens of rich ore show large flakes of greenish-yellow horn silver deposited in the cavities of dark quartz. Some of the iron-stained siliceous ore pans gold liberally.

The lodes are fractured and faulted, and locally the ore is reduced to a white sand, in which there are numerous small rounded fragments of quartz about the size of a hazel nut. The faults that cross the

lodes are mainly of the normal type, the hanging wall having dropped with respect to the foot wall.

MINE DESCRIPTIONS.

Protection mine.—The Protection mine, located three-fourths of a mile below Mountain City, was one of the early discoveries of the district and was worked in the early seventies, when considerable chloride ore is said to have been treated in a silver mill near by. In late years the mine has been reopened and considerable exploration work has been done. A 10-stamp amalgamating and concentrating mill was built near the portal of the tunnel to treat the ore. At present part of the mine is leased and is being worked in a small way, but the mill is shut down. The principal vein is a fissure filling in granodiorite and has a maximum width of about 4 feet. The sulphide ore is composed of quartz, pyrite, galena, zinc blende, gray copper, brittle silver, and ruby silver. The surface ore is stained with iron and manganese oxides and contains horn silver, a little copper carbonate, pyromorphite, and a yellowish-green mineral, said to be silver bromide. The sorted ore carries \$100 a ton in silver and gold. At some places near the vein the granite wall rock is but little altered; at others it is a light-colored decomposed rock, the ferromagnesian minerals having been leached out and the feldspar sericitized. A shaft is sunk to a depth of 62 feet and a level turned at the bottom. This is connected with an adit driven 80 feet below the bottom of the shaft, which gives a depth of 142 feet at this place.

The Protection vein strikes a few degrees west of north and has been followed into the hill on the adit level for a distance of some 750 feet to a point where it abuts against a fault that strikes eastward and dips about 40° N. A drift has been run on this fault for 400 feet, but no vein has been encountered on the hanging-wall side at this end of the drift. On the other side of the fault a vein with the same general dip and strike as the Protection vein is 35 feet farther south and is possibly the same vein, but if so the fault is reverse—a rare type of faulting in this part of Nevada. Along this fault there are stringers of quartz in place, and in the level above both the Protection vein and the faults are mineralized on both sides of their intersection. The relations indicate that the Protection fissure was displaced by faulting before deposition of the ore and that there has been considerable movement subsequently.

Resurrection mine.—At the Resurrection mine, a few rods north of Mountain City, a large amount of work has been done in tunnels, pits, and shallow inclines, but most of the workings were inaccessible when the mine was visited in 1908. The country rock is granodiorite, to the east of which are flows of rhyolite and basalt. The granodiorite, which is highly altered, is sheeted by closely spaced fissures that strike northeastward. Several narrow quartz veins cut the granodiorite

parallel to the sheeting. The surface ore is composed of quartz, horn silver, lead carbonates, and iron oxide; the sulphides are galena, gray copper, a little pyrite, and chalcopyrite. In the seventies considerable rich chloride ore was taken from the surface pits and worked in silver mills near by.

Nelson mine.—The Nelson mine is on a branch of the north fork of the Owyhee, about $1\frac{1}{4}$ miles above Mountain City. Some 4,000 feet of workings have been run, mainly on two adit levels driven at a difference in elevation of about 100 feet. When the mine was visited in 1908 only the lower adit was accessible. A mill recently built at the portal of the lower adit is equipped with Blake crusher, nine stamps, amalgamation plates, and three Wilfley tables, and has treated a small amount of ore. The country rock of the mine consists of granodiorite, limestone, and aplite. The granodiorite intrudes the limestone and causes contact metamorphism with the development of epidote, actinolite, garnet, and mica in the limestone. In places this rock is so rich in actinolite that it has the appearance of a basic igneous rock and has been mistaken for diabase. The granodiorite is cut by aplite, which occurs as dikes and irregular intrusive masses. The ore deposits are fissure fillings from 1 to 3 feet wide and occur in granodiorite, limestone, and aplite. Several veins outcrop boldly on the hill above the mine, cutting across beds of metamorphosed limestone. The veins cross the contact of igneous and sedimentary rocks unbroken, but have been developed mainly in the granodiorite. The sulphide minerals present are quartz, pyrite, galena, zinc blende, gray copper, chalcopyrite, and arsenopyrite, with here and there a small amount of ruby silver and argentite. Native silver and horn silver are present near the surface, where the ore is stained with copper carbonates, iron oxides, and manganese oxides. Free gold, some of it with the crystal form, is associated with quartz and brown iron oxide. The sorted ore carries good values in both silver and gold, some specimens containing a high percentage of horn silver.

The Standard vein, which is developed in the lower tunnel, strikes southeastward and has been followed for about 1,000 feet, with overhead stoping here and there. This vein is faulted at three places by faults that strike eastward and dip northward at various angles. One of the faults shows a horizontal displacement of about 150 feet, the other two of less than 15 feet. All are of normal type, the hanging wall having dropped with respect to the foot wall.

Mountain City mine.—The Mountain City mine is located about 1 mile southwest of Mountain City, at the top of a low, flat ridge that rises some 200 feet above Owyhee River. The country rock is a metamorphosed black, shaly limestone which strikes eastward and dips 50° N. The lode is a fissure vein which cuts across the limestone, striking N. 50° W. The ore is highly siliceous and is a simple fissure

filling, cementing angular fragments of the altered limestone. It carries silver chloride and native silver, and in the seventies, according to report, several hundred thousand dollars' worth of silver ore was taken from the deposit through a shaft now inaccessible. About 500 feet S. 75° E. of the principal workings of the old Mountain City vein and lower on the hill are a number of open pits, some of which have been sunk on a vein which strikes N. 32° W. Possibly it is the faulted continuation of the Mountain City vein, but this has not yet been determined. This ground has recently been acquired by J. Hall and others, of Mountain City, and is called the New Yorkeys claim. The country rock of the lower deposit is a dark-gray metamorphosed limestone flaked with tremolite crystals. A tunnel is driven 95 feet N. 70° W. to the vein, which it follows for 90 feet, and a winze is sunk on the ore body 60 feet below the adit level. The deposit is a fissure vein and at some places a sheeted zone composed of several narrow veins with slabs of limestone between. Much movement has occurred since deposition, for at places the quartz is brecciated almost to powder. The ore is composed of quartz, iron oxide, copper carbonates, and silicates, and a little pyrite is present at the bottom of the winze. The vein strikes N. 32° W. and dips from 56° to 85° S. It has a maximum width of 5 feet, and is said to carry good milling values.

VAN DUZER CREEK PLACERS.

Van Duzer Creek is a small stream which flows eastward, joining the north fork of the Owyhee some 6 miles south of Mountain City. About 2 miles above the point where the stage road crosses the stream and extending westward up the main fork for about a mile portions of the channel have been washed for placer gold. Two strips of the channel, from 20 to 60 feet wide, have been worked out, one of these for a distance of 1,200 feet and the other for about 1,000 feet. Two hydraulic plants with 10-inch pipe and monitors are installed along the stream about three-fourths of a mile apart. The depth of work is nearly everywhere less than 15 feet. The ground is mainly fine gravel with few small boulders, and the bed rock is presumably limestone. The gold is said to vary from fine dust to nuggets of 5 or 6 ounces and sells for \$17 an ounce. The mines, which were discovered by R. M. Woodward in 1893, have been worked in some seasons for about fifteen years, but in 1908 were idle. The source of the gold is presumably some undeveloped veins at the head of the stream.

LONE MOUNTAIN.

Lone Mountain, which is called Nannies Peak in the reports of the King Survey, is a striking topographic feature rising conspicuously above the main axis of the Seetoya Range to an elevation of 9,046

feet. The mountain mass is composed in the main of Carboniferous limestone—dark-blue or gray massive beds locally metamorphosed by intruding igneous rock. The prevailing dip of the limestone is from 30° to 80° W. The crest of the mountain is composed of quartz monzonite and quartz monzonite porphyry, which cut through the limestone and inclose small blocks of it. On the west slope of the mountain, from 200 to 700 feet below the summit, is a long, narrow intrusive mass of felsitic quartz porphyry which trends nearly due north. Certain phases of the porphyry are almost as dense as rhyolite; others resemble the common types of quartz porphyry. The quartz monzonite and the porphyry locally grade one into the other and are probably phases of the same intrusion. The quartz monzonite, which is of medium grain, is composed of feldspar, quartz, hornblende, and biotite. An analysis of this rock is given on page 26, where the granitic rocks are described. The porphyry has a finely crystalline groundmass composed of quartz and orthoclase, in which are embedded phenocrysts of acidic plagioclase and resorbed quartz, biotite, and hornblende.

Both the quartz monzonite and the quartz monzonite porphyry have caused contact metamorphism of the limestone. At some places the limestone is changed to a garnet-calcite rock, at others to a green actinolite rock, at others to a hard cherty hornstone, and at still others to a coarse-grained marble.

The Merrimac district is situated on Lone Mountain about 28 miles by wagon road northwest of Elko. More or less prospecting has been done in this district since 1879, and from various claims about 1,000 tons of ore has been shipped, with a total value of over \$30,000.

The ore bodies, so far as observed, are deposits of lead and copper ore in marbleized limestone and contact-metamorphic deposits of copper ore in garnetized limestone. When the camp was visited by the writer nearly all the workings were inaccessible. The lead and copper ores in marbleized limestone are near the intruding igneous rocks and are usually highly shattered and oxidized. Some of them are at the contact of quartz monzonite or porphyry with limestone, and some of them are several hundred feet from it. The common minerals are galena, pyrite, chalcopyrite, and their oxidation products. Most of the ore consists of iron-stained limestone and quartz containing lead carbonate and green and blue copper carbonates, with copper oxides and yellow pyromorphite. Such ore is exposed at several places at the south end of the mountain on the Baltimore group of claims, and also on the Floradora and Ajax claims, a mile or more northeast of the summit. This ore is said to carry high values in silver, with \$1 to \$2 in gold to the ton. A number of shafts have been sunk and short tunnels have been driven on the deposits, but the present accessible workings do not show

the extent or shape of the ore bodies. Some promising lead-silver deposits in limestone are said to be located on the north end of the mountain just outside of the area shown in the outline map (fig. 2).

The contact-metamorphic deposits occur in the garnet zones near the intrusive igneous rocks. The large dumps from workings now inaccessible on the Baltimore and Cuag claims show a wide zone of decomposed limestone, with some garnet rock stained with iron oxides and copper carbonates. A low-level tunnel was driven north-eastward for 850 feet to explore the ore bodies in depth. It cross-cuts westward-dipping limestone for 750 feet, to a point where it encounters quartz monzonite porphyry similar to that which forms a large part of the mountain crest. Near the contact the limestone is metamorphosed to garnet, actinolite, and other silicates, with which are intergrown iron and copper sulphides. Pyrite, arsenopyrite, and chalcopyrite are present, and locally copper carbonate.

A zone of metamorphosed limestone is situated on the east slope of the mountain and trends northward approximately parallel to the crest, and there are a number of pits and short tunnels which are driven in ore. On the Morgan claim and north of it for several hundred feet, the porphyry lies to the east and metamorphosed limestone to the west of the contact. Highly oxidized copper and lead ore, some of it associated with garnet gangue, is exposed at several places. On the Pacific claim, which lies to the south of the Morgan, there is a good-sized body of iron ore at the contact of limestone and porphyry. This ore is composed of magnetite, limonite, garnet, and copper carbonate. Some of the limonite is pseudomorphous after pyrite, indicating that the original deposit was composed of intergrown magnetite and pyrite. This ore is said to carry a small percentage of copper, and if it were accessible to smelters it would make a good fluxing rock.

CORTEZ RANGE BETWEEN THE CARLIN AND DALTON PEAKS.

GENERAL FEATURES.

The Cortez Range, north of Humboldt River, is a lofty ridge about 45 miles long and from 5 to 20 miles wide. The highest summits are the Carlin Peaks, which reach an elevation above 7,700 feet, and the Dalton Peaks, about 25 miles farther north, whose highest point is 9,232 feet above the sea. Rye Meadows, a broad area of gently rolling sagebrush plain, lie to the west of this portion of the range, and on the east side is the broad, gently sloping valley of Maggie Creek, trending southward approximately parallel to the range. The Carlin Peaks^a are composed in the main of extensive flows of rhyolite, which on the west slope of the highest summit surround a large mass

^a Emmons, S. F., U. S. Geol. Expl. 40th Par., vol. 2, 1877, p. 587.

of andesite. The rhyolite series is probably not so thick as would appear from the extent of the flow and from the differences in the elevations at which it is found, for here and there small areas of sedimentary rocks outcrop where the rhyolite has been eroded. These may be observed in the Richmond district and on either side of Maggie Canyon, where the stream cuts through the low ridge that joins Carlin with Maggie Peak. Granodiorite and other coarse-grained rocks which are probably intrusive in limestone were noted in the Richmond district on the east slope of the range, 8 miles north of the Carlin Peaks. The north end of the range, between the Lynn district and Soldier Gap and including the main mass of the Dalton Peaks,^a is made up of the Weber quartzite; which dips toward the west and is bordered on the east by rhyolite. In this area there are several mines and prospects, none of which are extensively developed or have produced more than a few tons of ore.

MINE DESCRIPTIONS.

Nevada Star mine.—The Nevada Star mine is located 9 miles northwest of the town of Carlin, on the ridge just west of Maggie Canyon. The deposit is a small replacement vein in limestone, dips steeply eastward, and contains bunches of galena and lead and copper carbonates said to carry good values in silver. A shaft is sunk on the lode to a depth of 60 feet, and at the bottom a drift is driven southward for 80 feet. Near by a two-compartment shaft has been sunk to a depth of 110 feet, but from it no crosscutting has been done.

Copper King claim.—At the Copper King claim, which is about 2 miles southwest of the Nevada Star, the country rock is banded, fissile rhyolite, which dips about 40° N. The rhyolite is shattered, leached, and cut by slickensided movement planes that carry bunches of oxidized copper ore here and there. A well-equipped two-compartment shaft has been sunk to a depth of 150 feet, with short levels turned at intervals of 50 feet.

Richmond district.—There are several small prospects of silver ore in the Richmond district, which is on the east slope of the Cortez Range, 14 miles northwest of Carlin. The rocks exposed in this district are limestones and quartzites intruded by granodiorite. In the limestones near the intruding rock there are several small outcrops of siliceous silver ore which contains a small proportion of galena, gray copper, and lead and copper carbonates. Shallow pits are sunk on several of these outcrops.

Lynn district.—The Lynn district, just east of a low divide at the crest of the Cortez Range, is about 20 miles northwest of Carlin. Gold was discovered in this district in April, 1907, and during the following summer the camp was the scene of one of the rushes which

^a Emmons, S. F., U. S. Geol. Expl. 40th Par., vol. 2, 1877, p. 608.

often follow discoveries in Nevada. The excitement had subsided in 1908, and the district was deserted by all except about a dozen miners who remained to develop their claims.

The country rock is bedded rhyolite, and in the vicinity of the principal deposits it dips westward at low angles. On the west side of the divide, a few hundred yards away, an outcrop of decomposed andesite was noted. At the Big Six mine the bedded rhyolite strikes S. 30° W. and dips from 20° to 35° W. Along a zone of shattering the rhyolite is silicified, is stained with iron oxide, and carries bunches of rusty ore which pans gold freely. Three inclines, 20, 40, and 65 feet long, are driven down the bedding planes of the rhyolite, and a few tons of ore has been shipped from these workings.

At the Gold Dollar mine, about 200 yards N. 10° E. of the Big Six inclines, a zone of shattering strikes northwestward across the rhyolite, and along this zone veinlets of quartz and silicified iron-stained rhyolite carry high values in gold. Four deep trenches are dug across the zone of shattered rhyolite.

Several placer claims are located in the gulches which drain eastward from the Big Six and Gold Dollar mines, and at a number of places the gravels carry placer gold. On the Hilltop claim, belonging to Hugh Jones, a block of gravel 60 feet long, 8 feet wide, and about 5 feet deep yielded \$150. During the summer of 1907 about \$1,000 was taken from this gulch. In the upper part of the gulch the bed rock is rhyolite, but this has been cut away lower down, locally exposing the underlying limestone. It is thought that water to be used for sluicing could be collected on the slopes of the Dalton Peaks, if the deposits prove extensive enough to warrant it.

PINYON RANGE.

GENERAL FEATURES.

The Pinyon Range is a long, narrow group of hills and mountains which extends southward from Humboldt River to a point near Eureka. Its highest point, called Ravens Nest, is 8,386 feet above sea level, or more than 3,000 feet above Pine Valley, which lies to the west of the range.

The rocks that constitute the range are in the main Paleozoic sedimentary rocks, including the section from the Ordovician to the Carboniferous. In the vicinity of the Bullion mining district these sedimentary rocks are intruded by granodiorite. At the north end of the range and at many places along its flanks the sedimentary rocks are covered over by rhyolite, basalt, and the Humboldt formation.

As shown by the analytical map of the Fortieth Parallel Survey (Pl. XI, vol. 1), the structure of the range is anticlinal from Ravens Nest to Pinyon Pass. From Mineral Hill to Diamond Valley the

rocks are compressed into an open syncline, the axis of which trends a few degrees west of north. South of this point the structure is that of an eastward-dipping monocline. In connection with the studies at Eureka, Nev., C. D. Walcott made a section across Pinto Peak for the purpose of comparison with the section at Eureka.^a This section (fig. 14) is given herewith and shows that the anticlinal structure of the range is modified to an important extent by faulting.




FIGURE 14.—Cross section through Ravens Nest, Pinyon Range. After C. D. Walcott.

Northwest of Pinto Peak, just above the valley flat, is a dark-blue limestone which carries Devonian fossils. This rock is faulted against Carboniferous quartzite that contains conglomerates and black siliceous pebbles. These dip northwestward and are underlain conformably by blue limestone which carries Upper Devonian fossils. High on the mountain, northwest of its summit, the limestones are faulted near the crest of the anticline. To the northwest of this fault the beds dip westward, but southeast of it they dip east. East of the fault is dark ferruginous Eureka quartzite, overlain by light-gray siliceous limestone, which contains *Halysites* and is probably of Ordovician age. The beds overlying the Ordovician carry an early Devonian fauna and above these Upper Devonian species. Above the Devonian is a great thickness of quartzites and sandstones with some argillaceous beds. The Upper Devonian and the Carboniferous dip away from the fault at the crest of the anticline on either side.

BULLION.

GEOLOGY AND ORE DEPOSITS.

Bullion, which is the principal settlement in the Railroad mining district, is about 28 miles southwest of Elko and 12 miles southeast of Palisade. It is situated at the base of the east slope of the Pinyon Range near the headwaters of the west fork of Dixie Creek, at an elevation of about 6,600 feet. The mines were discovered in the late sixties and were successfully worked in the seventies and eighties, when two small smelters were in operation. In 1906 some of the mines were reopened and since that time a moderate tonnage of ore has been taken from the old workings, together with several carloads

^a Hague, Arnold, *Geology of the Eureka district, Nevada*: Mon. U. S. Geol. Survey, vol. 20, 1892, p. 201.

of slag from the slag dump of the lead smelter at Bullion. The district is said to have produced about \$3,000,000 in silver, lead, copper, and gold. At present the principal holdings are owned by or are under option to three companies, each of which is actively engaged in mining or in exploration work. They are the Nevada Bunker Hill Mining Company, the Trimetal Mining Company, and the Delmas Copper Company. Two of these companies are driving long low-level tunnels below the old workings, giving vertical depths up to 1,500 feet below the surface. From these tunnels it is planned to prospect a considerable area of promising ground. At present the ore from the Standing Elk and near-by mines is hauled by wagon 28 miles to Elko. From the Delmas property on the west side of the range ore is hauled 12 miles to the Eureka and Palisade Railroad, to be reloaded at Palisade into broad-gage cars.

The mines are situated high on the slopes of Bullion Hill, a spur of Ravens Nest. This mountain is composed in the main of Ordovician limestone, which is for the most part a gray marbled limestone with a general westward dip. On the northeast slope of the mountain there is considerable faulting and the abrupt changes of dip indicate a complex geologic structure. The limestone is intruded by granodiorite and by quartz porphyry.

The granodiorite, which is of medium grain, is composed of quartz, feldspar, and biotite. Under the microscope the feldspars are seen to be oligoclase and andesine, with only a little orthoclase. The rock approaches quartz diorite in composition and is poorer in potash than most granodiorites. It forms extensive outcrops near the summit of the range and causes considerable contact metamorphism of the Ordovician limestone which it intrudes. Near the granodiorite the limestone is changed to a rock composed of garnet, tremolite, actinolite, calcite, andalusite, and white mica, which at some places are intergrown with pyrite, chalcopyrite, and zinc blende.

The intruding quartz porphyry is a light-colored rock which contains abundant rounded phenocrysts of quartz and a smaller number of feldspar phenocrysts, in a light-colored, fine-grained groundmass. Muscovite and some biotite are present, but the dark-colored silicates are very sparingly developed and are much less abundant than in the granodiorite. The quartz porphyry forms large intrusive masses on the northeast slope of Bunker Hill and is exposed at several places in the Standing Elk and Tripoli mines, where it is greatly altered by hot waters, white mica and carbonates being extensively developed in the groundmass. The limestone near the quartz-porphyry intrusions is somewhat marbled, but garnet zones are not developed along its contacts.

The most important ore bodies are replacement deposits of lead, silver, and copper ore in marbled limestone, and copper deposits

of contact-metamorphic origin. At some places the quartz porphyry is impregnated with copper minerals and possibly some of this rock could be worked under favorable conditions. There are also auriferous quartz veins in the granodiorite, but these, so far as developed, are of small economic importance. The replacement deposits of lead, silver, and copper ore in marbleized limestone are commercially of greatest importance, and most of the production from the district has come from ore bodies of this character. Most of these deposits were inaccessible when the mines were visited, but those which were exposed appear to be nearly vertical chimneys of ore situated at the intersection of two or more relatively narrow replacement veins. The maximum depth to which the ore is exposed is about 500 feet below the surface and the ore is almost completely oxidized. The principal ore minerals are lead carbonate, horn silver, pyromorphite, malachite, azurite, chrysocholla, cuprite, pyrite, chalcopyrite, galena, bornite, copper glance, and a copper-antimony mineral that is probably gray copper. The gangue is composed of quartz and calcite. The ore is nearly everywhere stained by red and brown iron oxides, which, together with calcite and quartz, form the most abundant minerals. A little manganese oxide is present.

The deposits outcrop as iron-stained gossans where the surface is strewn with craggy boulders composed of iron oxides, with copper carbonates here and there. Some of the gossans have not been explored in depth, and none of the deposits have been followed down to pure sulphide ores. Owing to the relief of the country and the shattered condition of the rocks, oxidation has extended to considerable depth.

The replacement deposits of copper ore of contact-metamorphic origin have received serious attention only within the last two years, probably because the mass of this ore is of too low grade to stand excessive freight and treatment charges. The Delmas Copper Company has lately sunk a number of shallow pits and has driven several short tunnels on these deposits, exposing a considerable quantity of this ore, some of which has been shipped to Utah smelters.

On the summit of the ridge which divides the drainage of Pine Creek from that of Dixie Creek the intruding granite porphyry is in contact with the Ordovician limestone. The line of contact stretches northeastward and may be followed for nearly 2,000 feet down the southwest slope of the ridge and for a considerable distance down the northeast slope. The zone of limestone along the margin for a width of 125 to 400 feet has been converted into a rock composed of garnet, actinolite, calcite, epidote, quartz, tremolite, zoisite, and pyroxene, and at many places copper-bearing sulphide and zinc blende are intergrown with the iron-bearing silicates. The principal ore minerals are pyrite, chalcopyrite, bornite, chalcocite, galena,

and zinc blende. These are locally altered to malachite, azurite, copper oxides, iron oxides, and chrysocolla. The ore of the better grade is said to carry from 3 to 8 per cent of copper with good values in silver. Although much of the garnet rock is barren of copper minerals, the work that has been done seems to show that some of the ore bodies are of considerable extent. The boundaries of the ore zone are very well defined. The contact between the garnet rock and the marble which cuts across the bedding is usually rather sharp and clean cut. Near the top of the ridge a vein of garnet rock about 1 foot wide cuts across the bedding of the marble, and just below this vein




FIGURE 15.—Sketch showing the relation of the ore zone to granodiorite and limestone on the Sweepstakes claim of the Delmas group, Bullion district.

small patches of the garnet rock are surrounded by the marble. The garnet, in its relations to the marble, resembles some deposits of siliceous ore in limestone and indicates that in this particular locality the garnetization follows fissures in the rocks. In the main, however, the garnet rock follows the contact of the limestone with the granodiorite, and no garnet veins were noted more than a few hundred feet away from the igneous rock. Figure 15 is a sketch showing the relations of granodiorite and garnet rock on the Delmas claims.

The deposits of quartz porphyry impregnated with copper minerals which have been developed through exploration for lead-silver

ores are too low in grade to work under present conditions. In the Standing Elk mine, in tunnel 5, the quartz porphyry intrudes the limestone. It contains much hydrous silica and is, for a distance of about 20 feet, extensively altered by hot solutions that have replaced the feldspars and the groundmass with sericite or white mica, which is now stained with copper carbonate. The feldspars have been leached out and their spaces are now filled with fibrous malachite and with chalcantite. The rock is said to carry 2 per cent of copper and a small amount of silver.

The auriferous quartz veins in granite porphyry are, so far as developed, small and of too low grade to work. On the Delmas ground one of these is exposed here and there for a distance of some 500 feet along the strike. It is from 2 to 4 feet wide, is composed of white quartz, limonite, and pyrite, and is said to carry about \$3 to the ton in gold. Several smaller veins of this character outcrop in the areas of granodiorite between the Delmas claims and the Copper Belle, but none of them have been extensively explored.

MINE DESCRIPTIONS.

Standing Elk mine.—The Nevada Bunker Hill Mining Company controls the Standing Elk, Tripoli, Red Bird, and other mines, and is driving a crosscut tunnel to intersect the several lodes in depth. This tunnel in July, 1908, was 1,500 feet long but was still several hundred feet from the nearest lode. The Standing Elk, the most important mine of this group, is opened on seven levels, mainly adits, having altogether a vertical range of 600 feet. There are several thousand feet of workings on this claim, the principal level being adit No. 5. When the camp was visited the workings above this level were not accessible.

The ore bodies are irregular replacement veins in limestone, which intersect to form chimneys of ore, the largest being about 50 feet in diameter. The country rock varies from a hard gray limestone to a massive marble which locally is very coarsely crystalline. The limestone is sheeted, brecciated, and filled with white calcite. In a few places garnet rock is developed in the mine, but none of the Standing Elk deposits consist of the typical garnet ore. The intruding quartz porphyry is much decomposed and is locally silicified, and a considerable mass of it carries copper. Nearly everywhere the ore is highly oxidized. The principal ore minerals are lead and copper carbonates, copper and iron oxides, bornite, pyrite, chalcopyrite, and a copper-antimony sulphide, which is probably gray copper. Calcite and quartz are the important gangue minerals; a little fluorite is present in microscopic crystals, intergrown with quartz.

Tripoli mine.—The Tripoli mine, about 800 feet northeast of the Standing Elk, is owned also by the Nevada Bunker Hill Mining Company. A tunnel is driven into the mountain for 100 feet to an engine

station, where a winze is sunk to a depth of 175 feet. Levels are turned at vertical intervals of about 50 feet and a second winze is sunk from the lowest level, giving a maximum depth of 300 feet below the surface. The country rock is marbled gray limestone which on the surface strikes N. 20° W. and dips from 85° SW. to 90°. The limestone is intruded by an acidic phase of granodiorite. The incline and drift are driven on a fissured zone, which strikes about N. 25° W. and is approximately vertical. The zone of crushed limestone is crossed by two fissures, approximately at right angles, and these dip steeply toward the northwest. Chimneys of rich silver-lead ore are formed at or near the intersections. Lead carbonate and pyromorphite are mixed with galena and copper carbonates, and the ore is highly oxidized as deep as development goes. In the bottom of the winze, 300 feet below the surface, garnet and tremolite are extensively developed. At this place the metamorphosed rock is crushed and greatly altered, and it is said to carry about \$25 in lead and silver.

At the surface to the east of the Tripoli claim the limestone is medium-grained marble containing reefs and patches of garnet rock, with which are intergrown small masses of pyrite and chalcopyrite, stained here and there with copper carbonates.

Red Bird mine.—The Red Bird mine, of the Bunker Hill group, is about half a mile northeast of the Standing Elk and nearest the portal of the low-level adit. A tunnel is driven for 250 feet on a steeply dipping lode, and 80 feet below this a second tunnel is driven for 200 feet. About 150 feet from the portal of the lower tunnel a raise connects the two levels. The lode, which at some places is 4 feet wide, carries good values in lead and silver.

Copper Belle mine.—The Trimetal Mining Company controls a large acreage which joins the Nevada Bunker Hill holdings on the northwest. This group includes the Copper Belle, Copper King, and Philippine claims, on each of which considerable development work has been done. The Copper Belle mine is about half a mile northeast of the Standing Elk. The principal tunnel is driven for 325 feet S. 60° E. to the ore body, which is a large irregular mass of oxidized ore carrying lead, silver, and copper. It is said to have produced about half a million dollars' worth of ore, which was smelted in the copper smelter at Bullion. The ore body, which is nearly everywhere inaccessible, resembles the deposits of the Standing Elk in that it is a replacement of marbled limestone. On the surface above the ore body and at several other places near by the rock is stained with iron oxides and copper carbonates.

Delmas mine.—The Delmas Copper Company owns several claims which are located near the crest of the mountain range and extend southwestward from it.

On the Sweepstakes claim, which is the most extensively explored, are the copper deposits of contact-metamorphic origin which have been mentioned above. These deposits, as already stated, are composed of garnet, tremolite, and other contact-metamorphic minerals, which at some places are intergrown with pyrite, chalcopyrite, bornite, galena, and zinc blende. Locally the copper-iron sulphides, which are unquestionably primary, are coated over with secondary copper glance. Here and there the copper sulphides have oxidized to copper carbonates and iron oxides, but the oxidation is not complete, even at the surface, and at some places less than 10 feet below the surface the sulphides are much more abundant than the carbonates and oxides. Shipments of 210 tons of ore from this claim averaged 70 ounces of silver to the ton, 10.4 per cent of copper, and 2.8 per cent of lead. There is a considerable tonnage of low-grade ore partly developed. It will probably be found that the high-grade ore is restricted to that which carries chalcocite-coated sulphides, or to that which has resulted from the oxidation of such ore. Figure 15 is a sketch of the Delmas property.

Other claims.—The Kenilworth claim, north of the Standing Elk; the Sylvania claim, west of the Standing Elk; and the Blue Belle, northwest of the Sylvania, have each produced considerable ore from workings which were inaccessible when the camp was visited by the writer. The principal deposits of these mines appear to resemble those of the Standing Elk rather than the contact-metamorphic deposits of the Delmas group.

MINERAL HILL.

LOCATION AND HISTORY.

Mineral Hill is a mining camp situated about 5 miles southeast of Mineral, a station on the Eureka and Palisade Railroad. It is at the north end of a small ridge of the same name which rises some 700 feet above the floor of Pine Valley and forms a foothill on the west slope of the Pinyon Range.

The deposits, which outcrop along the summit of Mineral Hill, were discovered by a party of prospectors from the Reese River district in 1868 and were sold soon afterward to a San Francisco company. In the early seventies the mines were sold again to a London corporation, which operated them with some success until 1878. Parker, Spencer & Co., who were mining at the south end of the hill, acquired the holdings of the London company in 1880 and operated the mines and mill until 1887. Since that time the mines have not been worked actively. The total production of Mineral Hill, so far as it can be estimated from various reports, is probably a little more than \$6,000,000, practically all of which is silver. The mines are now in the hands of the Mineral Hill Consolidated Mining

Company, with headquarters in New York, and this company plans to work the low-grade ore that remains in the dumps and mines.

The ore was treated in two silver mills, one of which was equipped with 15 and the other with 20 dry-crushing stamps. The 15-stamp mill was in operation for a number of years, but the larger mill was sold and removed after it had been running for only a short period. The mills were equipped with Stetefeldt roasters to work the ore by the Reese River process, but it was found that the additional cost of roasting was greater than the increased saving effected, and so the Washoe process with dry crushing and raw amalgamation was early adopted. The details of treatment are given by M. Eissler in the "Metallurgy of silver," page 154. The present owners plan to treat the ore by concentration and cyanidation.

GEOLOGY.

The Pinyon Range to the east of Mineral Hill is made up of steeply dipping sedimentary rocks consisting of Paleozoic limestones, quartzite, and shales. The ore deposits are in a gray crystalline limestone, which, along the crest of Mineral Hill, dips from 45° to




FIGURE 16.—Sketch illustrating the structure at Mineral Hill.

75° E. This limestone rests upon a dark shale that is exposed along the west slope of the hill and in the Taylor tunnel below the ore bodies. On the west slope of the Pinyon Range, east of the south end of Mineral Hill, the shales, which here are at least 600 feet thick, dip westward below the limestone and rest upon a great series of quartzites which are mapped as Ogden (Devonian) in the atlas of the Fortieth Parallel Survey. Figure 16 is a sketch in cross section drawn westward through Mineral Hill.

The structure appears to be synclinal, the axis of the syncline lying somewhere to the east of the crest of Mineral Hill and east of the outcrops of the ore deposits. The limestone that contains the ore bodies is about 400 feet thick, but it is probably only the lower part of a formation which may be thicker, the upper portion having been eroded away at this place. The limestone is cut by three narrow dikes of a decomposed intruding igneous rock, the least-altered specimens of which are composed of quartz, sericite, calcite, and limonite. These dikes are approximately parallel, strike eastward, and dip to the south at high angles.

ORE DEPOSITS.

The principal workings of Mineral Hill are open cuts and shallow stopes which are closely spaced along a zone from 200 to 300 feet wide and about 1,200 feet long. The open cuts are from 25 to 75 feet long and their width is somewhat less. The stopes range from 10 to 40 feet in width and do not extend downward more than 150 feet below the surface. The Queen tunnel is driven southward from the north end of the workings, exploring a large part of the ore zone, and ore chutes are raised to the ore bodies near the surface. About




FIGURE 17.—Cross section of Star Chamber stope, Mineral Hill.

150 feet below the Queen adit the Taylor tunnel is driven westward to intersect a winze from the Queen.

The deposits are very siliceous and are eroded less rapidly than the country rock; consequently they form bold outcrops, and the zone along which a maximum of mineralization has taken place has remained longer at the surface, the hard rock tending to monopolize the outcrop. The configuration of the hill is probably due to the resistance offered to erosion by the large masses of quartz. The ore bodies are chambers or irregular replacement deposits which cut

across the bedding of limestone. Along the line of the principal deposits the limestone dips steeply eastward, as shown in figure 17. In places near the ore the limestone is strongly sheeted or brecciated, the open spaces being filled with white calcite, which contrasts strikingly with the gray limestone bands of the original rock, producing gneisslike banding. At some places the limestone is replaced by the quartz and sulphides and the ore grades into the country rock; elsewhere the limestone is brecciated, and the small angular fragments of limestone, which are cemented by white quartz, show no evidence of having been dissolved, the sharp edges remaining intact. The relations indicate that the composition of the solutions changed while the deposits were being formed—that they had spent their power to replace the limestone before deposition ceased. The ore, deposited by solutions which were capable of replacing the limestone, is richer in the sulphides and in silver than the quartzose material that simply filled the open spaces.

The minerals of the ore are quartz, calcite, barite, silver chloride, argentite, gray copper, galena, zinc blende, copper carbonates, pyromorphite, lead carbonate, pyrite, and iron and manganese oxides. According to Eissler,^a polybasite, stephanite, bromide of silver, and molybdenite are also present. Some of the ore carries a considerable quantity of galena, but in most of it the proportion of sulphides is small, the gangue minerals constituting considerably more than 90 per cent of the bulk of the rock. A large proportion of the ore carried from 100 to 200 ounces of silver to the ton, although ore which carried as low as 25 ounces was worked. The ore bodies are closely spaced along a zone of fracturing that strikes northward and has been extensively explored for a distance of about 1,500 feet. So far as developed, the ore is in the main at the surface or less than 100 feet deep. Most of the ore bodies dip about 45° E.; some are vertical and one dips steeply westward. The silicified zone is cut by a number of fissures which strike east and dip about 60° N., and in two of the ore chambers these fissures form the south wall of the ore body. Possibly these fissures have faulted the ore, but the limestone has been silicified on both sides of them and the ore zone is not displaced by them to any great extent.

From the Star Chamber stope was removed a wide mass of ore, extending from the surface to the level of the main adit with a dip of about 40°. This stope, which is shown in figure 17, is practically continuous toward the south with the Giant stope, a huge cavity about 50 feet in diameter from which small tortuous stopes extend in several directions. Still farther south along the zone of silicification are the Live Yankee, the Austin, and several other stopes of smaller size. The localization of the ore bodies is due to the intensity

^a Eissler, M., loc. cit.

of fracturing and sheeting in the ore zone, the larger deposits having formed where there was a maximum amount of shattering.

The shales which underlie the limestone are crumpled, fissured, and cut by small veins of white quartz, but are not known to carry deposits of economic value. Below the ore bodies some development work has been done in the shales, but this prospecting was not thorough and the present owners plan further exploration at the lower level. Work to the east of that already done in the Taylor tunnel may prove more productive. To judge from the dip of the beds the limestone will probably be found at greater depth at that place than in the ground directly below the outcropping deposit.

ALPHA.

Alpha is a small camp about 15 miles south of Mineral Hill and 5 miles east of Alpha station on the Eureka and Palisade Railroad. The principal claims are the Arizona, Utah, Oregon, and Idaho. These claims have been developed by a number of shallow inclines and short tunnels, driven for the most part in silicified limestone which carries a considerable amount of barite and a small amount of the metalliferous sulphides. A concentration plant equipped with ten stamps and five vanners was built at Chimney station, about 3 miles west of the mines, but the treatment was presumably unsatisfactory, as only a small amount of ore was put through the plant. The country rock is Devonian limestone. It dips from 30° to 40° E. and presumably rests upon the quartzite which outcrops as a marginal band in the low hills west of the mines and is mapped as Ogden by the Fortieth Parallel Survey. The ore deposits outcrop boldly at the surface, some of them forming more or less noticeable reefs. The metalliferous minerals include freibergite, galena, zinc blende, pyrite, and copper carbonates, and in some places barite forms more than half of the ore. The lodes are sheeted zones and replacement deposits in limestone, of which some follow the stratification and some cut across the bedding.

CORTEZ RANGE SOUTH OF HUMBOLDT RIVER.

GENERAL FEATURES.

The Cortez Range extends from Carico Peak northeastward about 50 miles to Humboldt River and from this point northward some 40 miles to Independence Valley in the region of Tuscarora. The range north of Humboldt River has been described. That portion which lies south of the Humboldt includes a part of the Safford district, near Palisade, and the Mill Canyon and Cortez district, on the slopes of Tenabo Peak. The country between Mill Canyon and the Safford district was not traversed in this reconnaissance, and the

notes referring to that portion of the range are taken from the descriptions given in the report of the Fortieth Parallel Survey.^a

The south end of the range as outlined above is 50 miles long and about 12 miles wide. The highest summits are Railroad, Tenabo, Cortez, and Papoose peaks. Of these Tenabo Peak is the most elevated, its summit being 9,240 feet above sea level, or about 4,000 feet above the valley flats. The range is crossed by several low passes, among them Wagon Canyon, Agate Pass, and Cortez Pass. Crescent Valley, a great gravel-covered sagebrush plain, borders the range to the northwest; Pine Valley and Garden Valley lie to the southeast.

GEOLOGIC FEATURES.

The rocks of the Cortez Range south of Humboldt River include limestone, quartzites, and shales, which are intruded by a great variety of igneous rocks and overlain by basalts and rhyolites. No fossils have been found in this portion of the range, but on lithologic grounds the sedimentary rocks have been referred to the Carboniferous. The intruded rocks include granodiorites, quartz monzonites, quartz diorites, dacites, and andesites. It has been suggested that the granular rock of Agate Pass^b is probably Archean, but the granitic rock in Mill Canyon and at Tenabo Peak is certainly later than the sedimentary rocks called Carboniferous, for the latter are metamorphosed at the contact. Neither the granular nor the sedimentary rocks have the schistose structure commonly found in pre-Cambrian formations.

The sequence or relative age of the eruptive rocks in this part of the range has not been determined, but probably all are later than the Carboniferous. The granular rocks are older than the rhyolites and basalts. On the east flank the range is bordered by a wide belt of the Humboldt formation which trends southward from Palisade for about 30 miles. Above the Humboldt and extending high on the east slopes of the range, at some places almost to the summit, there is a wide belt of basalt. The rocks along the crest of the western slope of the range from the Safford district to Mill Canyon include limestones, quartzites, granodiorites, diorite, rhyolites, and dacites. The prevailing dip of the bedded rocks is eastward.

CORTEZ AND MILL CANYON DISTRICT.

LOCATION AND HISTORY.

Tenabo Peak, near the southwest end of the Cortez Range, is about 30 miles south of Beowawe, the nearest station on the Southern Pacific Railroad. Cortez, on the southwest slope of this peak, was

^a U. S. Geol. Expl. 40th Par., vol. 2, 1877, p. 570.

^b *Idem.* p. 575.

one of the most productive mining camps in this part of Nevada. There are also a number of mines and prospects in Mill Canyon on the north slope of Tenabo Peak and on the slopes of Bullion Hill at the head of Mill Creek. The mines of Cortez and Mill Canyon were discovered in 1863, and rich ore yielding several hundred dollars to the ton was hauled to Austin for metallurgical treatment. Simeon Wenban, one of the early locators, obtained control of the Garrison, St. Louis, Arctic, Fitzgerald, and other important claims and organized the Tenabo Mill and Mines Company, which operated the mines for a number of years and still owns them. A small stamp mill with roasters and silver pans was built in Mill Canyon in 1864 and for three years was employed in treating the ore from various mines near by. This mill was bought by Wenban in 1867 and the ore was hauled to it from Cortez by wagon and pack trains.

The mines on the Cortez side of the mountain were exploited by Wenban with some success for many years. In 1886 a leaching plant with a daily capacity of about 50 tons was built at Cortez and this plant was in constant service for eight years. Water was piped from the mountains to the south, a distance of 7 miles, and the fuel supply was pine timber, a scattered growth of which is found on the mountain slopes. From the best sources of information that are available the production of Cortez and Mill Canyon since discovery is estimated at \$10,000,000, the larger part of which was taken from the Garrison mine.

In the summer of 1908 cyanide tanks were built to treat the tailings from the Cortez mill, which are estimated to amount to about 120,000 tons. At this time the mines were not producing.

GEOLOGY.

Tenabo Peak is composed of limestone and quartzite which are intruded by granitic rocks and porphyries. The southwest slope of this mountain is one of the most striking features of this portion of Nevada. High on the slope a massive ledge of quartzite from 200 to 300 feet thick forms an abrupt wall, which for a distance of about 2 miles is practically vertical. This bed strikes a few degrees west of south and dips about 23° E. On the southeast slope of Tenabo Peak it descends steeply toward the valley flat and is covered by flows of Tertiary lavas. Underneath the quartzite is a series of gray limestones, presumably the Pogonip, which are conformable with the quartzite in dip and have a thickness of not less than 2,000 feet. At the top of the mountain above the massive ledge is a series of gray limestones, possibly the Lone Mountain, which are cherty near the base and pass upward into thinly bedded limestones with some siliceous bands. The relations are indicated by figure 18. The

massive wall of quartzite appears to be uninterrupted by faulting except at the St. Louis mine, where a short block of ground is faulted above the ledge about 100 feet. On the northwest slope of the




FIGURE 18.—Section drawn eastward through Tenabo Peak, showing underground workings at Garrison mine. The position of the ore chambers is taken from the maps of the Tenabo Mill and Mines Company, and some of the ore bodies are projected on the plane of the section.

mountain a great granite intrusive cuts across the sedimentary beds and forms a broad precipitous spur which is cut by a number of ravines that drain westward to Crescent Valley. This granitic mass

extends northeastward for several miles, forming the summit of the Cortez Range to the north of Tenabo Peak and occupying its western slope. In Mill Canyon, where the granitic mass is in contact with the sedimentary rocks, it sends out small apophyses into the limestone. In some places tremolite is formed in the country rock near the contact, but garnet zones seem not to have been developed. The granitic intrusive varies in composition from a rock composed of quartz feldspar and only a little biotite to one in which the dark silicates are present in considerable quantity. A thin section of a specimen taken from the western slope of Tenabo Peak and from the upper portion of the mass is composed of quartz, orthoclase, oligoclase, biotite, and muscovite, and is therefore quartz monzonite. In Mill Canyon a more basic phase of the intrusive rock contains considerable hornblende and less orthoclase and may properly be called granodiorite.

CORTEZ MINES.

Garrison mine.—The mines on the south and west slopes of Tenabo Peak belong to the Tenabo Mill and Mines Company. The Garrison mine, which is the most extensively developed, is located about a mile northeast of Cortez and in it centers a labyrinth of closely spaced intersecting workings from which a number of long crosscuts are driven to prospect the various claims near by. The level that is most extensively developed has an elevation of about 7,000 feet above sea level. A tunnel is driven eastward for 4,000 feet and from this several long crosscuts are run to the south. At 1,600 feet from the portal a raise inclined 85° N. connects the main tunnel with the seventh-level adit, and from this raise six levels are turned at unequal vertical distances. On the main adit, 1,500 feet east of the raise and just below the quartzite, an incline is driven eastward at an angle of 23° to a depth of 145 feet below the adit level, and three other short inclines are driven from crosscuts on the adit level. On level 5 the Garrison mine is connected by a long crosscut with the St. Louis mine and on level 6 with the Fitzgerald, but these connections had caved when the mines were visited.

Nearly all the workings of the Garrison mine are in limestone and below the quartzite cliff which forms the escarpment of the mountain. The usual strike is S. 11° W. and the dip is 24° to 27° E. Although there is some minor warping and puckering of the strata, their attitude is fairly constant over a considerable area.

A dike of decomposed igneous rock cutting through the limestone strikes S. 75° E. and dips northward at a high angle. On the adit level this is exposed here and there for 2,000 feet or more from a point west of the main upraise to the east incline. Three dikes, similarly decomposed and with approximately the same strike and dip, are

crosscut in the Boss tunnel on the adit level and small exposures of the same rock may be noted at several places on the surface above the mine. These dikes, which are everywhere highly altered, are now composed mainly of quartz, sericite, and calcite; the original mineral constituents can not be made out. The dike, which is exposed here and there along the main adit, is not continuous, and from the lack of continuity it presumably did not everywhere fill the fissure into which it was intruded but was injected here and there from below. At some places along the dike the limestone near the contact is metamorphosed to a rock composed of quartz, calcite, tremolite, actinolite, orthoclase, pyrite, sericite, and biotite. In certain phases enough sulphides are present to constitute an ore, but the developments along the dike, though they include altogether several thousand feet on the six levels, have not been profitable. After the dike solidified it was sheeted and shattered, the fissures being in a broad way parallel to the intrusive but extending into the limestone wall rock also. These fissures were filled with banded sulphide ore, but most of those which are exposed on the lower levels are of too low grade to work.

The principal deposits of the Garrison mine are very irregular chambers of ore in limestone. The minerals are quartz, calcite, galena, stibnite, pyrite, zinc blende, stromeyerite, gray copper, and other minerals containing antimony and arsenic. The oxidized ore is composed of silver chloride, copper carbonates, and iron and manganese oxides. The galena is very rich in silver, especially where it is coated with dark films of a sooty black sulphide that is presumably argentite. The ore in mill runs ranges from 30 to 80 ounces of silver and \$3 in gold to the ton. Where pyrite is abundant the mill runs range up to \$15 in gold to the ton. The completely oxidized ores were very much richer. According to J. D. Hague,^a 88 tons of ore taken from the St. Louis mine in 1868 yielded \$600 to the ton.

The ore body which led to the discovery of the mine outcropped at a place near the top of an air shaft, which is raised to the surface, 400 feet in from the portal of the main adit. From the point of discovery the ore, which carried galena and silver chlorides, was followed almost vertically along a vein striking N. 60° E., to a point near the tunnel level, where the ore made out in irregular chambers in the limestone. A long, flat ribbon of ore was followed not far above the tunnel level to the Red Breast stope, a distance of about 700 feet. From the top of the Red Breast stope the ore dipped about 12° S. and was followed down a low-angle incline for 275 feet, to a point 60 feet below the tunnel level.

Eastward, along a fissure which strikes N. 20° W. and dips 55° NE., this ore joined the northwest ore body, which dips about 38° SE., as

^a U. S. Geol. Expl. 40th Par., vol. 3, 1870, p. 406.

shown in figure 19. The northwest ore body is a ribbon of ore 350 feet long and from 25 to 100 feet wide, which cuts across the limestone. As shown by the maps of the company and indicated in figure 18, it extended upward almost vertically from level 5 to level 7, forming a large flat-lying mass, which made against the quartzite roof. Farther east the middle ore body, which was approximately in the same plane, joined the large mass near the base of the quartzite ledge. At the Capps breast, about 800 feet southeast of the main upraise, an ore body which pitches westward at a high angle is followed upward through very tortuous workings to level 5.

The fissure veins that occur in the zone of sheeting parallel to the dike are stoped here and there on the tunnel level and on two levels driven from the east incline, shown in figure 18. The large stopes




FIGURE 19.—Section through Garrison mine, Cortez. Line of section trends N. 15° W. along northwest ore body, about 10 feet north of main winze.

above the fifth level, which were inaccessible when the mine was visited, are said to be in the plane of the sheeted zone. They are above the intersections of this zone and the ore bodies, which in the lower part of the mine are called the northwest and northeast ore channels.

On the south and west slopes of Tenabo Peak there are several smaller mines which the Tenabo Company exploited many years ago. On the Arctic claim, above the wagon road between the Garrison mine and the mill, a short incline is driven down a bedding-plane deposit, making off from a narrow fissure vein which cuts across the bedding of the limestone. Figure 20 is a cross section of this deposit. On the mountain trail between Cortez and Mill Canyon there are a number of small deposits of highly siliceous silver ore in limestone and in quartzite.

Valley View mine.—The Valley View mine is 3 miles northwest of Cortez, on the border of the Crescent Valley. The country is an area of tilted limestone overlain by rhyolite and cut by porphyry dikes similar to those of the Cortez mine. Between Cortez and the Valley View mine are numerous small outcrops of ore showing copper carbonates, and in the vicinity of the Valley View are several small veins of lead ore. The Valley View vein follows in a general way a dike of altered porphyry. A shaft is sunk on this vein at an inclination of 58° to a depth of 103 feet, and at a depth of 50 feet a level is turned on the vein for 50 feet to the east; at the bottom of the shaft a second drift is run for 100 feet to the east. The vein is from 1 to 5 feet wide, strikes southeastward, and consists of white quartz, galena, pyrite, and silver chloride. The values are in lead, silver, and gold.

White Horse turquoise mine.—About half a mile south of the Valley View mine some small open cuts show a white decomposed rock which is presumably an altered rhyolite. This rock contains many small




FIGURE 20.—Cross section of Arctic mine, Cortez district, near top of incline, looking north.

veinlets of turquoise from one-sixteenth to one-quarter of an inch wide. Some good gem material is said to have been obtained from this mine.

MILL CREEK MINES.

General features.—Mill Creek, the largest of several small streams which flow northwestward from the crest of the Cortez Range, drains the northern slope of Tenabo Peak through a steep-walled canyon. The mouth of this canyon is deeply incised in limestone, which about a mile above is intruded by granodiorite. To the northeast of the canyon are massive beds of quartzite and near the head are limestones that are presumably of the same age as the beds which form the summit of Tenabo Peak. Although the veins on this side of the mountain have not been so productive as the mines near Cortez, the country is highly mineralized and there is a great variety of deposits, some of which have produced considerable ore. The ore bodies are fissure veins of silver ore in granodiorite, silver-lead replacement deposits in limestone, and ferruginous deposits of gold ore replacing limestone.

The fissure deposits, which are the most important of the canyon, include the Aurora, Benjamin Harrison, Rhoda, Empire State, and other veins on Bullion Hill, a lofty spur which extends northward from Tenabo Peak. Most of the veins strike a few degrees west of north and dip eastward at moderately low angles. The ore consists of banded quartz and sulphides and carries from 30 to 200 ounces of silver and considerable lead. The walls along the veins are strongly leached by the hot waters which deposited the ores. Pyrite, sericite, calcite, and quartz are formed in abundance, replacing the ferromagnesian minerals and feldspars. The minerals of the ore are quartz, calcite, galena, zinc blende, pyrite, argentite, stephanite, gray copper, and stibnite. Along the outcrops and in some places as deep as 100 feet below the surface the veins are altered to spongy ferruginous quartz carrying silver chloride, lead carbonate, and other minerals.

The silver-lead deposits in limestone occur in the main within 300 feet of the intrusive granodiorite and are of the irregular replacement type, forming in the limestone along zones of fissuring and at some places following the bedding of the limestone. The minerals are calcite, quartz, galena, pyrite, zinc blende, and chalcopryrite. The values are mainly silver, but some of the deposits carry considerable gold.

Bullion Hill mines.—The Aurora vein of the Bullion Hill group strikes a few degrees west of north and is opened here and there on the surface for a distance of about 2,000 feet. The Water tunnel on this vein follows it for about 600 feet along the strike. The lode dips eastward and in places is a sheeted zone from 6 inches to 5 feet wide, with slabs of country rock between the ore shoots. The Aurora tunnel, 375 feet long, is driven in decomposed granodiorite 58 feet above the level of the Water tunnel, and part of the workings are west of the Aurora vein. Here and there stopes are carried up from the Water tunnel to the level of the Aurora tunnel. The ore minerals are quartz, calcite, galena, zinc blende, pyrite, chalcopryrite, gray copper, and a sooty black powder on galena, which is probably silver glance. A little ruby silver was found in the bottom of the winze on the Water tunnel level. The vein is 2 feet wide and carries about 75 ounces of silver to the ton. It has produced \$50,000 since 1883, when it was first located.

The Rhoda fissure parallels the Aurora vein and lies 30 feet to the east on the Water tunnel level. It is a great zone of crushed granite with rounded friction fragments of granite and quartz. The Rhoda shaft, which is now inaccessible, was put down 250 feet on an incline 80° W. A shoot of rich ore found in this shaft is said to have been cut off by a fault, which is possibly the same as one exposed in the Water tunnel 30 feet east of the Aurora vein. The Bullion tunnel, 260 feet below the Water tunnel level, has been driven for about 1,000

feet westward to tap the Rhoda and Aurora veins. Several small veins parallel to the Aurora have been found, but nothing of great value has yet been discovered in this tunnel.

The Benjamin Harrison vein, which lies to the east of the Aurora vein, on an adjoining claim, has been developed in tunnels and pits for a distance of about 2,000 feet. This vein is in quartz monzonite, strikes N. 25° W., and dips toward the east. The ore and the alteration of the wall rock are similar to those of the Aurora, but the vein is narrower. It belongs to the Bullion Hill Company and is said to have produced \$8,000 in silver.

Empire State mine.—The Empire State mine, one-fourth mile above the Mill Canyon mill, was located in 1872 and has produced about \$2,500. The deposit is a fissure vein in granodiorite that strikes S. 55° W. and dips 35° SE. The lode is exposed in three tunnels, the highest one about 100 feet above the lowest. In the lower tunnel the vein, which is about 2 feet wide, is stoped here and there at several places and is offset by small normal faults which dip southwestward. About 50 feet from the face a fault nearly parallel to the vein offsets it a few feet toward the west. The ore is somewhat similar to the Bullion Hill ore, but contains less galena and has higher gold values. The minerals of the sulphide ore are pyrite, zinc blende, galena, stibnite, and gray copper.

Hidden Treasure mine.—The Hidden Treasure mine is one-half mile S. 30° E. of the Rhoda shaft of Bullion Hill. From a short tunnel a 60-foot incline is driven S. 23° W. at an angle of 25°, on a lode of highly oxidized iron-stained gold ore. The hanging wall is limestone and the foot wall a white decomposed igneous rock which is probably leached quartz monzonite. At the top of the winze the ore is several feet wide and carries \$35 in gold to the ton. It thins out at the bottom of the incline, but the face of a drift along the contact for 75 feet west from the bottom is in silicified iron-stained limestone, which is said to carry \$20 in gold to the ton. At 150 feet west of the portal of the tunnel a second incline 70 feet deep is driven in limestone and intersects the same contact at the bottom.

Lewis Canyon claims.—Near the contact the limestone carries bunches of oxidized gold ore. In Lewis Canyon, on the west slope of Bullion Hill, granodiorite cuts through limestone and sends off dikes into it. A number of fissure veins strike northward and dip from 50° to 80° E. They are in granodiorite, in limestone, and at contacts between the two rocks. The veins are up to 4 feet wide, are composed of quartz, calcite, galena, and pyrite, and carry values in silver. At the Isaacs mine a silver-bearing vein strikes north, dips 80° E., and is opened in several short tunnels. In the only accessible tunnel the vein is 3 feet wide and is composed of ribbons of quartz and sulphides forming bands parallel to the walls.

At the Cynthia mine, on the northwest slope of Bullion Hill, a lode which in places is 4 feet wide strikes south and dips 68° E. It is opened by short tunnels, shafts, and underhand stopes for several hundred yards along the strike. The ore is composed of quartz, calcite, galena, and pyrite and is said to carry 60 ounces of silver to the ton. The vein was one of the first discoveries in the camp and has produced a considerable quantity of ore, which in the sixties was sent to the Canyon mill.

Falconer and Berlin mines.—About one-fourth mile below the Canyon mill the contact of limestone and granite strikes N. 53° E. and dips northwest at a high angle. The limestone, which lies northwest of the granodiorite, is locally rather closely folded and in general dips away from it at a moderately steep angle. The granodiorite sends out small dikes, which cut across the bedding and form thin sheets in the bedding planes. Along the contact, extending for a




FIGURE 21.—Cross section of ore body of Falconer mine, Mill Canyon.

distance of several thousand feet, a number of small veins and irregular shoots of silver-lead ore are developed in the limestone within a few yards of the granite, and there is an almost continuous chain of small deposits up the canyon sides from the bottom of the gulch. The ore shoots are not large, but some of them are said to have produced ore of very high grade.

The Falconer mine is on the northeast side of the canyon about 300 feet above Mill Creek. The country rock is a dark-gray limestone which strikes S. 60° E. and dips 30° NW. The ore body is a replacement deposit that cuts the bedding at a small angle and here and there small masses follow bedding planes and the small fissures which cross them to join the main ore body. Figure 21 is a cross section of this mine. From a 40-foot tunnel near the apex of the ore a 120-foot incline is sunk on the ore and small stopes are carried up from the incline at several places in the deposit. At the bottom of the incline the ore shoot flattens and dips about 20° E. It is followed in a drift

for 40 feet along the strike, at the end of which a winze is sunk 40 feet on a dip of 30° N. A crosscut tunnel 125 feet below the upper tunnel is driven northeastward for 375 feet. At 330 feet from the portal a raise is turned to the ore body to connect with the incline from the upper tunnel. The minerals of the ore are quartz, calcite, galena, pyrite, chalcopyrite, and zinc blende. Much of the ore is a black sooty powder, probably pyrite in the main. The ore is said to carry \$35 to the ton in silver, lead, and gold.

The Berlin mine, on the southwest side of the canyon, has supplied about \$40,000 in gold, silver, and lead. Most of the ore bodies are small and some seem to be related to fissures which strike northeastward and dip flatly to the southwest.

On the New York claim, which joins the Berlin on the northwest, a shaft is down 28 feet on a vein striking northwest and dipping 80° SW. At the face of a short drift this vein shows a foot of lead and iron sulphides which run high in gold. A car of ore has been shipped from this level. A 200-foot tunnel about 90 feet below the collar of the shaft has been run to intersect the ore below, but has not encountered it.

Caledonia mine.—The limestones on the northeast slope of Mill Canyon near its mouth strike N. 65° W. and dip 44° NE. A shoot of ore trending approximately with the bedding has been followed for 50 feet down an incline. A carload of ore taken from this incline and from a short drift from the bottom is reported to have yielded 120 ounces of silver and \$20 in gold to the ton and 13 per cent of lead. Approximately 300 yards farther north, at about the same horizon in the limestone, two drifts are run on a thin bedding-plane deposit which is said to carry \$25 in gold to the ton.

SAFFORD DISTRICT.

GENERAL FEATURES.

The Safford district includes a number of small mines and prospects situated in the vicinity of Barth, a small camp on Humboldt River about 6 miles west of Palisade. The district takes its name from James Safford, who discovered the Onondaga mine in 1881. The mines of the district have produced about \$200,000 in silver, nearly all of which was taken from the Onondaga and Zenoli mines. The West Mining Company is exploiting an iron deposit at Barth and ships from 100 to 300 tons of iron ore daily to the Utah plants of the American Smelting and Refining Company, where the ore is used for flux.

In the report of the Fortieth Parallel Survey the rocks of the Safford district are represented as trachytes, which are said to overlie andesites. The rocks collected by the writer proved to be varied in composition and included fine-grained diorites that carry some orthoclase and quartz, glassy vesicular andesites, and dacite por-

phyries. These species are probably the eruptions or the intrusions from a center of Tertiary volcanism which has been deeply eroded by Humboldt River, but which in the higher country on both sides of the river is nearly everywhere surrounded by rhyolite. The deposits are fissure veins of silver ore, which carries among other minerals some antimony and arsenic compounds—a type of deposit which is common in the Tertiary eruptive rocks. The wall rock is somewhat altered near the veins, the feldspar having been partly changed to sericite and calcite, and the ferromagnesian minerals are nearly everywhere altered to chlorite. These changes, which are presumably the result of the action of the vein-forming solutions on the wall rock, are not so intense as the hydrothermal action at Tuscarora and at some other places in the later Tertiary lavas.

MINE DESCRIPTIONS.

Zenoli mine.—The Zenoli mine is about 1 mile southeast of Barth, in a small gulch which joins the Humboldt Valley. The mine was discovered by Italian prospectors in the eighties and was held by them for a number of years, yielding a small amount of shipping ore. In 1907 the Zenoli Silver Copper Company was organized and since that time has been producing ore steadily and employing a force of about ten miners. The values are in silver, with important amounts of copper and lead, and the ore shipped carries from \$60 to \$70 to the ton. An inclined shaft sunk on the principal deposit to a depth of about 100 feet below the surface is intersected by an adit 220 feet from its portal, and levels are turned from this shaft at vertical intervals about 10 feet apart, making altogether about 2,500 feet of underground workings. The country rock is an andesite, which carries here and there a crystal of resorbed quartz and approaches dacite in composition. The deposit is a well-defined fissure vein from 1 to 5 feet wide. The sulphide ore is composed of quartz, calcite, barite, stibnite, gray copper, galena, pyrite, and chalcopyrite. In the upper levels there is considerable iron oxide, copper carbonates, and horn silver. The ore, which is sorted in the stopes, is shipped to Utah smelters. The principal lode strikes about north and near the surface dips 23° E.; on the 50-foot level the dip steepens to 45°. Nearly everywhere on the foot wall of the vein there is a smooth, well-defined slickensided surface carrying more or less gouge. On the tunnel level this plane of movement departs from the foot wall, cutting across the vein and displacing it to an undetermined extent. A second vein, known as the Lead Stope vein, has also a north-south strike, but dips eastward at a higher angle, joining the main lode near the surface. Some high-grade ore was found above the junction of the two veins.

Onondaga mine.—The Onondaga mine is located on the south side of the gulch, about 500 yards southeast of the Zenoli mine. Two

lodes outcrop boldly on top of the hill. The principal one strikes northwestward and dips from 60° to 80° SW. It is developed for a distance of 700 feet by several short tunnels and underhand stopes driven downward from the grass roots, the workings having a vertical range of 250 feet. A tunnel is driven northwestward for 600 feet and intersects the bottom of a shaft 200 feet in depth about 565 feet from the portal. The country rock is andesite, which is altered near the vein, and the joint planes are covered with a dark coating that is probably a mixture of iron and manganese oxides. The ore consists of quartz, iron oxide, calcite, and barite, with copper carbonates and oxides, and is said to carry from 30 to 100 ounces of silver to the ton.

In the gulch below the mine on the Malachite claim a crosscut tunnel is driven northward for 350 feet toward the Onondaga vein.

Ruby claim.—The Ruby claim, which is a few rods east of the Zenoli mine, adjoins the Malachite claim on the northwest. Four tunnels, from 50 to 250 feet long, have been driven to explore three veins which strike northwestward. The principal lode is an iron-stained zone of crushed porphyry, which here and there carries banded ribbons of ore composed of quartz, chalcopyrite, pyrite, gray copper, and iron oxides, with a little ruby silver. A carload of selected ore, running about 100 ounces of silver to the ton, has been shipped to smelters.

Humboldt mine.—The Humboldt mine is located on the north side of Humboldt River, about $3\frac{1}{2}$ miles below Palisade. Andesite, which is locally vesicular, outcrops at the surface, and copper-bearing sulphides have been deposited in a zone of brecciation. Some ore taken from a shallow pit shows bornite, chalcopyrite, and chalcocite, partly altered to copper carbonates and iron oxides. A crosscut tunnel 200 feet long has been driven at the level of the river, but has not yet encountered the lode.

Bonanza mine.—The Bonanza mine is about three-fourths of a mile S. 70° E. of the Humboldt mine. The lode is a zone of crushed and altered eruptive rock, cemented by iron oxides. In the shattered zone, which can be traced for about 200 feet, thin sheets of galena, rich in silver, may be found here and there at the surface, filling the crevices in the shattered rock. Two shallow pits have been sunk on the deposit.

Pittsburg and Palisade tunnel.—On the Pittsburg and Palisade group of claims, which join the Humboldt on the west, a crosscut tunnel has been driven for 600 feet and is intended to intersect a small vein that outcrops on the hill above.

West iron mine.—The West iron mine is located at Barth, 6 miles below Palisade. An open cut 200 feet in diameter and 80 feet deep is dug in the red hematite ore, which is shipped to Salt Lake and used for flux. The rock is quarried from the pit, hoisted up an incline, and loaded into railroad cars.

The cut is approximately the size of the known area of iron ore. The rock to the north and east is covered by gravels, which at the pit are about 20 feet deep, and on the south and west the iron ore is in contact with a fine-grained diorite. About 100 feet south of the pit the fine-grained diorite is in contact with an altered rock, probably a metamorphosed limestone. The iron ore, which is in the main red hematite, carries a very small amount of magnetite and specularite and is cut by many small veinlets of apatite. The deposit is presumably of contact-metamorphic origin and not the capping of a sulphide ore. The sulphides alter mainly to hydrated iron oxide, and not to hematite and magnetite, like the ore of the West mine. Outcrops of iron deposits are reported to have been found also in the Cortez Range about 15 miles south of Palisade.

SHOSHONE RANGE.

GENERAL FEATURES.

The Shoshone Range extends southwestward from Humboldt River near Beowawe for 50 miles or more. The present reconnaissance covered only the northern part of the range, which includes a group of lofty mountains, among them Shoshone Peak, 9,760 feet above the sea, the most elevated portion of the mountain mass. The central portion of the north end of the range is a group of steep-walled mountains with deeply incised amphitheatres, which are clearly the result of mountain glaciation. Northeast of the group of elevated peaks is the Whirlwind Mesa, which extends about 15 miles to Humboldt River. This mesa, which is over 7,000 feet above sea level, slopes gently northward and is divided by the Whirlwind Valley, which joins the Humboldt Valley near Beowawe.

As the central mountain mass is higher than the neighboring mountains, it is better watered and it supports a scattered growth of timber useful for local needs. The mining camps of the north end of the range are Tenabo, Lander, Mud Springs, Grey Eagle, Hilltop, Maysville, Pittsburg, Dean, and Lewis; all of these are on the east and north slopes of the mountains and are included in a rectangular area 6 by 16 miles. In the eighties there was considerable mining in some of these camps, especially at Lewis, Dean, and Lander. The Battle Mountain and Lewis Railroad, a narrow-gauge line, was built from a point on the Nevada Central Railroad through Lewis to Quartz Mountain, at the north end of the range.^a This road was never profitable and it was dismantled many years ago. At present the transportation to the various camps is by wagon road from Beowawe and from Battle Mountain.

^a Bancroft, H. H., Works, vol. 25, Nevada, p. 238.

Development work was being carried on at several of these camps in 1908 and some of the mines were shipping a little ore. The Dean mill, the only one in working condition, was idle at that time.

GEOLOGIC FEATURES.

The most abundant rocks of the northern part of the Shoshone Range are quartzites, siliceous shales, and limestones. Interbedded with these are conglomerate beds which are exposed, among other places, on the north slope of Shoshone Peak. On lithologic grounds^a the sedimentary series has been termed Carboniferous, but a considerable thickness of other rocks may be represented. At most places the sedimentary rocks dip toward the east or southeast from 20° to 50°, but locally they dip at high angles in other directions. The structure, as shown in the atlas of the Fortieth Parallel Survey, is that of an eastward-dipping monocline crossed by normal faults which dip westward. The sedimentary rocks are intruded by large masses of granodiorite and granodiorite porphyry. Such intrusive rocks are exposed at Tenabo, Mud Springs, Grey Eagle, Hilltop, and Dean, and the largest one of them forms the central part of Shoshone Mountain. An analysis of this rock, made by R. W. Woodward, is given on page 25, where the granodiorites are described.

Some actinolite and doubtless other contact-metamorphic minerals are developed in the quartzite, but no garnet zones were found in the few places where the contacts were observed. The siliceous sedimentary rocks are the most abundant and these, as is well known, are not so favorable to contact metamorphism as the calcareous rocks. Many of the ore deposits are in or near the granodiorite.

Dense andesites outcrop at Tenabo and at Lander, where they are surrounded by sedimentary rocks. The andesites have a glassy groundmass and in some places they are vesicular. It is assumed that they represent a magma which consolidated as an intrusive near the surface or as a flow. A broad belt of rhyolite borders the range low down on the east slope. A small intrusive mass of dense quartz porphyry was noted on the north slope of the gulch below the Pittsburg mine.

TENABO.

GENERAL STATEMENT.

Tenabo is situated on the east slope of the Shoshone Range, near the edge of Crescent Valley, about 21 miles southwest of Beowawe. It is in the Bullion mining district, which was organized many years ago to cover locations at Lander, but when a number of ore deposits were discovered in 1907 at some distance from that camp the new

^a U. S. Geol. Expl. 40th Par., vol. 2, 1877, p. 619.

town Tenabo was established 2 or 3 miles away, more conveniently situated for them. The town is well laid out and has some substantial wooden buildings, but is without a regular water supply. Early in 1907, when there was a rush to the mines, about a thousand persons were living there, but within a year all except a few score had left. A stage is run thrice a week between Tenabo and Beowawe, and during times of prosperity automobiles and a steam traction line were operated. The name Tenabo is not well chosen, as it is likely to be confused with that of Tenabo Peak, on the opposite side of the Crescent Valley, where the Tenabo Mill and Mines Company carried on extensive operations for many years.

GEOLOGY.

The relief at Tenabo is slight and consequently the natural exposures are not so good as at camps which are situated in more commanding positions, but owing to the large number of location pits recently dug a fairly satisfactory knowledge of the geologic features may be obtained. The rock formation which occupies the greater part of the area is composed of quartzites and fine-grained siliceous shales that have a general eastward dip of 15° to 40° . There are no pure limestones in the area near Tenabo, but some of the sandy layers of the quartzitic formation contain calcium carbonate which cements the grains of quartz. Certain fissile layers on the ridge above the Gem mine are dark and very fine grained and in the field resemble carbonaceous shale, but under the microscope they are seen to be composed of very fine particles of quartz with a small amount of sericite between the grains. The series is termed Carboniferous in the reports of the Fortieth Parallel Survey. The sedimentary rocks are intruded by porphyritic granodiorite, which is best exposed in the Phoenix mine. This rock in hand specimen has a dense gray groundmass, in which are embedded phenocrysts of feldspar, quartz, biotite, and hornblende. Under the microscope the groundmass is seen to be composed of quartz and orthoclase crystals which are distinctly smaller than the phenocrysts but larger than the constituents of the groundmass of the ordinary porphyries. The feldspar phenocrysts are in the main oligoclase and andesine, and many of them are zonally built. At the Two Widows claim a quartz diorite porphyry, which is probably a phase of the granodiorite magma, intrudes siliceous shales. Near the contact small veinlets of actinolite are developed, presumably as a result of contact metamorphism, but in the sections studied no garnet or other contact minerals have been found.

A dark, dense rock with an altered glassy groundmass containing phenocrysts of orthoclase and acidic plagioclase is exposed at the Gem mine and on the hills above the shaft. This rock will be called

andesite, but its determination is unsatisfactory owing to the decomposed condition of the material at hand; the slides studied contain more orthoclase than the other andesites described in this report, and possibly the rock should be classed as latite. It is vesicular in places and is everywhere finely crystalline and obscurely porphyritic, and it is therefore probably a flow or an intrusive formed near the surface. The andesite is believed to be younger than the other porphyries and appears to have formed at less depth. Near the lodes the country rock is hydrothermally altered, but the altered rock is restricted to a relatively narrow zone. Sericite, calcite, and pyrite have formed in the granodiorite, and chlorite, sericite, and pyrite are developed in the andesite.

ORE DEPOSITS.

The ore deposits are sheeted zones in quartzose sedimentary rocks and in andesite and fissure fillings in granite. The lodes do not fall into well-defined groups or parallel systems, but dip to all points of the compass. On the surface the ore is iron-stained quartz, carrying some chloride, and is usually kaolinic. Some of the surface ore gives very high pannings in gold. Sulphides appear in depth from 50 to 100 feet below the surface, and include arsenopyrite, pyrite, chalcopyrite, bornite, galena, zinc blende, and chalcocite. The gangue is quartz and calcite. Thin films of molybdenite appear on the joint faces of the country rock in the Violet shaft. Since the deposition of the ore there has been considerable movement, and practically all the lodes are sheared by slickensided planes parallel to the vein or are faulted by cross faults. These are small as far as developed and do not cause loss of the vein in any of the mines visited.

MINE DESCRIPTIONS.

Little Gem mine.—The Little Gem mine, located about $1\frac{1}{2}$ miles west of Tenabo, was discovered in 1907 and is the most extensively developed property in the area. A shaft is driven southwestward on the lode at inclinations of 20° to 30° . This shaft is 400 feet long and on the four levels turned from it there are altogether about 900 feet of drifts. In 1907 eighteen carloads of ore was shipped and yielded about \$30 a ton.

The country rock is a dark, fine-grained andesite, showing very few phenocrysts even where most porphyritic. On the hill just above the mine it is vesicular. The andesite is strongly sheeted parallel to the lode and is altered near the lode, but at most places it is not strongly leached; chlorite and pyrite, with some sericite and calcite, are developed by secondary processes, chlorite being formed in the greatest abundance. The ore outcrops near the collar of the shaft and at the outcrop is composed of iron oxide and quartz carry-

ing copper carbonates, silver chloride, and free gold. The sulphide ore, which appears about 75 feet below the surface, is composed of quartz, arsenopyrite, pyrite, chalcopyrite, galena, and zinc blende, with a little bornite and a sooty black film which covers other sulphides and is probably chalcocite. On level 1 the lode strikes north of west, but on the second level, the most extensive in the mine, it bends and strikes south of west. It dips about 30° S., a little more steeply than the incline, which is driven on the lode but to the right of the line of steepest dip. The lode is from 1 to 6 feet wide and, in the main, is a zone composed of several closely spaced parallel sheets of quartz and sulphides, between which the country rock is highly fractured and seamed with veinlets crossing the general strike of the vein. Thin drusy cavities parallel to the walls are lined with sulphide coatings, and these are in turn covered with quartz crystals pointing to the center. Since it was formed the vein has been much crushed and shattered by movement, which in the main was parallel to the walls. It is also crossed by several small normal faults, the offsets of which are not great enough to throw the vein out of continuous view in regular workings. The mine is reported to have in sight 7,000 tons of ore, with an average value of \$2 in gold and 10 ounces of silver to the ton and 3 per cent of copper.

Phoenix mine.—At the Phoenix mine, $1\frac{1}{4}$ miles southwest of Tenabo, a two-compartment vertical shaft is 250 feet deep, and an adit driven northward for 280 feet intersects the shaft about 100 feet below the collar. This adit is continued northward for 220 feet beyond the shaft, and short drifts are run on two veins at this level. A second level is turned from the shaft 60 feet below the adit. The country rock is porphyritic granodiorite, cut by a mass of intruding quartz porphyry, which measured on the adit level is about 100 feet thick. The south contact of the quartz porphyry, where it is encountered in the main adit, 70 feet from the portal, dips about 20° S. Three veins are exposed, two of them in granodiorite and one at the north contact of the quartz porphyry and granodiorite. They are from 4 to 18 inches wide and all strike westward. The deposits are banded fissure fillings, and the country rock near the lode is strongly sericitized. The sulphide ore, 100 feet in depth, is composed of quartz, pyrite, arsenopyrite, chalcopyrite, galena, and blende. In the granite sericite, calcite, chlorite, and pyrite have been deposited by ore solutions. There has been much movement since the veins were formed, and in places the lodes consist of 2 feet of white sericitized decomposed granite, containing here and there broken masses of quartz and ore up to a foot thick.

Gold Quartz mine.—The Gold Quartz mine is 700 yards south of the Little Gem. A vertical two-compartment shaft is down 308 feet, with short levels turned at intervals of 35, 100, 150, and 275 feet

below the collar. The country rock is quartzite, cut by an intruding quartz-bearing porphyry and by basalt. In a shallow opening near the main shaft some rich gold ore was encountered. From this and from the main shaft a shipment of ore, with an average value of \$80 a ton in gold, has been made. The ore is highly oxidized; about 75 feet below the surface the sulphides appear, arsenopyrite and pyrite predominating. In the main shaft, between the 35-foot and 100-foot levels, there is a body of shattered, altered quartzite, which is cut by stringers of iron oxide carrying free gold. This ore body is 4 feet wide, strikes N. 20° W., and dips 35° NE. Between the 35-foot and 100-foot levels an incline follows it for 30 feet; at the bottom of the incline the lode breaks into several small stringers, and this deposit has not been encountered below the 100-foot level.

Violet claim.—The Violet shaft, about 1 mile southwest of Tenabo, is down 208 feet in siliceous shale and quartzite. A crosscut is being run below an outcrop which has been opened in a surface pit. In the jointing of the country rock there are some small seams of molybdenite.

Two Widows claim.—On the Two Widows claim, half a mile west of Tenabo, an incline is sunk 110 feet at an angle of 70°, and from the bottom short crosscuts are run east and west. The country rock is fine-grained quartzite cut by intruding quartz diorite porphyry. Veinlets of actinolite cut the quartzite near the intruding porphyry. The joint planes of the quartzite are filled with copper carbonates and iron oxides, which are said to carry 40 ounces of silver and several dollars in gold to the ton.

LANDER.

GENERAL FEATURES.

Lander, 2 miles northwest of Tenabo, is the oldest camp in the Bullion district and was the milling center for the district in the seventies and eighties, when the Lovie mines were being worked. The rocks of the area are siliceous and carbonaceous shales interbedded with quartzite and limestone and have a general southeastward dip. The sedimentary rocks are capped with andesite and cut by intrusive quartz porphyry. The principal mines are the Bonnie Jean, the Silver Prize, and the Silverside.

MINE DESCRIPTIONS.

Bonnie Jean mine.—The Bonnie Jean, known also as the Lovie mine, is 1½ miles northwest of Lander and is opened by four tunnels which have a vertical range of 250 feet. In the gulch below these workings a shaft was being sunk in 1907 to explore the vein in depth. This mine was worked in a small way for many years, and the ore was treated in the 5-stamp pan-amalgamation mill at Lander. It

is currently reported to have produced some \$300,000 in silver, the chloride ore mined near the surface of the deposits running several hundred ounces to the ton. The ore body is a fissure vein in siliceous shales and quartzite. Near the apex and northwest of the deposit is an outcrop of andesite similar to that of the Little Gem mine at Tenabo. The lode, which lies approximately with the bedding, strikes N. 65° E. and dips 20° to 50° SE. The outcrop of the deposit is exposed at the surface for about 400 feet along the strike and has been followed down the dip for about 300 feet. The oxidized ore is composed of quartz, iron oxides, silver chloride, and lead and copper carbonates. The values are principally in silver. There is a considerable tonnage of low-grade silver ore blocked out which the present owners consider concentrating ore.

Silver Prize vein.—On the Silver Prize claim, half a mile north of the Bonnie Jean, the country rock is decomposed andesite, and to the northwest, near the crest of the hill, are outcrops of massive quartzite. The lower vein, which is opened in three short tunnels, strikes northwest and is approximately vertical. The lode locally carries 2 feet of highly oxidized ore, and galena and zinc blende appear in depth. Higher on the hill and about 75 feet to the northeast of this lode a second lode strikes N. 50° W. and dips 58° SW. A 40-foot incline is driven on the lode, which carries 20 inches of rich silver-bearing galena and lead carbonate. The two veins probably join on the slope of the hill.

Silver Side mine.—At the Silver Side mine, in Lander, a tunnel is run 625 feet southward on a vein which dips 20° to 38° E., and from this stopes are carried here and there above and below. The vein, which is about 3 feet wide, is at most places approximately parallel to the bedding. The deposit with respect to its structural features resembles that of the Bonnie Jean, but the ore is not so rich.

MUD SPRINGS.

GENERAL FEATURES.

Mud Springs is on the east slope of the Shoshone Range, about 4 miles north of Lander. In the summer of 1907 a number of claims were located, and recently a few tons of ore has been shipped. The country rock consists of quartzite, with fine-grained, shaly beds, and is cut by granodiorite and andesite. The deposits are fissure veins in the sedimentary rock and in granodiorite. The minerals of the ore are quartz, limonite, lead carbonate, silver chloride, and gold.

MINE DESCRIPTIONS.

Triumph mine.—At the Triumph mine a narrow lode carrying oxidized gold and silver ore cuts across siliceous shales, strikes N. 60° W., and dips 45° SW. It has been followed for 40 feet down an incline from which a 50-foot drift has been run, and from this a

stope has been raised east of the incline. On the surface 300 yards farther west a vein which is presumably the same cuts through a mass of undetermined igneous rock, probably an altered granodiorite. From this mine in the summer of 1908 a shipment of 18 tons of ore was made. This ore carried silver, lead, and gold, and is said to have yielded \$60 a ton. The lode is from 1 to 3 feet wide, and the ore minerals include iron oxide, lead carbonate, and silver chloride.

Big Bug claim.—At the Big Bug claim two tunnels, each about 100 feet long, are driven to intersect a lode cutting across quartzite which dips steeply southeastward. The lode is about 6 inches wide, strikes N. 55° W., and dips 80° S. The decomposed iron-stained quartzose ore carries low values in gold and silver.

Bridal Wreath claim.—On the Bridal Wreath claim, which joins the Big Bug claim on the northeast, a 30-foot shaft is sunk on a crushed vein in quartzite which strikes S. 65° W. and dips 70° N.

Uncle Sam claim.—The Uncle Sam shaft, half a mile below the Big Bug, is sunk 50 feet on a silver lode which dips 76° S. A small vein of quartz and sulphides is located near a contact of granite and quartzite.

GREY EAGLE MINE.

The Grey Eagle mine, a mile or two west of Mud Springs and about 22 miles in an air line southeast of Battle Mountain, is located near the summit of a high granodiorite ridge, which is separated by a deep valley from the main axis of the Shoshone Range. The mine was worked in the seventies and eighties, lay idle some twenty years, and was worked again in 1905. It produced about \$25,000 in 1906 and 1907. The main shaft is down 250 feet, and levels are turned at 60, 115, and 215 feet in depth. When the mine was visited in 1908 only the 60-foot level was accessible. The deposit is a fissure vein of banded quartz and sulphide, strikes N. 70° E., and dips 70° N. The country rock is a coarse granodiorite composed of oligoclase, andesine, quartz, hornblende, and biotite. Near the vein the granodiorite is strongly altered by hydrothermal metamorphism. The feldspars, biotite, and hornblende are replaced by sericite and by numerous small crystals of pyrite. The minerals of the ore include quartz, zinc blende, galena, pyrite, and gray copper, which near the surface are altered to oxides, carbonate, and chloride. The mine is reported to have produced since its discovery several hundred thousand dollars in silver, gold, and lead. It is commonly regarded as a property which possesses considerable promise.

HILLTOP.

Hilltop, or Marble Canyon, is a small camp which has recently been established on the east fork of Rock Creek, about 18 miles southeast of Battle Mountain. The country rock is quartzite,

which includes fine siliceous shales with here and there some fine conglomerates containing small fragments of angular jasper. The sedimentary rocks are cut by dikes of green chloritic porphyry which under the microscope shows phenocrysts of quartz, orthoclase, and some oligoclase with ferromagnesian minerals altered to chlorite and sericite. This rock is presumably an altered granodiorite porphyry.

The principal mines are the Independence and Hilltop, which were located in 1906. Two 50-foot shafts and two short tunnels are driven on a zone of fractured quartzite, which trends northward and is exposed at intervals for about 2,000 feet. Small intrusive masses of leached porphyry cut this zone of quartzite at three or four places, the intrusive rock including fragments of the quartzite. At some places the shattered quartzite is seamed with stringers of quartz and iron oxide carrying a large amount of free gold. Here and there are small bodies of pyrite and galena in the fracture planes of quartzite and minute stringers of gray metallic mineral which is said to contain bismuth. Assays show the presence of silver, by weight equal to the gold. The quartzite near the ore is impregnated with pyrite and is locally decomposed to a soft white claylike substance which is said to carry high values in gold.

When the camp was visited the amount of high-grade ore in sight was small and developments were not sufficient to show the extent of mineralization in the fractured zone.

MAYSVILLE MINE.

The Maysville mine is on the east slope of Shoshone Peak about 1 mile west of Hilltop. The country rock is quartzite, which near the lode dips east of south about 35°. The principal vein cuts across the quartzite, striking N. 80° W. and dipping 80° S., and is opened by three short tunnels and a shaft sunk on the deposit. In places it is stoped to the grass roots, the surface ore consisting of yellowish-brown siliceous rock which carries silver chloride. The ore about 50 feet below the surface is composed of quartz, pyrite, galena, chalcoppyrite, and gray copper. The vein, where accessible, is about 2 feet wide and shows banded quartz and pyrite parallel to the walls. The quartz and sulphides are highly crushed and at places mixed with much pulverized quartzite. About 1,500 feet northwest of the shaft a crosscut tunnel is driven southward for 215 feet to a vein, which is presumably the same lode. From this tunnel drifts are run 150 feet east and 75 feet west. The lode here strikes N. 60° W. and dips southward at a high angle. On the east drift some stoping has been done, but at most places the vein is small. A four-pan silver mill in the canyon below the mine was used some thirty years ago to treat the ore.

LEWIS AND DEAN.

LOCATION AND HISTORY.

Lewis is situated about 12 miles southeast of Battle Mountain, on the north edge of the elevated portion of the Shoshone Range, at the mouth of Lewis Canyon. Dean is 4 miles above Lewis, in the bottom of Lewis Canyon, which at this place is one of the most rugged and picturesque mountain features in northeastern Nevada. Silver deposits were discovered in Lewis Canyon in the seventies, and elaborate preparations were made by a company well provided with capital to develop the Eagle, Starr Grove, and other mines. A railroad which has long been dismantled was built from Lewis Junction to Lewis, and afterward was extended up Lewis Canyon to the Starr Grove mine. At Lewis a small silver amalgamation mill was built on the east side of the canyon, and considerable silver ore is said to have been put through this plant. Subsequently a 40-stamp mill equipped with roasters, pans, and settlers was built on the west side of the canyon, but was not long in operation. Extensive development work was done in several mines and some ore was stoped, but it is doubtful whether the mining and metallurgical operations were ever on a paying basis.

The gold deposits of the Morning Star and Pittsburg mines were opened soon after the silver deposits at Lewis. According to reports of the Director of the Mint both of these mines were being developed in 1882. The Pittsburg mine is located near the summit of a ridge east of Lewis Canyon, and in the eighties a mill was built in the canyon east of this ridge and connected by a tramway with the mine. This mill reported a production of \$46,000 in 1887 and was in successful operation until 1891. The Morning Star mine, on the same ridge and southwest of the Pittsburg, was sold in 1892 to W. E. Dean, of San Francisco, who built a 10-stamp amalgamating mill in Lewis Canyon and ran an adit from the mill to the lode. This mine was worked for many years under the management of D. J. Bousfield. After tedious litigation with the owners of the Pittsburg mine that property was acquired in 1904 by the owners of the Morning Star and the two mines were consolidated as the Cumberland mines. The Morning Star mill has not been in operation since 1906 and the company is devoting its attention to a long adit which it is driving to develop the lodes below the present workings.

GEOLOGY AND ORE DEPOSITS.

The rocks in the vicinity of Lewis and Dean are in the main quartzites, limestones, and shales, which, on lithologic grounds, were termed Carboniferous by the geologists of the Fortieth Parallel Survey. The quartzite beds are more extensive than the other sedimentary rocks, although some of the limestone and shale members have a thickness

of several hundred feet. High on the north slope of Shoshone Peak there are some rather coarse conglomerates, which presumably are the Weber. The sedimentary rocks have a general northeastward dip, but locally dip in other directions. They are intruded by granodiorite porphyry, the somewhat altered specimens of which are grayish-green rocks composed of a dense greenish paste carrying phenocrysts of feldspar, quartz, hornblende, and biotite. Under the microscope the groundmass is seen to be a microcrystalline aggregate of colorless minerals and the phenocrysts are oligoclase, andesine, and quartz, with hornblende and biotite altered to chlorite, sericite; and other minerals. This porphyry may have the composition of the rock from Shoshone Peak analyzed by R. W. Woodward and described on page 25. The sedimentary rocks are also cut by quartz porphyries, some of which are glassy devitrified rocks which were formed presumably very near the surface. That below the Pittsburg mine has the dense pasty appearance of rhyolite.

The ore deposits are fissure veins of pyritic gold ore in quartzite and in granodiorite porphyry and replacement deposits of siliceous silver ore in limestone. The gold deposits are the Morning Star and Pittsburg lodes and the silver deposits include the lodes of the Eagle, Starr Grove, Betty O'Neil, and other mines which have long since caved.

MINE DESCRIPTIONS.

Morning Star and Pittsburg mines.—The Morning Star and Pittsburg mines of the Cumberland group are developed on different lodes. They are close together and were worked through the same adit. The Morning Star with connecting adits has over 2 miles of underground workings, which have a vertical range in elevation of 925 feet. The Concert and Mayo inclines are sunk on the Morning Star lode about 300 feet apart to the level 230 feet below the collar of the Concert incline, and from this level a crosscut adit connects with the surface on the east side of the ridge. A second adit is driven from the mill northeastward for 1,500 feet to the lode, which it intersects on the 460-foot level. A vertical winze, 1,335 feet from the portal of this adit, is sunk 465 feet to connect with the lowest adit, which is driven eastward for 3,500 feet to the winze. An incline, driven on the lode near the foot of the Mayo incline, connects the 460-foot level with the 230-foot level, and three levels are turned between.

The country rock is quartzite and granodiorite porphyry. The quartzite is of medium grain, gray or buff in color, and includes some thin-bedded siliceous shales. The granodiorite porphyry is a mass of large areal outcrop which intrudes the quartzite and sends off thin dikelike apophyses into it. The lode is a fissure vein which cuts both the quartzite and the porphyry and is mineralized in both. The country rock is greatly altered at some places for 50 feet from the

vein. The porphyry has suffered greatest alteration and is changed to a mass of sericite and pyrite with a little calcite. Farther away chlorite and calcite are formed as an alteration of hornblende, biotite, and other minerals. The quartzite also is altered by the vein-forming solutions, the change being in the main the development of pyrite and sericite. Small amounts of galena and chalcopryrite are intergrown with the pyrite of the porphyry.




FIGURE 22.—Cross section along Mayo incline, Morning Star lode, Dean.

The lode outcrops at the surface on the east slope of the ridge and the main fissure was followed downward to the 380-foot level. It splits below this level and both branches carry good ore. On the 460-foot level there are six veins approximately parallel and in places three of them carry ore shoots. On this level the lode is composed of branching and forking, anastomosing fissures, the whole series being confined within a relatively narrow zone which strikes

north and dips about 60° W. A vertical cross section of the lode in the plane of the Mayo incline is shown by figure 22. The ore is chiefly quartz, pyrite, and arsenopyrite, all of which carry free gold. Calcite and barite are present in subordinate quantities in the gangue and a little galena and zinc blende are associated with the pyrite. The quartz and sulphides are locally banded. In places comb structure is shown and drusy cavities with quartz crystals pointing to the center represent the filling of open spaces. At some places the wall rock near the ore carried working values, but the filled portion of the vein is much the most important part. Near the surface at the outcrop of the deposit, but not below a depth of 100 feet, the ore is oxidized to a brownish-yellow mass carrying considerable iron oxide and some lead carbonate. A little native copper was found on the 128-foot level in the oxidized ore, and a small amount of ruby silver is reported from the 380-foot level. The veins vary from the thickness of a knife-edge to 10 feet. The principal ore shoot which was opened between the Concert and Mayo inclines had a maximum length along the level of about 300 feet altogether and pitched approximately down the dip of the vein. It split between the 380-foot level, forming an inverted Y with ore on both legs.

The waters of the mine are acid, and this condition, together with the somewhat crushed state of the ore, would favor secondary enrichment, but little is evident from the exploration done. The sulphide ore on the 430-foot level is said to be as rich as the ore higher up and of about the same value as the oxidized ore, which did not extend more than 100 feet below the surface.

The Pittsburg mine, which joins the Morning Star on the northeast, is opened on seven levels, which have a vertical range of 550 feet and include about 4,500 feet of horizontal workings, principally drifts on the Pittsburg vein. The vein strikes west, dips 60° S., and is composed of quartz, calcite, pyrite, and arsenopyrite, with a little galena, blende, chalcopyrite, and gray copper. The country rock is quartzite and granodiorite porphyry, the latter predominating. The wall rock is altered by hot waters, as in the Morning Star. The ore carries free gold which is mainly associated with quartz and pyrite, and a small amount of silver is present, with less than 1 per cent of copper.

The principal ore shoot pitches westward in the vein at an angle of about 45° and has been stoped from near the grass roots to levels about 400 feet vertically below the surface. On the levels this ore shoot is from 50 to 150 feet long and from 2 to 10 feet wide. Another stope to the east of this ore body lies between the fifth and sixth levels and extends downward to level 7, having a vertical range of 175 feet and a length on the levels of about 140 feet.

As the Morning Star lode strikes about N. 30° W. and the Pittsburg lode strikes west, the two should intersect in the Morning Star workings on the adit level 460 feet below the Concert incline. Some small fissures which have about the same attitude as the Pittsburg vein have been exposed in the east end of the Morning Star workings, but it can not be shown now whether they represent the Pittsburg lode or sheeting parallel to it.

Starr Grove mine.—The Starr Grove mine is one-eighth mile southwest of the compressor plant of the Cumberland mines, which is one-fourth mile below the portal of the lowest adit of the Morning Star. A crosscut tunnel is driven for 150 feet to an ore body which is followed through irregular drifts and crosscuts, some 200 feet of which are still open. A deep shaft and other workings below the tunnel are inaccessible. The country rock is dark-gray limestone, which above the mine is overlain by massive quartzite beds. The deposit is a large flat body which lies approximately parallel with the bedding of the country rock, but locally cuts across it. Where developed it strikes southward and dips at a very low angle toward the west. The ore body is composed of barite and quartz, carrying here and there a small amount of pyrite, galena, and zinc blende, finely disseminated in the white gangue. At some places there is as much as 10 feet of nearly pure barite. The quartz and sulphides are in part of later origin than the barite, for at some places the crushed fragments of barite are surrounded by the quartzose ore and cracks of the ore cross the barite. On the dump some quartzose ore carrying ruby silver was found.

Betty O'Neil mine.—The Betty O'Neil mine, about half a mile west of Lewis, was worked through a 225-foot shaft which was under water when the mine was visited. A shallow adit is driven for 70 feet southwestward to the vein, which it follows for 300 feet. The vein strikes S. 30° E. and dips from 20° to 50° E. The country rock is quartzite with shaly black siliceous beds, and fragments of porphyry were noted on the dump. The vein is from a few inches to 3 feet wide and is composed of banded quartz and sulphides, chiefly pyrite, galena, and zinc blende. In 1908 a crosscut tunnel was being driven to drain the old workings.

INDEX.

A.	Page.
Acknowledgments to those aiding	11
Agate Pass, granite from, analysis of	26
quartz diorite from, analysis of	27
Ajax claim, ores of	85
Alluvial deposits, character and distribution of	23-24
Alpha, description of	99
ore deposits at	99
Andesite, age of	32-33
analysis of	33
character and distribution of	32
fissure veins in	44
Arctic claim, description of	105
Arizona claim, description of	99
Ash, volcanic, character and distribution of ..	21-23
Aura, description of	71
ore deposits at	42, 70, 71-74
placers at	45
production of	15
Aura King mine, description of	74
Aurora vein, description of	107
B.	
Baltimore group, description of	55, 65
Barth, mines near. <i>See</i> Safford district.	
Basalt, character and distribution of	31
Becker, G. F., on propylite	34
Belvoir claim, description of	57
Benjamin Harrison vein, description of	108
Berlin mine, description of	110
Betty O'Neil mine, description of	126
Bibliography of region	11-12
Big Bob vein, description of	79
Big Bug claim, description of	120
Big Four mine, description of	72
section in, figure showing	72
Big Six mine, description of	88
Blue Belle mine, description of	95
Blue Jacket Canyon, ore deposits in	69, 71
Bonanza mine, description of	112
Bonneville, Lake, location of	24
Bonnie Jean mine, description of	118-119
Bridal Wreath claim, description of	120
Buckeye and Ohio mine, ores of	65-66
plan of	66
Bullion, description of	89
geology of	90-93
intrusion near	25, 28, 29
mines of	93-95
ore deposits at	41, 90-95
production of	15
Bullion Hill mines, description of	107-108
Bull Run, deposits at	42
Bull Run mine, description of	79-80
section of, figure showing	79

	Page.
Burner Hills, description of	66-67
ore deposits at	44, 67
C.	
Caledonia mine, description of	110
California Hill, ore deposits of	81
California mine, description of	74
Cambrian rocks, occurrence and character of ..	16, 17-18
Carboniferous rocks, occurrence and character of	16, 17, 20-21
Carlin Peaks, rocks of	86-87
Centennial Range, description of	67-68
geology of	68-69
intrusions in	25, 28, 69
ore deposits in	41, 42, 69-70
<i>See also</i> Lime Mountain; Columbia; Aura; Edgemont; Mountain City.	
Chimney, mill at	91
Climate, description of	12
Coal measures, character and distribution of ..	16, 20-21
Columbia, description of	71
ore deposits at	42, 69, 71-74
Columbia Queen mine, description of	72
plan of, figure showing	73
Columnar section, description of	15-17
figure showing	16
Contact-metamorphic deposits, description of	40-41
Copper, occurrence of	43, 91-93
Copper Belle mine, description of	94
Copper King claim, description of	87
Cornucopia, geologic map of	63
ore deposits at	44, 62-65
production at	62
Cortez district, description of	100-101
geology of	101-103
Cortez, intrusion near	26, 28, 30
ore deposits at	41, 103-106
section of, figure showing	102
Cortez Range, andesite from, analysis of	33
description of	86-87, 99-100
geology of	86-87, 100
ore deposits of	87-88
mines of	87-88
<i>See also</i> Cortez; Mill Canyon; Safford districts.	
Cretaceous ore deposits, character and distri- bution of	40-43
minerals in	40
production from	40
<i>See also</i> Contact metamorphic deposits; Replacement deposits; Fissure veins.	
Crops, character of	12-13

K.	Page.
Kenilworth claims, description of.....	95
King, C., on geologic history.....	37-39
on glacial history.....	24
on Humboldt formation.....	22
on intrusive rocks.....	27
on Weber quartzite.....	21
L.	
Lahontan, Lake, location of.....	24
Lander, description of.....	118
ore deposits of.....	118-119
Lavas. <i>See</i> Eruptive rocks.	
Lead, production of.....	15
Lewis, description of.....	122
geology of.....	122-123
ore deposits at.....	41, 42, 123-126
Lewis Canyon claims, description of.....	108-109, 122
Lime Mountain, description of.....	70
ore deposits at.....	41, 70-71
Limestone, replacement deposits in.....	41-42
Lindgren, Waldemar, on age of intrusions...	29
on granodiorite.....	24-25
on ore deposits.....	44
Literature, list of.....	11-12
Little Gem mine, description of.....	116-117
Location of area.....	9
maps showing.....	10, 13
Lone Mountain, description of.....	84-85
geology of.....	85
ore deposits at.....	41, 85-86
intrusions at.....	25, 26, 28, 29, 31
quartz monzonite of, analysis of.....	26
section of.....	18-19
Lone Mountain limestone, character and distribution of.....	16, 18-19
Lucky Boy mine, description of.....	
sections of, figures showing.....	76, 77, 78
Lucky Girl group, description of.....	75-79
production of.....	15
Lynn district, ore deposits in.....	45, 70, 87, 88
placers in.....	45
M.	
Map, index, of Nevada.....	10
Map, outline, showing area.....	13
Maysville mine, description of.....	121
Maysville district, intrusives at.....	30
Merrimac district, ore deposits of.....	85
Mesozoic rocks, occurrence and character of..	41
Metamorphism, occurrence and character of..	15
<i>See also</i> Contact metamorphism; Hydro-metamorphism.	
Midas district. <i>See</i> Gold Circle district.	
Midas mine.....	57
Mill Creek, description of.....	100-101
geology at.....	101-103
intrusion near.....	26, 28
ore deposits at.....	41, 42, 43, 106-110
Mineral Hill, description of.....	95
geology of.....	96
intrusives at.....	30
ore deposits at.....	41, 97-99
production of.....	95
section of, figures showing.....	96, 97
Mining districts, description of.....	48-126
Mint mine, description of.....	67

	Page.
Miocene ore deposits, character and distribution of.....	43-45
minerals in.....	43
production from.....	43
Miocene rocks, character and distribution of..	22
Modoc Peak, section near.....	19-20
Morgan claim, ore of.....	86
Morning Star mine, description of.....	122, 123-125
section of, figure showing.....	124
Mountain City, description of.....	80
geology of.....	80-81
intrusions at.....	26, 28, 30
ore deposits at.....	42, 43, 70, 81-84
production of.....	15-80
Mountain City mine, description of.....	83-84
Mountain ranges, account of.....	47-126
Mud Springs, description of.....	119
ore deposits of.....	119-120
N.	
Nannies Peak. <i>See</i> Lone Mountain.	
Navajo lode, description of.....	60-61
Nelson mine, description of.....	83
ores of.....	81, 83
Nevada, index map of.....	10
Nevada limestone, character and distribution of.....	16, 19-20
section of.....	19-20
Nevada Star mine, description of.....	87
New York claim, description of.....	110
O.	
Onondaga mine, description of.....	111-112
Ordovician rocks, occurrence and character of.....	16, 18-19
Ore deposits, account of.....	39-47
association of, with eruptive rocks.....	39
<i>See also</i> Cretaceous deposits; Miocene deposits.	
Oregon claim, description of.....	99
Owyhee Bluffs, description of.....	47
geology of.....	47
mining in. <i>See</i> Gold Circle district.	
Owyhee Desert, description of.....	47, 57
P.	
Pacific claim, ore of.....	86
Pahute Lake, history of.....	38
Panther vein, mines on.....	64-65
Phoenix mine, description of.....	117
Pinyon Range, description of.....	88-89
geology of.....	88-89
mines of. <i>See</i> Bullion; Mineral Hill; Alpha.	
section of, figure showing.....	89
Pittsburg and Palisade group, description of..	112
Pittsburg mine, description of.....	122, 123, 125-126
Placer deposits, character and distribution of..	45
Pliocene lakes, deposition in.....	35
Pliocene rocks, character and distribution of..	22-23
Pogonip limestone, character and distribution of.....	16, 18
Polaris mine, description of.....	74
Porphyries, association of, with intrusive rocks.....	29-30
Production, estimate of.....	14-15
Prospecting, suggestions for.....	46-47

	Page		Page
Prospect Mountain limestone. <i>See</i> Eldorado limestone.		Starr Grove mine, description of	126
Prospect Mountain quartzite, character and distribution of	16, 17	Stratigraphy, description of	15-17
Protection mine, description of	82	Surprise group, description of	61-62
Q.		Sweepstakes mine	95
Quartz diorite, analysis of	27	Sylvania claim, description of	95
Quartzite, ore deposits in	46, 68	T.	
veins in	42	Tenabo, description of	114-115
Quartz monzonite, analysis of	26	geology at	115-116
Quaternary deposits, occurrence and character of	23-24	intrusions at	28, 30
Queen Canyon, geology in	48	ore deposits at	116, 118
R.		Tenabo Peak, geology at	100-101
Ransome, F. L., on age of intrusions	28-29	mines at	103-106
Ravens Nest, section through, figure showing	89	section of, figure showing	102
Red Bird mine, description of	94	Tertiary lakes, history of	24, 38-39
Replacement deposits, character and distribution of	41-42	Tertiary rocks, occurrence and character of	21-23
<i>See also particular mines.</i>		ore deposits in	46
Resurrection mine, description of	82-83	<i>See also</i> Eruptive rocks.	
Rex mine, description of	54-55	Tiger lode, description of	74
plan of, figure showing	55	Topography, description of	12
Rhoda fissure, description of	107-108	Trimetal Mining Co., mines of	90, 94
Rhyolite, character and distribution of	30-31	Tripoli mine, description of	93-94
fissure veins in	44-45	Triumph mine, description of	119-120
Richmond district, ore deposits of	87	Truckee formation, character and distribution of	22, 38
Richthofen, F. von, on Tertiary eruptives	33-34	Turquoise, mine of	106
Rose mine, description of	61	Tuscarora, geology of	59-60
Ruby claim, description of	112	gold of	61-62
S.		history of	58
Safford district, description of	110-111	intrusions near	25, 28
ore deposits of	111-113	location of	58
production of	110	ore deposits at	44-45, 58, 60-62
St. Paul-Banner lode, description of	56	placers at	45, 58
Salt, occurrence of	24	production of	14-15, 89
Scope of paper	9	Tuscarora Nevada Mines Co., work of	59
Secret Canyon shale, character and distribution of	16, 18	Two Widows claim, description of	118
Sedimentary rocks, description of	15-24	U.	
Shale, deposits in	42	Ulrich, E. O., on Pogonip limestone	17
Shoshone Peak, granodiorite from, analysis of	25	Uncle Sam claim, description of	120
Shoshone Lake, history of	39	Utah claim, description of	99
Shoshone Mesa, capping of	31	V.	
Shoshone Range, description of	113-114	Valley View mine, description of	106
geology of	114	Van Duzee Creek, placers at	45, 84
glaciation in	23	Vegetation, character of	12-13
intrusions in	25, 28, 114	Veins. <i>See</i> Fissure veins.	
mines of. <i>See</i> Tenabo; Lander; Mud Springs; Grey Eagle; Hilltop; Maysville; Lewis and Dean.		Violet claim, description of	118
ore deposits in	42	W.	
Sierra Nevada, birth of	38	Walcott, C. D., section by	18
Signal Peak, section near	19-20	section by, figure showing	89
Silver, production of	15	Walcott, C. D., and Hague, A., on sedimentary rocks	17
Silver mining, rise and fall of	14, 58	Water Witch mine, description of	56-57
Silver Prize vein, description of	119	Weber conglomerate, character and distribution of	16, 20-21
Silver Side mine, description of	119	West mine, description of	112-113
Spurr, J. E., on lavas	34	White Horse mine, description of	106
Squaw Valley, description of	47	White Pine shale, character and distribution of	16, 20
Stafford, ore deposits of	44	Woodward, R. W., analyses by	25, 33
Standard vein, description of	83	Z.	
Standing Elk mine, description of	90, 93	Zenoli mine, description of	111
		production of	15
		Zinc, production of	15
		Zirkel, F., on trachytes	34