

OIL SHALE IN NORTHWESTERN COLORADO AND ADJACENT AREAS.

By DEAN E. WINCHESTER.

PREFATORY NOTE.

By MARIUS R. CAMPBELL.

For several years it has been known that some of the shale of the Green River formation in northwestern Colorado and northeastern Utah would produce oil when subjected to destructive distillation, but the yield of petroleum from the oil fields was so great that production by distillation did not seem to be feasible, despite the fact that in Scotland such an industry has long been developed and is to-day paying dividends on a large investment.

The United States Geological Survey has regarded this oil shale as a great reserve—an undeveloped resource—and one that would be developed as soon as the demand for petroleum greatly exceeded the supply. In anticipation of such an event, E. G. Woodruff and David T. Day began, in 1913, an examination of the Green River formation in Colorado and Utah and made rough field tests to determine the richness of the shale. Although these tests were not entirely satisfactory, they tended to confirm the general impression that this shale constitutes a source of oil which, sooner or later, will be called into use. Of course, no prediction could be made as to the date when this additional supply would be needed, but the Survey felt justified in continuing the geologic investigation, in order that when the time of need arrived it would have first-hand information on the richness and quantity of the shale available for distillation. Accordingly the field examination was continued during the summers of 1914 and 1915 by Dean E. Winchester, who devised a more efficient and portable apparatus for determining not only the quantity of crude oil in the shale but also the amount of gas and of ammonium sulphate (fertilizer) that might be obtained as a by-product and sold. The experiments of Mr. Winchester confirmed the results of the work done in the previous year and indicated even more strongly that a great quantity of high-class fuel is locked up in this shale.

At the present time, owing to the great increase in the consumption of gasoline and the failure to discover large new oil fields, it

seems that the day is approaching when this additional supply will be needed and that the public will demand all the information in the possession of the Geological Survey on this subject. I feel confident that this report of Mr. Winchester will supply many of the data needed to establish and develop the oil-shale industry in this country. The report contains information showing the quantity and quality of the oil that may be distilled from the richer beds of shale, the number of such beds at the different localities examined, and the general distribution of the shale throughout northwestern Colorado and adjacent parts of Utah and Wyoming. Mr. Winchester's results, which have been corroborated by tests made in the laboratory of the Bureau of Mines, show that the quantity of oil that can be derived from such shale ranges from less than 1 gallon to 90 gallons to the ton of shale. The field tests, however, were not intended to determine the best method of utilizing the shale in commercial operations, but simply to provide data for fixing approximately the value of the shale as it is found to-day. Mr. Winchester, as a result of the field tests, estimates that in Colorado alone there is enough shale to produce 20,000,000,000 barrels of oil, and it seems probable that in actual practice a greater yield than this can be obtained. He also estimates that 300,000,000 tons of ammonium sulphate could be recovered as a by-product in the manufacture of the oil.

The results given in this paper show that although the Geological Survey has done only reconnaissance work in this region, it has demonstrated that the shale of the Green River formation will yield a vast quantity of oil, gas sufficient to carry on the process of distillation, and fertilizer enough to enrich most of the farms of the Middle West, and that this reserve is ready whenever the demand is sufficient to warrant the establishment of a new industry to supplement the failing supply of petroleum from the oil fields.

INTRODUCTION.

The economic study of the oil shale of Colorado and Utah was begun in 1913 by Woodruff and Day,¹ and the results of their preliminary work showed that the shale of the Green River formation in Colorado and Utah will yield oil in commercial quantities when subjected to destructive distillation. As a result of a more detailed examination along the north, east, and southeast sides of the Uinta Basin, in northwestern Colorado, made during the summer of 1914 and in eastern Utah in 1915 the writer finds that although the thickest and richest beds of shale are exposed along the southern margin of the basin, nearly every section examined contains beds of shale more than 3 feet thick that will yield considerable oil. An examination of

¹ Woodruff, E. G., and Day, D. T., Oil shale of northwestern Colorado and northeastern Utah: U. S. Geol. Survey Bull. 581, pp. 1-21, 1914 (Bull. 581-A).

the same formation in southwestern Wyoming in 1915 indicates that in that area there are no thick beds of rich shale.

The oil shale of the United States has received very little attention, chiefly because petroleum has been abundant. Before petroleum was discovered in Pennsylvania the Mormons distilled oil from shale near Juab, Utah, where the ruins of an old still can yet be seen. Many attempts have been made to distill cannel coal, and a few experiments have been made with the Devonian black shale of the East, but no shale-oil industry has been established in America. According to Baskerville¹ there were 55 oil companies in the United States in 1860. "Many of the companies were of small capacity and most of them were not more than fairly started when the discovery of petroleum paralyzed the industry." In Scotland, however, several well-established plants mine and distill shale of Carboniferous age. In 1908, according to Ells,² the oil-shale industry of Scotland employed about 8,300 men, of whom nearly 4,000 were miners; and in the preceding year the production of oil shale in Scotland amounted to 2,775,799 (long) tons, and the average yield of crude oil was 23 (Imperial) gallons to the long ton (24.6 United States gallons to the short ton).³ The operations have paid dividends in spite of this low yield, because of the cheapness of labor, the value of the by-products, and lack of competition with petroleum. The cost of mining shale in Scotland is reported by the same author to be \$1 a ton, the cost of distilling the crude oil from the shale is 40 cents a ton, and the cost of making ammonium sulphate (the principal by-product) from the shale is 46 cents a ton. All mining in Scotland is underground, and in many of the mines the shale beds dip at angles of 30° to 60° and there are numerous faults, which greatly increase the expense of mining. At many places in Colorado and Utah, however, the rich shale has only a light overburden (see Pl. XI) and could be mined with a steam shovel.

In Colorado alone there is sufficient shale, in beds that are 3 feet or more thick and capable of yielding more oil than the average shale now mined in Scotland, to yield about 20,000,000,000 barrels of crude oil, from which 2,000,000,000 barrels of gasoline may be extracted by ordinary methods of refining, and in Utah there is probably an equal amount of shale just as rich. The same shale in Colorado, in addition to the oil, should produce with but little added cost about 300,000,000

¹ Baskerville, Charles, Economic possibilities of American oil-shales: Eng. and Min. Jour., vol. 88, pp. 149-154, 195-199, 1909.

² Ells, R. W., Report on tests made in Scotland of oil shale sent from New Brunswick in the spring of 1908, with a view of ascertaining its economic value, especially as regards the yield of crude oil and sulphate of ammonia, pp. 24, 26, Canada Dept. Mines, 1910.

³ In this paper results of distillation are given in United States gallons (42 gallons to the barrel) and referred to the short ton (2,000 pounds). Care should be taken in comparing figures with those given in reports on oil shale of Scotland, in which results are given in Imperial gallons (35 gallons to the barrel) and referred to the long ton (2,240 pounds).

tons of ammonium sulphate, a compound especially valuable as a fertilizer. The industry requires a large equipment of retorts, condensers, and oil refineries, as well as of mining machinery, so that it can not be profitably handled on a small scale.


In Scotland, according to Ells,¹ the following valuable products are manufactured from the shale: (1) permanent gas, used principally for fuel in the retorts; (2) naphtha, gasoline, and motor spirit; (3) burning or lamp oil; (4) intermediate oil used for gas making; (5) lubricating oil; (6) solid paraffin; (7) still grease; (8) still coke, which contains some oil and is used for gas fires, smokeless fuel, and carbon for electrical purposes; (9) sulphate of ammonia [a fertilizer which in the United States is worth from \$50 to \$60 a ton]; and (10) liquid fuel, used in the refineries. The distillation of the oil shale of the Green River formation will probably yield different products, and the processes of distillation used in Scotland may not be well adapted to this shale.

In the 11 field tests made by Woodruff² in 1913, 1 sample of shale yielded as low as 10.4 gallons of oil to the ton of shale; 8 between 16 and 40 gallons, averaging 27.2 gallons; one 45.2 gallons; and one 61.2 gallons. Of the 57 samples tested in 1914 (Nos. 1-57, pp. 152-153), 17 samples yielded less than 10 gallons of oil to the ton of shale; 22 samples between 10 and 20 gallons; 11 samples between 20 and 30 gallons; 3 samples between 30 and 40 gallons; 2 samples 40.6 gallons each; 1 sample 65.3 gallons; and 1 sample 86.8 gallons. Seventy-five samples (Nos. 58-133) were tested in 1915. Of those obtained in Utah (34 samples, Nos. 58-91) 6 yielded less than 10 gallons of oil to the ton of shale; 7 between 10 and 20 gallons; 7 between 20 and 30 gallons; 9 between 30 and 40 gallons; and 5 more than 40 gallons. The maximum yield, 90 gallons to the ton, was obtained from a thin bed near Watson, Utah. Samples 93-132, from Wyoming, yielded less than 30 gallons, except 4, which yielded more than 30 gallons. One of these represented a 2-foot bed, which yielded 50 gallons to the ton.

As these quantities compare favorably with those obtained from the oil shale of Scotland, it seems probable that in time the distillation of oil from the Green River shale may become as important an industry in this country as the distillation of oil from Carboniferous shale has become in Scotland, or if richer raw material can be found here in abundance it may even exceed the present shale-oil industry abroad.

Ells, R. W., *op. cit.*, pp. 54-56.

²Woodruff, E. G., and Day, D. T., *op. cit.*, p. 4.



KEY MAP OF NORTHWESTERN COLORADO AND ADJACENT AREAS.

FIELD WORK.

Personnel and methods.—In order to estimate the value of the oil shale of the Green River formation as an economic resource, the writer, assisted by H. M. Robinson and Frank A. Elliott, in 1914, made a careful study of the entire exposed portion of the formation at a number of places along the north, east, and southeast sides of the Uinta Basin in Colorado (localities A to P, Pl. XVIII). During the later part of the season Mr. Robinson made a hasty examination of the shale of the same formation at several places in northeastern Utah (localities 1-4, Pl. X). During the season of 1915 Walter B. Wilson, John N. Massey, and Yong K. Lee assisted the writer in examining the southern part of the area of the Green River formation in southwestern Wyoming and the south side of the Uinta Basin, near the eastern line of Utah. A large number of samples of shale were distilled in the field and in the laboratory of the Bureau of Mines at Washington to determine the quantity of oil and other products which could be obtained from them. Much of the chemical work was done under the direction of David T. Day, of the Bureau of Mines. C. A. Davis, also of the Bureau of Mines, was detailed to the Survey for several months to make a microscopic study of the organic matter in the shale. To Messrs. Day, Davis, and David White, who visited the field early in September, 1914, the writer is indebted for many helpful suggestions concerning methods of work and interpretation of results.

The limits of the Green River formation were mapped in the field and the exact place at which each sample of shale was obtained was determined by the use of telescopic alidade, stadia rod, and plane table. Geologic sections were measured at several localities. The accompanying map of northwestern Colorado and eastern Utah (Pl. XVIII) is compiled from field sheets prepared by the writer, from published and unpublished data collected by Gale¹ in connection with an examination of coal fields in the adjacent areas, and from data shown on the township plats of the General Land Office. The land net was compiled from these plats, but owing to discrepancies between surveys and resurveys it was necessary to make numerous more or less arbitrary adjustments and the land net shown on the map therefore comprises only township lines. The resurvey of the area north of the base line in Colorado and along Douglas Creek south of it was only recently completed, and the corners are well marked, but the corner posts of the much older surveys elsewhere are very poorly preserved or entirely gone. The land survey of that part of eastern Utah shown on Plate XVIII is good. In many places the geologic boundaries shown are taken directly from maps published

¹ Gale, H. S., Coal fields of northwestern Colorado and northeastern Utah: U. S. Geol. Survey Bull. 415, 1910.

by Mr. Gale, inasmuch as wherever tested these maps proved to be entirely satisfactory for the purposes of this work. The places from which samples 1-4 were collected are shown on the smaller map (Pl. X).

The map of southwestern Wyoming (Pl. XIX) is compiled from maps by Veatch¹ and Schultz² and unpublished data by Schultz, together with the data collected by the writer during the field work and data shown on the township plats of the General Land Office. A large part of the area has been recently resurveyed, and the land lines are trustworthy, except in the extreme eastern part, where the old survey is known to be very poor.


Area examined.—The examination of 1914 was confined to a narrow strip about the north, east, and southeast sides of the Uinta Basin in Colorado, except that during a short trip into central Utah Mr. Robinson sampled shale beds at the four localities shown on Plate X. A portion of the outcrop of the formation along the Colorado-Utah line near White River was examined, but the greater part of the season was spent along the margin of the main basin to the east. The Green River formation occupies an area of about 1,900 square miles in Colorado and a larger area in northeastern Utah. The main area in Colorado is separated from the Utah part of the Uinta Basin by the Douglas Creek anticline, which extends in a general northerly direction along the valley of Douglas Creek near the State line. The limit of the oil-yielding shale in most places practically coincides with the boundary of the Green River formation, but northwest of Meeker only the lower part of the formation is present, and this is barren of oil. A similar condition prevails south of Grand River.

The eastern edge of the Uinta Basin in Utah, near Watson, was examined in considerable detail during August and September, 1915. Most of the season, however, was spent in a reconnaissance examination of southwestern Wyoming (Pl. XIX). The eastern margin of the Green River formation in Wyoming between the Union Pacific Railroad and the Colorado-Utah-Wyoming line was examined, and samples of shale from the localities indicated on the map were tested. A hurried trip was made across the Green River basin to Fossil, Wyo., and southward along the western part of the area known to contain the Green River formation.

Northwestern Colorado and northeastern Utah.—The surface of the part of the field represented by Plate XVIII consists mainly of deeply dissected uplands surrounded on all sides by more open valleys carved in the shale and soft sandstone of the Wasatch formation,


¹ Veatch, A. C., Geography and geology of a portion of southwestern Wyoming, with special reference to coal and oil: U. S. Geol. Survey Prof. Paper 56, pl. 3, 1907.

² Schultz, A. R., The southern part of the Rock Springs coal field, Sweetwater County, Wyo.: U. S. Geol. Survey Bull. 381, pp. 214-281, 1908.




A. OIL SHALE NORTHEAST OF WATSON, UTAH.

Shows thin bedding in this part of the Green River formation. Darker bands are richest beds. About 600 feet of rock exposed.




B. OIL SHALE ON EAST SIDE OF PICEANCE CREEK NEAR WHITE RIVER, COLO.

Most of the projecting ledges are rich shale.



A. BOOK CLIFFS WEST OF RIFLE, COLO.

Oil shale forms upper cliff. About 3,000 feet of rock exposed. (See fig. 13, p. 145.)



B. GREEN RIVER FORMATION NORTH OF WHITE RIVER, 9 MILES WEST OF RANGELY, COLO.

Dark projecting beds are oil shale. Cliff about 1,000 feet high.

which lies beneath the oil-yielding Green River shale and above the coal-bearing Mesaverde formation. The Book Cliffs near Rifle, Colo. (Pl. XII, *A*), stand about 9,000 feet above sea level. The Grand Hogback, which extends along the eastern margin of the field nearly parallel with the boundary of the Green River formation, reaches an altitude of about 9,500 feet and is formed by the steeply upturned resistant sandstones of the Mesaverde formation. The Green River formation, probably owing to the presence of oil-yielding shale, resists erosion to a marked degree and gives rise along the southern margin of the area to nearly perpendicular and at most places impassable cliffs, which in the vicinity of Rifle and De Beque rise to heights of 3,000 or 4,000 feet above the valley of Grand River, only a few miles distant. (See fig. 13.) West of the Grand Hogback and north of Rio Blanco post office the shale forms hills of considerable prominence, known as the Petrolite Hills. The outcrop of




FIGURE 13.—Section from Grand River near Rulison, Colo., to the Book Cliffs on the north, showing the position of the oil shale.

the oil shale along the east side of the Douglas Creek anticline is marked by nearly perpendicular cliffs known as the Cathedral Bluffs. Cliffs similar to these and equally impassable form the boundary of the formation along White River (Pl. XII, *B*) and southward along the Colorado-Utah State line to a point near Watson, Utah. (See Pl. XI, *A*.)

By far the larger part of this area drains northward to White River through Evacuation, Douglas, Yellow, and Piceance creeks, and to the Pacific Ocean through Green and Colorado rivers. That part of the area south of the Book Cliffs is drained by Grand River, which joins Colorado River in Central Utah. White and Grand rivers and Roan, Douglas, Yellow, and Piceance creeks are the only streams that carry water throughout the year.

The valley of Grand River is traversed by the main line of the Denver & Rio Grande Railroad. The narrow-gage line of the Uintah Railway between Mack, Colo., and Watson, Utah, crosses the Book Cliffs near the Colorado-Utah line and furnishes transportation facilities for a considerable part of the Uinta Basin. The

ranchers and farmers in this area do a very large part of their shipping by way of Meeker, Rifle, and the Denver & Rio Grande Railroad. The proposed extension of the Denver & Salt Lake Railroad ("Moffat road") westward from Craig down Yampa River, to the north of the field, will furnish an additional outlet.


The surface of the area is so rough that wagon travel is forced to follow certain long-used routes, such as the Government road from Rifle to Meeker and the well-kept road between Meeker and Rangely down the valley of White River. Roads also lead from Rangely northwest and southwest to Vernal and Dragon, Utah, respectively, and there is a fair wagon road up Piceance Creek connecting with the Government road at Rio Blanco post office. Aside from these wagon roads there are few routes that can be used to advantage.

The valleys of White and Grand rivers and of Piceance and Roan creeks are occupied by small but prosperous ranches which under irrigation produce alfalfa, timothy, grain, and vegetables. Considerable fruit is raised in the valley of Grand River. The area outside of these valleys is used only for stock range.


Rifle is the largest railroad town in the area and is a shipping point for nearly the entire region. It is the starting point of the mail stage and freight lines which supply Meeker and the settlements in that vicinity. Meeker, a thriving town of 800 inhabitants on White River in T. 1 N., R. 94 W., is the distributing point for a large area to the north, east, and west. Rangely, 60 miles down White River, consists of a store and post office and is connected with Meeker by stage that makes three trips a week. Supplies for Rangely and vicinity are freighted from Dragon, Utah, a station on the Uintah Railway, and Watson, at the end of the same railway 9 miles north of Dragon, is the shipping point for ore from the gilsonite mines of the Uinta Basin and for produce from Ashley Valley, 50 miles to the northwest. A toll road from Watson to Vernal is used by daily automobile mail and passenger stages that connect with the trains of the Uintah Railway. Rio Blanco, halfway between Rifle and Meeker, consists of a small store and post office. Piceance and Sulphur are merely post offices located at ranch houses. De Beque and Grand Valley are small but prosperous towns on the Denver & Rio Grande Railroad south of the Book Cliffs.

Southwestern Wyoming.—That portion of southwestern Wyoming which is shown on the map (Pl. XIX) includes the southern part of the Green River basin and the western rim of the Southern Red Desert Basin. The surface is in most places rolling and covered with vegetation characteristic of arid regions.


Trees grow only along streams and in a few small upland areas. The region is drained through Green River and its tributaries except in the area adjacent to the west line of the State, where the run-off



A. GREEN RIVER FORMATION EAST OF GREEN RIVER CITY, WYO.
Samples of oil shale 127 and 128 taken from beds at base of upper cliff.




B. GREEN RIVER FORMATION IN T. 10 S., R. 15 E., ABOUT 25 MILES NORTH OF SUNNYSIDE, UTAH.



A. SAMPLING BED OF OIL SHALE SOUTH OF GREEN RIVER CITY, WYO.

Shows characteristic weathering of bed containing alternating rich and poor seams. The richer seams are more resistant.



B. SAMPLING BED OF OIL SHALE NEAR WATSON, UTAH.

A, Narrow channel in weathered face, from which sample 65 was taken. B, Larger cut in same bed, from which sample 66 was taken.

finds its way to Great Salt Lake by the way of Bear River. Steep cliffs mark the eastern margin of the main basin, and Green River valley is bordered by precipitous walls for several miles near the town of Green River. (See Pl. XIII, A.) Altitudes range from 5,900 feet above sea level on Green River near Linwood to 8,750 feet in the southwestern part of the area shown on the map.

Green River has its beginning far to the north, in the Wind River Mountains, but its principal tributaries in this area, Blacks Fork and Henrys Fork, rise in the Uinta Mountains of northern Utah. Aside from the three streams mentioned above, together with Hams Fork, a tributary of Blacks Fork, and Bear River, there are in this area very few streams that carry water the year round.

The main line of the Union Pacific Railroad crosses the area in a general westerly direction and is joined at Granger by the Oregon Short Line, which connects with points to the northwest. Rock Springs, Kemmerer, and Evanston, coal-mining towns, and Green River, a railroad division point, are the principal towns of the region. Several villages have been established in the irrigated district around old Fort Bridger and along Henrys Fork near the southern line of the State. Many of the towns on the railroad consist of only a few houses, a store, and a post office. The Lincoln Highway follows in general the line of the Union Pacific Railroad, and good automobile roads connect many of the smaller towns with the railroad and this highway.

TESTS OF THE SHALE.

APPARATUS FOR FIELD DISTILLATION.

A large number of samples of shale were collected (see Pl. XIV) and tested in the field in order that a definite idea of its value as a source of oil might be obtained. Several large samples were also shipped to the chemical laboratory of the Geological Survey so that different methods of treatment might be devised to increase if possible the yield of crude oil and other valuable products. In the early part of the work (1914) the apparatus designed and used by Woodruff and Day¹ during the previous season, described below, was used to make the tests.

The retort into which the shale was charged consisted of a section of 12-inch iron casing pipe 4 feet long, having flanges screwed on the ends and a removable iron plate with asbestos gaskets fitted to each end of the retort. On one side of the retort there was fitted a small steam dome, a pressure gage, and a safety valve. From the top of the dome a pipe led to a block-tin condensing coil in a small water-filled tank. The coil discharged into Wolff bottles set in series and provided with stopcocks so that the liquids could be drawn off without interfering with the operation of the condenser. During the operation the retort was suspended from iron supports in a narrow trench,

¹ Woodruff, E. G., and Day, D. T., *op. cit.*, p. 4.


covered with iron plates and earth, and a flue erected at the back. Heat was obtained from a wood fire placed under the retort.

The operation consisted of removing the head, charging the retort with shale broken into pieces not larger than 4 inches in diameter, and replacing the head. Fire was started to give a gentle heat at first and was gradually increased until the lower part of the retort became red hot; then the fire was held constant until near the close of the process, when it was increased for a short time and then allowed to subside. Water vapor, gas, oil and gas, and finally only gas was the order in which the products were obtained. From seven to eight hours' heating was required for a charge.


This apparatus necessitated the mining of a large quantity of shale (100 pounds or more) for each test. The apparatus, being large and not easily transported and eventually becoming unfit for use through leakage, was abandoned and replaced by a small still which required a sample weighing only about 1 pound and which was heated by gasoline torches. With two of these small distilling outfits four samples of shale were tested easily in one working day, whereas only one sample of shale a day could be tested with the larger apparatus. The distilling apparatus used in 1915 (see Pl. XV), which is similar to that used during the later part of the 1914 season, but much more compact and lighter, consists of the following essential parts:

- Two gasoline blast lamps (Barthel's).
- One iron mercury retort (one-half pint) with delivery tubes.
- One brass condenser.
- Two ring stands.
- One 3½-inch ring.
- One large condenser clamp.
- One receiver for condensed liquids (50 cubic centimeter glass graduate).
- One ammonia scrubber (8-ounce bottle filled with glass beads).
- Two pairs combination pliers.
- One postal balance.
- Six feet of rubber tubing.
- Glass tube for connecting condenser, receiver, and ammonia scrubber.
- One glass separatory funnel.


Because of its simplicity and because its flame can be adjusted to any desired angle or length, the Barthel blast lamp was chosen to furnish heat for the still. This lamp consists of a small spherical gasoline tank with burner, mounted on an iron base in such a way that the burner may be turned at any angle. To manipulate the burner the tank is first filled nearly full of gasoline and the cap securely screwed down. Gasoline is placed in the small cuplike depression around the burner and lighted. When this gasoline is burned out, sufficient heat will have been produced to generate gas under pressure, which may be lighted at the end of the burner on opening the burner valve. If the flame is yellow or sputters the burner is not sufficiently hot and must be reheated. The gasoline tank of each burner holds sufficient fuel to keep the blast burning about 2 hours. Inasmuch as each distillation of shale requires from



FIELD APPARATUS FOR DISTILLING OIL SHALE.



A. CHARACTERISTIC WEATHERING OF RICH, MASSIVE OIL SHALE.
Near Welch ranch, T. 1 N., R. 100 W., Colorado.



B. CHARACTERISTIC WEATHERING OF OIL-YIELDING PAPER SHALE.
West side of Piceance Creek near White River, Colo.

3 to 4 hours, two burners are used with each retort. The second torch may be most easily lighted by playing the flame of the first on the burner of the second.

The vessel in which the shale is placed to be heated is an ordinary iron mercury retort (capacity, one-half pint), which is equipped with close-fitting lid and clamp and an iron delivery tube. The delivery tube is fastened to the inner tube of the condenser by a small brass plumber's union, which provides a very easily disconnected joint. The retort is held in place above the flame of the burner by a ring and ring stand.

The condenser used in the outfit consists of an inner tube of thin brass three-eighths of an inch in diameter and 15 inches long, to one end of which is soldered a small brass plumber's union. This inner tube is surrounded by a second thin-walled brass tube $1\frac{1}{4}$ inches in diameter, 11 inches long, which is provided with a two-hole rubber stopper at each end, one hole being for the inner tube. A small brass tube 2 inches long is inserted through the second hole of each stopper to provide connection for the entrance and waste of the water which is circulated between the inner and outer tubes to keep the inner tube cool. The delivery tube from the retort is so bent that when the retort is in an upright position the condenser will be turned at an angle of about 40° from the horizontal. The condenser is held in position by a single clamp, attached to a ring stand.

The receiver for the condensable products of the distillation consists of a 50 cubic centimeter glass graduate, provided with a two-hole rubber stopper through which are thrust two glass tubes, one for the entrance of the liquids and permanent gases from the condenser and the other for the escape of the permanent gases to the ammonia scrubber. The glass tubes have a diameter of about a quarter of an inch and are bent at the proper angles to make connections with the condenser and scrubber. The tubes should barely penetrate the cork.

The ammonia scrubber consists of an ordinary glass cylinder or 8-ounce wide-mouthed bottle, provided with a glass tube reaching nearly to the bottom of the bottle for the entrance of permanent gas from the receiver. The bottle is filled with glass beads, which provide additional surface and a means of breaking up the gas into small bubbles as it passes up through a 10 per cent solution of sulphuric acid.

The water for cooling the condenser may be had from waterworks, or if no running water is at hand a tank or tub may be stationed near the condenser, at a slightly higher level, and the water conveyed over the top by a siphon entering the condenser at the lower end and wasting at the upper end.

The pliers are used for handling the retort.

The postal balance has a capacity of 4 pounds and is used to weigh the sample of shale.

The glass separatory funnel is used to separate the oil from the water derived from the shale.

In order to determine the quantity of oil and ammonia that may be derived from a sample of shale, the shale is first pulverized to pass through a screen of $\frac{1}{2}$ -inch mesh. After thorough mixing a sample weighing $8\frac{1}{2}$ ounces is so selected as to represent the entire quantity. This sample is placed in the iron retort and the cover securely fastened. In order to prevent leaks the joint between the cover and retort bowl is plastered with a thick paste made of a mixture of powdered graphite and glycerine. The delivery tube from the retort is then coupled with the inner tube of the condenser. The ammonia scrubber is filled approximately two-thirds full with a 15 per cent solution of sulphuric acid, and cool water (not ice water) is started circulating through the condenser. The blast lamp is then lighted and placed beneath the retort, with the flame turned as low as possible. After heating about 10 minutes water and oil will begin to condense and be delivered into the receiver. The permanent gas will pass into the ammonia scrubber and bubble up through the sulphuric acid, which will combine with any ammonia contained in it, producing soluble ammonium sulphate. Gentle heat should be applied to the retort as long as any oil is delivered to the receiver; then the flame of the burner may be lengthened until at the end of three or four hours the burner will be at full blast, the retort will be red hot, and the shale will cease to yield either oil or gas. The products of the distillation are then measured; the quantity of oil in the receiver is recorded, as well as the quantity of water in the same receiver. The liquid products of the distillation are then transferred to the separatory funnel and the water drawn off from the oil and added to the liquid contained in the ammonia scrubber. The material contained in the ammonia scrubber is then placed in a glass-stoppered bottle and transferred to a chemical laboratory for the determination of the quantity of ammonium sulphate. The yield of oil in United States gallons to the short ton of shale is equal to the number of cubic centimeters of oil in the receiver, provided the sample of shale used weighs $8\frac{1}{2}$ ounces. The oil obtained from the distillation should be placed in a small bottle for determination of its specific gravity, which can best be done in the laboratory. In order to compute the number of pounds of ammonium sulphate which may be derived from a short ton of shale it is necessary only to multiply the number of grams of ammonium sulphate found in the sample by 8.8.

RESULTS OF TESTS.**SALIENT FEATURES.**

The results of the tests made in the field and in the laboratory at Washington are given below and show variation from a maximum yield of 90 gallons of oil to the ton of shale to a minimum of 0.31 gallon. The yield of ammonium sulphate was not determined for samples tested during the early part of the first season's work, but those which were tested show a range from 18.3 pounds by dry distillation or 34 pounds by steam distillation to 0.4 pound to the ton of shale. Likewise the yield of inflammable gas ranges in the samples for which the amount was recorded from 4,549 cubic feet to the ton of shale to less than 500 cubic feet. Many of the samples that were subjected to destructive distillation were selected not because they were supposed to be rich in oil but in order that the field men might be better able to judge from the appearance of the shale how much oil it would yield when subjected to distillation. Several samples were selected in order to determine the geologic range of oil-yielding shale. All samples were taken near the outcrop, and it is probable that some of the more volatile constituents of the oil had been lost by evaporation, hence the results of distillation tests do not show the maximum amount of oil that may be obtained even by the method of distillation used. The stratigraphic position of nearly all the beds sampled is given in the sections (pp. 170-189).

Results of distillation of samples of oil shale collected in 1914.

Sample No.	Location.			Thickness of shale sampled.	Weight of shale used.	Crude oil obtained (cubic centimeters).		Equivalent yield of oil per short ton of shale (United States gallons).	Gravity of oil at 60° F.		Yield of gas per short ton of shale (cubic feet).	Yield of ammonium sulphate per short ton of shale (pounds).	Remarks.
	Sec.	T.	R. W.			Dry.	With steam.		Specific gravity	Degrees Baumé.			
1.	Juab.	Utah.		<i>Ft. in.</i> 2 9	6 ounces.	8.5		11.9	0.8995	25.6	1,198	2.5	Laboratory test.
2.	Mt. Pleasant.	Utah			6 ounces.	8.5		11.9	.8866	27.9	3,352	2.2	Do.
3.	Tucker.	Utah.		3 0	6 ounces.	8.5		11.9	.8965	26.2	2,525	6.5	Do.
4.	Soldiers Summit.	Utah.		6 3	6 ounces.	12.0		16.8	.8937	26.6	2,864	3.5	Do.
5.	Green River.	Wyo.	a		6 ounces.	21.0		29.4	.9130	23.3	2,978	9.0	Do.
6.	Elko.	Nev.	b		6 ounces.	62.0		86.3	.8850	28.2	3,891	6.0	Do.
7.	9	1 N.	103	2 3	85 lbs.	2,040.0		12.6+	.9255	21.6			Field test; retort leaked.
8.	9	1 N.	103	4 8	117 lbs.	946.2		4.26	.9371	19.4			Field test.
9.	9	1 N.	103	3 10	6 ounces.	8.5		11.9	.9010	25.4	2,395	7.1	Laboratory test.
10.	9	1 N.	103	3 10	104 lbs.	2,217.7		11.3	.9138	23.2			Field test.
11.	9	1 N.	103	1 11	130 lbs.	2,040.3		8.22					Field test; retort leaked.
12.	9	1 N.	103	2 3	150 lbs.		2,454.3	8.64	.9290	20.7			Field test.
13.	26	1 N.	100	5 0	6 ounces.	24.0		33.6	.8919	27.0	3,034	9.7	Laboratory test.
14.	15	2 N.	98	5	6 ounces.	9.5		13.3	.9165	22.7	3,034	12.8	Do.
15.	15	2 N.	98	1 2	1 lb.	7.4		3.0					Field test; retort leaked.
16.	15	2 N.	98	1 4	1 lb.	3.7		1.9					Field test.
17.	15	2 N.	98	6	6 ounces.	15.0		21.0	.9059	24.8	3,123	18.2	Laboratory test.
18.	16	2 N.	98	3 4	142 lbs.		6,150.6	22.88	.9290	20.7			Field test.
19.	16	2 N.	98	3 4	142 lbs.	1,685.5		6.27	.9327	20.0			Do.
20.	10	2 N.	98	3 0	6 ounces.	9.0		12.6	.9030	25.0	3,512	16.4	Laboratory test.
21.	10	2 N.	98	4 6	115 lbs.	1,685.5		26.8					Field test.
22.	15	2 N.	98	3 5 1/2	6 ounces.	11.0		15.4	.9310	20.4	3,592	9.0	Laboratory test.
23.	15	2 N.	98	2 2	150 lbs.	2,158.6		7.61	.9321	20.2			Field test.
24.	14	2 N.	98	1 6	135 lbs.		1,537.6	6.25+	.9352	19.7			Field test; retort leaked.
25.	19	2 N.	97	8 1	150 lbs.		1,360.0	4.78+	.9109	23.7			Do.
26.	19	2 N.	97	8 1	1 lb.	7.4		3.85	.9050	24.7			Field test.
27.	11	1 N.	97	18 1	6 ounces.	6.0		8.4	.8946	26.5	1,926	18.3	Laboratory test.
28.	11	1 N.	97	4 0	6 ounces.	10.0		14.0	.9090	24.0	2,075	1.0	Do.
29.	11	1 N.	97	4 0	1 lb.	22.2		12.5	.9082	24.1			Field test.
30.	11	1 N.	97	4	1.5 lbs.	25.9		9.1					Do.
31.	11	1 N.	97	17 0	1 lb.	29.6		15.5	.8880	27.6			Do.
32.	11	1 N.	97	3 0	6 ounces.	29.0		40.6	.8838	28.4	3,832	8.5	Laboratory test.
33.	11	1 N.	97	3 0	1 lb.	48.1		25.5	.8920	26.9			Field test.
34.	11	1 N.	97	5 6 1/2	1 lb.	42.5		23.0	.8884	27.6			Do.
35.	11	1 N.	97	4 2	1 lb.	27.7		14.7	.8864	27.9			Do.
36.	11	1 N.	97	3 3 1/2	8 ounces.	29.6		31.0	.8831	28.5			Do.

37	11	1 N	97	5	8 ³	1 lb.	44.4	23.25	.8877	27.7			Do.
38	14	1 S.	97	5	11	6 ounces.	18.0	25.2	.8955	26.4	4,294	6.3	Laboratory test.
39	36	1 N.	96	3	0	1 lb.	25.9	13.7	.8986	25.8		8.2	Field test.
40	36	1 N.	96	3	8	1 lb.	22.2	12.5	.8959	26.3		4.48	Do.
41	36	1 N.	96	1	4	1 lb.	25.9	13.7	.8603	32.7		4.91	Do.
42	29	1 N.	97			10 ounces	3.7	.31					Do.
43	2	3 S.	95	4	9	1 lb.	17.7	9.4	.8897	27.4		5.72	Do.
44	2	3 S.	95	13	6	1 lb.	11.8	6.2	.9025	25.1		3.25	Do.
45	32	4 S.	94	9	1	6 ounces.	4.5	6.1	.9205	22.1	639	2.8	Laboratory test.
46	16	4 S.	94	5	2	1 lb.	22.2	12.5	.9037	23.2			Field test.
47	27	4 S.	94	2	0	6 ounces.	5.0	7.0	.9030	25.4	479	1.7	Laboratory test.
48	27	4 S.	94	4	0	1 lb.	29.6	15.5	.9331	20.0		.4	Field test.
49	27	4 S.	94	3	0	6 ounces.	7.5	10.5	.8955	26.4	4,071	7.4	Laboratory test.
50	6	6 S.	94	10	0	6 ounces.	29.0	40.6	.8790	29.2	1,916	4.3	Do.
51	6	6 S.	94	4	3	6 ounces.	20.0	28.0	.9126	23.4	4,294	7.3	Do.
52	22	6 S.	95	8	8	6 ounces.	17.5	24.5	.8636	32.1	1,198	4.4	Do.
53	22	6 S.	95	8	8	6 ounces.	25.0	35.0	.8802	29.0	3,500	8.7	Do.
54	22	6 S.	95	8	0	1 lb.	39.5	20.7	.8585	33.1			Field test.
55	22	6 S.	95	13	0	6 ounces.	11.0	15.4	.9070	24.3	1,443	3.2	Laboratory test.
56	22	6 S.	95	15	0	6 ounces.	8.0	11.2	.9265	21.1	1,198	4.0	Do.
57	1	7 S.	98			6 ounces.	46.5	65.3	.9115	23.6	4,549	7.0	Do.

^a Hand specimen collected by W. T. Lee.

^b Hand specimen collected by D. T. Day.

Results of distillations of samples of oil shale in the field in 1915.

Sample No.	Location.				Thick-ness of shale sampled.	Gravity of oil at 60° F.		Yield of oil per short ton of shale (U. S. gallons).	Yield of ammonium sulphate per short ton of shale (pounds).
	Sec.	T.	R.	State.		Specific gravity.	Degrees Baumé.		
58	17	11 S.	25 E.	Utah.	<i>Ft. in.</i> 3 11	0.8989	25.74	23	5.04
59	17	11 S.	25 E.	do.	1 0	.9327	20.10	9	4.59
60	17	11 S.	25 E.	do.	2 2	.9019	25.22	12	4.38
61	17	11 S.	25 E.	do.	6 7	.9041	24.85	10	3.92
62	20	11 S.	25 E.	do.	3 11	.8983	25.85	18	5.37
63	20	11 S.	25 E.	do.	4 3	.8998	25.59	32	6.96
64	20	11 S.	25 E.	do.	6 2	.8870	27.83	15	4.09
65	20	11 S.	25 E.	do.	6 6	.9090	24.01	32	5.45
66	20	11 S.	25 E.	do.	6 6	.9052	24.66	55	9.65
67	20	11 S.	25 E.	do.	6 6	.8745	30.09	90	6.89
68	20	11 S.	25 E.	do.	6 2	.9112	23.64	31	6.99
69	20	11 S.	25 E.	do.	3 2	.9021	25.19	19	5.04
70	20	11 S.	25 E.	do.	5 9	.9260	21.18	14	4.98
71	20	11 S.	25 E.	do.	7 0	.9098	23.88	7	3.48
72	20	11 S.	25 E.	do.	6 2	.8775	29.54	7	2.25
73	20	11 S.	25 E.	do.	4 8½	.9263	21.13	6	2.61
74	20	11 S.	25 E.	do.	5 9	.8887	27.53	32	7.05
75	22	11 S.	25 E.	do.	2 4	.9036	24.93	35	5.14
76	22	11 S.	25 E.	do.	4 6	.9034	24.97	31	5.20
77	26	10 S.	25 E.	do.	6 9	.8727	30.42	37	7.81
78	24	11 S.	25 E.	do.	6 3	.8833	28.49	48	9.76
79	27	10 S.	25 E.	do.	4 1			1	2.11
80	15	10 S.	25 E.	do.	4 7	.9094	23.94	33	5.87
81	15	10 S.	25 E.	do.	1 11	.9073	24.30	24	6.72
82	15	10 S.	25 E.	do.	6 0	.9163	22.78	20	6.49
83	15	10 S.	25 E.	do.	4 0	.8975	25.98	8	5.32
84	15	10 S.	25 E.	do.	7 8	.8966	26.14	21	3.77
85	15	10 S.	25 E.	do.	1 8	.8879	27.07	22	5.39
86	15	10 S.	25 E.	do.	3 2½	.8934	26.70	37	6.52
87	15	10 S.	25 E.	do.	2 1½	.8866	27.90	54	5.51
88	15	10 S.	25 E.	do.	2 1	.8914	27.05	25	4.05
89	15	10 S.	25 E.	do.	4 0	.9059	24.54	17	5.48
90	15	10 S.	25 E.	do.	4 8	.8976	25.97	45	9.22
91	15	10 S.	25 E.	do.	5 1	.8953	26.37	29	5.35
92	9	13 N.	99 W.	Wyoming.	5 0	.8709	30.79	30	3.94
93 ^a	19	13 N.	103 W.	do.	4 0	.8760	29.81	11	4.88
94 ^a	25	13 N.	104 W.	do.	2 0	.8937	26.65	15	5.91
95	26	13 N.	108 W.	do.	5 0	.9496	17.43	13	7.49
96	26	13 N.	108 W.	do.	5 4	.9277	20.91	4	12.69
97	26	13 N.	108 W.	do.	4 10	.9062	24.49	6	4.71
98	27	13 N.	108 W.	do.	5 0			3	7.18
99	27	13 N.	108 W.	do.	3 3½			19	9.32
100	27	13 N.	108 W.	do.	5 0			3	9.52
101	13	13 N.	108 W.	do.	2 7	.8994	25.65	34	5.70
102	13	13 N.	108 W.	do.	4 0	.9060	24.52	15	8.50
103	23	14 N.	108 W.	do.	2 11	.8818	28.77	32	6.62
104	23	14 N.	108 W.	do.	11	.8892	27.44	20	4.39
105	9	14 N.	108 W.	do.	4 2	.8885	27.56	7	2.47
106	11	14 N.	108 W.	do.	7 2	.9183	22.45	9	11.14
107	36	16 N.	108 W.	do.	5 3	.9022	25.18	21	5.69
108	36	16 N.	108 W.	do.	6 0	.8925	26.86	13	5.06
109	17	17 N.	106 W.	do.	5 9	.8798	29.12	11	5.51
110	17	17 N.	106 W.	do.	6 3	.9190	22.34	19	9.82
111	17	17 N.	106 W.	do.	5 4	.9111	23.66	19	8.81
112	17	17 N.	106 W.	do.	5 1	.9075	24.26	9	7.59
113	17	17 N.	106 W.	do.	5 3	.9050	24.69	10	5.10
114	17	17 N.	106 W.	do.	4 6	.9143	23.12	11	3.86
115	17	17 N.	106 W.	do.	4 6	.8848	28.22	9	2.28
116	17	17 N.	106 W.	do.	4 10			4	3.02
117	27	16 N.	106 W.	do.	8 1	.9003	25.50	19	8.68
118	27	16 N.	106 W.	do.	5 6	.9120	23.50	14	5.50
119	19	17 N.	106 W.	do.	5 0	.8963	26.19	12	7.17
120	19	17 N.	106 W.	do.	5 0	.8702	30.88	14	7.93
121	16	17 N.	106 W.	do.	7 4	.9456	18.05	14	11.19
122	9	17 N.	106 W.	do.	7 3	.9077	24.23	14	4.27
123	9	17 N.	106 W.	do.	6 6	.9197	22.22	19	2.74
124	8	18 N.	107 W.	do.	5 9	.9027	25.09	13	5.80
125	8	18 N.	107 W.	do.	3 3½	.8800	29.09	6	3.73
126	8	18 N.	107 W.	do.	1 8	.9182	22.47	29	11.71
127	24	18 N.	107 W.	do.	6 3	.9148	23.03	18	7.27
128	24	18 N.	107 W.	do.	2 4	.8862	27.97	7	5.65
129	5	21 N.	107 W.	do.	1 7	.8705	30.82	8	.69
130	5	21 N.	107 W.	do.	5 0	.8837	28.42	10	.86
131	5	21 N.	107 W.	do.	2 0	.8889	27.49	50	1.99
132			(b)	do.	2 4	.8449	35.70	50	4.50

^a Collected by A. R. Schultz from fissile shale described as Wasatch in U. S. Geol. Survey Bull. 381, p. 222, 1910.
^b Collected by David White from a point near Elko, Nev.

OIL.

Most oils obtained from the distillation of shale are reddish brown, and at ordinary temperatures range from semisolid vaseline-like products to a thin liquid.

Samples 7, 8, 10, 11, 12, 18, 19, 21, 23, 24, and 25 were distilled in the large field retort, and the resulting oils have a specific gravity ranging from 0.9109 (23.7° Baumé) to 0.9371 (19.4° Baumé). The oils obtained from samples distilled in the small apparatus range in specific gravity from 0.8449 (35.7° Baumé) to 0.9496 (17.4° Baumé), though by far the larger number of samples show a specific gravity of less than 0.90 (25.0° Baumé). The difference in the oils obtained by using the large and small apparatus is probably due to conditions of heating rather than to differences in original material, but the oil from all samples distilled in a particular apparatus may be assumed to have been obtained under conditions approximately similar. About one-fourth of the distillations were made in the Washington laboratory of the Bureau of Mines, with an apparatus similar to the small outfit used in the field. In order to compare the tests made in the laboratory with those made in the field with the small apparatus, shale from two samples tested in the field was tested in the laboratory, and the results were found to check very closely.

It was suggested at the beginning of the oil-shale investigations that distillation of shale with the injection of steam into the heated retort might result in a product of increased value, but before 1914 no sample had been treated in such a way that a comparison of the value of dry and steam distillation could be made. The writer therefore during the first field season selected a sample of shale and, after careful mixing, divided the sample into two equal parts (142 pounds each). One part was distilled dry and the other part distilled in the same apparatus on the following day with the addition of steam to the heated mass. From the dry-distilled sample (No. 19) oil was obtained at the rate of 6.27 gallons to the ton of shale; the steam-distilled sample (No. 18) yielded 22.88 gallons of oil to the ton. It is not likely that the larger yield of oil obtained from the second sample could be entirely due to the effect of the steam, and it is probable that to other factors, such as leakage, may in part be due the difference in quantity of oil obtained under the two methods of distillation. That the oil produced by steam distillation is not radically different from that produced by dry distillation is shown by the similar proportion of products which the oils yield on fractionation. (See table, p. 156.) It is unfortunate that in this test, the only one made on a large sample, there was no equipment available to ascertain the effect of steam distillation on the yield of ammonium sulphate. Later, six samples of oil shale were distilled with steam at the Washington laboratory of the Bureau of Mines.

For these samples the quantity of oil derived by steam distillation exceeded that by dry distillation only slightly or in some samples not at all. At the same time the oil produced by steam distillation has a uniformly heavier gravity. The detailed results of these tests are given on page 160.

Oils from nine samples treated in the field and in the laboratory were redistilled with the following results:

Results of fractionation of shale oil.

Sample No.	4	6	10	18	19	27	32	51	57
Begins to boil at °C.	80	52	50	70	72	65	80	70	54
Distillation (cubic centimeters):									
To 100° C.	6	10	2	2.5	2	4	2.5	7	4
100 to 125° C.	2	.5	3.5	1	1	1.5	3.5	1	2
125 to 150° C.	2	1.5	6.5	2.5	8	1.5	4.5	1	3
Total gasoline.	10	12	12	6	11	7	10.5	9	9
150 to 175° C.	2	2	8	6	6.5	3	6	2	5
175 to 200° C.	2	4	6	5	5	3	5	4	4.5
200 to 225° C.	5	4	7.5	5	5	4	6	4	5
225 to 250° C.	7.5	6	7	5	5	5	8	6	5
250 to 275° C.	6	6	7.5	7	7	7	8.5	6.5	7
275 to 300° C.	6	10	13	7	7	17	9	13	12
Total kerosene.	28.5	32	49	35	35.5	39	42.5	35.5	38.5
Total distillate.	38.5	44	61	41	46.5	46	53	44.5	47.5
Total residuum.	61.5	56	39	59	53.5	54	47	51.5	52.5
Amount of oil used.	100	100	100	100	100	100	100	96	100
Specific gravity at 60° F.:									
Crude.	0.8937	0.8850	0.9138	0.9290	0.9327	.08946	0.8838	0.9126	0.9126
Gasoline.7947	.7769	.8090	.7974	.8202	.7849	.7568	.7838	.7605
Kerosene.8602	.8466	.8260	.8742	.8876	.8722	.8524	.8682	.8538
Residuum.9695	.9643	.9884	.9894	1.0160	.9684	.9368	.9695	.9628
Asphalt, per cent by weight.	1.35	.82	2.82	4.10	3.62	2.49	.47	1.40	1.03
Paraffin.	7.70	6.93	2.22	3.72	1.63	4.56	4.70	9.21	4.00
Sulphur.54	1.06				.73	1.42	.41	.69
Nitrogen.	1.848	.887	2.198	1.549	1.643	1.267	1.849	1.820	2.135
Unsaturated hydrocarbons:									
Crude. per cent.			82	86	81.6		72		
Kerosene. do.	55	55	64	71	71	61	57	62	58

A study of the table reveals a fairly uniform quantity of products from the different samples, the gasoline ranging from 6 to 12 per cent, the kerosene from 28.5 to 49 per cent, the paraffin from 1.63 to 7.70 per cent, and the sulphur from 0.41 to 1.42 per cent. The samples for fractionation were chosen to illustrate both range in physical character of the shale and wide geographic distribution. Sample 4 was from massive shale obtained near Soldier Summit, Utah (for location see Pl. X); sample 6, from brown shale near Elko, Nev.; sample 10, from massive shale in sec. 9, T. 1 N., R. 103 W., Colo. (see Pl. XVIII); samples 18 and 19, from massive shale in sec. 16, T. 2 N., R. 98 W., Colo. (No. 18, steam distilled; No. 19, same sample, dry distilled); sample 27, from papery shale in sec. 11, T. 1 N., R. 97 W., Colo.; sample 32, from massive shale in sec. 11, T. 1 N., R. 97 W., Colo.; sample 51, from massive shale in sec. 6, T. 6 S., R.

94 W., Colo.; sample 57, from massive black shale in sec. 1, T. 7 S., R. 98 W., Colo.

The samples, except No. 27, were obtained within a few inches of the outcrop and probably were slightly weathered, although physical evidence of weathering in the type of shale sampled extends only a very short distance back from the outcrop. As the papery shale weathers to considerable distance back from the outcrop a drift 18 feet long was driven for the purpose of obtaining a sample (No. 27), but the zone of weathering had not been passed at the point of sampling.

In order to determine the behavior of shale oil when subjected to the Rittman process¹ of refinement, two samples were tested at the Pittsburgh laboratory of the Bureau of Mines. The only samples of shale oil available for these tests were two obtained by Woodruff² in 1913 from field distillations of shale described below. The samples had remained sealed since the date of original distillation and were therefore probably not greatly altered. Each sample was fractionated in the ordinary way with the following results:

Fractionation of shale oil by ordinary method.

Distillation—	Sample A: Specific gravity, 0.882; boiling point, 32° C.		Sample B: Specific gravity, 0.925; boiling point, 25° C.	
	Percentage by weight.	Specific gravity.	Percentage by weight.	Specific gravity.
To 75° C.	3.4	0.691	2.5	0.700
75° to 100° C.	2.4	.738	1.6
100° to 125° C.	4.8	.754	3.6	.772
125° to 150° C.	5.6	.775	5.5	.792
150° to 175° C.	6.4	.795	5.8	.814
175° to 200° C.	6.6	.821	6.7	.842
200° to 225° C.	6.8	.844	6.1	.862
225° to 250° C.	7.2	.866	6.4	.884
250° to 275° C.	9.4	.892	7.2	.913
275° to 300° C.	8.7	.901	9.0	.929
Residuum	38.5	Solid.	43.0	Solid.

The residuum over 175° C. of each sample (A and B) was then divided into two parts (A and A', B and B') and run in a Rittman furnace with 150 pounds pressure, different temperature being used for each part, as indicated below.

A. Subsurface sample from north side of White River, 5 miles east of White River stage station, T. 9 S., R. 25 E., Utah. Bed, 3 feet 6 inches thick. Yield, 33.3 gallons of oil to the short ton of shale.

B. Sample obtained 4 miles north of Ninemile Creek, approximately in sec. 12, T. 11 S., R. 16 E., Utah. Bed, 6 inches thick. Yield, 39 gallons of oil to the short ton of shale.

¹ Rittman, W. F., Dutton, C. B., and Dean, E. W., Manufacture of gasoline and benzene-toluene from petroleum and other hydrocarbons: Bur. Mines Bull. 114, 1916.

² Woodruff, E. G., and Day, D. T., op. cit. pp. 4, 19, 20.

Rittman furnace tests on residuum obtained from distillations of shale oil at temperature over 175° C. Pressure 150 pounds (gauge reading).

	Sample A.	Sample A'.	Sample B.	Sample B'.
Temperature used (°C.).....	525	550	525	600
Specific gravity of residuum.....	0.920	0.920	0.957	0.957
Specific gravity of recovered oil.....	.901	.902	.929	.959
Percentage recovery.....	79	79	82	70

The oil recovered from the treatment in the Rittman furnace was then distilled with the following results:

Distillation of oil recovered from Rittman tests of shale oil.

	Sample A.	Sample A'.	Sample B.	Sample B'.
Specific gravity of recovered oil.....	0.901	0.902	0.929	0.959
Distillation (per cent by weight)—				
To 75°.....		4.2		3.8
To 100°.....	2.2	a 6.2	2.8	a 1.8
100° to 125°.....	1.1	8.5	1.1	2.5
125° to 150°.....	1.4	10.7	1.6	2.7
150° to 175°.....	2.7	13.8	3.0	4.2
175° to 200°.....	7.0	20.0	6.3	6.9

^a Distillation, 75° to 100° C.

An examination of the above tables shows that by ordinary methods of fractionation Samples A and B will yield 22.6 and 19.0 per cent, respectively, of distillate up to 175° C. Treatment of the residuum above 175° C. in the Rittman furnace with 150 pounds pressure and temperatures of 525° to 600° C. produces an oil which will yield additional fractions below 175° C. according to the temperature used in the furnace. Tests of two samples of shale oil do not furnish adequate data for generalizations but merely indicate that under proper treatment shale oil may be made to yield a much larger percentage of gasoline than that shown in the tables on pages 156 and 157.

AMMONIUM SULPHATE.

Inasmuch as one of the valuable products derived from the distillation of shale in Scotland is ammonium sulphate, it is interesting to compare the amount of nitrogen in the shale with the amount that is accounted for in the products of distillation. The following table is based on the assumption that all the nitrogen in the shale can be converted to ammonium sulphate:

Relation between total amount of nitrogen in shale and amount accounted for in products of distillation.

Sample No.	Nitrogen in shale.	Theoretical equivalent of nitrogen in ammonium sulphate (per ton of shale).	Yield of oil per ton of shale.	Nitrogen in oil.	Equivalent ammonium sulphate (per ton of shale).	Ammonium sulphate extracted from gas (per ton of shale).	Total nitrogen accounted for in ammonium sulphate (per ton of shale).	Total nitrogen unaccounted for in ammonium sulphate (per ton of shale).
	<i>Per cent.</i>	<i>Pounds.</i>	<i>Gallons.</i>	<i>Per cent.</i>	<i>Pounds.</i>	<i>Pounds.</i>	<i>Pounds.</i>	<i>Pounds.</i>
4.....	0.39	36.6	16.8	1.848	10.9	3.5	14.4	22.2
6.....	.39	36.6	86.8	.887	26.7	6.0	32.7	3.9
27.....	.46	43.2	8.4	1.267	3.7	18.3	22.0	21.2
32.....	.54	50.8	40.6	1.849	26.1	8.5	34.6	16.2
51.....	.46	43.2	28.0	1.820	18.2	7.3	25.5	17.7
57.....	1.22	115.0	65.3	2.135	50.0	7.0	57.0	58.0
63.....	.53	49.9	32.0	(a)	(a)	7.0	(a)	b 42.9
64.....	.35	33.0	15.0	(a)	(a)	4.1	(a)	b 28.9
65.....	.73	68.8	32.0	(a)	(a)	5.4	(a)	b 63.4
66.....	.80	75.4	55.0	(a)	(a)	(a)	(a)	(a)
67.....	1.30	122.5	90.0	(a)	(a)	6.9	(a)	b 115.6
68.....	.70	66.0	31.0	(a)	(a)	7.0	(a)	b 59.0
69.....	.32	30.1	19.0	(a)	(a)	5.0	(a)	b 25.1
74.....	.43	40.5	32.0	(a)	(a)	7.1	(a)	b 33.4
77.....	.68	64.1	37.0	(a)	(a)	7.8	(a)	b 56.3
92.....	.60	56.7	30.0	(a)	(a)	3.9	(a)	b 52.8
101.....	.68	64.1	34.0	(a)	(a)	5.7	(a)	b 58.4
129.....	.10	9.4	8.0	(a)	(a)	.7	(a)	b 8.7
131.....	.50	47.1	50.0	(a)	(a)	2.0	(a)	b 45.1
132.....	.85	80.1	50.0	(a)	(a)	4.5	(a)	b 75.6

^a Not determined.

^b Nitrogen unaccounted for is much greater in these samples because the amount in oil was not determined.

It will be noticed that in no test has the entire amount of nitrogen present in the shale been accounted for in the products of distillation, and except for one sample (No. 6) there is a larger amount of nitrogen unaccounted for than the amount which was extracted from the fixed gas in the form of ammonium sulphate. In the samples tested the amount of ammonium sulphate extracted from the permanent gas averages 11.1 per cent of the theoretical total amount of ammonium sulphate available from the shale, and an average of 41 per cent of the theoretical total available ammonium sulphate is accounted for by the nitrogen in the shale oil itself. Therefore the theoretically available ammonium sulphate unaccounted for averages more than twice as much as is extracted from the gas produced in the distillation of the shale. In the shale works of Scotland distillation is carried on with the injection of steam, which results, according to Stuart,¹ in practically doubling the amount of ammonium sulphate that is extracted from the permanent gas by the dry method. The table given above indicates that there is ample nitrogen in the shale to allow the extraction of double the amount of ammonium sulphate obtained in the dry distillation, provided the right process is used.

¹ Stuart, D. R., The oil shale of the Lothians, 2d ed., p. 144, Scotland Geol. Survey Mem., 1912.

During May, 1916, six samples of oil shale were tested at the Bureau of Mines, with an apparatus similar to that used in the field, but so arranged that superheated steam was injected into the retort during the entire process of distillation. The samples were selected to represent a wide geographical distribution, as well as differences in richness and physical character, and the results of the tests are extremely interesting. Each of the samples had been tested previously in the field apparatus without steam, and the results therefore furnished factors that may be used to convert the results of field tests into what are very probably close approximations to results to be expected from commercial practice.

Comparison of steam and dry distillation.

Sample No.	Oil.				Ammonium sulphate.		
	With steam.		Without steam.		Theoretical yield, equivalent of nitrogen in shale (pounds per ton).	Yield as determined.	
	Yield (gallons per ton).	Specific gravity.	Yield (gallons per ton).	Specific gravity.		With steam (pounds per ton).	Without steam (pounds per ton).
4.....	23.0	0.9346 (19.7° B.)	16.8	0.8937 (26.6° B.)	36.6	13.4	3.5
27.....	10.0	.9135 (23.2° B.)	8.4	.8946 (26.5° B.)	43.2	29.9	18.3
32.....	44.0	.9630 (15.3° B.)	40.6	.8838 (28.4° B.)	50.8	34.0	8.5
51.....	39.0	.9234 (21.6° B.)	28.0	.9126 (23.4° B.)	43.2	15.8	7.3
66.....	55.0	.9286 (20.7° B.)	55.0	.9052 (24.6° B.)	75.4	23.1	9.6
132.....	50.0	.9109 (23.7° B.)	50.0	.8449 (35.7° B.)	80.1	8.4	4.5

The average amount of ammonium sulphate produced from the shale by steam distillation was about two and one-half times the amount obtained from the same samples by dry distillation, thus providing a factor for the conversion of the figure for ammonium sulphate by dry distillation to ammonium sulphate which may be obtained with steam distillation (the method practiced in the oil-shale industry of Scotland and France).

In the six samples tested an average of 37.8 per cent of the nitrogen in the shale was accounted for in the ammonium sulphate obtained by steam distillation, compared with an average of 15.7 per cent recovered by dry distillation. During the two seasons the yield of ammonium sulphate was determined for 57 of the samples that yielded more than 15 gallons of oil to the ton of shale. In these samples an average of 6.7 pounds of ammonium sulphate to the ton was obtained. This multiplied by 2.5, the factor mentioned above, gives an average of 16.7 pounds of ammonium sulphate to the ton, which seems to be a fair estimate of the quantity that may be produced in commercial practice from shale of the area examined in 1914 and 1915.

CHEMICAL ANALYSES OF OIL SHALE.

Shale samples 4, 6, 27, 32, 51, and 57 were analyzed by methods used in making ultimate analyses of coal, with the following results:

Analyses of samples of shale from the Green River formation in Colorado, Utah, and Nevada.

[Made at the Washington laboratory of the Bureau of Mines; J. D. Davis, chemist in charge.]

Sample No.	Location.			Form of analysis. ^a	Proximate.				Ultimate.					Heating value (British thermal units).
	Sec.	T.	R.		Moisture.	Volatile matter.	Fixed carbon.	Ash.	Sulphur.	Hydrogen.	Carbon.	Nitrogen.	Oxygen.	
4.....	(b)			A	1.05	33.55	(c)	65.43	0.27	1.80	13.37	0.39	18.74	2,266
				C	33.91	(c)	66.12	.28	1.70	13.51	.39	18.00	2,290
6.....	(d)			A	1.05	45.04	(c)	45.73	1.07	5.19	36.76	.39	10.86	7,714
				C	45.52	(c)	46.21	1.08	5.13	37.15	.39	10.04	7,796
27.....	11	1 N.	97 W.	A	3.18	19.55	(c)	79.00	1.08	1.75	8.34	.46	9.37	1,157
				C	20.19	(c)	81.59	1.12	1.44	8.61	.48	6.76	1,195
32.....	11	1 N.	97 W.	A	.45	37.90	(c)	62.65	.55	2.76	22.48	.54	11.02	4,012
				C	35.07	(c)	62.93	.55	2.72	22.58	.55	10.67	4,030
51.....	6	6 S.	94 W.	A	.43	39.85	(c)	59.95	.30	2.24	18.87	.46	18.18	3,055
				C	40.02	(c)	60.21	.30	2.20	18.95	.46	17.88	3,068
57.....	1	7 S.	98 W.	A	.85	51.60	(c)	46.23	.95	4.32	36.40	1.22	10.88	6,976
				C	52.04	(c)	46.63	.96	4.26	36.71	1.23	10.21	7,086

^a Analysis A represents the composition of the sample as it comes from the ground. Analysis C represents the theoretical condition of the shale after all the moisture has been eliminated.

^b Soldier Summit, Utah.

^c The conditions of heating in the volatile matter determination are different from those in the ash determination, and owing to different reactions the quantity of inorganic residue is not the same in both. As a result, the value of the fixed carbon is for some examples negative.

^d Elko, Nev.

In addition to the ultimate analyses shown above, the percentages of nitrogen and sulphur in 14 samples of shale tested for oil in 1915 were determined, with the following results:

Partial analyses of samples of shale from Utah and Wyoming.

[Made at the Washington laboratory of the Bureau of Mines; J. D. Davis, chemist in charge.]

Sample No.	Location.				Nitrogen.	Sulphur.
	Sec.	T.	R.	State.		
63.....	20	11 S.....	25 E.....	Utah.....	<i>Per cent.</i> 0.53	<i>Per cent.</i> 0.58
64.....	20	11 S.....	25 E.....	do.....	.35	.31
65.....	20	11 S.....	25 E.....	do.....	.73	.70
66.....	20	11 S.....	25 E.....	do.....	.80	.87
67.....	20	11 S.....	25 E.....	do.....	1.30	1.59
68.....	20	11 S.....	25 E.....	do.....	.70	.58
69.....	20	11 S.....	25 E.....	do.....	.32	.81
74.....	20	11 S.....	25 E.....	do.....	.43	.42
77.....	26	10 S.....	25 E.....	do.....	.68	.56
92.....	9	13 N.....	99 W.....	Wyoming...	.60	.71
101.....	13	13 N.....	108 W.....	do.....	.68	1.66
129.....	5	21 N.....	107 W.....	do.....	.10	.77
131.....	5	21 N.....	107 W.....	do.....	.50	2.78
132.....	(a)				.85	5.15

^a Elko, Nev.

GEOLOGY.

THE SHALE.

The Green River formation, of early Tertiary (Eocene) age, consists predominantly of shale. It exhibits on the weathered outcrop a more or less white color, but closer examination reveals an alternation of gray, bluish gray, and white bands. (See Pls. XI and XIII.) The shale that yields the most oil when subjected to distillation is that which weathers into massive benches of grayish-blue color but which is dark brown to black on a freshly broken surface. After this tough rich shale, which appears to be without bedding planes or laminations, is heated and the oil driven off it crumbles easily and exhibits true shale structure. Where thin benches of rich shale are interbedded with lean or barren shale, the former, being resistant, weathers to projecting ledges. (See Pl. XIV, *A*.) Some of the very rich beds show a vitreous luster similar to that of coal. The massive shale (Pl. XVI, *A*) is exceedingly tough, resists erosion to a remarkable degree, and as it weathers to a bluish-white surface and will burn when ignited the ranchers of some parts of the region refer to it as "white rock that will burn." When freshly broken, the shale gives off an odor of petroleum. All gradations exist between this hard, tough, massive rock and the papery shale (Pl. XVI, *B*), which weathers to curly forms. The papery shale is in a few places black, but usually light brown, and the thin plates of weathered shale are remarkably flexible, a characteristic which distinguishes it from ordinary carbonaceous shale. Weathering affects the papery shale to a distance of more than 20 feet back from the outcrop, but, except along joint planes, the hard, massive shale shows little evidence of weathering for more than a quarter of an inch from the exposed surface. Oil has been distilled from the papery shale as well as from the hard, massive variety, although in smaller quantity.

The hard, rich shale that crops out as projecting ledges and weathers to a gray or grayish-blue color is dark brown to black on the unweathered surface, and in all probability weathering does not affect the oil-yielding capacity of the shale to any considerable depths. Most of the samples for distillation were taken after chipping away the part of the bed that had been changed from dark to light color, and for most of the samples this required the removal of only an inch or two of weathered shale. In order to determine the difference between the shale just back of this obviously weathered zone and that a foot or so deeper samples 65 and 66 were taken from a single bed near Watson, Utah. Plate XIV, *B*, shows the relative position of the materials for the two samples, as well as the manner in which the fresh shale breaks when mined. Sample 66 was obtained after blasting away an average of about 18 inches of outcrop shale; sample

65 was taken within an inch or two of the weather-exposed face. The yield of oil for sample 65 was at the rate of 32 gallons to the short ton, but sample 66 gave nearly twice as much, or 55 gallons of oil to the ton.

Inasmuch as the samples of shale for the tests listed on pages 152-154 were obtained near the outcrop, like sample 65, it is safe to assume that the results given in this paper do not represent the maximum yield that may be obtained from the same shale when it is mined.

Although oil may be obtained from the shale by destructive distillation, it does not appear that more than a small percentage exists in the shale as oil; at least oil has not been found in commercial quantity in two wells that have been drilled into the shale in northwestern Colorado. The "gas well" in sec. 22, T. 1 N., R. 97 W., is about 400 feet deep, and develops a strong flow of artesian water, together with considerable gas, but no oil. This well does not reach the richest shale beds. Another well, drilled to a depth of 1,345 feet in sec. 30, T. 2 S., R. 94 W., apparently passes through the Green River formation and penetrates the underlying Wasatch formation, but develops only a show of oil, and that apparently near the contact between the two formations. A spring in sec. 14, T. 1 N., R. 97 W., east of Piceance Creek, issues from between two rich beds of shale. The water is of excellent quality and does not show any evidence of oil or gas.

The following description of the microscopic structure of the shale was furnished by Charles A. Davis, who at the time of his death was engaged in making a careful study of the rocks with special attention to their fossil content:

The shale as seen under the microscope varies as greatly in appearance and structure as it does when examined with the unaided eye, according to the locality and bed from which the material examined originally came. The darker specimens yield sections showing characteristic opacity, density, and compactness of structure and fineness of grain. The lighter-colored samples show less opacity and appear more granular, even when casually examined. Specimens from the same sample also present quite different appearances, according to the direction in which the section is cut.

Under low powers of the compound microscope sections cut transverse to the bedding planes show a more or less distinct lamination or banded structure, the dark-brown to light-yellow bands alternating irregularly with colorless or light-grayish ones. Usually the dark bands of the section are longer and more continuous than the light bands.

Under higher magnifications these bands resolve themselves into a series of more or less discontinuous thin laminæ, of which the brown and yellow or dark ones are certainly longer and more continuous than the gray or colorless ones. Careful inspection shows that the dark laminæ are usually not easily resolvable into granules, while the light ones are obviously made up chiefly of particles of crystals of mineral matter; much of the mineral matter is very finely divided, although some sand is present. Moreover, most of the laminæ containing mineral matter are lens-shaped, and in the samples in which they were carefully examined many of them were found to

be from two to eight or ten times as long as they are thick, and the darker laminae in the same sections are characteristically many times longer than they are thick.

In addition to these general characteristics shown in the cross sections the samples exhibit the following features:

1. Well-marked openings, in places of polygonal or even square outline, but more commonly of irregular lens shape or the shape of a flattened sphere. The larger of such openings occur usually in the lenses of mineral matter that is in the form of coarse particles. Some of these openings still contain bits of crystalline minerals, and others contain fine silty residues. It is thus evident that the openings themselves may have been formed by the breaking up of the original filling, which corresponds to that of the shorter, lens-shaped laminae, during the cutting of the section.

2. Irregularly distributed black opaque bodies, which glisten by reflected light. These fill lenslike openings that seem identical with those just described, as well as other openings, including the irregularly polygonal matrices of minerals. The material forming the opaque bodies only partly fills some of the openings in which it occurs. Such partly filled spaces show by their irregular filling and by the relation of the opaque matter to included minerals that they have been filled by local intrusion from the surrounding walls, as the material has not yet been observed running from one cavity to another, and unfilled cavities are numerous near and between those that are filled.

3. Very dark brown or black lines, which seem to represent the edges of vegetable structures such as fibers, filaments, or films that are of different composition or are more fully carbonized than the general mass of material with which they were laid down. These dark lines are in places interlaminated with light-colored organic matter and clearly can not be considered as intrusions into the laminated material after it was laid down, as the whole series of laminae is in places very finely wrinkled and each individual layer is of extreme thinness, many being less than a micron thick. As these very thin light-colored laminae are entirely conformable with the thinnest dark ones, they must have been laid down with them. The thicker dark laminae, being more opaque than the light ones, can not be so readily resolved for measurement, but where they have been carefully studied they show minute secondary lamination as well as the inclusion of thin lines of inorganic sediment, a structure which clearly proves their sedimentary origin and entirely undisturbed condition.

The wide range of variation in the characteristics of the structure, bedding, and grain of the different samples of shale examined makes it impossible to discuss all the variations which have been found, but the points brought out are the most striking ones noted in examining the cross sections.

A study of sections made parallel with the bedding planes of the oil-yielding shale shows also a large number of variations, even in the same fragment. The dark samples in general give the most interesting sections, the lighter-colored ones showing chiefly finely divided amorphous particles of mineral matter, mixed with crystalline particles, which are possibly of secondary origin, and with scattered fragments which from their color and structure are easily recognizable as of organic origin—that is, derived from the remains of plants or animals.

The examination of a large number of horizontal sections shows that the laminae are not generally flat or perfectly horizontal, but rather occur in undulating bands, apparently formed while the material was still in the process of deposition. Some horizontal sections of darker samples show only slightly granular or nearly homogeneous texture, with no inclusions. Other sections show a large number of minute plants and plant remains that are surprisingly well preserved. In general, these plant remains are embedded in the same sort of material of which the seemingly structureless sections are composed. Some of this structureless material can be resolved by careful manipulation to show a granular structure, and under such conditions the shadowy outlines of plant remains can be seen in some sections.

The inference is clear that the structureless material probably originated in a collection of plant débris which has by decomposition and the activities of bacteria and other microscopic organisms passed into a jelly-like phase such as is to be found in certain kinds of modern peat deposits. The plant remains that have been found in the shale from every locality which has furnished samples are those of microscopic algae mixed in smaller percentage with pollen and similar parts of higher plants. Animal remains are rare in the material studied, and those noted were chiefly the remains of insects in a very fragmental state.

It seems apparent, therefore, that the study of the microscopic structure of this shale as seen in vertical and horizontal sections leads to the conclusion that the material was laid down originally in water and that it passed through a series of stages of decomposition before consolidation and lithification had taken place. The remarkably well preserved state of the delicate plant structures which have been examined indicates very slight disturbances of the original material and an almost entire lack of changes produced by the action of the usual metamorphosing agencies since lithification.

STRATIGRAPHY.

The oil-yielding shale is confined almost entirely to the middle part of the Green River formation, and during the present examination little attention was paid to rocks of other formations. In northwestern Colorado the Green River formation is the youngest present, but north of White River and only a few miles west of the Colorado-Utah line the Bridger formation rests unconformably on the Green River as well as older formations, and along the northern edge of the Uinta Basin in Utah the Bridger obscures the entire outcrop of the oil shale. The Bridger formation also occupies the central part of the Green River basin in southwestern Wyoming and west of Burnt Fork and south of Carter overlaps the Wasatch formation, covering the outcrop of the Green River.

The Green River formation is underlain by the Wasatch and this in turn by the Mesaverde, which in Colorado and Utah is coal bearing. The Wasatch in Wyoming contains valuable beds of coal, which may be needed for fuel when the shale-oil industry is developed.

The Green River formation has a maximum thickness near the mouth of Piceance Creek of about 2,600 feet and may be separated there on the basis of the presence or absence of oil-yielding shale into three fairly distinct parts. The upper and lower parts of the formation are practically barren, but the middle part of the formation contains, at every locality examined, beds of shale that will yield considerable oil. A single test was made of shale from the upper division of the formation, which yielded 0.31 gallon of oil to the ton of shale (sample 42, p. 153). The section measured near the mouth of Piceance Creek (H, pp. 177-178) shows the oil-yielding part of the formation to be 1,550 feet thick, whereas the lower barren part is only 342 feet thick and the upper barren part 716 feet thick. According to measurements made near Morris station (O, p. 182) on the Book Cliffs, the upper 595 feet of the section there exposed is

oil yielding, but the underlying part, 1,487 feet thick, includes no beds that will yield any appreciable quantity of oil. The great thickness of lower barren beds near Morris corresponds very closely with the thickness of the lower part of the formation as described by Woodruff¹ in a section measured along the Mount Logan trail in sec. 26, T. 7 S., R. 97 W., only a few miles to the west and on the same general cliff. In general the lower member of the Green River formation is extremely variable both in thickness and character. Along Evacuation Creek, near Dragon, Utah (sec. Q, pp. 183-184), this member includes about 600 feet of coarse sandstone, oolite, and shale with no persistent bed and some very remarkable lenticular beds. Only a few miles away, in Hells Hole Canyon, northwest of Watson (sec. T, pp. 185-186), the lower part of the Green River formation includes largely shale with comparatively thin sandstone beds and only a little oolite.

In contrast to this extreme irregularity in the lower part of the formation the thin beds of the oil-yielding portion are remarkably persistent. Sections Q, S, and T, representing the strata exposed at three places in eastern Utah, separated from one another by 5 and 7 miles (see Pl. XVIII), show three thin beds of sandstone which are remarkably regular in thickness, and at the same time the interval between them varies only slightly from place to place.

The line between the Wasatch and Green River formations is very difficult to follow because of lack of exposures and very hard to identify accurately from place to place. There seems to be a general gradation from the upper part of the Wasatch formation into the lower part of the Green River, and it is possible that the correlations shown on Plate XVII are not absolutely correct.

The Green River formation consists principally of shale but contains, especially in its lower part, beds of sandstone, many of which are ripple marked. Most sections show one or more beds of oolite and some conglomerate or conglomeratic sandstone. Near the old Black Dragon mine, Utah, however, the lower part of the formation, according to measurements made by Woodruff,² contains oolite and sandstone equal to more than half of the exposed thickness of beds (529 feet). There is at the base of the upper part of the formation on Yellow and Piceance creeks a bed of massive brown sandstone which may be equivalent to the Tower sandstone of Powell³ in southwestern Wyoming. As is shown by the sections given below, there are in nearly every section many beds of shale that will yield at least 15 gallons of crude oil to the ton, but the correlation of beds from one measured section to another, although the sections may be

¹ Woodruff, E. G., and Day, D. T., *op. cit.*, p. 14.

² *Idem*, pp. 16, 17.

³ Powell, J. W., *Geology of the eastern portion of the Uinta Mountains*, pp. 40-45, U. S. Geol. and Geog. Survey Terr., 2d div., 1876.

only a few miles apart, is very uncertain. A careful study of the strata exposed in a continuous cliff face a mile or more in extent shows that although the formation appears to be remarkably regular in thickness, individual beds vary greatly from place to place and that a single massive bed 5 feet thick at one place may change to comparatively thin-bedded shale within less than half a mile. Study of any single bed at several places along its outcrop to determine its variability in thickness, bedding, mode of weathering, and value as a source of oil was made impossible by lack of exposures except near the mouth of Piceance Creek in Colorado and along the west side of Green River in southwestern Wyoming. In the Colorado locality a zone of oil-producing shale was examined, carefully measured, and sampled at three localities within a distance of approximately 1,100 feet along its outcrop with the following results:

Sections of oil-shale zone along the west side of Piceance Creek, Colo., in sec. 11, T. 1 N., R. 97 W.

Location 34.¹

	Ft.	in.		Ft.	in.
Shale, hard, black.....	1	2	Shale, brown, lean ²	2	
Shale, light brown.....	3		Shale, hard, dark.....	3	
Shale, dark brown.....	2		Shale, brown.....		$\frac{1}{2}$
Shale, light brown.....	1		Shale, hard, dark.....	1	
Shale, hard, black.....	7		Shale, lean.....		$4\frac{1}{2}$
Shale, light brown.....		$\frac{1}{2}$	Shale.....		1
Shale, hard, black, in beds 2 inches thick.....	6		Shale, hard, black.....	1	2
Shale, brown.....		$2\frac{1}{2}$			
Shale, hard, dark.....	2			5	4

Location 35, 100 feet N. 7° E. of location 34.

	Ft.	in.		Ft.	in.
Shale, dark brown.....	2		Shale, hard, brown.....	2	
Shale, hard, dark brown.....	2		Shale, rich.....	3	
Shale, hard, black.....		$1\frac{1}{2}$	Shale, hard, brown.....		$1\frac{1}{2}$
Shale, sandy, lean.....	1	4	Shale, brown, lean.....	3	
Shale, brown, rich.....	5				
Shale, hard, brown.....	3			4	2
Shale, brown, rich.....	11				

Location 36, about 1,000 feet N. 10° E. of location 35.

	Ft.	in.		Ft.	in.
Shale, hard, dark.....	1		Shale, hard, black.....		$8\frac{1}{2}$
Shale, brown, thin bedded.....	1		Shale, brown.....		$\frac{1}{2}$
Shale, hard, dark.....		$4\frac{1}{2}$	Shale, hard, black.....		$4\frac{1}{2}$
Shale, lean.....		$\frac{1}{2}$	Shale.....		1
Shale, hard, black.....		$5\frac{1}{2}$	Shale, hard, black.....		3
Shale, brown.....		$\frac{1}{4}$	Shale, brown, lean.....		4
Shale, hard, black.....	4				
Shale, brown.....		$1\frac{1}{2}$		3	$3\frac{1}{4}$

¹ Location numbers correspond to those used on the maps. Stratigraphic sections are arranged so that the youngest beds are described first and successively older beds follow.

² In this paper the term "lean" is applied to shale that will yield less than 15 gallons of oil to the short ton and "rich" to shale that will yield more than 15 gallons.

Samples from these localities when subjected to distillation gave the following results:

Results of distillation of samples from three localities on a single bed of shale on the west side of Piceance Creek, sec. 11, T. 1 N., R. 97 W.

Location No.	Total thickness sampled.	Yield of oil per ton of shale.	Gravity of oil.
	<i>Ft. in.</i>	<i>Gallons.</i>	
34.....	5 4	23.0	0.888 (27.6° B.)
35.....	4 2	14.7	.887 (27.9° B.)
36.....	3 3½	31.0	.883 (28.5° B.)

This zone contains shale which on weathering resembles somewhat closely a massive bed but which may be subdivided upon close examination into a number of very thin units differing from one another only in minor particulars. The gravity of the oil derived from these samples is fairly uniform, but the quantity differs widely. It is possible that part of this difference in yield may be due to changes produced by weathering, although if such were the case it would seem that the gravity of the oil in sample 35 would show a corresponding increase. However, the data at hand are not sufficient to make generalizations.

Along both sides of Green River, Wyo., in Tps. 13 and 14 N., R. 108 W., a single bed or zone of rich oil-yielding shale is exposed in almost continuous outcrop for several miles. The bed is made up of alternating thin benches of rich and lean shale which weather into a most characteristic form, so that the bed can be easily identified from place to place. The richer benches weather to grayish-blue ledges which project beyond the softer lean shale, as shown in Plates XIII, B, and XIV, A. In places slabs 3 or 4 feet square and only an inch thick have weathered out and lie scattered over the surface. The four sections given below illustrate the variability of the different parts of the bed. The results of the field distillations of samples taken at three of the places are given in a separate table following.

Sections of shale bed along Green River, Wyo.

Location 99, sec. 27, T. 13 N., R. 108 W.

Shale, hard, black, rich.....	Ft. in.	Shale, soft, brown, lean.....	Ft. in.
	½		1 8
Shale, soft, brown, lean.....	5	Shale, hard, black, rich.....	3
Shale, hard, black, rich.....	1		
Shale, soft, brown, lean.....	8		3 3½
Shale, hard, black, rich.....	2		

Sections of shale bed along Green River, Wyo.—Continued.

Location 101, sec. 13, T. 13 N., R. 108 W.

	Ft.	in.		Ft.	in.
Shale, hard, black, rich.....	5		Shale, thin bedded, brown, lean...		4
Shale, thin bedded, brown, lean...	2		Shale, hard, black, rich.....		5
Shale, hard, black, rich.....	10				
Shale, thin bedded, brown, lean...	3				2
Shale, hard, black, rich.....	2				7

Location 103, sec. 23, T. 14 N., R. 108 W.

	Ft.	in.		Ft.	in.
Shale, thin bedded, brown, lean...	5		Shale, hard, black, rich.....	1	5
Shale, hard, black, rich.....	1	1			
Sandstone.....	1				3
					0

Location 105, sec. 9, T. 14 N., R. 108 W.

	Ft.	in.
Shale, brownish, black, thin bedded, lean; not sampled.....	2	4

Results of distillation of samples of shale from three localities along the outcrop of a single bed in Tps. 13 and 14 N., R. 108 W., Wyoming.

Location No.	Total thickness sampled.	Yield of oil per short ton of shale.	Gravity of oil.	Yield of ammonium sulphate per short ton of shale.
	Ft. in.	Gallons.		Pounds.
99.....	3 3½	19	Not determined.	9.3
100.....	2 7	34	0.8994 (25.65° B.).	5.7
103.....	3 0	32	.8818 (28.77° B.).	6.6

In many places massive beds of dark, tough, rich shale contain lenses of coarse sand that show no free oil. In other places small masses (some of them mere films between beds) of solid black hydrocarbon are found in the shale. Hydrocarbon occurring in this way in a small gulch east of Piceance Creek near its mouth possesses all the properties of elaterite, but in most places the material is similar to gilsonite. In sec. 14, T. 1 N., R. 97 W., this elaterite may be seen at a number of places between two beds of rather rich shale. In some places, such as Hay Gulch, in sec. 36, T. 1 N., R. 96 W., there are pockets of black material which have the shape of partly compressed stems but which show no woody structure, as would be expected if they were carbonized wood. The material contained in these pockets is not soluble in ether, chloroform, gasoline, or turpentine, the ordinary solvents of hydrocarbons.

Fossil remains, except those of microscopic size, are scarce but include leaves, fresh-water shells, insects, and fish remains. None of the fossils collected were found to be especially diagnostic, although several were identifiable. The shells were collected near the base of

the formation, as were most of the fish and leaf remains, but excellently preserved remains of insects (Diptera larvæ, etc.) were found at a number of horizons in the oil-yielding portion of the formation.

The following stratigraphic sections were measured at places indicated on the map (Pl. XVIII), and illustrate the character of the rocks exposed in different parts of the field. The beds of shale that are known by testing or are estimated to yield 15 gallons of oil or more to the ton of shale are indicated by heavy type in the sections.

Sections in northwestern Colorado.

Location A, T. 2 N., R. 104 W.

	Ft. in.		Ft. in.
Shale, gray, with a few hard sandstone beds each a few inches thick.....	350 0	Shale, sandy.....	24 0
Sandstone, white.....	1 0	Sandstone, ripple marked; ripples 4 inches from crest to crest and three-quarters of an inch deep.....	3 0
Shale, gray.....	60 0	Shale, sandy.....	10 0
Sandstone, clayey.....	15 0	Sandstone, friable, with about 10 per cent of shale; sandstone is oolitic....	3 0
Shale, thin bedded, with a few thin beds of sandstone.....	15 0	Sandstone, friable, with about 33 per cent of shale.....	54 0
Shale, brown to black, contains thin beds of rich oil shale.....	2 0	Sandstone, thin bedded.....	15 0
Shale, thin bedded, slightly carbonaceous, but is supposed to yield very little oil.....	30 0	Shale, gray.....	25 0
Shale, brown, thin bedded, will probably yield some oil.....	1 0	Sandstone, shaly.....	1 0
Shale, brown, will probably yield some oil.....	2 0	Shale, sandy.....	15 0
Shale, gray, thin bedded.....	18 0	Sandstone, coarse grained.....	3 0
Shale, thin bedded, brown, contains thin laminae of oil shale.....	10 0	Shale.....	75 0
Shale, sandy.....	40 0	Oolite with grains as large as one-sixteenth of an inch; this stratum is a massive ledge maker, the most resistant rock of the formation.....	5 0
Shale, dark brown, rich in oil.....	4	Shale, sandy.....	62 0
Shale, light gray, sandy.....	7 0	Sandstone, thin bedded.....	5 0
Shale, will yield a little oil.....	1 0	Shale, gray, mostly covered.....	114 0
Shale, gray.....	2 0	Sandstone, thin bedded, cross bedded, and slightly conglomeratic; the grains are mostly silica and well rounded....	10 0
Shale, dark brown, rich in oil.....	4	Shale, gray.....	50 0
Shale, slightly sandy.....	7 0	Sandstone, conglomeratic; largest pebbles observed have a maximum diameter of one-half inch.....	3
Sandstone and shale; sandstone shows ripple marks as much as 6 inches from crest to crest.....	15 0	Shale, gray.....	26 0
Sandstone, coarse, containing concretions.....	3 0	Sandstone, shaly.....	3 0
Sandstone, clayey.....	5 0	Shale, gray.....	20 0
Shale, sandy.....	50 0	Surface covered, supposed to be mostly shale; tan-colored shale exposed at base.	280 0
Sandstone and shale, about 60 per cent sandstone; sandstone for the most part ripple marked; one thin bed of carbonaceous shale.....	90 0	Sandstone, white, lenticular.....	3 0
Sandstone, conglomeratic at the base; most of the pebbles are flat; some are 4 inches across.....	2 0	Shale, drab, contains some sand.....	204 0
Sandstone, thin bedded, not resistant....	15 0	Sandstone, tan-colored.....	6
		Shale, sandy.....	120 0
		Sandstone, tan-colored, definitely Watsch; bottom of section.	1,872 5

Sections in northwestern Colorado—Continued.

Location B, T. 1 N., R. 103 W.

	Ft.	in.		Ft.	in.
Shale, evenly thin bedded, with very little sandstone.....	250±		Shale, sandy.....	7	0
Shale, dark, thin bedded, estimated that at least 50 per cent is oil-bearing shale (sample 10 from bed 3 feet 10 inches thick near top; 11.3 gallons).....	50	0	Sandstone, even bedded.....	2	0
Shale, sandy, thin bedded, lean.....	11	0	Shale, grading into sandstone at the top.....	7	0
Sandstone.....	4		Shale, dark brown, with disseminated iron sulphide, rich.....	1	0
Shale, thin bedded, sandy in places, bituminous in others, will yield some oil.....	7	0	Shale, sandy at base, thin bedded at top.....	9	0
Shale, dark brown, weathers bluish gray, rich.....	7		Sandstone, thin bedded.....	3	0
Sandstone, shaly.....	5	0	Shale, sandy.....	9	0
Sandstone, friable, weathers to round forms.....	1	0	Shale, thinly laminated, dark brown on fresh surface; contains beds of rich shale.....	7	0
Shale, hard, dark brown, (sample rich.....) } 8; 4:26	1	0	Shale, sandy.....	2	0
Shale, light brown..... } gallons). (3	8	Shale, thinly laminated, dark brown on fresh exposure; probably will yield some oil.....	8	0
Shale, alternating beds of rich oil-bearing shale (estimated 10 per cent) and lean shale.....	7	0	Shale, dark brown, thin bedded, rich.....	1	0
Shale, clayey, containing thin beds of rich oil shale.....	4	0	Shale, sandy.....	13	0
Shale, thin bedded.....	6	0	Sandstone, thin bedded, ripple marked.....	2	0
Shale and sandstone, containing some oil-bearing layers; the entire member is colored red by burning.....	45	0	Shale, sandy; in places will yield oil.....	4	0
Shale, sandy; contains some thin sandstones.....	5	0	Shale, dark brown; weathers bluish gray; rich.....	2	
Shale, sandy.....	10	0	Shale, sandy.....	7	0
Shale, dark brown, hard; weathers bluish gray; rich.....	1	0	Sandstone, thin bedded, with some shale.....	7	0
Shale, clayey, thin bedded; contains a few thin layers of rich shale.....	19	0	Shale, lean.....	18	0
Shale, sandy.....	3	0	Sandstone, cross-bedded at top.....	3	0
Shale, thin bedded, clayey.....	8		Sandstone and shale.....	25	0
Shale, sandy; contains some bituminous matter.....	1	0	Sandstone, in beds having a maximum thickness of 4 inches.....	5	0
Shale, with thin beds of rich oil-bearing shale; estimated that 50 per cent is rich rock.....	4	0	Shale, with some sandstone; shale is dark and carbonaceous; probably will yield some oil.....	33	0
Shale, sandy.....	11	0	Conglomerate; maximum size of pebbles half an inch.....	8	
Shale, dark brown; weathers bluish gray; rich.....	2	6	Shale, drab.....	25	0
Shale, sandy, thin bedded, calcareous.....	4	0	Shale, brown; slight oily odor.....	4	
Shale, carbonaceous; contains beds of rich shale as thick as three-quarters of an inch (sample 12; 8.64 gallons).....	2	3	Shale, dark brown; weathers blue.....	1½	
Sandstone, ripple marked at top, thin bedded.....	7	0	Shale, light brown; weathers platy..... (sample 7; 12.6 gallons).	5	
Shale.....	3	1	Shale, dark brown, hard; weathers blue.....	2	
Shale, lean.....	1	11	Shale, soft, brown; weathers into fine laminae and curls on surface.....	1	0
Shale, dark brown; contains disseminated iron sulphide (sample 11; 8.22 gallons).....	1	11	Shale.....	50	0
Shale, thin bedded, brown.....	9		Oolite.....	4	
Shale, sandy.....	6	0	Sandstone and shale.....	25	0
Shale, dark brown, massive, rich.....	4	0	Oolite.....	6	
Shale, gray, lean; contains some bituminous matter.....	4	0	Shale and sandstone in layers as thick as 6 inches.....	40	0
Sandstone, thin bedded.....	5	0	Shale, finely laminated; gives slight oily odor when broken.....	8	0
			Sandstone, calcareous.....	6	
			Shale; lower part drab; upper part weathers curly.....	13	0
			Limestone.....	2	
			Shale, mostly drab, partly carbonaceous, finely laminated.....	33	0
			Shale, thin bedded, rich; gives slight oily odor when freshly broken.....	5	0

Sections in northwestern Colorado—Continued.

Location B, T. 1 N., R. 103 W.—Continued.

	Ft.	in.		Ft.	in.
Sandstone, containing clay balls.....		2	Shale, thick bedded, rich; gives oily		
Shale, drab.....	25	0	odor when broken.....	3	0
Shale, sandy.....	3	0	Shale.....	12	0
Sandstone, containing pebbles as large as			Sandstone, coarse grained.....		2
half an inch in diameter.....		2	Shale.....		10
Shale, sandy.....	1	6	Sandstone.....		5
Shale, dark brown, rich; gives oily			Shale, thinly laminated, dark brown...	3	0
odor when broken.....		4	Talus slope at bottom.		
Shale, thick bedded, rich.....	2	10			
Shale, finely laminated, brown, carbon-					
aceous.....	2	6			
				929	1½

Location C, on north side of White River, T. 1 N., R. 104 W.

	Ft.	in.		Ft.	in.
Shale, sandy; weathers to round forms...	50	0	Shale.....	5	0
Shale, light brown, lean.....		6	Shale, dark brown; weathers bluish gray;		
Interval, probably sandy shale.....	20	0	rich.....	1	0
Shale, dark brown; weathers bluish gray;			Shale.....	9	0
rich.....		6	Shale, sandy; about 33 per cent rich, dark		
Shale.....	7	0	brown.....	13	0
Shale, dark brown; weathers bluish gray;			Shale, sandy.....	3	0
rich.....	1	0	Shale, dark brown; weathers bluish		
Shale.....	3	0	gray; rich.....	2	0
Shale, dark brown; weathers bluish gray;			Shale, dark brown; weathers bluish		
rich.....		6	gray; rich; and sandy shale.....	1	0
Sandstone, thin bedded.....	15	0	Shale, sandy.....	3	0
Shale, dark brown; weathers bluish			Shale, dark brown; weathers bluish gray;		
gray; 75 per cent rich shale and 20			rich.....	1	0
per cent lean shale.....	8	0	Shale, drab, with thin layers of rich dark-		
Shale, dark brown; weathers bluish			brown shale.....	3	0
gray; rich.....		8	Shale, dark brown; weathers bluish gray;		
Shale, brown, thin bedded.....	1	4	rich.....		6
Shale, dark brown; weathers bluish			Interval, probably mostly shale.....	13	0
gray; 75 per cent rich shale and 25			Shale, dark brown; weathers bluish gray;		
per cent lean shale.....	5	0	rich.....	1	0
Shale, dark brown; weathers bluish			Shale, sandy.....	5	0
gray; about 10 per cent lean shale,			Sandstone.....	3	0
remainder rich shale.....	5	0	Shale.....	7	0
Shale, dark brown; weathers bluish gray;			Shale, dark brown; weathers bluish gray;		
about 40 per cent rich shale.....	35	0	rich.....	1	
Shale, dark brown; weathers bluish gray;			Shale.....	4	0
about 10 per cent lean shale, remainder			Shale, dark brown; weathers bluish gray;		
rich shale.....	1	0	rich.....	1	2
Shale.....	3	0	Shale.....	35	0
Shale, dark brown; weathers bluish gray;			Sandstone, shaly.....	4	0
about half rich shale and half lean shale.	3	0	Shale, light gray, sandy.....	4	0
Shale, sandy.....	4	0	Shale; black; contains thin beds of rich		
Shale, dark brown; weathers bluish gray;			dark-brown shale.....	1	0
rich.....		6	Shale, drab, with some thin beds of rich		
Sandstone, massive.....	1	6	dark-brown shale near top.....	20	0
Shale, dark brown; weathers bluish			Shale, dark brown; weathers bluish		
gray; rich and lean shale inter-			gray; rich; interbedded with lean		
bedded.....	5	0	shale.....	3	0
Shale, sandy.....	23	0	Shale, thin bedded; 25 per cent rich dark-		
Sandstone, thin bedded.....	2	0	brown shale.....	6	0
Shale.....	20	0	Shale, dark brown; weathers bluish		
Shale, dark brown; weathers bluish gray;			gray; rich.....	2	2
rich; contains an abundance of iron			Shale, with thin beds of rich dark		
pyrite.....	1	0	brown shale.....	6	
Shale; about 25 per cent rich, dark brown	2	0	Shale, dark brown; weathers bluish		
Shale, dark brown; weathers bluish gray;			gray; rich.....	1	0
rich.....	2	0	Sandstone, shaly, grading to shale in the		
			lower part.....	8	0

*Sections in northwestern Colorado—Continued.***Location C, on north side of White River, T. 1 N., R. 104 W.—Continued.**

	Ft.	in.		Ft.	in.
Shale, dark brown; weathers bluish gray; rich		1	Oolite, with grains as large as one-eighth inch in diameter.....		5
Shale, drab.....	27	0	Shale, with some rich dark-brown shale..	5	0
Sandstone, shaly.....	5	0	Oolite.....		1
Shale, sandy.....	23	0	Shale, sandy, with some sandstone.....	13	0
Shale, drab.....	20	0	Oolite.....		1
Shale, thin bedded, black on fresh surface, rich	4	0	Shale, upper part tan-colored, lower part gray.....	100	0
Shale, sandy.....	5	0	Sandstone, light colored, with shaly layers and some rich dark-brown shale....	4	0
Shale; weathers curly; will yield some oil.....		6	Shale, light colored at bottom, dark at top; some rich dark-brown shale.....	27	0
Shale.....	18	0	Sandstone, including clay balls; a single fossil gastropod was found in this bed..		4
Sandstone, shaly.....	5	0	Shale, drab.....		2
Shale.....	8	0	Sandstone, at some places oolitic and at others slightly conglomeratic.....		8
Sandstone, brown.....	2		Shale, dark.....		20±
Shale.....	20	0	River.....		
Shale, dark brown; weathers bluish gray; rich	1	0			
Shale, gray, sandy at base, thin bedded at top.....	75	0			
				765	3.

Location D, T. 1 N., R. 100 W.

	Ft.	in.		Ft.	in.
Shale, drab; contains sandy layers.....	40	0	Shale, with a few streaks of rich dark-brown shale.....	5	0
Shale, dark brown, rich.....	6		Shale, drab.....	30	0
Sandstone.....	6	0	Sandstone, thin bedded.....	2	0
Shale, dark brown, rich	1	0	Shale and thin sandstone; contains beds as thick as 2 feet which probably will yield some oil.....	45	0
Sandstone, thin bedded.....	5	0	Shale, thin bedded, lean.....	10	0
Shale, drab.....	20	0	Shale, drab, slightly sandy.....	15	0
Sandstone, thin bedded.....	3	0	Shale, thin bedded, lean.....	2	0
Shale, thin bedded, lean.....	15	0	Shale.....	30	0
Sandstone; contains much crystalline quartz; weathers like an oolite.....	4		Shale, thin bedded, slightly carbonaceous; in places contains thin sandy beds.....	25	0
Shale, drab, with sandy layers.....	30	0	Covered.....	65	0
Shale, dark brown, rich	3	0			
Shale, drab.....	35	0			
Shale, lean.....	6				
Shale, sandy.....	4	0			
Shale; about 30 per cent rich shale in thin layers.....	4	0			
Shale, drab.....	3	0			
				399	4

Location E, T. 1 N., Rs. 99 and 100 W.

	Ft.	in.		Ft.	in.
Sandstone, yellow, slightly friable; contains concretions of pyrite.....	15	0	Shale; estimated 15 per cent rich oil shale.	20	0
bedded.....	20	0	Shale, gray on outcrop; top sandy, in beds 2 inches thick.....	20	0
Shale; upper part slightly sandy; contains some mica; lower part not well exposed.....	120	0	Shale, thin bedded; probably 15 per cent rich oil shale.....	10	0
Shale, dark gray to brown; contains a few beds of bituminous shale about half an inch thick.....	65	0	Shale, thin bedded, papery; estimated 75 per cent rich oil shale	7	0
Shale, gray, calcareous, in beds about 2 inches thick.....	8	0	Shale, sandy; contains few carbonaceous layers.....	5	0
Shale, gray to brown, slightly bituminous.....	20	0	Shale, thin bedded, estimated 50 per cent rich oil shale.....	1	6
Shale, dark brown, rich	8		Shale; contains little rich bituminous shale, but the whole is brown on fresh surface and probably will yield oil.....	1	0
Shale, thin bedded, gray on weathered outcrop but contains thin beds of brown and black rich bituminous shale; estimated that 10 per cent of the rock will yield oil.....	18	0	Shale, sandy, thin bedded.....	4	0
Shale, dark brown, thin bedded, 75 per cent rich and 25 per cent lean	1	0	Shale, dark brown, rich	1	
			Shale, thin bedded, lean.....	6	
			Shale, thin bedded, dark brown, rich	2	0
			Shale, sandy.....	2	0

Sections in northwestern Colorado—Continued.

Location E, T. 1 N., Rs. 99 and 100 W.—Continued.

	Ft.	in.		Ft.	in.
Shale, for the most part thin bedded, gray on weathered outcrop; probably 20 per cent of the whole is shale which will yield oil.....	13	0	Shale , thin bedded (estimated yield, 15 gallons).....	1	3
Shale , dark brown, massive; resists weathering; rich.....	2	0	Shale, drab.....	7	0
Shale , massive (estimated yield, 20 gallons).....	4	0	Sandstone, shaly.....	2	0
Shale, light gray, slightly sandy and bituminous.....	45	0	Shale , dark brown (estimated yield, 25 gallons).....	2	6
Shale , dark brown, rich.....	1	0	Shale, friable, yellow, lenticular.....	1	0
Shale, thin bedded, lean.....	18	0	Shale, drab.....	3	0
Shale (estimated yield, 20 gallons).....	6	0	Shale, very thin bedded (probable yield, less than 15 gallons).....	2	0
Shale, gray on weathered surface, thin bedded, lean.....	43	0	Shale, sandy.....	3	0
Sandstone, with numerous dark specks..	2	0	Shale, thin bedded, with a few bituminous layers.....	10	0
Shale, sandy.....	7	0	Shale , dark brown, rich, thin bedded (estimated yield, 35 gallons).....	3	0
Shale , dark brown; about 67 per cent rich and 33 per cent lean.....	1	0	Sandstone, contains quartz grains cemented with iron oxide.....	4	4
Shale, drab.....	1	0	Shale, thin bedded.....	4	4
Shale , dark brown; appears to be very rich (sample 13; 33.6 gallons)..	5	0	Shale, drab; contains sandy beds.....	5	0
Shale, about 50 per cent oil shale.....	2	0	Shale, gray, sandy.....	10	0
Shale, sandy.....	33	0	Shale , thin bedded, with some rich oil shale (estimated yield, 30 gallons).....	5	0
Shale , dark brown, rich.....	1	3	Shale, sandy.....	8	0
Shale.....	2	6	Shale, thin bedded; will yield some oil..	2	6
Shale, thin bedded, slightly carbonaceous, but will probably yield some oil.	6	0	Shale, sandy.....	7	0
Shale, gray, thin bedded.....	15	0	Shale, thin bedded, lean.....	5	0
Shale , dark brown, rich.....	6	0	Shale, drab; some layers 1 foot thick; will yield oil.....	10	0
Shale.....	8	0	Shale , dark brown, rich.....	4	0
Shale , dark brown, rich.....	6	0	Shale, thin bedded.....	4	0
Shale, thin bedded.....	4	0	Shale , brown (estimated yield, 25 gallons).....	3	0
Shale , dark brown, rich.....	2	0	Shale, about 50 per cent bituminous.....	2	0
Shale, thin bedded; contains possibly about 10 per cent rich shale.....	8	0	Shale, thin bedded, bituminous.....	7	0
Shale , dark brown, rich.....	1	0	Shale, thin bedded; contains rich bituminous layers; also sandstone lenses and lean shale.....	15	0
Shale.....	6	0	Sandstone, thin bedded.....	5	0
Sandstone, fine grained, massive.....	4	0	Shale , thin bedded (estimated yield, 15 gallons).....	3	0
Shale.....	8	0	Sandstone, light-colored.....	2	0
Shale , dark brown, rich.....	2	0	Shale , dark brown; part rich and part lean.....	1	0
Shale, gray, thin bedded.....	7	0	Interval largely covered, but probably shale.....	40	0
Sandstone, shaly.....	2	0		874	9
Shale, sandy.....	7	0			
Shale , dark brown, rich.....	8	0			
Shale; contains some sandy beds.....	40	0			
Sandstone, yellow, shaly.....	15	0			
Shale and sandstone in about equal quantities.....	50	0			
Shale , dark brown, rich.....	6	0			
Sandstone, shaly.....	8	0			
Shale , dark brown, rich, with some sandstone lenses.....	1	0			
Sandstone, shaly.....	8	0			
Shale , very thin bedded (estimated yield, 25 gallons).....	1	6			
Shale , 25 per cent rich.....	1	0			
Shale, dark brown, interbedded with papery shale.....	6	0			
Shale, drab.....	3	0			
Shale, lean.....	3	0			
Shale, brown, thin bedded.....	6	0			

Between the last bed mentioned above and the base of the Green River formation there is a distance of about 300 feet in which the rocks are largely concealed by surface material. It is probable, however, that there is considerable oil shale in this interval, but that it has been burned so as to lose its ordinary characteristics. The burning is indicated by loose fragments of red burned rock on the surface, by slaglike masses of fused material, and by black burnt-out shale exposed near the base of the formation.

Sections in northwestern Colorado—Continued.

Location F, T. 2 N., R. 98 W.

	Ft.	in.		Ft.	in.
Sandstone (about 75 per cent) and slightly bituminous shale.....	150	0	Sandstone, weathers rusty tan.....	9	0
Shale, gray, sandy; some layers brown on fresh surfaces.....	45	0	Shale (80 per cent) and sandstone; some beds of shale 1 foot thick may yield as much as 15 gallons of oil to the ton....	40	0
Shale; about 20 per cent rich bituminous shale (samples 18, 19, 22, 23, and 24; 6.25 to 22.88 gallons)...	5	0	Shale (estimated yield, less than 10 gallons).....	1	0
Shale, thin bedded, slightly bituminous, and sandstone.....	150	0	Oolite.....	4	
Sandstone, minutely cross-bedded, massive.....	1	0	Shale, slightly bituminous.....	1	1
Shale, drab; contains a few thin beds of sandstone.....	70	0	Oolite.....	2	
Sandstone, massive; contains a few thin layers of conglomerate.....	2	4	Sandstone, thin bedded.....	5	0
Shale, in part slightly bituminous, with a few thin beds of sandstone.....	90	0	Oolite.....	2	
Sandstone, in part chertlike.....	6		Shale, slightly bituminous.....	1	1
Shale, thin bedded.....	28	0	Oolite.....	3	
Shale, dark brown, rich, and thin-bedded lean shale (estimated yield, 25 gallons).....	5	0	Sandstone, thin bedded, with some bituminous shale.....	58	0
Shale, sandy, lenticular.....	3	0	Shale, brown, lean, interbedded with thin sandstone.....	50	0
Shale, thin bedded (estimated yield, 15 gallons).....	4	0	Shale, brown to black, thin bedded, slightly bituminous.....	50	0
Sandstone.....	6		Shale, brown, very thin bedded; weathers curly (estimated yield, 15 gallons)....	1	0
Shale, dark brown, rich (sample { Sandstone, cherty..... } 21; 26.8	11		Sandstone, shaly.....	15	0
Shale, dark brown, rich (gallons). {	3	7	Shale, brown, thin bedded; weathers curly (estimated yield, 15 gallons)....	1	0
Sandstone and shale.....	1	0	Sandstone, shaly.....	8	0
Conglomerate with pebbles half an inch in diameter.....	2		Shale, drab to gray, interbedded with thin beds of sandstone.....	90	0
Sandstone and shale, with layers of rich oil shale 1 inch thick.....	8	0	Shale, brown to black, thin bedded (estimated yield, less than 15 gallons)...	25	0
Shale and some sandstone.....	8	0	Shale, drab.....	10	0
Sandstone.....	1	0	Shale, brown on fresh surface, thin bedded, slightly bituminous.....	8	
Shale, gray, thin bedded.....	34	0	Shale, tan-colored; weathers white; many of the joint planes, which are at right angles to the bedding, are filled with a siliceous deposit.....	2	0
Sandstone, slightly conglomeratic and oolitic.....	8		Shale, gray; weathers almost white; upper part thin bedded; some of the shale is slightly carbonaceous.....	20	0
Shale, thin bedded (estimated yield, 25 gallons).....	5	0	Shale, drab and tan-colored.....	100	0
Shale, dark brown (sample 20; 12.6 gallons).....	3	0	Sandstone, light gray, fine grained, calcareous, ripple marked, lenticular....	8	0
Sandstone, irregular in thickness, weathering light yellow.....	9		Shale, drab and tan-colored.....	200	0
Shale, dark brown; weathers to papery shale (sample 14; 13.3 gallons).....	5		Sandstone, ripple marked, lenticular....	1	0
Sandstone, shaly.....	8		Shale, sandy.....	15	0
Shale, light brownish drab; weathers to sheets one-eighth to three-eighths inch thick (sample 15; 3 gallons).....	1	2	Sandstone, tan-colored, thin bedded....	1	0
Sandstone, coarse, yellow.....	2		Shale, drab.....	28	0
Shale, light brown.....	1	4	Shale, tan-colored.....	50	0
Shale, dark brown, fairly rich (sample { weathers papery..... } 16; 1.9	8		Oolite.....	6	
Shale, hard, dark brown; weathers to blue resistant ledges; this contains lenses of rock which weather yellow and resemble sandstone (sample 17; 21 gallons).....	6		Shale, tan-colored.....	20	0
			Sandstone, thin bedded.....	6	
			Shale, drab, thin bedded.....	40	0
			Sandstone, tan-colored, ripple marked, thin bedded.....	1	0
			Shale, with a few sandstone beds, tan colored; Wasatch formation.....	200	0
				1,677	11

Sections in northwestern Colorado—Continued.

Location G, T. 2 N., R. 97 W.

	Ft.	in.		Ft.	in.
Top of hill.			Shale, 10 per cent dark brown, rich and 90 per cent lean.....	2	0
Sandstone, massive, tan-colored, slightly friable, fairly coarse grained, weathers into nodular forms.....	5	0	Shale, thin bedded (possible yield, 15 gallons).....	10	0
Shale, sandy.....	300±		Shale, carbonaceous.....	3	0
Shale, dark brown, rich.....		½	Covered, probably mostly shale.....	100	0
Shale, sandy.....	30	0	Shale, curly; contains lenses of bituminous sandstone (samples 25 and 26, lower 8 feet of this bed; 4.78 and 3.85 gallons).....	15	0
Shale, dark brown (probable yield, 10 gallons).....	1	0	Shale, carbonaceous and sandy.....	2	0
Shale, drab, thin bedded.....	50	0	Sandstone.....	4	
Shale (estimated yield, 20 gallons).....	3	0	Shale, thin-bedded, carbonaceous; will probably yield a small quantity of oil.	3	0
Shale, thin bedded, sandy.....	25	0	Interval, probably mostly lean shale but with a few thin beds which are rich in oil.....	60	0
Shale, dark brown, rich.....	1	0	Shale, thin bedded (estimated yield, 15 gallons).....	3	0
Shale, sandy.....	45	0	Sandstone.....		½
Sandstone.....	3	0	Shale, thin bedded, black on fresh surface (estimated yield, 25 gallons).	8	
Shale.....	5	0	Interval, covered, probably shale.....	6	0
Shale, dark brown, rich.....		½	Shale, dark gray to black, thin bedded (estimated yield, 25 gallons).....	3	0
Shale, sandy.....	100	0	Interval, mostly covered but probably sandy shale.....	20	0
Shale (estimated yield, 25½ gallons).....	3	0	Oolite.....	10	
Shale, drab, with some sandy beds, lean.	60	0	Shale, thin bedded, brown on fresh surface, with sandstone beds as thick as 1 foot near the base (top includes beds estimated to yield less than 15 gallons)	15	0
Shale, generally lean, but with some rich layers.....	1	0	Shale, thin bedded, brown on fresh surface; contains lenses of sandstone....	4	0
Shale (80 per cent) and sandstone (20 per cent).....	20	0	Interval, covered but probably shale...	15	0
Shale, thin bedded, containing thin layers of rich shale (probable yield of the whole, 15 gallons).....	5	0	Sandstone and sandy shale; the sandstone, especially in the upper part, is oolitic.....	13	0
Shale, drab, with some sandy layers....	20	0	Sandstone, mostly friable; contains beds which weather rusty.....	15	0
Shale, sandy.....	10	0	Shale.....	3	0
Shale, dark brown, with some lean beds (estimated yield of the whole, 25 gallons).....	2	0	Sandstone and oolite.....	3	0
Sandstone, shaly.....	1	0	Sandstone, friable.....	2	0
Shale, dark brown (estimated yield, 20 gallons).....	2	0	Shale, upper part sandy, lower part slightly carbonaceous in places.....	15	0
Shale, lean, oolite, and sandstone, intimately interbedded; oolite and sandstone lenticular.....	3	0	Oolite and sandy shale.....	3	0
Sandstone, shaly.....	3	0	Sandstone, shaly.....	8	0
Shale, drab.....	4	0	Sandstone, with oolitic phases.....	3	0
Sandstone.....	7	0	Sandstone, shaly, friable.....	3	0
Sandstone, shaly.....	10	0	Oolite.....	17	0
Shale, with some thin beds of sandstone; part of the shale is carbonaceous; other parts will probably yield a little oil...	40	0	Sandstone, friable.....	10	0
Interval, mostly covered but probably shale.....	50	0	Oolite.....	20	0
Shale, thin bedded (estimated yield, 15 gallons).....	2	0	Sandstone, shaly.....	15	0
Shale, carbonaceous, containing beds as thick as 6 inches which will yield oil..	20	0	Shale and sandstone, folded and faulted to such an extent that detailed measurements are not possible. The folding and faulting are believed to be confined to this member. Some of the shale will probably yield oil.....	100	0
Shale, sandy.....	15	0	Sandstone, tan-colored, slightly cross-bedded; contains oolitic layers near the top.....	17	0
Shale (estimated yield, 25 gallons).....	5	0			
Shale (estimated yield, 20 gallons).....	2	0			
Shale, thin bedded, mostly carbonaceous; some parts are flexible and probably will yield oil; benches as thick as 2 feet are estimated to yield 25 gallons.	30	0			
Shale, sandy, in places carbonaceous....	25	0			
Shale, for the most part drab and thin bedded; contains beds as thick as 1 foot which will probably yield 20 gallons.....	30	0			

Sections in northwestern Colorado—Continued.

Location G, T. 2 N., R. 97 W.—Continued.

	Ft. in.		Ft. in.
Sandstone, friable; contains a few thin lenses of oolite near the base.....	15 0	Oolite, containing numerous gastropods at the base.....	8 0
Sandstone, with lenses of oolite averaging about 2 inches long.....	1 0	Sandstone, shaly at the base, gray on unweathered surface.....	3 0
Sandstone, shaly; cut by normal faults having throws of 6 to 8 feet; fault planes dip about 50° S.; ripple marks were noted in the highest and lowest layers of the bed.....	3 0	Shale, slightly gray, sandy.....	4 0
Interval, mostly covered, probably sandy shale.....	40 0	Sandstone, with numerous gastropods..	1 0
Sandstone; contains oolitic phases.....	5 0	Sandstone, massive, slightly cross-bedded, fairly well rounded grains of quartz.....	4 0
Sandstone, shaly.....	1 0	Sandstone, calcareous, contains gastropod remains.....	6
Sandstone, oolitic.....	5 0	Sandstone, massive, slightly cross-bedded.....	15 0
Sandstone, shaly.....	3 0	Sandstone; weathers buff; resistant, in layers 1 inch to 6 inches.....	1 6
Sandstone, minutely cross-bedded; top oolitic.....	5 0	Shale, sandy, and friable sandstone; the sandstone predominates but does not form ledges. This is supposed to be typical Wasatch.	
Sandstone, shaly.....	1 0		
Oolite.....	15 0		
Sandstone, oolitic.....	2 0		1,605 1½
Shale, sandy, but contains carbonaceous beds, also sandstone layers as thick as 2 inches.....	25 0		

Section near location H, from gas well to mouth of Piceance Creek, T. 1 N., R. 97 W.

	Ft. in.		Ft. in.
Sandstone, coarse, yellowish brown, friable, cross-bedded.....	169 0	Sandstone, clayey.....	5 0
Shale, light gray.....	10 0	Shale, hard, brown, lean.....	3 0
Sandstone, tan-colored, friable.....	20 0	Sandstone, clayey, shaly.....	17 0
Sandstone, brown, friable.....	160 0	Shale, sandy (80 per cent), and hard, rich shale (20 per cent).....	2 6
Shale, light gray, sandy (sample 42; 0.31 gallon).....	45 0	Shale, hard, brown (sample 38; 25.2 gallons).....	5 11
Shale, gray, thin bedded.....	12 0	Shale, slightly sandy.....	10 0
Shale, hard, brown, rich, lenticular.....	1 0	Sandstone, calcareous.....	3 0
Shale, gray.....	73 0	Shale, partly rich and partly curly.	3 0
Sandstone, yellowish brown at base, contains vegetable remains.....	55 0	Shale, top sandy, tan-colored.....	80 0
Shale, with few lenses of sandstone, much distorted.....	40 0	Shale, hard, brown, rich (90 per cent) and lean shale (10 per cent).....	2 0
Shale, sandy, very lean.....	5 0	Shale, thin bedded, carbonaceous, sandy.	3 0
Shale, sandy at base.....	60 0	Shale, curly, lean (75 per cent), and hard, rich shale (25 per cent).....	3 4
Shale, sandy at top.....	24 0	Shale, sandy.....	5 0
Shale, hard, brown, rich.....	3	Shale, curly, lean.....	2 0
Shale, sandy.....	25 0	Shale, drab.....	6
Shale, sandy, thin bedded.....	2 6	Sandstone, irregularly bedded.....	2
Shale, lean (90 per cent) and hard, rich shale (10 per cent).....	1 6	Shale, thin bedded, lean.....	10 0
Shale, sandy, thin bedded, gray.....	3 0	Shale, thin bedded, lean (90 per cent), and hard, rich shale (10 per cent).....	6
Sandstone, massive, yellowish brown, coarse grained.....	17 0	Shale, thin bedded, sandy.....	5 0
Sandstone, yellowish, thin bedded.....	12 0	Shale, thin bedded (90 per cent), and hard, rich shale (10 per cent).....	1 0
Shale and thin beds of rich, hard shale..	32 0	Shale, slightly curly, lean.....	5 0
Shale, sandy, thin bedded.....	17 0	Sandstone, clayey.....	1 0
Shale, hard, rich, and thin-bedded shale.....	3 0	Shale, thin bedded at base, sandy at top.	25 0
Shale, sandy.....	10 0	Shale, hard, rich.....	6
Shale, hard, brown, rich.....	1	Shale, sandy.....	2 0
Shale, sandy, thin bedded.....	30 0	Sandstone, calcareous.....	5 0
Shale, for the most part barren but including beds as thick as 1 foot which will yield considerable oil.....	132 0	Shale, thin bedded, curly.....	1 0
		Shale, hard, brown, variable thickness (sample 37; 23.25 gallons).....	5 8½
		Shale, thin bedded.....	20 0

Sections in northwestern Colorado—Continued.

Section near location H, from gas well to mouth of Piceance Creek, T. 1 N., R. 97 W.—Continued.

	Ft.	in.		Ft.	in.
Shale, curly, and hard, rich shales.....	10		Shale, curly, papery; will yield oil.....	40	0
Shale, brown, thin bedded, lean.....	3	0	Shale, dark brown, thin bedded.....	3	4
Shale, sandy, thin bedded.....	15	0	Shale, sandy.....		
Shale, thin bedded, curly.....	1	0	Shale, brown, curly, with lenses of sandstone.....	5	0
Shale, sandy.....	5		Sandstone, lenticular.....	1	2
Shale, thin bedded, slightly curly.....	1	0	Shale, thin bedded.....		
Shale, hard, brown, rich.....	2	0	Shale, light brown, lean.....	8	8
Shale, brown, lean.....	10		Shale, very dark brown, curly, thin bedded.....		
Shale, thin bedded, slightly carbonaceous.....	10	0	Shale, thin bedded brown.....	8	
Shale, hard, rich.....	6		Shale, curly.....	4	4
Shale, black, carbonaceous, lean.....	2	0	Shale, curly papery; will yield oil.....	20	0
Shale, thin bedded, drab.....	18	0	Covered, mostly shale.....	30	0
Shale, hard, rich (50 per cent), and lean shale (50 per cent).....	2	0	Shale, curly, papery, will yield oil.....	3	0
Shale, slightly curly (95 per cent), and rich shale (5 per cent).....	1	0	Covered, probably mostly shale.....	15	0
Shale, sandy.....	100	0	Shale, dark brown, rich.....	1	
Shale, hard, rich, and lean shale (estimated yield, 30 gallons).....	3	0	Shale, much of it papery, curly; will yield a small amount of oil.....	154	0
Shale, thin bedded, lean.....	5	0	Shale, curly, papery, fairly rich (samples 27, 28, 29, and 30; 8.4 to 12.5 gallons)...	15	0
Shale, massive, rich, weathers bluish.....	5	0	Shale, much of it curly and will yield oil.....	67	0
Shale, thin bedded, lean (90 per cent), and rich shale (10 per cent).....	52	0	Shale, curly, papery; will yield oil.....	8	0
Shale, dark brown, rich, and lean shale (50 per cent each).....	16	0	Shale and clay, tan.....	48	0
Shale, hard, dark brown, rich.....	5	0	Oolite.....	1	0
Shale, drab.....	2	0	Shale, sandy at base.....	40	0
Shale, thin bedded, lean.....	3	0	Oolite.....	3	0
Shale, weathers blue; massive, rich (samples 32, 33, 34, 35, and 36; 14.7 to 40.6 gallons).....	5	0	Sandstone.....	42	0
Shale, thin bedded, lean.....	31	0	Sandstone, ripple marked at top, with lenses of oolite.....	22	0
Shale, hard, dark brown, rich.....	3	0	Shale, sandy.....	5	0
Shale, hard, black, rich.....	1	0	Oolite, massive; weathers brown.....	3	0
Shale, thin bedded, lean.....	28	0	Sandstone.....	8	0
Shale, hard, dark brown; estimated yield, 15 gallons.....	1	0	Sandstone, oolitic at base, thin bedded at top.....	7	0
Shale, light gray.....	2	0	Oolite.....	2	0
Shale, curly; will yield oil.....	1	6	Sandstone.....	41	0
Shale, dark brown, hard, rich.....	3		Sandstone, brown, fossiliferous (gastropods).....	2	0
Shale, sandy, carbonaceous.....	3	4	Sandstone, massive, irregular base, ripple marked on top.....	5	0
Shale, curly, papery, and small lenses of solid hydrocarbons.....	2	0	Covered, but probably tan-colored shale.....	83	0
Shale, dark brown, hard, rich.....	4		Sandstone, resistant and cross-bedded at top, friable near base.....	27	0
Shale, drab.....	37	0	Sandstone, light gray to white, coarse... Sandstone, clay, and varicolored lenticular shale.....	3	0
Shale, partly papery, curly, with a few thin one-half inch beds of hard, dark-brown, rich shale.....	35	0		200	0
				2,496	6½

Location I, north of White River, T. 1 N., R. 96 W.

	Ft.	in.		Ft.	in.
Sandstone, ripple marked.....	25	0	Shale, lean (25 per cent), and rich shale (75 per cent).....	1	0
Sandstone and shale.....	189	0	Shale, sandy.....	4	0
Sandstone, conglomeratic, friable, yellow.....	15	0	Shale, dark brown, massive, rich.....	1	0
Shale, yellow.....	58	0	Sandstone, friable.....	6	
Shale, drab.....	1	0	Shale, thin bedded, and lean brown shale.....	4	0
Shale, brown, lean.....	1	0	Sandstone, oolitic, tan-colored.....	1	0
Shale, drab, with thin beds of rich shale.....	2	0	Shale, gray, thin bedded.....	15	0
Shale, thin bedded, curly.....	1	0	Sandstone, oolitic and conglomeratic....	3	0
Shale, drab.....	1	0			

Sections in northwestern Colorado—Continued.

Location I, north of White River, T. 1 N., R. 96 W.—Continued.

	Ft.	in.		Ft.	in.
Shale, gray, thin bedded.....	35	0	Sandstone, oolitic and oolite.....	10	0
Shale and sandstone, lean.....	77	0	Shale and thin beds of sandstone.....	85	0
Sandstone, friable.....	4	0	Shale, brown, curly.....	5	0
Shale, sandy.....	2	0	Sandstone, friable, lenticular.....	10	0
Shale, curly (estimated yield, not more than 10 gallons).....	5	0	Shale, sandy, drab.....	47	0
Interval.....	24	0	Shale, thin bedded.....	3	0
Sandstone, oolitic, ripple marked.....	3	0	Shale, sandy, drab.....	75	0
Shale and sandstone, lenticular.....	167	0			
				874	6

Location J, T. 1 N., R. 96 W.

	Ft.	in.		Ft.	in.
Shale, gray; probably will yield little or no oil.....	200		Sandstone, tan-colored.....		4
Shale, containing a few thin beds of rich black shale.....	20	0	Shale; probably will yield a small amount of oil.....		10
Shale; contains a large number of rich layers interbedded with sandy shale.....	20	0	Shale, gray.....	6	0
Shale, thin bedded; contains a large number of thin beds of rich shale; the whole slightly richer than the unit next above.....	15	0	Shale, hard, dark brown, rich.....	2	0
Shale, hard, dark brown, rich.....	6		Shale, papery, lean.....	3	0
Shale; contains a few thin beds of rich shale.....	8	0	Shale, sandy.....	5	
Shale, hard, dark brown, rich.....	2		Shale, dark brown, rich.....		3
Shale, thin bedded, slightly sandy.....	2	0	Shale, lean.....		7
Shale, hard, dark brown, rich (sample 41; 13.7 gallons).....	1	4	Shale, rich.....		8
Shale, lean.....	2	0	Shale, gray, slightly sandy.....	10	0
Shale, hard, brown, massive (sample 40; 12.5 gallons).....	3	8	Shale, hard, dark brown, rich.....		6
Shale, interbedded with layers of rich shale; the whole probably lean.....	6	0	Shale, lean.....		6
Shale, thin bedded, slightly sandy.....	15	0	Shale, very lean.....		1
Shale, hard, dark brown, rich.....	4		Shale, sandy.....		5
Shale, thin bedded.....	5	0	Shale, dark brown, rich.....		1
Shale, hard, dark brown, rich.....	10		Shale, lean.....		1
Shale, sandy.....	5	0	Shale, hard, dark brown, rich.....		3
Shale, hard, dark brown, rich.....	1	0	Shale, lean.....		10
Sandstone.....	8		Shale, dark gray, thin bedded, lean....		6
Shale, hard, dark brown, rich with some lean layers.....	3	0	Shale, gray, slightly sandy.....		5
Shale, thin bedded, slightly sandy.....	50	0	Shale, brown to black, thin bedded, rich.....		3
Shale, hard, dark brown, rich.....	8		Shale, brown to black, thin bedded, probably lean.....		2
Shale, thin bedded, slightly sandy; containing caverns in places filled with solid hydrocarbon.....	5	0	Shale, sandy, thin bedded.....		3
Shale, sandy.....	7	0	Shale, dark brown to black, thin bedded, lean.....		2
Shale, lean.....	1	0	Shale, dark brown to black, thin bedded to almost papery, lean.....		1
Shale, hard, dark brown, rich.....	1	0	Shale, hard, dark brown, rich.....		2
Shale, thin bedded, slightly sandy.....	5	0	Shale, drab, thin bedded, sandy.....		5
Shale, hard, dark brown, rich.....	1	0	Sandstone, massive, showing slight ripple marks.....		3
Shale, lean, slightly sandy.....	2	0	Shale, sandy.....		8
Shale, sandy, thin bedded.....	11	0	Sandstone, massive, resistant.....		2
Shale, sandy; contains a few thin layers of rich shale.....	5	0	Shale, gray, thin bedded.....		3
Shale, gray, slightly sandy.....	4	0	Sandstone.....		4
Shale, hard, dark brown, rich, with lean shale interbedded.....	1	10	Shale, gray, thin bedded.....		45
Shale, gray, slightly sandy.....	8	0	Sandstone, massive, cross-bedded.....		15
Shale, lean.....	6		Shale, gray, thin bedded.....		5
Shale, sandy.....	3	4	Shale, dark gray to black, probably lean.....		1
			Shale, gray, thin bedded, with thin beds of sandstone.....		25
			Shale, hard, dark brown, rich.....		3
			Shale, weathers gray, lean.....		1
			Shale, gray, thin bedded.....		12
			Shale, thin bedded, dark brown to black on fresh surface (sample 39; 13.7 gallons).....		3
			Shale, hard, brown, probably rich.....		6

Sections in northwestern Colorado—Continued.

Location J, T. 1 N., R. 96 W.—Continued.

	Ft.	in.		Ft.	in.
Shale, thin bedded, lean.....	8		4 inches and as thick as one-quarter inch.....	3	0
Shale , hard, dark brown, rich.....	3		Sandstone, tan-colored, massive, slightly friable, minutely cross bedded.....	35	0
Sandstone, with numerous fragments of vegetable remains.....	2	0	Shale, drab, sandy.....	55	0
Shale, gray, sandy, with thin beds of lean shale.....	10	0	Sandstone.....	1	0
Sandstone, coarse grained, conglomeratic, containing flat pebbles of shale.....	1	0	Sandstone and shale, fossiliferous.....	30	0
Shale and sandstone; much of the surface covered.....	45	0	Sandstone, tan-colored on weathered surface, minutely cross-bedded.....	2	0
Sandstone, conglomeratic, tan-colored; contains flat pebbles of shale as long as			Shale, sandy; one fossil leaf was found in this member.....	5	0
					1,666 10½

Location K, on north side of Fourteenmile Creek, T. 3 S., R. 95 W.

	Ft.	in.		Ft.	ins
Sandstone, brown, massive.....	50	0	Shale, thin bedded, lean.....	11	0
Shale, light brown.....	75	0	Sandstone, coarse grained, irregular bedded.....	6	
Shale, thin bedded, light brown, lean (sample 44; 6.2 gallons).....	64	0	Shale, bluish gray.....	11	0
Sandstone, shaly.....	19	0	Sandstone, fine grained.....	8	
Shale, thin bedded; weathers light gray; probably as rich as the bed from which sample 43 was taken.....	40	0	Shale, bluish gray; contains vegetable remains.....	12	0
Sandstone, shaly.....	17	0	Sandstone, brown, shaly.....	10	0
Shale, thin bedded; weathers light gray; probably as rich as the bed from which sample 43 was taken.....	59	0	Shale, gray, with a few thin beds of rich shale.....	117	0
Shale, dark brown, but probably not rich.....	3		Sandstone, grayish brown.....	8	
Shale, thin bedded, very hard, black.....	1	5	Shale, drab, platy; contains stems of wood and pockets of solid hydrocarbon.....	65	0
Shale, brown, hard, probably lean.....	7		Sandstone, gray.....	4	0
Shale , very dark brown, hard.....	1	6	Shale, gray.....	17	0
			Sandstone, brown.....	1	0
Shale, dark brown, containing small lenses of sandstone.....	1	0	Shale, gray.....	2	0
Sandstone, irregular in thickness.....	2		Sandstone, brown.....	1	0
Shale, lean (80 per cent), and rich, brown shale (20 per cent).....	35	0	Shale, gray, clayey.....	77	0
Shale, light brown; weathers white; lean.....	7	0	Sandstone, coarse, yellow.....	7	0
Interval, probably mostly shale.....	40	0	Shale, drab; contains wood fragments.....	80	0
Shale, gray.....	110	0	Sandstone, coarse, angular grains, yellow.....	2	0
Shale, thin bedded, light colored.....	10	0	Shale, gray.....	5	0
Shale, dark drab, thin bedded.....	28	0	Sandstone, brown, coarse.....	1	0
Shale, very thin bedded, with thin layers of rich, dark-brown shale.....	1	0	Shale, light drab.....	15	0
Shale, gray, lean; weathers to small plates.....	5	0	Sandstone, coarse, yellowish brown.....	3	0
			Shale, light drab, very clayey.....	53	0
			Sandstone, yellowish brown, ripple marked.....	1	6
			Shale, drab, sandy, with three thin beds of brownish sandstone.....	32	0
			Sandstone, yellowish brown, upper part oolitic; contains rounded quartzite pebbles.....	19	0
			Top of Wasatch formation.....		
					1,113

Location L, on Piceance Creek, T. 4 S., R. 94 W.

	Ft.	in.		Ft.	in.
Sandstone, brown, massive.....	150±		Shale, dark brown, hard; weathers gray; lean.....	40	0
Shale, sandy; weathers light gray.....	95	0	Shale , hard, dark brown (estimated yield, 30 gallons).....	1	0
Shale, dark brown, hard; weathers gray; lean.....	112	0	Shale, dark brown; weathers gray; lean.....	168	0
Shale , hard, dark brown, rich.....	5		Shale , dark brown, rich, and thin-bedded shale.....	2	0
Shale, dark brown, hard; weathers gray (probable yield, less than 15 gallons)....	11	0	Shale, brown; weathers gray; with a few thin beds of hard, dark-brown, rich shale.....	38	0
Shale , hard, black; weathers bluish (estimated yield, 25 gallons).....	2	0			

Sections in northwestern Colorado—Continued.

Location L, on Piceance Creek, T. 4 S., R. 94 W.—Continued.

	Ft.	in.		Ft.	in.
Sandstone, shaly; weathers reddish brown.....	90	0	Sandstone, brown.....	10	0
Shale, drab.....	100	0	Shale, drab, somewhat sandy, interbedded with about 20 per cent of sandstone in beds 8 inches thick.....	65	0
Sandstone.....	4	0	Sandstone, brown.....	8	0
Shale.....	18	0	Shale, drab, sandy, with about 40 per cent of sandstone in beds 1 foot thick.....	45	0
Sandstone.....	3	0	Sandstone, gray.....	15	0
Shale, drab, slightly sandy.....	523	0	Sandstone in beds averaging 6 feet thick, interbedded with gray sandy shale (shale probably 50 per cent of the whole)	200	0
Shale, bluish drab, with about 5 per cent of beds of sandstone 5 inches thick.....	102	0	Sandstone, massive, brown.....	12	0
Shale, drab, slightly sandy, interbedded with layers of sandstone 1 foot thick....	120	0	Sandstone and gray shale, Wasatch.....	2,002	5
Sandstone, brown, coarse grained.....	8	0			
Shale, drab, slightly sandy, with about 8 per cent of sandstone in beds 4 inches thick.....	60	0			

Location M, on north side of Pole Gulch, T. 4 S., R. 94 W.

	Ft.	in.		Ft.	in.
Shale, light brown, weathers gray; contains a few thin beds of rich shale.....	75	0	Shale, brown, probably not quite as rich as the bed from which sample 46 was taken.....	4	6
Shale, black, massive; weathers dark blue; rich.....	1	0	Shale, light brown; will yield a small quantity of oil.....	5	0
Shale, light brown; weathers gray; contains several thin beds of sandstone, also thin beds of rich shale.....	17	0	Shale, brown, weathers thin bedded, curly (estimated yield, 15 gallons)....	2	0
Shale, dark brown, rich.....	8		Shale, light brown, with beds of rich shale (probable yield less than 10 gallons)...	11	0
Shale, light brown, weathers gray; will yield a small quantity of oil.....	25	0	Shale, dark-brown, weathers blue.....	1	1
Shale, brown (sample 46; 12.5 gallons)....	5	2	Shale, very light brown; weathers yellowish gray; will yield very little oil.....	15+	
Shale, gray, lean.....	5	0		167	5

Location N, on the face of a cliff southwest of Cook's ranch, T. 4 S., R. 94 W.

	Ft.	in.		Ft.	in.
Top of cliff approximately 1,400 feet above the base of the Green River formation.			Shale, dark brown, rich.....	1	
Shale, thin bedded; weathers platy; lean.	15	0	Sandstone, irregularly bedded.....	2	
Shale, dark brown, rich; weathers bluish gray and thin bedded.....	6		Shale, dark brown, rich, containing much pyrite.....	2	
Shale, light brown; weathers yellow; probably will yield a small quantity of oil.....	3	0	Shale, gray and thin sandstone.....	10	0
Shale and thin sandstone bed.....	5	0	Shale, dark brown, rich; weathers bluish gray.....	1	
Shale, lean.....	3	6	Shale, lean.....	11	0
Shale, yellowish gray; will yield but little oil.....	5	0	Shale, hard, dark brown; weathers bluish gray; contains a large amount of pyrite.....	10	0
Shale, light brown; weathers to thin laminae (sample 49; 10.5 gallons).....	3	0	Shale, gray (may yield as much as 10 gallons).....	11	0
Shale, yellowish gray, probably very lean.	7		Shale, thin bedded (may yield as much as 15 gallons).....	8	
Shale, dark brown, rich.....	6		Shale, weathers white.....	16	0
Shale; weathers gray; probably will yield a small quantity of oil.....	7	0	Shale; weathers bluish (will not yield more than 10 gallons).....	10	
Shale, dark brown, rich (sample 48; 15.5 gallons).....	4	0	Shale, dark brown, rich; weathers blue; contains a large amount of pyrite.....	3	
Shale and thin-bedded sandstone; will probably yield a little oil.....	5	0	Shale and sandstone.....	6	0
Shale, brown, lean.....	2	0	Shale; weathers white.....	1	
Shale, light gray.....	6	0	Sandstone.....	3	
Shale, dark brown, rich; weathers bluish gray.....	5		Shale; weathers white; lean.....	1	6
Shale, lean (estimated yield, 10 gallons)..	2	0	Sandstone.....	3	
Shale, light brown, very lean.....	4	0	Shale; weathers white; will yield a little oil.....	1	8
Shale, dark brown, rich.....	1		Shale; weathers bluish, thin bedded (sample 47, 7 gallons).....	2	0
Shale.....	3		Shale; will yield a little oil.....	2	6
			Talus slope concealing the lower rocks.....	141	4

Sections in northwestern Colorado—Continued.

Location O, in Book Cliffs north of Morris station, T. 6 S., R. 94 W.

	Ft.	in.		Ft.	in.
Shale, brown, makes the upper part of the cliff; probably will yield considerable oil but could not be examined.....	210±		Shale (90 per cent) and sandstone (10 per cent); this member contains beds of sandstone as thick as 8 feet; shale is in places slightly carbonaceous.....	100	0
Shale , weathers thin bedded; black to bluish gray (sample 50; 40.6 gallons).....	10	0	Sandstone, light tan-colored to gray, very persistent in thickness.....	5	0
Shale (probably will yield less than 15 gallons).....	25	0	Sandstone (25 per cent) in beds not over 8 inches thick, and drab shale (75 per cent).....	25	0
Shale , hard, black, rich Sandstone (excluded from sample).....	2	0	Oolite (50 per cent) and thin-bedded carbonaceous shale (50 per cent); contains fish scales.....	2	0
Shale , light brown, very hard.....	1	10	Sandstone (10 per cent) and shale (90 per cent); some of the sandstone beds are as thick as 5 feet and are conglomeratic...	60	0
Shale , black, hard; appears to be very rich.....	5		Shale, drab; contains a few thin beds of sandstone.....	60	0
Shale; weathers white; thin bedded, slightly carbonaceous; contains a few thin beds of rich shale, also sandstone lenses.....	40	0	Shale (90 per cent), in places thin bedded and slightly carbonaceous, and sandstone, in part ripple marked (10 per cent).....	70	0
Shale, light yellow, thin bedded, sandy..	30	0	Sandstone (50 per cent) and gray shale, in part clayey (50 per cent); sandstone at top is even bedded and appears persistent; that in the lower part is irregular in thickness.....	100	0
Sandstone, coarse grained.....	2	0	Shale, for the most part gray (75 per cent) and thin-bedded sandstone (25 per cent)	300	0
Shale; weathers white; contains some sandy beds.....	40	0	Sandstone and shale; sandstone beds are massive, tan-colored for the most part and cross-bedded; shale in the lower part of the member is yellow and is similar to Wasatch. At the base of the member is a bed of massive sandstone 10 feet thick with 15 feet of thin-bedded sandstone above. These beds are irregular in thickness and look like the typical Wasatch.....	750	0
Shale , rich, dark brown.....	1		Variegated clays and shales with sandstone lenses at irregular intervals.....		
Shale, slightly sandy, thin bedded.....	1	0			
Shale , rich, dark brown.....	3				
Shale, slightly sandy, gray.....	1	0			
Shale , dark brown, rich.....	8				
Shale, lean.....	6				
Shale, mostly gray, partly sandy, contains a few beds of curly shale; the whole is supposed to be very lean.....	75	0			
Shale, with some sandstone beds as thick as 2 feet.....	120	0			
Sandstone, tan-colored, coarse grained...	25	0			
Shale , dark brown, rich.....	3				
Sandstone (90 per cent) and shale (10 per cent); sandstone coarse grained, cross-bedded, and conglomeratic in lower part.....	25	0			
				2,082	2

Location P, in upper part of Book Cliffs north of Rulison, Colo., T. 6 S., R. 95 W.

	Ft.	in.		Ft.	in.
Shale, massive, probably will yield considerable oil; forms at this locality impassable cliff.....	100±		Sandstone, shaly.....	10	0
Shale, light brown (sample 56; 11.2 gallons).....	15	0	Shale , probably fairly rich in oil....	35	0
Shale brown, probably richer than that of the bed above.....	100	0	Shale , thin bedded; weathers black; inclined to be curly (to be correlated with the bed from which sample 50 was taken; see above)...	6	0
Shale , dark brown (sample 55; 15.4 gallons).....	13	0	Shale; weathers yellowish; lean.....	25	0
Shale , probably as rich as that above	20	0	Shale , dark brown; weathers gray (sample 54; 20.7 gallons).....	8	0
			Talus slope.....		
				332	0

Sections in northeastern Utah.

Location Q, along Evacuation Creek between Temple switch and Dragon, Utah.

	Ft.	in.		Ft.	in.
Shale, thin bedded, lean to barren.....	40	0	Shale, sandy; weathers greenish gray; lean to barren.....	2	0
Shale, hard, dark (estimated yield, 20 gallons).....	1	0	Shale, hard (sample 59; 9 gallons).....	1	0
Shale, lean to barren, thin bedded; a few rich layers less than 1 inch thick (Diptera larvæ).....	155	0	Sandstone, rough, coarse, containing asphalt; top and bottom surfaces irregular, with shale conforming to the irregularities.....	1	10
Sandstone.....	1	0	Shale.....	7	0
Shale, platy, lean to barren; two or three rich beds about 1 inch thick.....	36	0	Sandstone, persistent.....	3	
Shale, thin bedded, rich.....	1	0	Shale, lean, sandy, gray to reddish, with several thin layers of sandstone.....	13	8
Shale, lean.....	6	0	Shale, hard, rich..... (sample 74; 2 0		
Sandstone, persistent.....	5		Shale, soft..... (sample 74; 2 0		
Shale, lean, thin bedded.....	4	0	Shale, hard, rich..... (sample 74; 1 9		
Shale, hard, dark brown, rich.....	1	10	Sandstone.....	3	
Shale, hard, light brown, rich.....	2	1	Shale, hard.....	3½	
Shale, hard, dark brown, rich.....	6		Sandstone (not included in sample).....	1½	
Shale, sandy (not in sample) (sample 68; 31 gal- lons).	2		Shale, hard.....	1	7
Shale, hard, dark brown, rich.....	3		Sandstone, persistent (not included in sample)..... (sample 73; 6 gallons).	2	
Shale, hard, light brown, rich.....	10		Shale, hard.....	11	
Shale, hard, dark brown, rich.....	8		Shale, clayey.....	5	
Shale, dark, tough..... (sample 67, from richest part of upper bench; 90 gal- lons. Sam- ple 65, from whole bed at surface; 32 gallons.)	1	4	Shale, hard, mostly lean, with thin beds of richer shale.....	1	6
Shale, dark, platy.....	1	5	Sandstone, persistent.....	3	
Shale, hard, dark, rich.....	2½		Shale, hard, rich.....	2	9
Shale, soft, dark brown.....	3		Sandstone..... (sample 72; 7 gallons).	1	
Shale, hard, dark, rich.....	3		Shale, hard, rich.....	2	6
Shale, soft, dark brown.....	5		Sandstone.....	2	
Shale, hard, dark, rich..... (sample 66 from whole bed 1½ feet back from outcrop; 55 gallons.)	3½		Shale, hard, dark (sample 71; 7 gal- lons)	7	0
Shale, soft, dark brown.....	10		Sandstone, persistent.....	6	
Shale, hard, dark, rich.....	1	6	Shale, hard, rich to lean.....	3	3
Shale, thin bedded, platy..... (sample 66; 15 gallons.)	4	2	Horizon of sandstone lenses, none of which came where sample was taken..... (sample 70; 14 gallons).		
Shale, rather lean and papery.....	2	0	Shale, hard, rich.....	2	6
Shale, hard, dark brown to black (sample 63; 32 gallons).....	4	3	Sandstone.....	1	
Shale, hard, lean, some thin sandstone layers.....	3	10	Shale, probably, lean. Sandstone, bearing gypsum (not in- cluded in sample)...	7	
Shale, hard, rich (samples 58 and 62; 23 and 18 gallons, respectively)...	3	11	Shale, hard, dark, rich; some gypsum near top..... (sample 69; 19 gallons).	5	
Shale, minutely banded, some rich layers (sample 61; 10 gallons).....	6	7	Shale, with consider- able gypsum.....	1	10
Shale, lean to barren, with two bands of small dark sandstone lenses.....	4	0	Shale, very dark brown, rich.....	4	
Sandstone, hard, quartzitic, persistent.....	5		Sandstone, brownish, shaly.....	23	0
Shale, sandy, barren; thin beds of sandstone.....	3	1	Shale, papery, lean.....	1	6
Shale, brown and black, rich..... (sample 60; 12 gallons).	7		Shale, rich; weathers blue.....	6	
Shale, hard; weathers green.....			Shale, sandy, and barren shaly sand- stone.....	10	0
			Shale, rich, papery.....	6	
			Shale, sandy, barren.....	17	6
			Sandstone, brown, massive.....	11	

Sections in northeastern Utah—Continued.

Location Q, along Evacuation Creek between Temple switch and Dragon, Utah—Continued.

	Ft.	in.		Ft.	in.
Shale, lean to barren.....	2	6	Shale, drab.....	11	0
Sandstone, massive, ledge making.....	5	0	Sandstone, poorly cemented.....	4	6
Sandstone, brownish, shaly.....	50	0	Shale, drab to green.....	17	0
Shale , forming ledge, lean to rich..	4	6	Oolite, much distorted.....	2	8
Shale, papery, lean to barren.....	4	0	Shale, gray.....	1	0
Shale , hard, rich.....	3		Oolite, much distorted.....	8	6
Shale, sandy, lean to barren; two or three sandstone ledges less than 1 foot thick.....	43	0	Shale, sandy.....	19	0
Shale , papery; numerous thin blue rich bands.....	7	0	Oolite.....	3	0
Shale, barren, with several brownish sandstone layers 2 to 4 inches thick..	20	0	Sandstone, massive.....	7	0
Shale ; weathers bluish; rich, ledge forming.....	1	0	Oolite.....	1	2
Shale; weathers bluish; barren.....	2	7	Sandstone, massive for the most part; some oolitic members near base.....	61	0
Sandstone, light brown.....	1	0	Shale, drab; upper surface very irregular.....	4	0
Shale; weathers bluish; barren.....	1	5	Sandstone, massive.....	8	0
Sandstone, light brown.....	4	0	Sandstone, shaly.....	18	0
Shale; weathers bluish; barren.....	1	6	Shale, gray, sandy.....	7	0
Sandstone, light brown.....	8		Sandstone, yellow, massive.....	5	6
Shale; weathers bluish; barren.....	1	3	Shale, gray, sandy.....	17	0
Sandstone, massive.....	2	6	Sandstone, massive.....	3	0
Sandstone, gray, shaly.....	3	0	Shale, drab.....	2	0
Sandstone, yellowish brown, not well cemented.....	48	0	Sandstone, massive.....	14	0
Shale, gray.....	35	0	Shale, drab to dark, sandy and concretionary.....	5	6
Sandstone.....	4	0	Shale, platy, gray and drab.....	11	0
Shale, gray.....	7	0	Oolite.....	6	
Sandstone.....	2	0	Sandstone, finely cross-bedded, poorly cemented.....	5	0
Shale, drab.....	13	0	Oolite.....	10	
Oolite.....	6	0	Sandstone, massive.....	2	0
Shale, drab.....	42	0	Shale, drab, sandy.....	3	10
Oolite.....	2	4	Sandstone, massive.....	1	6
Shale, gray.....	4	0	Shale, drab.....	3	3
Oolite.....	3	10	Sandstone, massive.....	3	0
Shale, gray.....	14	6	Shale, gray and drab.....	102	0
Sandstone, massive cross-bedded.....	14	0	Sandstone, minutely cross-bedded.....	8	
Shale, drab.....	6	0	Shale, gray and drab.....	97	0
Sandstone, massive.....	8	6	Oolitic sandstone, forming ledge; lower 4 inches conglomeratic and may well be considered basal member at Green River.....	1	4
Shale, gray and drab.....	28	0	Sandstone, yellow, poorly cemented....	28	0
Sandstone, resistant.....	10		Shales, red and green, undoubtedly Wasatch.....	1,306	10½
Shale, drab.....	4	0			
Sandstone, massive.....	23	0			

Location R, north side of Saddle Post Canyon, sec. 22, T. 11 S., R. 25 E.

	Ft.	in.		Ft.	in.
Shale, hard, rich, weathers papery.....	2	11	Shale, lean; weathers white and platy..	2	4
			Shale , hard, rich.....	10	
Shale, rather soft, white on surface, platy.....	1	7	Shale, lean.....	12	0
			Shale , hard, rich.....	4	
Shale , hard, rich.....	10		Shale, lean, platy.....	4	8
Shale, white, platy.....	9		Sandstone, persistent, quartzitic (identical with 8-inch sandstone at location Q; see p. 183).....	5	
Shale , hard, rich.....	9				
Shale, lean to rich; weathers papery.....	1	6			
Shale, lean; weathers white and platy..	1	3			
Shale , rich; weathers papery.....	6½				
				30	8½

Sections in northeastern Utah—Continued.

Location S, sec. 24, T. 11 S., R. 25 E.

	Ft.	in.		Ft.	in.
Sandstone.....		8	Shale, lean.....	2	0
Shale, lean, sandy.....	20	0	Shale, rich.....	1	0
Shale, rich.....	3	0	Shale, rich, and papery shale.....	4	0
Shale, lean.....	6	0	Shale, medium rich.....	3	0
Shale, rich.....	1	6	Shale, rich, hard.....	2	0
Shale, lean to barren.....	6	6	Shale, lean.....	4	0
Sandstone.....		5	Shale, medium rich.....	9	0
Shale, lean.....	4	6	Shale, lean.....	4	0
Shale, rich.....		8	Sandstone, ferruginous.....		5
Shale, lean.....	4	0	Shale, lean to barren.....	4	6
Shale, rich.....		10	Shale, hard, rich.....		6
Shale, lean, papery.....	1	6	Shale, lean to barren.....	5	0
Shale, rich.....	3	6	Sandstone, coarse, asphaltic (same as 22-		
Shale, soft.....		2	inch asphaltic sandstone at location Q;		
Shale, rich.....		4	see p. 183).....	1	1
Shale, soft.....	1	7			
Shale, rich.....		8			
				91	9

Location T, on north side of Hells Hole Canyon, sec. 22, T. 10 S., R. 25 E.

	Ft.	in.		Ft.	in.
Shale, platy, sandy, lean to barren.....	15	0	Shale, hard.....		
Shale, mostly lean, with rich beds too thin to sample.....	11	0	Sandstone.....		
Shale, thin, platy.....	1	11	Shale, hard, rich.....	2	11
Shale, hard, rich.....	3	2	Shale, hard, richest in upper 2 feet (sample 82; 20 gallons).....	6	0
Shale, lean.....	2	1	Sandstone, asphaltic, variable in thickness (same as 22-inch asphaltic sandstone at location Q, see p. 183).....	1	4
Sandstone, persistent.....		5	Shale, hard, rich (sample 81; 24 gallons).....	1	11
Shale, lean, platy, containing one 6-inch bed of rich shale.....	6	5	Sandstone.....		1
Shale, hard, very rich.....	3	2	Shale, sandy, platy, lean to barren.....	28	0
Shale, very sandy, lean.....		11	Shale, rich.....		3
Shale, hard, rich (sample 90; 45 gallons).....	4	8	Shale, lean to barren, sandy.....	2	4
Shale, hard.....	1	3	Shale, rich.....		2
Shale, soft.....	1	4	Shale, lean to barren.....	2	6
Shale, hard, rich.....	1	5	Sandstone.....		1
Shale, lean to barren.....	1	9	Shale, gray, lean to barren.....	18	6
Shale, rich (A).....		1	Shale, hard, rich.....		2
Shale, soft (B).....		1½	Shale, gray, lean to barren.....	2	10
Shale, rich (A).....		3½	Shale, alternating hard and soft layers.....	2	4
Shale, soft (B).....		3½	Shale, soft.....	1	0
Shale, rich (A).....		½	Shale, alternating hard and soft layers.....	1	3
Shale, soft (B).....		1	Shale, lean.....	5	0
Shale, rich (A).....		5	Shale, rich.....		1
Shale, soft (B).....		7	Shale, papery, or sandy and platy, lean.....	34	0
Sandstone (discarded).....		1	Shale, rich.....		10
Shale, soft (B).....		6	Shale, sandy, platy, lean to barren.....	37	0
Shale, rich (A).....		1	Shale, rich.....		10
Shale, soft (B).....		2	Shale, barren.....		6
Shale, rich (A).....		½	Shale, hard, rich.....		1
Shale, soft (B).....		5	Shale, platy, barren.....	5	0
Shale, rich (A).....		1½	Shale, hard, rich.....		2
Shale, lean.....	1	8	Shale, for the most part gray and lean, but a few rich layers less than 1 inch thick.....	85	0
Shale, hard, rich (sample 85; 22 gallons).....	1	8	Shale, papery, lean.....	8	0
Shale, lean mostly, with rich layers too thin to sample.....	7	6	Shale, platy, lean.....	3	6
Shale, hard, rich (sample 84; 21 gallons).....	7	8	Shale, hard, rich.....		3
Shale, hard, lean to rich, cliff forming.....	7	4	Shale, sandy, barren.....	3	0
Sandstone, asphaltic.....		3	Shale, hard, rich.....		2
Sandstone, coarse.....		1½			

Sections in northeastern Utah—Continued.

Location T, on north side of Hells Hole Canyon, sec. 22, T. 10 S., R. 25 E.—Continued.

	Ft.	in.		Ft.	in.
Shale, thin, platy, lean to barren.....	30	0	Sandstone.....	3	0
Oolite.....	5	0	Shale, gray, lean to barren.....	19	0
Shale, papery, lean.....	3	8	Oolite.....	6	6
Shale, hard, rich.....	1½	0	Sandstone, twisted and deformed.....	1	10
Shale, barren.....	6	0	Oolite.....	6	6
Shale, hard, rich.....	2	0	Shale, gray to drab, barren.....	19	0
Shale, thin, platy, barren.....	16	0	Sandstone, massive, yellow.....	12	6
Oolite.....	4	0	Shale, gray.....	5	0
Shale, thin, platy, barren.....	1	2	Oolitic sandstone.....	4	0
Sandstone.....	7	0	Shale, sandy.....	13	0
Shale, thin, platy, barren.....	16	0	Sandstone, massive, yellow, cross-bedded.....	10	6
Cherty layers, distorted, concretionary..	1	0	Shales, gray and drab, barren.....	45	0
Oolite, conglomerate at base.....	2	4	Sandstone, mostly yellow and massive, but with two somewhat shaly zones..	41	0
Shale, gray and drab, barren.....	2	4	Shale, sandy, barren.....	21	0
Sandstone.....	8	0	Shale, rich.....	6	6
Shale, gray and drab, barren.....	1	10	Shale, gray and drab, barren.....	35	0
Sandstone, concretionary.....	3	6	Sandstone and barren, sandy shale....	37	0
Shale, gray and drab, barren.....	20	0	Shale, largely masked, mostly drab, papery, with sandy zones; not sufficiently cemented to form ledges.....	95	0
Sandstone, with oolitic bands.....	4	0	Sandstone, coarse, yellow.....	1	10
Oolite.....	3	9	Shale, lean to barren, papery.....	7	0
Shale, gray and drab, barren.....	23	0	Sandstone, yellowish.....	6	6
Oolite, top distorted and sandy.....	3	0	Shale, soft, sandy (sample 79; 1 gallon)..	4	1
Sandstone, oolitic.....	5	0	Sandstone.....	3	3
Shale, gray.....	28	0	Shale, masked, but for the most part a barren greenish shale.....	26	0
Sandstone, shaly.....	15	0	Sandstone, yellowish brown, poorly cemented.....	10	0
Sandstone, massive.....	4	0	Shales, red and green (Wasatch).....		
Shale, barren, gray, sandy.....	18	6			
Oolite.....	3	6			
Shale, barren, gray sandy.....	10	0			
Oolite.....	2	0			
Sandstone.....	2	0			
Shale, barren, gray, sandy.....	65	0			
Oolite.....	1	0			
				1,047	5

The following stratigraphic sections were measured in southwestern Wyoming and show the general character of the Green River formation in the southern parts of the Green River and Southern Red Desert basins:

Sections of parts of Green River formation in southwestern Wyoming.

T. 14 N., R. 99 W.

	Ft.	in.		Ft.	in.
Sandstone, coarse grained, not massive..	50	0	Shale, thin bedded, lean.....	14	6
Sandstone, containing fossil shells.....	4	0	Shale, thin bedded; weathers blue; rich.....	2	0
Sandstone, coarse grained, thin bedded..	10	0	Shale, gray, sandy (not included in sample).....	1	7
Covered, probably sandy shale.....	35	0	Sandstone, yellow (not included in sample).....		
Sandstone, coarse.....	8	0	Shale, thin bedded; weathers blue; rich.....	3	0
Covered, mostly shale.....	30	0	Shale, yellow, sandy.....	28	0
Shale, papery, drab, lean.....	5	0	Shale, papery, lean.....	40	0
Shale, thin, barren and sandstone.....	72	0	Shale, drab, fissile.....	10	0
Shale, drab, thin, lean.....	3	0	Sandstone, concretionary.....	1	0
Shale, thin, drab, barren.....	20	0	Shale, drab, papery.....	13	0
Shale, thin, lean.....	30	0			
Sandstone, concretionary.....	1	0			
Shale, thin, lean.....	14	0			
Oolite and chert.....	0	0			

Sections of parts of Green River formation in southwestern Wyoming—Continued.

T. 14 N., R. 99 W.—Continued.

	Ft.	in.		Ft.	in.
Oolite.....		6	Sandstone, shaly, yellowish.....		1 0
Shale, drab, papery.....	10	0	Shale, drab, papery, barren.....		5 0
Sandstone, oolitic.....		4	Sandstone, shaly, yellowish.....		1 6
Shale, drab, fissile.....	12	6	Shale, greenish drab.....		37 0
Sandstone, micaceous.....		1 0	Maroon clay shale (probably Wasatch).....		489 10
Sandstone, yellowish.....		3 0			
Shale, drab, thin sandstone lenses.....	26	0			

T. 13 N., R. 108 W.

	Ft.	in.		Ft.	in.
Sandstone, ferruginous, containing fossil shells.....		6	Shale, light brown, lean (sample 98; 3 gallons).....		5 0
Shale and sandstone.....	500±		Interval.....		25 0
Shale, thin, brown.....		5±	Shale, lean (?).....		10±
Shale, thin, brown (sample 100; 3 gallons).....		5 0	Shale, dark brown (sample 95; 13 gallons).....		5 0
Shale, thin, brown.....		2 6	Shale, thin bedded, tough.....		10 6
Shale, hard, black, rich.....		3	Shale, thin bedded, tough (sample 96; 4 gallons).....		5 4
Shale, brown, soft.....		5	Shale, thin bedded, tough.....		2 6
Shale, hard, black, rich.....		1	Shale, massive, light brown.....		2 6
Shale, brown, soft.....		8	Shale, massive, light brown (sample 97; 6 gallons).....		4 10
Shale, hard, black, rich.....		2	Shale, lean.....		
Shale, brown, soft.....		1 8			
Shale, hard, black, rich.....		3			586 1½

Sec. 27, T. 17 N., R. 106 W.

	Ft.	in.		Ft.	in.
Shale, drab.....		45 0	Covered, mostly light-colored shale or clay.....		75 0
Sandstone, platy.....		16 0	Clay, mostly gray, with some red.....		80 0
Shale, lean.....		22 0	Clay, gray, yellow, green, and red, with beds of yellow sandstone. All beds very lenticular. A 40-foot bed of massive sandstone at one place splits and within 100 yards along its outer top is represented by variegated clay beds, with a few thin sandstones.....		75 0
Sandstone, brown, platy.....		7 0			
Shale, hard, not so rich.....		24 0			
Shale, hard, thin, platy ^a (sample 118; 14 gallons).....		5 6			
Shale, hard, dark ^a		2 3			
Shale, hard, dark, rich ^a (sample 117; 19 gallons).....		8 1			
Shale, weathers to thin plates; part of member will yield small amount of oil..		120 0			479 10

Secs. 17 and 19, T. 17 N., R. 106 W.

	Ft.	in.		Ft.	in.
Sandstone, massive, brown, coarse (Tower sandstone of Powell).....		125 0	Shale, hard, thin, medium rich.....		1 6
Sandstone, thin bedded.....		35 0	Shale, barren.....		15 0
Shale, papery, gray.....		25 0	Shale, medium, with large gypsum crystals.....		1 6
Sandstone, shaly, gray.....		32 0	Shale, thin, barren.....		80 0
Shale, sandy, lean.....		65 0	Shale, medium, with gypsum crystals.....		8
Shale, hard; contains fish remains (sample 120; 14 gallons).....		5 0	Shale, gray, sandy.....		26 0
Shale, lean.....		20 0	Shale, medium, with gypsum crystals.....		1 6
Shale, thin with lenses of very rich waxy shale.....		55 0	Shale, hard, rich.....		10
Shale, hard.....		15 0	Sandstone, thin, gray.....		8 0
Shale, hard (sample 119; 12 gallons).....		5 0	Shale, hard, rich.....		10
Shale, hard, lean.....		12 0	Shale, gray, sandy.....		17 0
Shale, gray, sandy.....		20 0	Shale, hard, rich.....		2
Shale, hard, rich.....		1 4	Shale, thin, gray, sandy.....		9 2
Shale, gray, sandy, thin sandstones, and a few 1 to 3 inch beds of rich shale.....		90 0	Shale, hard, rich.....		0
			Shale, sandy.....		118 0
			Sandstone, gray.....		4 0
			Shale, sandy, greenish.....		6 0

^a Probably same zone as lower 22 feet 8 inches of section measured in secs. 17 and 19 of this township.

Sections of parts of Green River formation in southwestern Wyoming—Continued.

Secs. 17 and 19, T. 17 N., R. 106 W.—Continued.

	Ft. in.		Ft. in.
Sandstone, gray, thin bedded.....	1 0	Shale, lean to barren.....	75±
Shale, sandy, green.....	27 0	Shaly sandstone, barren.....	15 0
Shale, sandy, thin bedded, gray.....	21 0	Shale, sandy, forming slope, lean.....	47 0
Sandstone and shale, green, in beds 2 feet thick; sandstone, concretionary.....	58 0	Shale, hard (sample 114, lower 4½ feet; 11 gallons).....	5 6
Shale, sandy, gray, slope.....	97 0	Shale, hard (sample 113; 10 gallons).....	5 3
Sandstone, massive, cross-bedded, forming ledge and capping hill.....	5 0	Shale, hard.....	1 4
Shale, forming slope.....	43 0	Sandstone.....	3
Sandstone, rather massive, forming ledge.....	10 0	Shale, fairly soft, thin-bedded.....	2 9
Shale, soft, thin, platy, barren.....	30 0	Shale, hard.....	1 0
Shale, medium hard, rather thin, very lean.....	4 11	Shale, hard, rich (sample 111; 19 gallons).....	5 4
Shale, medium hard (sample 116; 4 gallons).....	4 10	Shale, hard, rich (sample 110; 19 gallons).....	6 3
Shale, sandy, lean to barren.....	70 0	Shale, soft.....	7
Shale, medium hard, very lean.....	10 0	Shale, hard, rich.....	2 5
Shale, lean.....	4 6	Shale, soft.....	6
Sandstone, brown, persistent.....	8	Shale, hard, rich.....	2 3
Shale, lean.....	3 6		
Shale, hard to medium hard (sample 115; 9 gallons).....	4 6		1,360 10

Along Bitter Creek, T. 18 N., R. 107 W.

	Ft. in.		Ft. in.
Sandstone, massive, brown.....	135 0	Shale, greenish gray.....	21 6
Shale, lean to barren.....	3 0	Sandstone, thin bedded.....	11 0
Shale, hard.....	6 0	Covered, mostly barren gray sandy shale, with a few ledges of gray shaly sandstone.....	128 0
Sandstone, brown, massive.....	1 0	Sandstone, platy.....	3 0
Shale, lean to rich.....	1 0	Shale, greenish.....	7 0
Sandstone, brown, massive.....	5 0	Covered, mostly barren gray sandy shale, with a few ledges of gray shaly sandstone.....	110 0
Shale, lean, papery.....	11 0	Shale; weathers papery.....	1 6
Sandstone.....	2	Sandstone.....	6
Shale, hard, rich.....	6	Shale; weathers papery.....	2 0
Sandstone.....	3	Shale, gray, sandy, with layers of shaly sandstone.....	21 0
Shale, hard, dark.....	8	Shale, greenish.....	20 0
Shale, brown, tough.....	4	Shale, gray, sandy, with layers of shaly sandstone.....	17 0
Shale, hard, rich, dark.....	1 4	Shale, greenish, with brown sandstone lentils.....	20 0
Sandstone.....	3	Sandstone, brown, with some clay shale.....	35 0
Shale, hard, rich.....	2 6	Sandstone, thin, platy.....	4 0
Shale, hard, rich (sample 127; 18 gallons).....	-6 3	Sandy shale and shaly sandstone, gray, barren.....	85 0
Sandstone.....	6		755 5
Shale, hard, rich.....	6 2		
Sandstone, hard, massive.....	6		
Shale, hard, gray, sandy, lean to barren.....	13 0		
Shale, soft, greenish, lean.....	15 0		
Partly masked, barren gray shale and sandstone, with some lean papery shale.....	55 0		
Sandstone, platy.....	2 0		
Shale, lean, soft.....	2 6		

White Mountain, sec. 36, T. 19 N., R. 106 W.

	Ft. in.		Ft. in.
Sandstone, brown, coarse (tower sandstone of Powell).....	245 0	Shale, sandy, gray, and shaly sandstone.....	22 0
Shale, gray, sandy, and shaly sandstone, with three beds of rich shale each 3 inches thick in lower part.....	265 0	Sandstone, shaly, yellow.....	2 0
Shale (estimated yield, 12 to 15 gallons).....	3 0	Shale, sandy, and clay, with a few thin sandstone beds; color predominantly white.....	133 0
Shale, gray, sandy, and thin sandstone with two or three 1-inch beds of rich shale.....	37 0	Sandstone, green, shaly.....	33 0
Sandstone, gray, ripple marked.....	1 0	Shale, gray, sandy, and thin sandstone.....	58 0
		Shale, green, sandy, and green sandstone.....	17 0
		Shale, greenish drab, sandy.....	35 0
		Sandstone, chalky, cross-bedded, brown.....	2 0

Sections of parts of Green River formation in southwestern Wyoming—Continued.

White Mountain, sec. 36, T. 19 N., R. 106 W.—Continued.

	Ft.	in.		Ft.	in.
Shale, drab, sandy.....	95	0	Shale, sandy, gray.....	24	0
Sandstone, ferruginous.....	4		Sandstone, shaly, gray, fossiliferous.....	2	0
Shale, sandy, gray-green, and shaly sandstone.....	75	0	Shale, carbonaceous.....	8	0
Sandstone, shaly.....	2	0	Clay, sandy, gray.....	40	0
Shale, papery, lean, with 2-inch beds of rich shale and some thin beds of sandstone.....	87	0	Sandstone, coarse, gray, possibly base of Green River formation.....	1	0
Sandstone, with clay balls.....	6		Clay, somewhat sandy, gray.....	55	0
Shale, papery, lean.....	26	0	Shale, clay, variegated, red at top.....	33	0
			Sandstone, yellowish green, friable.....	30	0
				1,331	10

Fossil Butte.

	Ft.	in.		Ft.	in.
Sandstone, shaly.....	95	0	Shale, white, chalky.....	10	
Sandstone, coarse, brown.....	3		Shale, hard, brown.....	6	
Shale, hard, rich, dark; weathers blue (sample 131; 50 gallons).....	2	0	Shale, white, chalky; contains abundant fish remains.....	129	8
Sandstone, shaly.....	7	0	Shale, lean, thin bedded.....	18	0
Shale, hard, rich; weathers blue.....	1	0	Sandstone, drab.....	10	0
Alternating bands of coarse sandstone and shale.....	8	0	Shale, sandy.....	5	0
Sandstone, cherty.....	10		Sandstone, coarse.....	65	0
Shale, lean, chalky, thin bedded.....	15	0			
Sandstone, reddish brown.....	8			250	1½
Shale, lean, chalky, thin bedded.....	2	6			
Sandstone, massive.....	8	6			
Chalk.....	3				
Sandstone, massive, yellow.....	3	2			
Shale.....	10				
Sandstone, gray (discarded).....	5				
Shale, hard, light brown.....	1	7			
Sandstone, light brown (discarded).....	4				
Shale, hard, brown, rich; weathers blue.....	1	1			
Sandstone, yellow.....	½				
Shale, hard, brown, rich; weathers blue.....	11				

STRUCTURE.

Northwestern Colorado and northeastern Utah.—The Uinta Basin is a broad, shallow syncline whose central portion is occupied by the Green River and younger formations. The Green River beds are only slightly tilted at most places around the margin of the basin, but the older formations dip at much greater angles only a few miles beyond the limits of the Green River formation. The Douglas anticline, which extends southward from Rangely, brings up the Mesa-verde and Wasatch formations and perhaps entirely separates the main Colorado area of the Green River formation from the much larger area in northeastern Utah, and it is almost certain that the areas of oil-yielding shale are completely separated. Although the Green River formation dips sharply (maximum observed dip 28°) toward the interior of the basin at its margin along the north and east sides of the Colorado area, the dip decreases very rapidly,

so that a few miles back of the margin the shale is horizontal or dips only slightly.

Small faults were observed at a few places, although it is probable that the broad movement which produced the great synclinal structure gave rise to fracture zones or cracks and faults at many other places. The great veins of gilsonite in the vicinity of Dragon, Utah, are cracks filled with the asphaltic material. West of the Petrolite Hills, in the eastern part of the area studied, a fracture zone cutting the Green River formation is filled with a very light yellowish-brown hydrocarbon (specific gravity 1.06) which does not answer the description of any of the ordinary asphaltites. The deposit may not be sufficiently extensive to be of economic importance, but it is of considerable scientific interest. The fracture zone in which the material is found is 2 or 3 feet wide and has been traced for several miles in a




FIGURE 14.—Diagram showing structure in the Green River formation near the mouth of Yellow Creek, Colo.

general northwest-southeast direction. Several small prospects have been opened along it.

On Yellow Creek, sec. 15, T. 2 N., R. 99 W., a heavy brown sandstone rests on a very irregular surface, and the shale beneath (A, fig. 14) is very much distorted, exhibiting in places very definitely overturned folds, as well as faults on a small scale. It is not at all impossible that the brown sandstone marks the base of a formation which should be separated from that below, but in this paper it is included in the upper part of the Green River formation. The massive sandstone appears to fill channels, yet the distorted condition of the shale beneath would suggest that the shale was still in a plastic condition after the deposition of part of the sand and that the sand settled unevenly, giving rise to the many peculiar forms which are exhibited in the shale beds beneath. After this settling the deposition of sand was continued over the edges of the portions of the sand which had been displaced. This appears to represent a change in conditions of deposition which may have more than local importance.

Southwestern Wyoming.—That part of southwestern Wyoming (see Pl. X) lying between the Uinta Mountains on the south and the Wind River Mountains on the north is in a broad structural basin which is separated into several smaller basins by north-south uplifts. Oil-yielding shale is present in three distinct areas; the central area, to which the name Green River basin is applied, is by far the largest. The Rock Springs uplift, on the east, described by Schultz,¹ and related features near the Colorado-Wyoming State line, to the southeast, separate the main area from the much smaller Southern Red Desert Basin, only the rim of which is shown on Plate XIX (in pocket). On the west the Green River basin is separated from the area of the Green River formation in the vicinity of Fossil, described by Veatch² as the Fossil syncline, by an anticlinal fold which the same author called the Meridian anticline.

Beds of the Green River formation are only slightly tilted in any part of the area shown on Plate XIX. Along the western rim of the Southern Red Desert Basin the oil shale at its outcrop dips as much as 11° NE., but it flattens rapidly toward the center of the basin. Along the east and west sides of the main Green River basin the oil-yielding shale shows dips not exceeding 3° or 4°, but in some places along the north flanks of the Uinta Mountains the beds of the Green River formation are tilted at greater angles. The oil shale of the Fossil syncline is practically horizontal at every point examined.

Faults in the Green River shale were noted in only one area, but they may be present at many other places within the area here described. West of Green River, near the center of T. 15 N., R. 108 W., there is an area perhaps half a mile wide and extending an unknown distance to the west, where the rocks are crumpled and cut by east-west trending faults. On both sides of this disturbed zone the strata are apparently unaffected.

The channel sandstone of Lee,³ which is the Tower sandstone of Powell,⁴ forms vertical cliffs in the vicinity of Green River, Wyo., where it rests on an irregular surface of shale and is itself very much distorted, whereas the shale beneath is not deformed.

BIBLIOGRAPHY.

The literature of oil shale is confined largely to reports dealing with the shale of foreign countries, very little having been published on that of the United States. The reports that treat of the methods of

¹ Schultz, A. R., The southern part of the Rock Springs coal field, Sweetwater County, Wyo.: U. S. Geol. Survey Bull. 381, pp. 218-220, 1910.

² Veatch, A. C., Geography and geology of a portion of southwestern Wyoming, with special reference to coal and oil: U. S. Geol. Survey Prof. Paper 56, pp. 103-110, 1907.

³ Lee, W. T., and others, Guidebook of the western United States, Part B, The Overland Route: U. S. Geol. Survey Bull. 612, p. 74, 1915.

⁴ Powell, J. W., Geology of the eastern portion of the Uinta Mountains, pp. 40, 45, U. S. Geol. and Geog. Survey Terr., 2d div., 1876.

mining and distilling oil shale in Scotland, France, and New South Wales are especially interesting and instructive in view of the probable development in the near future of an oil-shale industry in the United States. The following list of reports includes a large number of rather extensive articles on oil shale of different localities, as well as numerous brief notes on the occurrence of shale in more or less isolated areas. The list includes most of the important reports that have been published.

- ALLEN, A. H., On the relative proportions of olefines in shale and petroleum products: *Analyst*, vol. 6, pp. 177-180, 1881.
- ANDERSON, F. M., Oil shales of Elko, Nev.: *Am. Inst. Min. Eng. Bull.*, 1914, pp. 1402-1403. Notes great percentage of organic matter in oil shale from Elko, Nev. States that no bitumen can be extracted by solvents such as chloroform and ether.
- ANDERSON, G., Description of the bituminous rocks which occur in Ross-shire and in the neighborhood of Inverness: *Edinburgh Jour. Sci.*, vol. 4, pp. 93-95, 1826. Brief note only.
- On the occurrence of a bituminous substance near Mountgerald (Scotland): *London Geol. Soc. Quart. Jour.*, vol. 19, p. 522, 1863. Brief mention of oil shale near Mountgerald (Scotland).
- ANDRÄ, C. J. [Fossil fuel from New South Wales (kerosene shale)]: *Naturhist Ver. preuss. Rheinl. Verh.*, vol. 32, *Sitzungsber.*, p. 5, 1875. Gives results of distillation of sample of kerosene shale sent from Sydney, New South Wales.
- ARON, A., Note sur l'industrie française des schistes bitumineux: *Annales des mines*, 10th ser., vol. 9, pp. 47-75, 1906. Discusses oil-shale industry of France, giving statistics of production, 1893-1904. Also gives relation of French to Scotch retorting practice as regards results obtained.
- ATKINSON, A. A., Reports of chief inspector of coal and [kerosene] shale mines: *New South Wales Dept. Mines Ann. Repts. for 1897-1910, 1898-1914*. Statistics of shale production in New South Wales in tons of shale mined and value for each yield from 1865 to date.
- BAKER, O. H., Australian kerosene shale—notes preparation being made to work the New South Wales deposits: *U. S. Cons. Rept.* 333, pp. 107, 108, 1908.
- BALL, L. C., Tertiary oil shales of Baffle Creek, Port Curtis district, Australia: *Queensland Govt. Min. Jour.*, vol. 15, pp. 19-25, 1914. Describes occurrence of oil shale and gives a few tests of the shale.
- Tertiary oil shales of the Narrows-Port Curtis district, Australia: *Queensland Govt. Min. Jour.*, vol. 15, pp. 73-76, 1914. Describes occurrence and mining possibilities of oil shales.
- Oil-shale industry, an introduction to a report on the oil shales of Lowmead now being prepared: *Queensland Govt. Min. Jour.*, vol. 16, pp. 608-616, 1915. Good description of geographic distribution of oil shales, with notes on their origin, characteristics, treatment, and products.
- BASKERVILLE, C., Economic possibilities of American oil shales: *Eng. and Min. Jour.*, vol. 88, pp. 149-154, 195-199, 1909. Gives history of oil-shale industry (especially American); discusses methods of mining and treatment in the light of the practice in Scotland.
- Oil shales of Canada: *Cong. internat. chim. appl. compt. rend.*, 7th sess., vol. 4, pp. 22-31, 1910.
- American oil shales: *Jour. Ind. and Eng. Chemistry*, vol. 5, p. 73, 1913. Mentions oil shales in Nevada, Montana, and California.

- BASKERVILLE, C., and HAMOR, W. A., Oil shales of America: Jour. Ind. and Eng. Chemistry, vol. 1, pp. 507-511, 1909. Gives history of oil-shale industry and describes the occurrence and tests of Canadian oil shales.
- BEILBY, GEORGE, On the production of ammonia from the nitrogen of minerals: Soc. Chem. Ind. Jour., vol. 3, pp. 216-224, 1884. Describes shale-distilling retorts and discusses nitrogen obtained from oil shale.
- BEILBY, G. T., The production of ammonia from nitrogen of minerals: Soc. Arts Jour., vol. 33, pp. 313-320, 1885.
- The nitrogen of crude petroleum and paraffin oils: Soc. Chem. Ind. Jour., vol. 10, p. 120, 1891. Note on nitrogen obtained from Scottish shale oil.
- Thirty years of progress in the shale-oil industry: Soc. Chem. Ind. Jour., vol. 16, pp. 876-886, 1897. Discusses evolution of modern shale still and ammonia scrubber, with tables of cost.
- BERTHIER, P., Analyse de quelques mineraux de bitume (de France): Annales des mines, 3d ser., vol. 13, pp. 605-616, 1838.
- Analyse des schistes bitumineux d'Autun: Annales des mines, 3d ser., vol. 13, pp. 616-618, 1839.
- BLAKE, G. S., Oil shale from Natal (analysis): Imp. Inst. Bull., vol. 1, pp. 74-76, 1903. Thin shale between coal beds yields less than 1 gallon of oil per ton.
- BOURGOR, —, Note sur le gisement des schistes bitumineux à impressions de végétaux de Ménat; sur la chaleur présumable d'un dyke basaltique, non venu au jour, au point où une partie de ces schistes a été transformée en tripoli par la chaleur souterraine; considérations générales sur ce sujet: Soc. géol. France Bull., 2d ser., vol. 8, pp. 39-44, 1850.
- BRANNER, J. C., The oil-bearing shales of the coast of Brazil: Am. Inst. Min. Eng. Trans., vol. 30, pp. 537-554, map, 1901. Describes the occurrence of Eocene oil shales along the east coast.
- BROWN, J. F. K., The working of South African oil shales: South African Min. Jour., vol. 9, pt. 1, pp. 352, 396, 1911. Likens oil shales of Transvaal to carbonaceous shale and concludes that there is little chance of their development.
- Oil-shale mining: Petroleum World, vol. 11, pp. 164-168, 1914. Good general discussion of occurrence, methods of mining, and distillation of oil shale, with costs.
- CADELL, H. M., The oil shales of the Scottish Carboniferous system: Jour. Geology, vol. 2, pp. 243-249, 1894. Describes geology briefly.
- Geology of the oil-shale fields of the Lothians: Edinburgh Geol. Soc. Trans., vol. 8, pp. 116-162, 1901. Good description, with map.
- The oil-shale fields of the Lothians: Inst. Min. Eng. [London] Trans., vol. 22, pp. 314-371, pls. 7, 9, 1902. Good description, with map and short discussion of the oil-shale industry.
- Scottish shale industry: Petroleum World, vol. 10, pp. 228-236, 1913. Good description of oil shale and processes of distillation and refining.
- CADELL, H. M., and GRANT WILSON, J. S., The geology of the oil-shale fields: The oil shales of the Lothians, 1st ed., pt. 1, pp. 1-97, Scotland Geol. Survey Mem., 1906. Describes geology and give data relative to value of different beds.
- CALDWELL, WILLIAM, Methods of working the oil shales: The oil shales of the Lothians, 1st ed., pt. 2, pp. 98-132, Scotland Geol. Survey Mem., 1906; 2d ed., pt. 2, pp. 95-135, 1912.
- The working of oil shale at Pumpherston (Scotland): Inst. Min. Eng. [London] Trans., vol. 36, pp. 581-589, 1909. Describes methods used.
- CAMERON, —, The bituminous deposits of the Camamu Basin, Province of Bahia, in the Brazilian Empire, London, 1884.

- CANAVARI, M., Gli schisti a fucoidi, e gli schisti bituminosi che spesso li accompagnano nell' Appennino centrale; Una radiolite del suavicino: Soc. toscana sci. nat. Proc. verb., vol. 3, pp. 6, 7, 1881.
- CARNE, J. E., On certain coal and shale lands in the Capertee Valley district, New South Wales: New South Wales Geol. Survey Rec., vol. 4, pp. 39-48, 1894. Describes geology of kerosene shale in Capertee Valley district.
- The kerosene-shale deposits of New South Wales: New South Wales Geol. Survey Mem., Geol. ser., No. 3, 1903. Describes development, geology, and method of treatment.
- CARRUTHERS, R. G., The geology of the oil-shale fields: The oil shales of the Lothians, 2d ed., pt. 1, pp. 1-94, Scotland Geol. Survey Mem., 1912. Describes geology.
- CATLIN, R. M., Oil shales of Elko, Nev.: Am. Inst. Min. Eng. Bull., 1914, p. 1402. Attributes Elko oil shale to saturation of shale by wax from oil which has escaped.
- CHESNEAU, G., Rapport sur l'industrie des huiles de schiste en France et en Angleterre, Paris, 1892.
- L'industrie des huiles de schiste en France et en Écosse: Annales des mines, 9th ser., vol. 3, pp. 617-673, 1893. Outlines oil-shale industry of France and compares distillation and refining practices of France with those of Scotland.
- CHOSSON, —, De la situation de l'industrie des schistes bitumineux du bassin d'Autun: Annales des mines, 6th ser., vol. 20, pp. 347-428, 1871. Detailed discussion of distillation and refining processes used in different fields of France.
- CLARKE, W. B., On the occurrence and geological positions of the oil-bearing deposits of New South Wales: London Geol. Soc. Quart. Jour., vol. 22, pp. 439-448, 1866. Short discussion of age of oil shale and cannel coals of New South Wales.
- COSTE, EUGENE, Oil shales of Elko, Nev.: Am. Inst. Min. Eng. Bull., 1914, pp. 1403-1404. Concludes from evidence of Elko oil shales that the wax or oil in them can be there only by impregnation and is of inorganic origin.
- CRAIG, CUNNINGHAM, Petroleum prospects in South Africa: Petroleum World, vol. 11, pp. 265-271, 1914. Describes geology of oil-shale fields of Transvaal and Natal and tests of shale and gives estimates of cost.
- DAWKINS, W. B., On the kerosene shale of Mount Victoria, New South Wales: British Assoc. Rept., 1886, p. 643, 1887. Mentions kerosene shale and its relations.
- DELAHAYE, N. B., Sur les schistes de Muse (Saône-et-Loire): Soc. géol. France Bull., 2d ser., vol. 5, pp. 304-308, 1848.
- Faits et observations pour servir à l'histoire des schistes bitumineux: Rev. sci. ind., vol. 38, pp. 1-17, 49-63, 97-100, 161-174, 225-232, 321-329, 337-352, 440-447, 1850.
- DESBIEF, P., Schistes bitumineux des environs d'Autun: Annales des mines, 5th ser., vol. 14, pp. 44, 45, 1858. Brief note on the yield of oil from oil shales of Autun, France.
- DIETRICH, P. F. DE, Description des gîtes de minéral et des bouches à feu de la France, Paris and Strasburg, 1786-1799.
- DUNLOP, R., Oil-shale deposits at Orepuki: New Zealand Dept. Mines Rept. C³, pp. 52-54, 1900. Describes plant being installed to get oil and ammonia from the oil shales at Orepuki, New Zealand.
- ELLS, R. W., Oil shales of eastern Canada: Canada Dept. Mines Summ. Rept., 1909, pp. 200-216. Describes occurrence and gives large number of tests of oil shales in eastern Canada.
- Bituminous shales in Nova Scotia and New Brunswick, with notes on the oil shales in Scotland: Canada Geol. Survey Summ. Rept., 1908, pp. 132-142, 1909. Describes geology of oil shales of Scotland and of Nova Scotia and New Brunswick and gives results of distillation tests of the latter.

- ELLS, R. W., Joint report on the bituminous or oil shales of New Brunswick and Nova Scotia also on the oil-shale industry of Scotland, Canada Dept. Mines, 1909. Discusses at length the oil-shale industry of Scotland. Describes geology and mining possibilities of oil shales of New Brunswick and Nova Scotia, giving results of distillation tests.
- Notes on the geology of the oil shales of Scotland and their relation to somewhat similar oil shales in eastern Canada: Roy. Soc. Canada Trans., 3d ser., vol. 3, sec. 4, pp. 35-44, 1909. Short sketch of geology of oil shales of eastern Canada.
- The commercial value of the oil shales of eastern Canada, based on their contents by analysis in crude oil and ammonium sulphate: Nova Scotia Min. Soc. Jour., vol. 15, pp. 29-56, 1910. Describes in detail tests on shales from different fields of Canada and discusses costs and values. Notes oil shale from Melville Island and area along lower part of Mackenzie River, also from Island of Spitzbergen.
- The oil shales of the Maritime Provinces (Canada): Min. Soc. Nova Scotia Jour., vol. 14, pp. 1-12, 1910. Outlines possibilities of oil shales of Canada and describes briefly the results of tests on these shales made in the shale works of Scotland.
- ELSNER, L., Ueber die Zusammensetzung des Schieferöl: Ver. Beford. Gewerbl. Preuss. Verb., Band 24, pp. 195, 196, 1845.
- FERRERO, O., Analisi chimica degli scisti bituminosi della valle di Setarolo: R. ist. Lombardo Atti, vol. 3, pp. 15-18, 1862.
- GILLET DE LAUMONT, F. P. N., Observations sur les schistes bitumineux, sur les bitumes ou matières bitumineuses, et sur les Grès Psammites cités dans le mémoire de M. Leschevin: Jour. Mines, vol. 33, pp. 46-52, 1813.
- GRÄFE, E., Ueber canadischen Oelschiefer: Braunkohle, Band 9, pp. 424-426, 1910. Brief description of Canadian oil shales.
- Die schottische Schiefertheerindustrie: Petroleum Zeitschr., Band 6, p. 69, 1910. Gives history of Scotch oil-shale industry, beginning with 1694, also good description of processes of distillation and products.
- GRAY, THOMAS, The phenols from shale oil: Soc. Chem. Ind. Jour., vol. 21, pp. 845-847, 1902. Gives chemical analysis of "green naphtha" from shale oil.
- GOODCHILD, J. G., Some of the modes of origin of oil shales, with remarks upon the geological history of some other hydrocarbon compounds: Edinburg Geol. Soc. Trans., vol. 7, pp. 121-131, 1897.
- GREEN, BURTON, Kimmeridge shale, its origin, history, and uses: London, 1886.
- GRIFFITHS, A. B., Paraffin shale from Servia: Chem. News, vol. 49, pp. 107, 108, 1884. Notes a shale in western Servia which will burn with a smokeless flame and yields paraffin on distillation.
- HART, C. F., Geology and physical geography of Brazil, pp. 263-264, Boston, 1870. Mentions shale ("turba") from Bahia Province, Brazil, that yields large amount of oil when distilled.
- HELLSING, GUSTAF, Skifferoljeindustrieen i Skottland och Frankrike: Sveriges geol. Undersökn. Årsb., 1907, No. 2, pp. 1-92, 1906. Discusses at length the oil-shale industry, giving geologic as well as economic data.
- HENDERSON, N. M., History of shale retorts at Broxburn: Soc. Chem. Ind. Jour., vol. 18, pp. 246-248, 1899. Discusses effect of different retorts on the percentage and character of products obtained by the distillation of oil shale.
- HEUSLER, F., Ueber die Zusammensetzung der schottischen Schiefertheeröle, ein Beitrag zur Theorie der Theerbildung: Deutsche chem. Gesell. Ber., Band 30, pp. 2743-2752, 1897.

- HUNT, T. S., Contributions to the chemical and geological history of bitumens and of pyroschists or bituminous shales: *Am. Jour. Sci.*, 2d ser., vol. 35, pp. 157-171, 1863.
- JENE, H. L., Oil-shale deposits, Blue Mountains, New South Wales: *Eng. and Min. Jour.*, vol. 90, p. 407-408, 1910. Describes briefly occurrence and working of deposits.
- LAHORE, —, Appareil de distillation des schistes, asphaltes et des matières sèches ou liquides: *Gen. Industr.*, vol. 24, pp. 304-308, pl. 322, 1862.
- LAURENT, A., Sur les schistes bitumineux et sur la paraffine: *Annales chimie et phys.*, 2d ser., vol. 54, pp. 392-396, 1833.
- De l'huile des schistes bitumineux et de quelques produits qu'on en obtient: *Compt. Rend.*, vol. 4, pp. 909-912, 1837.
- LEE, G. W., The occurrence of oil shales among the Jurassic rocks of Rassay and Skye: *Nature*, vol. 92, pp. 169-170, 1913. Notes the occurrence of Jurassic oil shale in northern Scotland.
- LEFEBVRE, —, Notice sur les mines de bitume de Bastennes et de Gaujacq (Landes): *Annales des mines*, 3d ser., vol. 13, pp. 585-594, 1838.
- MCGRATH, J. W., Oil shales of Newfoundland: *Petroleum Rev.*, vol. 33, No. 686, p. 209, 1915.
- MAGNIN, J., L'industrie des schistes bitumineux: *Jour. pétrole*, vol. 8, pp. 20-23, 35-37, 51, 52, 68, 69, 1908. Describes processes of distillation and refining used in Scotland.
- MANSFIELD, W. H., Oil shales and their occurrence: *Petroleum Rev.*, vol. 34, pp. 159-160, 199-201, 1916. Good general statement on distribution of oil shales and detailed report on oil shale of Dorsetshire, England, and their treatment.
- MASTON, —, and PARISOT, —, Sur le depot des schistes bitumineux de Froibe-Fontaine, 1862.
- MAYER, J., The mineral-oil industry of Scotland: *Min. Mag. Rev.*, vol. 1, pp. 118-128, 1872.
- MINING WORLD, Transvaal oil-shale deposits: *Min. World*, vol. 34, pp. 74-75, 1911. Describes briefly oil shales at Ermelo and Wakkerstroom. Reprinted in *Petroleum Rev.*, vol. 24, pp. 147-148, 1911.
- Ponthierville oil shales: *Min. World*, vol. 34, p. 1182, 1911. Brief mention of oil shales of the Congo, Africa.
- Bituminous and oil shales in Canada: *Min. World*, vol. 35, p. 202, 1912. Also calls attention to oil shales in Book Cliffs region of Colorado.
- Prospecting the Transvaal oil shales: *Min. World*, vol. 35, p. 714, 1912. Describes occurrence and prospecting of oil shale near Ermelo, Transvaal, Africa.
- MOORE, R. T., The mineral oil industry of Scotland: *Fed. Inst. Min. Eng. Trans.*, vol. 4, pp. 36-47, 1893. Describes briefly geology and method of mining of oil shale, also refining of shale oil and value of different shale beds.
- NEWBERRY, J. S., On the discovery of mineral wax in Utah: *Am. Jour. Sci.*, 3d ser., vol. 17, pp. 340-341, 1879. Mentions oil shales high in paraffin in central Utah.
- OLIVEIRA, E. P. DE, Sobre alguns affloramentos de carvão no Parana: *Escola de minas de Ouro Preto Annaes*, No. 9, pp. 91-94, 1907; reviewed in *Inst. Min. Eng. [London] Trans.*, vol. 36, p. 759, 1907. On Carboniferous oil shale near Canoinhas and Iraty.
- PAYEN, ANSELME, Rapport * * * au nom du Comité des arts chimiques, sur divers ouvrages en mastic bitumineux, de MM. Pillot et Eyquem: *Soc. encour. bull.*, vol. 23, pp. 169-173, 1824.
- Note sur le schiste bitumineux et le lignite d'Ardres: *Annales chimie et phys.* 2d ser., vol. 29, p. 335, 1825.

- PETRIE, J. M., The mineral oil from the torbanite of New South Wales: Soc. Chem. Ind. Jour., vol. 24, pp. 996-1002, 1905. Describes torbanite and oils obtained by its distillation.
- REDWOOD, BOVERTON, A treatise on petroleum, 3 vols., 3d ed., London, 1913. Vol. 2, pp. 83-139, gives general treatise on oil shale, its occurrence and development, with a history of the industry. Vol. 3 contains an extensive bibliography of petroleum and oil shales.
- REDWOOD, BOVERTON, and TOPLEY, WILLIAM, Report on the Riacho Doce and Camaragibe shale deposits on the coast of Brazil near Maceio, London, 1891.
- REDWOOD, I. I., A practical treatise on mineral oils and their by-products, London and New York, 1898. Includes a short history of the Scotch shale-oil industry, the geologic and geographic distribution of Scotch shales, recovery of acid and soda used in oil refining, and a list of patents relating to mineral oils.
- REISSIG, W. [Analysis of shale-oil gas]: Jour. Gasbeleuchtung, Band 5, pp. 131-133, 1862.
- RICCIARDI, LEONARDO, Ricerche chimiche sopra una lignite e alcuni scisti bituminiferi di Giffoni, Valle e Piana, provincia di Salerno: Accad. gioenia sci. nat. Atti, 3d ser., vol. 16, pp. 123-128, 1882.
- ROWAN, F. J., On the physical conditions existing in shale-distilling retorts: Soc. Chem. Ind. Jour., vol. 10, pp. 436-443, 1891. Good discussion of temperatures obtained in different shale retorts and their effect on the products.
- ROYER, E., Essai sur la constitution chimique de l'huile de schiste: Soc. sci. phys. nat. Bordeaux Mém., vol. 2, pp. 285-292, 1862.
- ST. ÈVRE, ÉDOUARD, Note sur divers hydrocarbures provenant de l'huile de schiste: Compt. Rend., vol. 29, p. 339, 1849.
- SELIGUE, —, Emploi de l'huile extraite des schistes bitumineux pour la fabrication du gaz d'éclairage: Compt. Rend., vol. 4, pp. 969, 970, 1837.
- Procédé pour la fabrication d'un gaz d'éclairage au moyen des huiles provenant de la distillation de certains schistes bitumineux: Compt. Rend., vol. 10, pp. 861-865, 1840.
- SELWYN-BROWN, ARTHUR, Fuel oil from shale: Eng. Mag., vol. 50, pp. 913-920, 1916. Good general description of oil shale and its possibility as a source of oil.
- SIMONIN, L., Sur les schistes bitumineux de Vagnas (Ardèche): Compt. Rend., vol. 64, pp. 1138-1185, 1867. Brief description of the occurrence and distillation of oil shale in France.
- SKEY, WILLIAM, Oil shales of New South Wales: New Zealand Colonial Mus. Lab. Ann. Rept., vol. 23, p. 50; vol. 25, p. 56; vol. 31, p. 10. Gives analyses of oil shales.
- STEUART, D. R., On the occurrence of petroleum in a shale mine at Broxburn: Soc. Chem. Ind. Jour., vol. 6, pp. 128-130, 352, 1887. Notes semisolid oil (specific gravity 0.842) from bore hole 600 feet deep in vicinity of oil-shale beds.
- Refining shale oil: Soc. Chem. Ind. Jour., vol. 18, pp. 248, 249, 1899. Describes refining processes.
- The chemistry of the oil shales: The oil shales of the Lothians, 1st ed., pt. 3, pp. 133-188, Scotland Geol. Survey Mem., 1906; 2d ed., pt. 3, pp. 136-194, 1912. Gives history of oil-shale industry and detailed treatise on the chemistry of the oil shales and statistics of production.
- The shale-oil industry of Scotland: Econ. Geology, vol. 3, pp. 573-598, 1908. Discusses geology and methods of distillation of oil shales of Scotland.
- STONE, G. H., Asphaltum of Utah and Colorado: Am. Jour. Sci., 3d ser., vol. 42, pp. 148-159, 1891. Describes oil shales of Colorado and Utah and discusses their possible relation to the origin of asphalt veins.

- TAYLOR, ANDREW, On the bituminous shales of Linlithgowshire and Edinburghshire: Roy. Phys. Soc. Edinburgh Proc., vol. 3, pp. 16-24, 1867. Discusses geologic relations of Scotch oil shales.
- Scotch and Welsh mineral-oil trades: Edinburgh Geol. Soc. Trans., vol. 1, pp. 19, 20, 1868. Mentions experiments by Clayton in 1728-29 distilling carbonaceous minerals at low red heat.
- On bitumen, oil shales, and oil coals: Edinburgh Geol. Soc. Trans., vol. 2, pp. 187-189, 1873. Brief mention of Scotch oil shales.
- THOMPSON, A. B., Petroleum mining, pp. 118-121, London, 1910. Short general discussion of oil shales.
- TINKLER, C. K., and CHALLENGER, F., The chemistry of petroleum and its substitutes, pp. 182-192, London, Crosby Lockwood & Son, 1915. Describes briefly the distillation of oil shales and the refining of the products.
- VALENTINE, G., A carbonaceous mineral or oil shale from Brazil: South Wales Inst. Eng. Proc., vol. 17, pp. 20-28, 1890. Describes oil shale from Bahia, Brazil, and theorizes as to its origin.
- VOHL, E. H. L., Ueber Destillationsproducte des Blätterschiefers und der Braunkohle Jour. prakt. Chemie, Band 67, pp. 418-420, 1856.
- Ueber die Producte der trockenen Destillation des Rheinischen Blätterschiefers (Schiste bitumineux) der Sächsischen sowie der Thüringischen Braunkohle, und die Anwendung derselben als Beleuchtungsmaterialien: Annalen Chemie Pharm., Band 68, pp. 504-508, 1856; Band 75, pp. 289-298, 1858.
- WAGENMANN, P., Ueber neue, in Schottland vorkommende Rohmaterialien zur Photogen- und Paraffinfabrication: Polytech. Jour., Band 151, pp. 116-119; Band 152, p. 113, 1859.
- WEED, R., Oil shales in Scotland: Iron and Coal Trade Rev., vol. 79, pp. 205, 206, 1909.
- WILLIAMS, C. G., On the presence of pyridine in naphtha among the volatile bases in the naphtha from the bituminous shale of Dorsetshire: Philos. Mag., 4th ser., vol. 8, pp. 209-212, 1854. Describes chemical work on pyridine in naphtha of shale oil of Scotland.
- On the volatile bases produced by destructive distillation of the bituminous shale of Dorsetshire. Chem Soc. Jour., vol. 7, pp. 97-107, 1854. Gives analysis of shale oil.
- WINDAKIEWICZ, E., Ueber die Wichtigkeit des Vorkommens von bituminösem Schiefer in Galizien: Oesterr. Zeitschr. Berg- u. Huttenw., Band 23, p. 196, 1875.
- WOODRUFF, E. G., and DAY, D. T., Oil shales of northwestern Colorado and northeastern Utah: U. S. Geol. Survey Bull. 581, pp. 1-21, 1914. Describe occurrence and geology of oil shales in Colorado and Utah and give results of distillation tests.
- WRIGHT, W. J., Geology of the Moncton map area, New Brunswick: Canada Geol. Survey Dept. Mines Summary Rept., 1913, pp. 223-227, 1914. Describes oil shales of Albert area and gives distillation tests.
- YOUNG, G. A., Geology of the Moncton map area, Westmoreland and Albert counties, New Brunswick: Canada Geol. Survey Summary Rept., 1911, p. 316, 1912. Mentions oil shale and attempted development near St. Johns.