

**Table 2.** Results of sediment contamination literature searches conducted from May 15 to June 6, 2014, for listed contaminants in Barnegat Bay-Little Egg Harbor, Newark Bay-New York Harbor-Raritan Bay-Passaic River Estuary, Jamaica Bay, Great South Bay, South Oyster Bay, and Peconic Bay, New Jersey and New York.

[Refer to table 1 for terms used in literature search. PAHs, polycyclic aromatic hydrocarbons; PBDEs, polybrominated diphenyl ethers; PCBs, polychlorinated biphenyls]

Contaminants and contaminant indicators	Barnegat Bay–Little Egg Harbor		Newark Bay–New York Harbor–Raritan Bay–Passaic River Estuary		Jamaica Bay		Great South Bay		South Oyster Bay		Peconic Bay	
	Search results	Relevant articles	Search results	Relevant articles	Search results	Relevant articles	Search results	Relevant articles	Search results	Relevant articles	Search results	Relevant articles
Antibiotics	31	0	32	2	149	17	39	1	0	0	61	0
Detergents	53	1	105	1	149	21	85	0	2	0	18	0
Fragrances	6	0	13	0	40	0	18	0	0	0	11	0
Hormones	58	0	153	7	134	19	57	0	2	0	17	0
PAHs	60	6	478	19	155	17	31	0	0	0	15	0
PBDEs/flamm retardants	14	1	66	3	18	1	3	0	1	0	0	0
PCBs	171	8	826	42	273	26	94	0	6	0	44	0
Personal-care products	112	1	127	2	65	13	88	0	0	0	45	0
Pesticides	213	4	580	69	308	26	214	0	9	0	74	0
Pharmaceuticals	42	1	120	0	60	7	67	0	3	0	39	0
Sediment toxicity	449	5	900	34	559	25	438	3	23	1	164	0
Trace metals	329	6	610	34	412	21	365	8	29	1	128	3
Wastewater indicators	266	1	385	18	2	0	295	0	16	0	105	0

