

DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY

SURFICIAL GEOLOGIC MAP OF THE
LAS VEGAS 1° x 2° QUADRANGLE, NEVADA, ARIZONA,
CALIFORNIA, AND UTAH

By


David L. Weide

Open-File Report 82-706

1982

This report is preliminary and has not been reviewed for conformity with U. S. Geological Survey editorial standards and stratigraphic nomenclature.

CORRELATION OF SURFICIAL MAP UNITS



DESCRIPTION OF SURFICIAL MAP UNITS
EOLIAN DEPOSITS

Windblown sand deposits in this area are reworked from alluvium in stream channels and (or) older eolian deposits. Commonly sand deposits occur as thin sheets and do not form dunes, suggesting the lack of a prevailing, unidirectional wind. Both active and stabilized sand deposits support some vegetation, but active deposits are marked by very sparse patches of immature soil and numerous wind-deflated hollows.

DUNE DEPOSITS

- Qda Active (Holocene)--Fine to medium quartz sand in thin sheets and very low dunes without significant vegetation covering. Locally may exceed 6 m in thickness
- Qds Stabilized (Holocene to late Pleistocene)--Fine to medium quartz sand in deposits stabilized by vegetation. Locally may include deposits formed of sand-size clay aggregates. Commonly includes inceptisols. As much as 6 m in thickness

LACUSTRINE AND PALUDAL DEPOSITS

Lacustrine deposits are subdivided into two major groups, clay and saline playa deposits that occur in closed basins occupied by ephemeral lakes, and lacustrine clay deposits that represent eroded remnants of older lake beds now exposed in areas not periodically inundated. Playa sediments overlie older deposits of interstratified clay and saline minerals that may exceed 400 m in thickness. Although no cores are available from this area, data from similar deposits in a similar setting at Searles Lake, Calif., suggest intermittent lacustrine deposition for at least 120,000 years (Smith, 1979, p. 113).

- Qm MARSH DEPOSITS (HOLOCENE)--Interbedded organic-rich silt and clay containing abundant plant fragments. Deposited in intermittent to perennial bodies of fresh, brackish, or saline water that support aquatic vegetation. May reach a maximum of 15 m in thickness
- Qps SALINE PLAYA DEPOSITS (HOLOCENE)--Silty sand capped by crusts of mud mixed with halite and gypsum. Salt-rich units as much as 8 m in thickness
- Qpc CLAY PLAYA DEPOSITS (HOLOCENE TO LATE PLEISTOCENE)--Thin-bedded silt and clay deposited by ephemeral flooding on playas. Commonly forms flat surfaces containing desiccation polygons and patches of easily deflated mud curls. Thickness 5-25 cm
- Qlc LAKE CLAY (LATE PLEISTOCENE TO PLEISTOCENE)--Thin-bedded silt and clay having well-developed stratification. Commonly restricted to areas inundated by Pleistocene lakes. Easily eroded to badland topography having as much as 15 m of relief. Contains numerous buried paleosols and volcanic-ash layers. Commonly marked by buried desiccation features and sand-filled paleochannels. As much as 40 m in thickness

ALLUVIAL DEPOSITS IN CHANNELS

In the Las Vegas quadrangle, major alluvial deposits are restricted to through-flowing streams that are perennial, such as the Virgin and Muddy Rivers, or streams that carried large volumes of water during the late Pleistocene, such as the White River or Meadow Valley Wash. In the eastern part of the quadrangle, channel and terrace gravels reflect earlier periods of increased stream capacity. Along drainages of the Muddy and White Rivers and Meadow Valley Wash, for instance, the lithology of the gravels does not reflect adjacent bedrock but is composed of material derived far upstream. In addition to geomorphic evidence, younger and older terrace gravels may be mapped separately because older deposits do not contain limestone or dolomite components.

Qal CHANNEL ALLUVIUM (HOLOCENE)--Silt, fine to coarse sand, and gravel in beds and along flood plains of ephemeral or permanent streams. Intermittently deposited as bars or other channel deposits, then eroded and redeposited. Unit occupies almost all drainage channels, but only the larger deposits are shown. Along major streams, such as the Virgin River, alluvium is inset in older flood-plain and lacustrine deposits. Alluvial units contain numerous cut-and-fill lenses of silt, sand, or gravel. Locally may exceed 20 m in thickness

TERRACE GRAVELS

Qtg Younger terrace gravels (Holocene to late Pleistocene)--Gray well-rounded gravel composed of volcanic rock, overlain by patches of white very fine sand and silt, and red angular calcareous gravel. Matrix is medium to coarse, angular to subangular sand composed of equal amounts of volcanic and calcareous detritus that commonly is cemented by 2 m of white powdery calcium carbonate. Unit caps terraces 18-25 m above present drainages. Thickness 8-21 m

Qtg₁ Older terrace gravels (late to middle Pleistocene)--Well-rounded gravel composed of volcanic rock and having no overlying sand or calcareous gravel. Matrix composed of medium to fine rounded volcanic sand. Upper 1-1.5 m very strongly cemented with calcium carbonate, in places having the texture, grain size, and degree of induration of dense limestone. Unit forms terraces 60-70 m above present drainages. Thickness 4-10 m

Qvf VALLEY FILL (LATE TO MIDDLE PLEISTOCENE)--Mixture of interbedded lake silt, fluvial sand and gravel, buried eolian deposits, and sand-to silt-size material deposited at the distal edges of alluvial fans. Commonly does not exhibit well-defined stratification but locally may contain buried paleosols and lenses of volcanic ash. Surface characterized by well-developed soils as thick as 1.5 m. Ranges from 10 to 60 m in thickness

ALLUVIAL FAN DEPOSITS

In this quadrangle, alluvial-fan deposits are subdivided into two broad categories, unconsolidated and cemented. Cemented fan deposits are further grouped by degree and amount of calcium carbonate (caliche) cementation. Degree of cementation is a highly subjective criterion because it results from many interacting causes including lithology of the contributing bedrock, ground-water and pedogenetic factors, climatic history, and time. Nevertheless, the practice adopted when separating fan units in the field suggests a relative age classification based on the number of cemented horizons within any given fan deposit. For instance, a fan deposit containing three cemented horizons, of probable pedogenetic origin, separated by layers of unconsolidated fan debris, is assumed to have had a longer depositional history than a fan having caliche cementation only on its surface. On this map, where sequential and superimposed (or inset) fan deposits that contain caliche can be recognized, they are numbered from youngest to oldest, Qfc, Qfc₁, Qfc₂, and Qfc₃. No data leading to an absolute fan chronology are presently available in this area. Where approximate ages are suggested, they are inferred from information collected at the Nevada Test Site (Hoover and others, 1981).

- Qfu UNCONSOLIDATED FAN DEPOSITS (HOLOCENE TO LATE PLEISTOCENE)--Mixture of uncemented, well-rounded, poorly sorted sand- to boulder-size material. Surface characterized by large areas of gravel pavement interspersed with pockets of weakly developed sandy soil. Particle size of surface gravels decreases downslope where fan deposits merge into valley-fill units (Qvf). Although surficial pavements and soils range in thickness from 0.5 to 3.0 m, the unconsolidated fan deposits commonly exceed 250 m in total thickness
- Qfc CEMENTED FAN DEPOSITS (HOLOCENE TO LATE PLEISTOCENE)--Mixture of sand- to boulder-size material containing as much as 1 m of surficial calcium carbonate cementation. Surfaces commonly are cemented with stage II to III carbonate (caliche partly coats pebbles or fills interstices between pebbles). Surfaces are highly resistant to erosion but support widely spaced, headward-cutting arroyos as deep as 10 m. Age of these deposits is poorly known, but deposition may have begun at least by middle Pleistocene. Deposit beneath calcium carbonate cement may exceed 100 m in thickness

- Qfc₁ Well-cemented deposits (late to middle Pleistocene)--Gray to red, well-rounded pebble to cobble gravel composed of calcareous, volcanic, and quartzite clasts in a matrix of coarse to medium angular to subrounded sand composed of calcareous and volcanic detritus. Commonly contains a single, well-developed caliche-cemented horizon within 1 m of the surface. Caliche commonly ranges from 1 to 2 m in thickness and attains stage III or IV in development (interstices between pebbles filled with carbonate, upper layers of carbonate massive, relatively pebble free, and having the texture, structure, and degree of induration of fine-grained limestone). Fans are undated but may have begun to form during the early or middle Pleistocene. Surficial cemented horizon ranges from 1 to 3 m in thickness. Deposit may exceed 150 m in thickness
- Qfc₂ Multiple cementation layers (middle Pleistocene to pre-Pleistocene)--Gray to red, well-rounded pebble to cobble gravel composed of calcareous, volcanic, and quartzite clasts in a matrix of coarse to medium sand comprised of calcareous and volcanic sand. Mapped on the basis of multiple carbonate-cement (caliche) horizons, two to three in number, 1-3 m in thickness. Upper caliche horizon commonly stage II to III development Lower caliche horizons commonly stage III or IV development. Age unknown, but formation of fan deposits may have begun during the late Tertiary. Surfaces commonly support vesicular, silty soils and desert-pavement deposits as much as 0.5 m thick. The caliche-cemented part ranges from 5 to 10 m in thickness. Deposit may exceed 150 m in total thickness
- Qfc₃ Multiple cementation layers, very well cemented (early Pleistocene to pre-Pleistocene)--Gray to red, well-rounded pebble to cobble gravel composed of calcareous, volcanic, and quartzite clasts in a matrix of coarse to medium sand composed of calcareous and volcanic detritus. Mapped on the basis of multiple superimposed caliche horizons that are 2-5 m thick and commonly attain stage III or IV development. These fan deposits appear to be the oldest in this quadrangle and may have formed between late Tertiary and early Pleistocene time. Surfaces commonly stripped of soil and (or) desert-pavement deposits, exposing large areas of caliche. Caliche-cemented part ranges in thickness from 10 to 40 m. Deposit may exceed 150 m in total thickness

MASS MOVEMENT DEPOSITS

- Qc COLLUVIUM (HOLOCENE TO PRE-PLEISTOCENE)--Cobble- to boulder-size angular fragments of underlying bedrock on slopes greater than 5°. Matrix is generally silty; may exhibit weakly developed soil as thick as 1 m. Subject to sheet erosion and rill wash. Marked in many places by small bedrock outcrops upslope from thick lobes and patches of cobble- to boulder-size concentrations without sand or silt matrix. Commonly cemented by carbonate throughout the quadrangle regardless of the lithology of the underlying bedrock. Unit includes sheetwash alluvium, discontinuous patches of talus and scree on steep slopes, and solifluction deposits at elevations above 8,500 ft (2600 m). Rarely exceeds 5 m in thickness
- Q1s LANDSLIDE DEPOSITS (HOLOCENE TO PRE-PLEISTOCENE)--Angular cobbles to boulders having poorly sorted sand to pebble matrix, unconsolidated. Unit includes other mass-movement deposits such as earthflows and rockfalls. Where landslide deposits exceed 1 km² in area, they generally include patches of other colluvial and residual surficial deposits that have been moved downslope. Thickness 1-30 m

RESIDUAL DEPOSITS

- Qr RESIDUUM DEPOSITS (HOLOCENE TO PRE-PLEISTOCENE)--Sparse angular cobbles of underlying bedrock in a silty matrix. May contain pockets of clay-rich, well-developed soil as thick as 1 m. Commonly forms on slopes of less than 5° and reflects chemical and mineralogical properties of the underlying bedrock. In the map area, residuum deposits include both solution residuum derived from calcareous bedrock, and decomposition residuum derived from silicious bedrock. Where developed on basic volcanic rocks, residuum is rich in smectitic swelling clays. Where bedrock is specifically basaltic in composition (unit Rv1), residuum is commonly indurated with a hard silica duricrust. Where developed on calcareous rock, residuum may be cemented with caliche. Rarely exceeds 3 m in thickness
- Qpg PEDIMENT GRAVEL (LATE PLEISTOCENE TO PRE-PLEISTOCENE)--Subrounded to angular pebbles to cobbles in a matrix of medium to coarse quartz sand. In east half of quadrangle, unit is developed on beveled surfaces adjacent to mountain ranges formed of intrusive igneous or metamorphic rock. Commonly has a well-developed desert pavement and desert varnish. Age is unknown, but unit may be at least as old as the older consolidated alluvial fans (units Qfc₂ and Qfc₃). As much as 4 m in thickness

BEDROCK UNITS

Rock units pose a particularly vexing problem to the constructor of a map of surficial deposits, especially in arid areas marked by steep slopes, numerous outcrops, and a lack of weathered debris. For the purposes of this map, an area was designated as a rock unit if structural information, such as an attitude on a bedding plane, joint, fracture, or fault, could be obtained within a circle of 100-ft (30-m) radius from any given point.

In general, areas designated as "rock" range from 85 to 100 percent outcrop. Any such outcrop area contains numerous patches of colluvium, residuum, talus, or scree, and small areas of coarse entisols and other immature azonal soils developed under discontinuous, highly variable plant communities.

The lithology of parent bedrock influences surficial deposits in terms of their soil development, ground-water characteristics, and engineering properties. On this map a simple classification is used to show the dominant lithology of a bedrock outcrop.

ROCK--Lithologic nomenclature follows that of Stewart and Carlson (1978)

Rv1 Basaltic volcanic

Rv2 Andesitic volcanic

Rv3 Rhyolitic volcanic

Rs Siliceous sedimentary rock--Sandstone, shale, and quartzite

Rc Calcareous rocks--Including limestone, dolomite, and marble

Rpm Plutonic, metamorphic, and ultramafic rocks

CONTACT

FAULT CUTTING QUATERNARY DEPOSITS--Bar and ball on downthrown side
where sense of movement is known

PLEISTOCENE VERTEBRATE-FOSSIL LOCALITY

AREA DISTURBED BY SURFACE MINING

METHOD OF COMPILATION

This map constitutes a portion of the Quaternary Geologic Map of the Mount Whitney 4° x 6° Quadrangle, in preparation for the National Quaternary Atlas of the United States at a scale of 1:1,000,000. The map units shown on the Las Vegas 1° by 2° quadrangle map are scaled for reduction from 1:250,000 to 1:1,000,000 and are, of necessity, generalized. Surficial deposits that form a linear pattern, for instance, are limited by cartographic constraints to a minimum width of 0.5 mile (800 m); those that occur as patchy deposits are restricted to a minimum area of 1 mi² (2.6 km²). Because of these size limitations, many specific deposits and groups of deposits are not shown or are generalized to reflect the dominant component. Use of this 1:250,000-scale map for specific information about a particular locality must be supplemented by field investigation.

The map was derived from three primary sources: (1) published geologic maps, (2) recent Landsat images corrected for a scale of 1:250,000, and (3) fieldwork. The first step was to compare specific surficial deposits that appeared on published maps with Landsat imagery. Such deposits were then visited and described. In areas where little or no published information was available, the Landsat signature (color hue, tone, and intensity, albedo or reflectivity, and degree of surface roughness) was compared with a known locality and a tentative unit description assigned to the deposit. The area was then field checked and the Landsat description verified or revised. Large areas, such as coalesced alluvial fans covering as much as 75 mi², were spot checked. If the signature was constant over the entire surface, the deposit was assumed to be uniform throughout and equivalent to adjacent areas that displayed the same signature.

Relative age determinations for deposits, such as alluvial fans or terrace gravels, were first made from tonal differences on Landsat images. These were then verified or revised in the field using such criteria as degree of calcium carbonate (caliche) development, soil profiles, and extent of weathering.

On this map, as is the case for the Quaternary Atlas of the United States in general, certain surficial deposits imply specific processes and materials. For the purposes of this map, "solifluction deposit" is a general term for material transported and deposited by viscous flow of unconsolidated debris saturated with water. Colluvium is material transported and deposited by mass-wasting processes, chiefly by creep. Sheetwash alluvium is material transported and deposited by running water, chiefly by sheetflow or rill wash. Decomposition residuum is material derived in place by decomposition of bedrock, and has not been moved laterally. Solution residuum is residuum derived primarily by solution of calcareous rock.

SOURCES OF INFORMATION

- Anderson, R. E., 1974, Large-magnitude late Tertiary strike-slip faulting north of Lake Mead, Nevada: U.S. Geological Survey Professional Paper 794, 18 p.
- Bell, J. W., 1981, Subsidence in Las Vegas Valley, Nevada: Nevada Bureau of Mines and Geology Bulletin 96, 21 p.
- Bell, J. W., and Smith, E. I., 1980, Geologic map of the Henderson quadrangle, Nevada: Nevada Bureau of Mines and Geology Map 67, scale 1:24,000.
- Bingler, E. C., 1977, Geologic map of the Las Vegas SE quadrangle, Clark County, Nevada: Nevada Bureau of Mines and Geology Las Vegas Area Urban Geology Folio, scale 1:24,000.
- Bohannon, R. G., 1977, Geologic map and sections of the Valley of Fire region, North Muddy Mountains, Nevada: U. S. Geological Survey Miscellaneous Field Studies Map MF-849, scale 1:25,000.
- _____, 1978, Preliminary geologic map and sections, White Basin and Bitter Springs Valley, Muddy Mountains, Clark County, Nevada: U.S. Geological Survey Miscellaneous Field Studies Map MF-922, scale 1:25,000.
- _____, 1978, Preliminary geologic map of the Las Vegas 1° x 2° degree quadrangle, Nevada: U.S. Geological Survey Open-File Report 78-670, scale 1:250,000.
- Burchfiel, B. C., and others, 1974, Geology of the Spring Mountains, Nevada: Geological Society of America Bulletin, v. 85, p. 1013-1022.
- Gardner, L. R., 1972, Pediments and terraces along the Moapa Valley, Clark County, Nevada: Geological Society of America Bulletin, v. 83, p. 3479-3486.
- Hoover, D. L., Swadley, W. C., and Gordon, A. J., 1981, Correlation characteristics of surficial deposits with a description of surficial stratigraphy in the Nevada Test Site region: U.S. Geological Survey Open-File Report 81-512, 44 p.
- Longwell, C. R., Pampeyan, E. H., Bowyer, Ben, and Roberts, R. J., 1965, Geology and mineral deposits of Clark County, Nevada: Nevada Bureau of Mines and Geology Bulletin 62, 218 p.
- Maxey, G. B., and Jameson, C. H., 1948, Geology and water resources of Las Vegas, Pahrump, and Indian Springs Valleys, Clark and Nye Counties, Nevada: Nevada State Engineer Water Resources Bulletin 5, 124 p.
- Miffelen, M. D., and Wheat, M. M., 1979, Pluvial lakes and estimated pluvial climates of Nevada: Nevada Bureau of Mines and Geology Bulletin 94, 47 p.
- Smith, G. I., 1979, Subsurface stratigraphy and geochemistry of late Quaternary evaporites, Searles Lake, California: U.S. Geological Survey Professional Paper 1043, 130 p.
- Stewart, J. H., 1980, Geology of Nevada--A discussion to accompany the Geologic Map of Nevada: Nevada Bureau of Mines and Geology Special Publication 4, 136 p.
- Stewart, J. H., and Carlson, J. E., 1978, Geologic Map of Nevada: U.S. Geological Survey map, scale 1:500,000.
- Tschanz, C. M., and Pampeyan, E. H., 1970, Geology and mineral deposits of Lincoln County, Nevada: Nevada Bureau of Mines and Geology Bulletin 73, 187 p.