

DEPARTMENT OF INTERIOR  
U.S. GEOLOGICAL SURVEY

**Latest Cretaceous and Cenozoic magmatism in mainland Alaska**

by

Elizabeth J. Moll-Stalcup<sup>1</sup>

Open-File Report 90-84

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards and stratigraphic nomenclature. Any use of trade names is for descriptive purposes only and does not imply endorsement by the U.S. Geological Survey.

<sup>1</sup>U. S. Geological Survey  
Menlo Park, CA 94025

## CONTENTS

Introduction .....	2
Latest Cretaceous and early Tertiary magmatism.....	3
Alaska Range-Talkeetna Mountains belt.....	3
Petrogenesis.....	5
Kuskokwim Mountains belt.....	6
Petrogenesis.....	8
Yukon-Kanuti belt.....	12
Petrogenesis.....	13
Late Cretaceous and early Tertiary tectonic implications.....	15
Middle Tertiary magmatism.....	20
South-central Alaska.....	21
Western interior Alaska.....	21
Petrogenesis.....	37
Late Cenozoic volcanism.....	23
The Bering Sea region.....	23
Petrogenesis.....	26
Eastern and central interior Alaska.....	31
Late Cenozoic tectonic implications.....	32
Summary and conclusions.....	34
Acknowledgements.....	36
References cited.....	37
Tables.....	53
Figures .....	66

## INTRODUCTION

Continental Alaska has been the site of widespread magmatism throughout much of the late Mesozoic and Cenozoic, but until recently most of this magmatism was unrecognized due to the lack of modern geologic maps or isotopic age data for large tracts of Alaska. Although parts remain unmapped, progress in reconnaissance mapping and dating have enabled workers to identify major late Mesozoic and Cenozoic magmatic provinces outside the well-known Aleutian arc and to speculate as to their tectonic implications and origin (Wallace and Engebretson, 1984).

This paper defines major Late Cretaceous and Cenozoic magmatic provinces in Alaska outside the Aleutian arc (Marsh, in press; Miller and Richter, in press) and southeast Alaska (Brew, in press) and discusses their distribution, age, petrology, and tectonic implications. The paper will eventually be published as a chapter in the DNAG volume on Alaskan geology (Plafker et al., in preparation), but is being open-filed because of long publication delays. The available data suggest that Late Cretaceous and Cenozoic magmatism in continental Alaska can be roughly divided into three periods: (1) latest Cretaceous and early Tertiary (73 to 50 Ma), (2) middle Tertiary (43 to 37 Ma), and (3) late Tertiary and Quaternary (6 Ma to the present). Late Cretaceous and early Tertiary calc-alkalic volcanism and plutonism was widespread over much of western, central, and southern Alaska and on the Bering Sea shelf. Middle Tertiary magmatism was characterized by the eruption of small volumes of calc-alkalic rocks in interior Alaska, contemporaneous with the inception of a major pulse of magmatism in the Aleutian arc. Late Tertiary and Quaternary volcanism has been characterized by the eruption of voluminous basaltic magma at numerous sites along the western margin of Alaska and on the Bering Sea shelf.

A map showing Cenozoic volcanic and plutonic rocks for the entire state at a scale of 1:2.5 million (Moll-Stalcup and others, in press) will be available as part of the DNAG volume. Place names used in this chapter appear on that map, but are also shown on 1:250,000 scale maps of the area published by the U. S. Geological Survey. Although only major belts of regional significance are discussed in this chapter, tables summarizing age, lithologic, and chemical data for all the Late Cretaceous to Quaternary volcanic and plutonic rocks outside the Aleutian arc and southeast Alaska are found on Table 1.

Nomenclature used in this paper generally follows that of Streckeisen (1980), Gill (1981) and Morrison (1980). On an anhydrous basis, basalts have less than 53% SiO<sub>2</sub>, andesites have 53 to 63% SiO<sub>2</sub>, dacites have 63 to 70% SiO<sub>2</sub>, and rhyolites have more than 70% SiO<sub>2</sub>. Fe<sub>2</sub>O<sub>3</sub>/FeO was set to 0.15 for the late Cenozoic basalts in order to calculate Mg numbers and normative mineralogies. The late Cenozoic rocks are classified as follows: Basalt having normative nepheline is called alkali basalt or alkali olivine basalt if it contains more than 10% normative olivine. Basalt having normative hypersthene is called tholeiite or olivine tholeiite if it contains more than 10% normative olivine. Basanites have 10 to 20% normative nepheline; nephelinites have more than 20% normative nepheline. Hawaiites have more than 5% total alkalis

( $\text{Na}_2\text{O} + \text{K}_2\text{O}$ ) and less than 5% MgO. The Late Cretaceous, early Tertiary, and middle Tertiary suites are classified using the Peacock index and then divided into low-, moderate-, or high-K after Gill (1981) or shoshonitic after Morrison (1980). An upper limit for  $\text{Fe}_2\text{O}_3$  was set by the formula  $\% \text{Fe}_2\text{O}_3 = \% \text{TiO}_2 + 1.5$  (after Irvine and Baragar, 1971). All ages were obtained by K/Ar methods unless otherwise noted.

All the Late Cretaceous and early Tertiary volcanic and plutonic rocks, and some of the middle Tertiary rocks are hydrothermally altered and weathered, and I therefore have relied heavily on trace elements for interpretation of the geochemical data. Typical plutonic samples have about 1%  $\text{H}_2\text{O}_t$  and 0.2%  $\text{CO}_2$  and typical volcanic rocks have 1-4%  $\text{H}_2\text{O}_t$  and 0.2%  $\text{CO}_2$ . Pyroxenes and feldspars are generally fresh, but olivine, biotite, and hornblende are altered in some samples. Alteration of rocks from five volcanic fields, which are typical of much of the magmatic province are described in more detail in Moll-Stalcup (1987). The late Cenozoic volcanic rocks are very fresh, and even olivine is well preserved in most samples.

### LATEST CRETACEOUS AND EARLY TERTIARY MAGMATISM

Latest Cretaceous and early Tertiary magmatic activity occurred in a vast region of Alaska stretching from the southern continental margin north to the Arctic Circle and west to the Bering Sea shelf (Fig. 1)--possibly extending as far west as contemporaneous magmatic belts in eastern Siberia. Hudson (1979) and Wallace and Engebretson (1984) group the widespread volcanic and plutonic rocks in southern, western, and central Alaska into volcano-plutonic belts. From south to north these are: the Sanak-Baranof belt, the Alaska Range-Talkeetna Mountains belt, and the Kuskokwim Mountains belt. An additional, previously unnamed, belt occurs farther to the northwest and is herein named the Yukon-Kanuti belt. Little-known volcanic and plutonic rocks of latest Cretaceous and early Tertiary age also occur in the Yukon-Tanana area of east-central Alaska (Foster and others, in press), but their correlation and tectonic affinities are unknown. Data on the Late Cretaceous and early Tertiary rocks in the Yukon-Tanana upland are summarized in Table 1, but those rocks are not discussed further. The Sanak-Baranof belt, which consists of early Tertiary granitic plutons emplaced into the Gulf of Alaska accretionary wedge, extends along the southern margin of Alaska to Baranof Island in southeast Alaska and is described by Hudson (in press).

### ALASKA RANGE-TALKEETNA MOUNTAINS BELT

The Alaska Range-Talkeetna Mountains belt consists of numerous coalescing plutons and subordinate volcanic rocks that extend in a broad belt, about 150 km wide, from the central Alaska Range, west and south to the Iliamna Lake region (Fig. 1). Most of the rocks occur south of the Denali fault, except for a few small bodies north of the fault near Farewell Lake and the Tonzona River (Reed and Nelson, 1980). Plutonic and volcanic rocks in the Alaska Range-Talkeetna Mountains belt intrude and overlie the Dillinger, Kahiltna and Peninsular terranes as well as a number of smaller

terrane in the Mt. McKinley area (Jones and others, in press). Aeromagnetic anomalies on the Bering Sea Shelf (Godson, 1984; Cooper and others, 1986) and the presence of compositionally similar contemporaneous volcanic rocks on St. Matthew Island (Patton and others, 1975) suggest that the belt may continue southwest of Iliamna Lake under Bristol Bay, curving west and north along the submerged continental shelf of Alaska (Fig. 1).

Plutonic activity in the Alaska Range-Talkeetna Mountains belt is divided into an early stage (75-60 Ma), which occurred chiefly on the the south-southeast flank of the belt; and a late stage (65 to 50 Ma), which occurred chiefly on the north-northwest flank of the belt. The early stage consists of dominantly intermediate to felsic plutons and includes many of the rocks of Summit Lake (Reed and Lanphere, 1972), the Mount Susitna pluton (Magoon and others, 1976), and a large tonalite pluton in the southern Talkeetna Mountains (Csejtey, 1974). The late stage consists generally of felsic plutons and includes the quartz monzonite of Tired Pup, the Crystal Creek sequence, the McKinley sequence (Reed and Lanphere, 1972), and numerous granitic plutons in the northern Talkeetna Mountains and the adjacent southern Alaska Range (Csejtey and others, 1978; 1986).

Volcanic rocks in the Talkeetna Mountains consist of several small fields and one large field approximately 90 by 25 km that trends southeast perpendicular to the belt. The main volcanic field is over 1,500 m thick and is composed of rhyolite and dacite stocks, irregular dikes, lenticular flows, and thick pyroclastic rocks at the base grading up into gently dipping interlayered basalt and andesite flows at the top (Csejtey and others, 1978). Three rocks from about midsection give ages of 56.5 to 50.4 Ma, indicating that the lower part of the section is Paleocene and Eocene in age. The stratigraphically high mafic and intermediate flows are thought to be equivalent in age to Miocene lava flows in the Wrangell Mountains (Csejtey, and others, 1978).

The volcanic rocks of the Cantwell Formation crop out north of the Talkeetna Mountains volcanic rocks in the central Alaska Range, covering about 165 km<sup>2</sup> in the eastern part of Mount McKinley National Park. They consist of at least 3,750 m of mostly andesite and rhyolite flows and subordinate basalt flows, felsic pyroclastic rocks, and related intrusive rocks (Gilbert and others, 1976). Gilbert and others considered ages of 60.6, 57.2, and 41.8 Ma to be minimum ages and interpret the formation as Paleocene in age.

Little known volcanic rocks near Lake Clark consist of undivided Paleocene and Eocene volcanic and associated plutonic rocks that crop out discontinuously over more than 3,000 km<sup>2</sup> in the area between Lake Clark and the Mulchatna River (Nelson and others, 1983). The volcanic rocks are 62.7 to 56.2 Ma (Eakin and others, 1978) and 44.4 to 39.7 Ma (Thrupp and Coe, 1986); and adjacent shallow plutons are 71.3 to 60.5 Ma (Nelson and others, 1983). The volcanic rocks are described as rhyolite breccia, lava flows and ash-flow tuffs, and subordinate mafic to intermediate flows; the intrusive rocks as granite, granodiorite, and diorite (Nelson and others, 1983).

The east Susitna batholith (Moll-Stalcup and others, in press) occurs at the east end of the Alaska Range-Talkeetna Mountains belt on the east limb of the bend in the

Denali fault. Although the batholith yields Late Cretaceous and early Tertiary minimum ages, it is not considered to be part of the Alaska Range-Talkeetna Mountains belt because it consists of regionally metamorphosed and penetratively deformed diorite, granodiorite, and quartz monzonite that give a wide range of K/Ar ages (Nokleberg and others, 1982; Table 1), and because it is thought to have been 400 km from its present position at the time of its emplacement (Nokleberg, 1985). Tertiary displacements on regional strike-slip faults along the east side of the bend in the Denali fault, where the east Susitna batholith occurs, are thought to be much greater than displacements on the west side, where most of the Alaska Range-Talkeetna Mountains belt occurs. Because the batholith probably is allochthonous relative to the Alaska Range-Talkeetna Mountains belt and thus is not part of this belt, and because its age is ambiguous, it is not discussed further in this chapter.

**PETROGENESIS:** The calc-alkalic plutons and volcanic rocks in the Alaska Range-Talkeetna Mountains belt are typical of continental-margin arc rocks, and are characterized chemically by low  $\text{TiO}_2$ , moderate  $\text{K}_2\text{O}$  and lack of Fe-enrichment (data from Reed and Lanphere, 1972: 1974; Csejtey, 1976; 1974; Csejtey and others, 1978; Gilbert and others, 1976; Lanphere and Reed, 1985). The early-stage plutons in the Alaska Range are dominantly intermediate to felsic (54.5-70%  $\text{SiO}_2$ ) and are compositionally equivalent to medium-K orogenic andesites and dacites. Plots of  $\text{SiO}_2$  vs.  $\text{Na}_2\text{O}$  and Ca-Na-K distinguish two suites of rocks--a diorite, tonalite, trondhjemite suite similar to the calc-alkalic-trondhjemite suite of southwest Finland (Arth and others, 1978) and a "normal" calc-alkalic suite. I was not able to determine from the published data (above references) whether the two suites are temporally or geographically distinct.

The late-stage plutons in the Alaska Range are generally more felsic and are divided into two groups on the basis of mineralogy and chemistry. One group is a normal calc-alkalic suite similar to the early-stage plutons and consists of the early Tertiary Crystal Creek pluton and numerous small granitic bodies in the northern Alaska Range. The other group is represented by the McKinley sequence and the quartz monzonite of Tired Pup (Lanphere and Reed, 1985; M.A. Lanphere, oral comm., 1984), which consist of siliceous peraluminous granites that plot at minimum-melt compositions on a Q-Ab-Or diagram and have moderately high strontium initial ratios ( $^{87}\text{Sr}/^{86}\text{Sr}=0.7054$  to  $0.7085$ ; Lanphere and Reed, 1985). Lanphere and Reed (1985) interpret the chemical and isotopic data of the McKinley sequence as the result of mixing of mantle-derived magmas with upper Mesozoic flysch into which the plutons were intruded. They believe that this mixing took place in the early Tertiary during the collision between northern Alaska and southern accreted terranes. Paleomagnetic data, however, suggest that the terranes were accreted, by Late Cretaceous time (Hillhouse and Coe, in press). K-Ar ages (Lanphere and Reed, 1985; Reed and Lanphere, 1972: 1974) suggest that these plutons were emplaced at the end of a long period of arc magmatism and that they may simply mark the end of this event. Chondrite-normalized multi-element diagrams for the McKinley sequence show large depletions in Nb and Ta, which are diagnostic of arc magmatism (Fig. 2; Perfit and others, 1980; Gill, 1981; Thompson and others, 1984).

Volcanic rocks having ages between 58 to 50 Ma crop out in the northern Alaska Range and Talkeetna Mountains at the east end of the belt. Limited petrologic data on the lower sequence of the Talkeetna Mountains volcanic rocks suggest they are typical of orogenic belts and are similar in composition to the older Summit Lake plutons. Analyzed rock samples from the Cantwell Formation have higher  $\text{TiO}_2$  and lower  $\text{Al}_2\text{O}_3$  (Gilbert and others, 1976) --both immobile elements--than the other clearly orogenic suites. Granitic plutons in the same region also have ages from 58 to 50 Ma, but their affinities are not known. The lack of muscovite and presence of rare hornblende suggest that they are similar to the Crystal Creek sequence, but the association of some plutons with tin mineralization suggests instead that they may be correlative with the peraluminous McKinley sequence.

The age and composition of the Talkeetna Mountains volcanic rocks suggest that arc volcanism occurred later in the eastern part of the belt than in the western or southern parts. Arc volcanism appears to have migrated gradually north between about 65 Ma and 58 Ma, then east, until it shut off at about 50 Ma.

### KUSKOKWIM MOUNTAINS BELT

The Kuskokwim Mountains belt is a Late Cretaceous and early Tertiary volcanic-plutonic belt that extends along the entire length of the Kuskokwim Mountains for over 800 km, from Bristol Bay to about latitude 64 degrees (Fig. 1). Studies in the Medfra quadrangle (Patton and others, 1980; Moll and others, 1981), in the Iditarod and McGrath quadrangles (M.L. Miller, oral comm., 1986; Bundtzen and Laird, 1982, 1983a, b, c), in the Sleetmute quadrangle area (Robinson and others, 1984; Decker and others, 1984; 1985; 1986; Reifentuhl and others, 1984; 1985), and in the Tikchik Lakes-Bristol Bay region (Hoare and Coonrad, 1978b; Wilson, 1977; Globberman, 1985) have led to the recognition of a volcanic and plutonic belt of Late Cretaceous and early Tertiary age (Moll and Patton, 1982; Wallace and Engebretson, 1984). Compilation of more than 90 K/Ar ages from volcanic and plutonic rocks along the belt (E.J. Moll, unpub. data, 1985) suggests that the main magmatic pulse occurred between 72 Ma and 60 Ma, contemporaneous with the older-stage plutonism in the Alaska Range. It is not yet clear if the Kuskokwim Mountains belt is separate from the Alaska Range-Talkeetna Mountains belt. Much of the area separating the two belts is covered by Quaternary surficial deposits on the north side of the Farewell fault and in the vicinity of the Mulchatna fault. Exposures of plutonic and volcanic rocks in low hills in the Farewell Lake area and on both sides of the Mulchatna fault suggest that the two belts may be continuous. The volcanic and intrusive rocks in the Kuskokwim Mountains belt are divided into six groups: (1) northern volcanic fields; (2) northern volcanoplutonic complexes; (3) northern plutons, dikes, and sills; (4) volcanic and plutonic rocks in the Sleetmute-Nyac area; (5) plutonic rocks of the southern Kuskokwim Mountains region; and (6) the Bristol Bay volcanic sequence. The northern part of the belt overlies the Ruby, Innoko, Nixon Fork, and Minchumina terranes; the southern part overlies the Dillinger, Tikchik Lake, Nyac, Kilbuck, Togiak, and Goodnews terranes of Jones and others (1987, and Plate 3, in press).

The northern volcanic fields include the Sischu, Nowitna, Dishna, and Yetna volcanic fields (Fig. 1). The Sischu volcanic field (71-66 Ma) consists of a narrow belt of poorly exposed rhyolite and dacite domes, flows, and tuff that extends from the Sischu Mountains to the southern Chitanatala Mountains (Moll and others, 1981; Moll-Stalcup and Arth, 1989). The main volcanic field covers an area of more than 725 km<sup>2</sup>, is at least 500 m thick, and is fault bounded on the southeast side. A felsic pluton that crops out just east of the volcanic rocks has an age of 64 Ma (Silberman, and others, 1979).

The Nowitna volcanic field consists of more than 1500 m of chiefly andesitic flows preserved in a gently folded northeast-trending syncline that is fault-bounded on the southeast side. The field covers more than 2700 km<sup>2</sup>, and overlaps the suture between the Innoko and Nixon Fork terranes. At least seven highly altered rhyolite domes overlie the andesite flows. K-Ar ages on three whole-rock andesite samples collected near the top of the section are 64 to 63 Ma (Silberman and others 1979).

The Dishna volcanic field consists of calc-alkalic dacite, rhyolite, and minor andesite poorly exposed in a series of isolated ridges and hills that rise above the alluvium in the Innoko-Dishna Rivers area (Chapman and others, 1985). These undated rocks are presumed to be Late Cretaceous or early Tertiary in age.

Volcanoplutonic complexes occur in the McGrath area at Page Mountain, Cloudy Mountain, Candle Mountain, Takotna Mountain, Mount Joaquin, the Beaver Mountains, and in the Lonesome Hills. These complexes have circular-shaped outcrop areas that consist of andesite flows and shallow hypabyssal rocks intruded by small granitic stocks. Most of the volcanic rocks are highly altered by the intrusions. The margins of many of the complexes appear to be fault-bounded and the complex at Page Mountain is down-faulted against the surrounding sedimentary rocks. Date volcanic and intrusive rocks from these complexes yield K-Ar ages ranging from 73 to 65 Ma (Moll and others, 1981; Bundtzen, and Laird, 1982; 1983a; 1983b; 1983c) and are interpreted as being deeply eroded volcanic centers.

Widespread intrusive rocks, many too small to be shown on published maps, occur throughout the Kuskokwim Mountains belt. In the northern Kuskokwim Mountains numerous dikes, sills, and small stocks, usually 1 to 9 km in diameter, give similar K-Ar ages (72 to 62 Ma) and are compositionally similar to the volcanic rocks. Most of the intrusive rocks are compositionally homogeneous monzonite, monzodiorite, quartz monzodiorite, quartz monzonite, or granite. Plutons at Von Frank and Stone Mountain, however, are compositionally zoned, and commonly grade inward from gabbro or monzogabbro to quartz monzonite in the center of the pluton.

Volcanic and plutonic rocks in the Sleetmute-Nyac area have K-Ar ages ranging from 75 to 61.7 Ma (Robinson and others, 1984; Decker and others, 1985, 1986; Reifentstahl and others, 1984). The volcanic sequence, named the Holokuk Basalt by Cady and others(1955), consists in fact of more than 1000 m of flows and lahars composed chiefly of andesite and minor amounts of rhyolite vitric tuff and breccia (Decker and others, 1986). Most of the andesites are older (74.5 to 64.3 Ma) than the

rhyolites. Intrusive rocks include the Chuilnuk and Kiokluk granodiorite plutons dated at 68.7 to 67.5 Ma, intermediate stocks and dikes dated at 69.8 Ma, and a number of small rhyolite porphyries dated at 70.5 to 67.9 Ma and 61.5 Ma. Similar biotite and biotite-muscovite rhyolite porphyries occur to the north along the Nixon Fork-Iditarod fault and contain garnet (Bundtzen and Swanson, 1984).

More than 30 plutons, usually small stocks 3 to 15 km in diameter, occur in the southern Kuskokwim Mountains in the area between the Nushagak and the Kuskokwim Rivers (Wilson, 1977; Hoare and Coonrad, 1978b). Hoare and Coonrad (1978b) describe them as monzonite, granodiorite, and quartz diorite stocks, "mafic" dikes and sills; and felsic dikes, sills, tuffs, and breccias. K-Ar ages for all the rock types in the Goodnews-Hagemeister quadrangles range from 72.5 to 60.7 Ma. Poorly known granitic stocks to the north and east of the the Goodnews-Hagemeister quadrangles appear to be contemporaneous with, and compositionally similar to, those in the quadrangle (Wilson, 1977; Hoare, oral comm., 1980).

A thick sequence of volcanic rocks called the Bristol Bay volcanic sequence (Globerman, 1985) is exposed on Hagemeister, Walrus, and Summit Islands in Bristol Bay and on the adjacent mainland. The rocks are dated at 68.7 to 64.5 Ma (Globerman, 1985; Box, 1985). The volcanic rocks consist of andesitic lava flows interbedded with tuffs, breccias, and volcanogenic sedimentary rocks that are exposed in a section more than 2 km thick.

**PETROGENESIS:** The Kuskokwim Mountains belt consists of moderate-K calc-alkalic to shoshonitic suites that range in composition from basalt to rhyolite. Present exposures suggest that andesite, followed by rhyolite, are the overwhelmingly dominant volcanic rock types. Dacite and basalt are relatively uncommon, and rocks having less than 52 percent SiO<sub>2</sub> are rare. Most of the intrusive rocks have intermediate to felsic compositions, and many are compositionally equivalent to dacites, plotting in the silica gap (63-70% SiO<sub>2</sub>) defined by the volcanic rocks. Mineralogies vary considerably according to rock type and K<sub>2</sub>O content (Table 1).

Major-element data on the volcanic and plutonic rocks show trends typical of most igneous calc-alkalic suites: MgO, FeO\*, TiO<sub>2</sub>, Al<sub>2</sub>O<sub>3</sub>, and CaO decrease with increasing SiO<sub>2</sub>; K<sub>2</sub>O and Na<sub>2</sub>O increase with increasing SiO<sub>2</sub>. TiO<sub>2</sub> is low (less than 1.75%), and Al<sub>2</sub>O<sub>3</sub> is moderate (12 to 17%). None of the suites shows Fe enrichment. K<sub>2</sub>O varies from moderate (1.3% at 56% SiO<sub>2</sub>) to very high values (4% at 56% SiO<sub>2</sub>). Moderate to high-K suites plot in the subalkaline field of Irvine and Baragar (1971) on a total alkalis vs. SiO<sub>2</sub> diagram and are calc-alkalic. Very high-K suites (Von Frank and Whirlwind on Fig. 5) plot in the alkalic field and are classified as shoshonitic (Morrison, 1980). Shoshonitic and calc-alkalic suites are similar in all major elements except K and P, which correlates with K. In the northern Kuskokwim Mountains the high-K calc-alkalic and shoshonitic suites tend to be older (71 to 65 Ma) than the moderate-K suites (68 to 62 Ma), although there is considerable overlap.

Major- and trace-element data suggest that the volcanic and plutonic rocks are highly enriched in Ba, Rb, Th, K, and Sr and depleted in Nb and Ta relative to La (Fig.

3). These features are characteristic of subduction-related arc rocks (Perfit and others, 1980; Gill, 1981; Thompson and others, 1984). All the rocks are LREE (light-rare-earth-element) enriched but the degree of enrichment varies, correlating with K-content and the abundance other incompatible elements: shoshonitic rocks have La about 150; high-K rocks have La about 100; and moderate-K rocks have La about 75 x chondritic abundances. In contrast andesites from the entire belt have similar HREE (heavy rare-earth-element) contents (6-13 x chondrites). There is also a rough correlation between geographic area and degree of incompatible-element enrichment. Andesites from the Bristol Bay volcanic sequence in the southernmost part of the belt have lower incompatible element contents than andesites from Sleetmute, 360 km to the north (Table 2), which have lower incompatible element contents than andesites and intermediate plutonic rocks from the northern Kuskokwim Mountains. LIL (large ion lithophile) elements (K, Rb, Ba, Th, LREE) show the greatest increase from south to north; high-field-strength (HFS) elements (Zr, Hf, Nb, and Ta), which are also incompatible, increase to a lesser degree (Table 2). Alkali element contents in the highly altered volcanoplutonic complexes are high and variable, but REE data from the complexes are similar to data from the less altered Nowitna volcanic field, suggesting that the two suites are chemically similar (compare Cloudy Mountain and Nowitna on Fig. 4D). Trace-element ratios (Ba/Ta, Ba/La, La/Nb) in andesites from all three K-groups are similar to arc andesites (Gill, 1981) despite the overall higher than "typical" arc abundances of these elements in the high-K calc alkalic and shoshonitic groups.

REE patterns for rhyolites, dacites, quartz monzonites, and granites from the Kuskokwim Mountains belt are highly variable. Most of the rhyolites and dacites from the Sischu volcanic field have patterns that have very high LREE and low HREE (3-5 x chondrites), but some have extremely high LREE and moderate HREE (Fig. 4A-4B). The samples with low HREE contents show a weak correlation between silica and decreasing HREE, which suggests that these magmas either fractionated garnet or hornblende, or formed from partial melting of garnet- or hornblende-bearing schist (Moll-Stalcup, 1987). Garnetiferous schist underlies the volcanic field and garnet xenocrysts were found in one thin section of rhyolite from the field. However, trace-element models (Moll-Stalcup and Arth, 1989) suggest that hornblende, rather than garnet, is the controlling phase. Rhyolites from the Sischu field that have extremely high LREE and very large negative Eu anomalies are probably highly fractionated high-silica rhyolites. Rhyolites and granites from the Sleetmute area have more moderate REE patterns (Fig. 4C) similar to those of the granitic intrusive rocks in the northern Kuskokwim Mountains.

Even the most mafic rocks in the Kuskokwim Mountains belt have compositions that suggest that they have undergone significant fractionation, and many show evidence for interaction with continental crust. In the northern Kuskokwim Mountains, where the basement is Precambrian and Paleozoic schist and/or sedimentary rocks, andesites from the Nowitna volcanic field have initial Sr isotope ratios ( $^{87}\text{Sr}/^{86}\text{Sr}=0.7045$  to  $0.7053$ ) and trace-element abundances that suggest that the magmas have assimilated small amounts of continental crust during crystal fractionation (Moll and Arth, 1985; Moll-Stalcup and Arth, 1989). Rhyolites from the

nearby Sischu volcanic field have high Sn, Be, U, W, and F contents and initial  $^{87}\text{Sr}/^{86}\text{Sr}$  greater than 0.7080 (Moll and Arth, 1985; Moll and Patton, 1983), which suggests that they either were strongly contaminated by continental crust or were partial melts of the crust.

In the Sleetmute area initial  $^{87}\text{Sr}/^{86}\text{Sr}$  varies from 0.704 to 0.706 (M. Robinson and R. Reifenhohl, written commun., 1985). Initial  $^{87}\text{Sr}/^{86}\text{Sr}$  does not correlate with  $\text{SiO}_2$ , Rb/Sr, or 1/Sr, indicating that crustal contamination of isotopically homogeneous magmas is not the only cause of this isotopic variation. Andesites, rhyolites, granites, and quartz monzonites have a similar range of initial  $^{87}\text{Sr}/^{86}\text{Sr}$  (andesites: 0.70403 to 0.70601; felsic rocks: 0.70488 to 0.70626; M. Robinson and R. Reifenhohl, written commun. 1985). The volcanic and plutonic rocks overlie sedimentary rocks of the Cretaceous Kuskokwim Group, but the nature and age of the older basement rocks is uncertain (S. Box, oral commun. 1986). The presence of andesites having initial  $^{87}\text{Sr}/^{86}\text{Sr}$  as high as 0.706 suggests that old (Precambrian and Paleozoic) radiogenic continental crust or lithosphere, or sedimentary rocks derived from old continental crust occur in the Sleetmute area, although the mechanism for interaction with the crust is not yet understood. Initial  $^{87}\text{Sr}/^{86}\text{Sr}$  on andesites from the Bristol Bay volcanic sequence are uniformly low (0.7037 to 0.7041) (Globerman, 1985), indicating that old continental crust does not underlie this area .

St. Matthew Island, on the Bering Sea shelf, is composed entirely of Late Cretaceous and early Tertiary volcanic rocks ranging in composition from basalt to rhyolite (Patton and others, 1975). The island may be part of the Alaska Range-Talkeetna Mountains belt, part of the Kuskokwim Mountains belt, or neither, but the  $\text{K}_2\text{O}$  contents are most like those in volcanic rocks in the Alaska Range-Talkeetna Mountain belt (Fig. 5). No trace-element data on the rocks from Alaska Range-Talkeetna Mountains belts have been published except for those on the anomalous McKinley sequence. Comparisons of incompatible-element contents between rocks from the Kuskokwim Mountains belt (Table 2) and rocks from St. Matthew Island show that the St. Matthew Island rocks have slightly lower abundances of incompatible elements than the Bristol Bay volcanic sequence, which has the lowest contents in the belt. REE data for St. Matthew andesites have lower LREE than andesites from either Sleetmute or the northern Kuskokwim Mountains (Fig. 4D) but have La contents similar to that of the Bristol Bay volcanic sequence (Table 2).

The unusually large range in incompatible element contents of rocks in the Kuskokwim Mountains belt is not well understood. K and incompatible elements vary geographically and temporally; along the strike of the belt K increases from south to north, and in the northern Kuskokwim Mountains the K content decreases with time, from shoshonitic to moderate-K. The volcanic and plutonic rocks in the Kuskokwim Mountains belt are compositionally similar to arc volcanic rocks which are thought to be generated by partial melting in a mantle wedge that has been modified by alkali-enriched fluids derived from the subducted slab and/or subducted sediments (Arculus and Powell, 1986). In arc magmas, K contents are separately influenced by the composition of the crust and the depth to the slab. K has been empirically

observed to increase with distance from the trench, or depth to the slab. Additional factors that can influence the degree of K and other incompatible-element enrichment include greater degrees of crustal contamination or the presence of old, thicker, or more enriched crust or lithosphere. Estimates of crustal thickness from gravity studies (Barnes, 1977) suggest that the present crust is 30 to 35 km thick under the Kuskokwim Mountains, except in the southernmost part near Bristol Bay where it is 25 to 30 km thick. However, these present-day estimates are not well constrained because of the lack of seismic refraction data for the entire region. Crustal thicknesses in the early Tertiary were probably about the same as at present because sedimentation and deformation has been minor since the early Tertiary. The basement in the northern Kuskokwim Mountains consists of Precambrian and Paleozoic continental crust of the Ruby, Nixon Fork, and Minchumina terranes, which contrasts sharply with the late Paleozoic and Mesozoic mafic and intermediate oceanic rocks of the Togiak terrane in the south. Isotopic data suggest that the igneous rocks in the northern part of the belt have interacted with old continental crust. Thus, the increase in incompatible-element contents from south to north along the belt is probably related to the presence of old, possibly thicker, continental crust in the northern Kuskokwim Mountains.

Increasing incompatible-element contents along the strike of a continental arc has been documented by Hildreth (1986) in the Chilean Andes. He attributes the increase in incompatible elements, from south to north, to continental influence because other factors, such as sediment subduction, or depth to the Benioff zone, do not vary geographically and the basement in the north is both thicker and older than the basement in the south.

The range in K from moderate-K to high-K to shoshonitic within the northern Kuskokwim Mountains cannot be attributed to the presence of old thick continental crust because all three suites overlie the same type of basement. Furthermore, there is no correlation between initial  $^{87}\text{Sr}/^{86}\text{Sr}$  and K-type (E. J. Moll and M.L. Silberman, unpub. data, 1985), which suggests that the shoshonitic K contents are not the result of greater degrees of crustal contamination nor the result of local crustal inhomogeneities. There is, however, a weak correlation between age and K-type, because the shoshonitic to high-K suites tend to be older than the moderate- to high-K suites. This correlation suggests that some of the variation in K in the northern Kuskokwim Mountains is tectonically controlled by depth to the Benioff zone and that the dip of the slab may have become shallower over time. This model would account for both the higher K-content of the rocks and the occurrence of a narrower magmatic belt prior to 65 Ma. On the other hand, the older rocks may be more enriched in incompatible elements simply because those elements were concentrated in the first partial melts.

## YUKON-KANUTI BELT

The Yukon-Kanuti belt (Fig. 1) consists of an aligned group of early Tertiary volcanic fields that form a northeast-trending belt located north and west of the Kuskokwim Mountains belt. The belt extends from the Arctic Circle near the town of

Bettles for over 300 km southwest to the Kaltag fault and continues south of the fault on the west side of the Yukon River (Moll-Stalcup and others, in press). There is no clear boundary between the Kuskokwim Mountains and Yukon-Kanuti belts in the region south of the Kaltag fault. Late Cretaceous and early Tertiary volcanic rocks are divided into the two belts because the Yukon-Kanuti belt contains some younger rocks (66 to 47 Ma) are found in the Kuskokwim Mountains belt (72 to 60 Ma), and because the Yukon-Kanuti belt is within the Yukon-Koyukuk province (Patton and others, in press b). The Yukon-Koyukuk province consists chiefly of late Paleozoic and Mesozoic mafic and intermediate volcanic rocks and associated sedimentary rocks, which contrast sharply with the Precambrian and lower Paleozoic schist and carbonate rocks that underlie the northern Kuskokwim Mountains belt. A further distinction is that the depth of post-early Tertiary erosion in the Yukon-Koyukuk province is shallower; as a result there are more volcanic rocks and fewer plutonic rocks exposed in the Yukon-Kanuti belt than in the Kuskokwim Mountains belt.

Three areas in the Yukon-Kanuti belt have been studied in detail: the Kanuti volcanic field in the northern part, the Yukon River area located south of the Kaltag fault, and the Blackburn Hills volcanic field about 50 km farther southwest (Moll-Stalcup and Arth, 1989; in prep.). Numerous small volcanic bodies, plutons, dikes, and sills, having K/Ar ages chiefly between 65 and 53 Ma, occur in the northern and central parts of the belt. Extensive volcanic and plutonic rocks of Tertiary age are shown on the geologic map of Alaska farther south (Beikman, Plate 1, in press), but no detailed map, age, or petrologic data are available.

The Kanuti field lies within the Yukon-Koyukuk province near its southeast margin (Fig. 1). The field consists of dacite, andesite, and rhyodacite flows, domes, and tuffs exposed in a broad syncline that trends northeast and covers an area of more than 550 km<sup>2</sup>. The base of the volcanic section is dated at 59.5 and 59.7 Ma, and the top at 55.9 Ma (Patton and Miller, 1973; Moll-Stalcup and Arth (1989).

A moderate size (275 km<sup>2</sup>) volcanic field is located about 75 km southwest of the Kanuti field near Tokatjikh Creek (Patton and others, 1978). The rocks are undated but consist of andesite and dacite flows and tuffs petrologically similar to those in the Kanuti field and probably contemporaneous. Another small volcanic field, located to the west, just south of the village of Huslia, near Roundabout Mountain, is also undated and described as andesitic (Patton, 1966; Moll-Stalcup and others, in press). Both of these fields are probably part of the early Tertiary volcanic belt, although they could be part of the younger middle Tertiary (40 Ma) province discussed below.

Isolated domes, cones and flows and one large volcanic field crop out in the dense vegetation along the west banks of the Yukon River south of the village of Kaltag (no. 2 on Fig. 1). These rocks are the youngest volcanic rocks (53-47 Ma) in the Late Cretaceous and early Tertiary province of western Alaska and occur in three main locations. From north to south they are (1) a large (400 km<sup>2</sup>) poorly exposed volcanic field near Poisen Creek and Stink Creek that consists of basalt, andesite, dacite, and rhyolite (50.6 and 47.6 Ma: W.W. Patton, Jr. and E.J. Moll, unpub. data, 1981); (2) a large composite rhyolite dome and associated olivine basalt and pyroxene andesite

flows at Eagle Slide (53.2 Ma: Patton and Moll, 1985); and (3) a morphologically well preserved andesite cone located 6 km west of Bullfrog Island (53.8 Ma: Patton and Moll, 1985). Harris (1985) reports similar K-Ar ages on rocks from the same area.

A well exposed volcanic field (150 km<sup>2</sup>), located in the Blackburn Hills, about 10 km west of the Yukon River and 70 km south of the Kaltag fault (no. 1 on Fig.1), consists of a thick section of andesite flows exposed in a northeast-trending syncline that is bounded by the Thompson Creek fault on the west flank and by an unnamed fault on the northwest flank. The flows are interlayered with rhyolite domes and basalt flows near the top of the section. The core of the syncline consists of a thick section of highly altered green tuff intruded by a granodiorite pluton. The base of the volcanic pile is dated at 65 Ma; the rhyolite domes and granodiorite pluton at 56 Ma (Patton and Moll, 1985; Moll-Stalcup, 1987). The green altered tuff is interpreted as intracaldera tuff, and the granodiorite pluton is interpreted as an eroded resurgent dome (Moll-Stalcup, 1987).

A thick sequence of east-dipping andesite flows crops out in low hills about 20 km south of the Blackburn Hills on the opposite side of the Thompson Creek fault. Available data suggest that these flows are part of the west-flank of the Blackburn Hills volcanic field, which in turn suggests that 20 km of left lateral movement has taken place along the Thompson Creek fault since the early Tertiary.

**PETROGENESIS:** The Yukon-Kanuti belt is generally similar to the Kuskokwim Mountain belt in that it consists of moderate- to high-K calc-alkalic suites of mafic to felsic compositions. Basalt, however, is more abundant in the Yukon-Kanuti belt, although still much less common than andesite. A further distinction between the two belts is that most rocks in the Yukon-Kanuti belt having 63-68% SiO<sub>2</sub> are volcanic rather than intrusive.

The Blackburn Hills volcanic field consists of a thick pile (1 km) of chiefly andesite flows at the base, and interlayered rhyolite, basalt, and andesite at the top. The lower andesite section is divided into two groups on the basis of REE patterns: One group consists of one- and two-pyroxene andesites and basalts that have REE patterns similar to the andesites in the northern Kuskokwim Mountains (Fig. 4D, G). The second group consists of hornblende and pyroxene andesites that have higher LREE and lower HREE than the first group (Fig. 4H). Rocks in the second group also have lower FeO<sup>t</sup> and TiO<sub>2</sub>, higher MgO at a given SiO<sub>2</sub>, and higher initial <sup>87</sup>Sr/<sup>86</sup>Sr than the first group, and they occur only in the northwest part of the volcanic field. The rhyolites at the top of the section have a wide variety of mineral assemblages, some of which are typical calc-alkalic and some of which are mildly alkalic. Common mineral assemblages include anorthoclase+hedenbergite, oligoclase+biotite, and oligoclase+orthopyroxene.

The occurrence of both calc-alkalic and mildly alkalic suites at the top of the Blackburn Hills section suggests that there was a chemical and mineralogical transition

at about 56 Ma. Rocks that are older than 56 Ma are compositionally similar to those in the Kuskokwim Mountains belt in that the andesites are enriched in K, Rb, Ba, Th, U, and Sr and depleted in Nb and Ta, relative to the LREE (Fig. 6A). These andesites contain minerals typical of calc-alkalic rocks such as plagioclase, orthopyroxene, clinopyroxene, and hornblende. Rocks between 56 and 47 Ma constitute a mixed assemblage of calc-alkalic and mildly alkalic rocks which occur at the top of the Blackburn Hills section and in the nearby Yukon River area. Some of the post-56 Ma rocks show the characteristic enrichments and depletions and have the typical calc-alkalic mineralogy (rhyolites: plag+biot; andesites: plag+opx+cpx) of the earlier suite. The post-56 Ma assemblage differs, however, in that: (1) there is a higher proportion of basalt and these basalts have smaller Nb-Ta depletions and lower alkali/ LREE ratios than typical arc rocks (Fig. 6A; Thompson and others, 1982); and (2) some of the rhyolites have mildly alkalic mineral assemblages, such as anorthoclase+hedenbergite (Moll-Stalcup, 1987). This chemical and mineralogical transition is not strongly reflected in the major-element variation that shows a typical calc-alkaline affinity for all the rocks.

The shift in mineralogy and chemistry upsection is found only in the Blackburn Hills field. The rocks in the Yukon River area (53 to 47 Ma) post date the transition, whereas the rocks in the Kanuti field (59 to 56 Ma) predate the transition. The volcanic rocks in the Yukon River area include a mixed assemblage of typical calc-alkalic andesites and dacites (andesite: plag +opx+cpx; dacite:plag+horn+biot) and mildly alkalic latites and basalts (latites:anorthoclase+plag+biot; basalt:ol+ plag+cpx and groundmass ol+biot). Rocks that have calc-alkalic mineralogy have moderate Nb-Ta depletions ( $La/Nb$  greater than 2), and those with mildly alkalic mineralogy have smaller Nb-Ta depletions. No correlation between age and rock types has been found in the Yukon River area and it appears that both calc-alkalic "arc-type" and mildly alkalic "non arc-type" rocks erupted between 53 and 47 Ma.

The Kanuti volcanic field is 59 to 56 Ma and predates the transition noted at the southern end of the belt. All the analyzed rocks from the Kanuti field have a high alkali and LREE content and deep troughs at Nb and Ta on chondrite-normalized multi-element plots (Fig. 6C), similar to the older suite of rocks in the Blackburn Hills. The volcanic rocks in the Kanuti field range from high-silica andesite to rhyodacite. Most contain hornblende and have REE patterns similar to hornblende-bearing andesites in the Blackburn Hills (Fig. 5A).

Samples of andesite, basalt, and dacite that give K-Ar ages of 59.8 to 50.2 Ma have been dredged from the edge of the Bering Sea shelf (Fig. 1; Davis and others, 1987). These rocks are altered but appear to have typical calc-alkalic compositions and may be correlative with the Yukon-Kanuti belt. Whether they are part of the Yukon-Kanuti or some other belt, however, is not known. They may be rocks that were accreted to the continental margin after eruption.

Volcanic rocks that have K-Ar ages from 64 to 62 Ma also occur on St. Lawrence Island in the Bering Sea. Patton and Csejtey (1980) describe them as basalt, soda rhyolite, trachyandesite, and andesite. Chemical analyses (W.W. Patton, Jr., unpub. data, 1971) suggest the rocks are mildly alkalic and possibly a bimodal basalt-rhyolite

assemblage ( $\text{SiO}_2$  48% and 68 to 71%). The K-Ar ages and mildly alkaline compositions suggest that the rocks are probably not part of any of the Late Cretaceous and early Tertiary belts, but they could be part of the Yukon-Kanuti belt if the transition from subduction-related calc-alkalic rocks to mildly-alkalic post-subduction magmatism occurred earlier in that part of the Bering Sea.

#### LATE CRETACEOUS AND EARLY TERTIARY TECTONIC IMPLICATIONS

The Alaska Range-Talkeetna Mountains, the Kuskokwim Mountains, and the Yukon-Kanuti belts constitute a Late Cretaceous and early Tertiary magmatic province that extends over a vast region of Alaska, from the Alaska Range north to the Arctic Circle and west to the Bering Sea shelf. The volcanic and plutonic rocks in this province overlap a number of tectonostratigraphic terranes that lie between the Alaska Range, Bristol Bay, and the Yukon-Koyukuk province, which suggests that these terranes were assembled by late Mesozoic time (Moll and Patton, 1982).

Paleomagnetic studies of Late Cretaceous and early Tertiary volcanic rocks at Bristol Bay (Globerman, 1985), Lake Clark (Thrupp and Coe, 1986), the northern Kuskokwim Mountains (Nowitna field: Coe and others, 1985; Blackburn Hills field: Thrupp and Coe, 1986), the Talkeetna Mountains (Hillhouse and others, 1985), and the east-central Alaska Range (Cantwell Formation, Hillhouse and Gromme', 1982) indicate about 30 to 55 degrees of counterclockwise rotation, but no major latitudinal displacement relative to North America since eruption (Hillhouse and Coe, in press; Thrupp and Coe, 1986).

Igneous rocks from all three belts have chemical compositions typical of subduction-related magmatism that occurred between 75 and 56 Ma. Dominant compositions are intermediate and felsic, and rocks from all three belts have high K, Ba, Sr, Rb, and Th, and low Nb, Ta, and Ti relative to LREE --the most diagnostic characteristics of arc magmatism. Most workers (for example Hudson, 1979; Reed and Lanphere, 1973) agree that the Alaska Range-Talkeetna Mountains belt is a continental volcanic arc related to subduction of the Kula plate under southern Alaska during the Paleocene. The tectonic environment of the Kuskokwim Mountains and Yukon-Kanuti belts has not been so well documented. Wallace and Engebretson (1984) outline three possible relationships between the Alaska Range-Talkeetna Mountains and Kuskokwim Mountains belts: (1) the Kuskokwim Mountains belt formed in an extensional environment (or intracontinental back-arc setting) behind the Alaska Range-Talkeetna Mountains belt (Bundtzen and Gilbert, 1983; Gemuts and others, 1983); (2) The Alaska Range-Talkeetna Mountains and Kuskokwim Mountains belts constitute an unusually wide arc analogous to Oligocene volcanism in the western U.S. (Gill, 1981 p. 39 and references therein); or (3) the Alaska Range-Talkeetna Mountains and the Kuskokwim Mountains belts were two separate arcs juxtaposed by subsequent tectonic activity. Of the three possibilities, the third is the least likely because no major Tertiary suture has been mapped between the two belts, and geologic and paleomagnetic data suggest that the two belts were essentially in place during formation--not accreted at at some later time. Both belts may have been brought into their current position by strike-slip faults, but Tertiary offset along faults in western Alaska are thought to be relatively small, probably less than 150 km (Grantz, 1966).

Finally, the abundance of high-K calc-alkalic and shoshonitic rocks in the Kuskokwim Mountains suggests that it was not a piece of the Alaska Range-Talkeetna Mountains belt that was originally along strike and was later strike-slip faulted to its present position.

The chemical composition of the plutonic and volcanic rocks in the Kuskokwim Mountains and Yukon-Kanuti belts strongly supports their formation in an arc rather than back-arc environment. Other workers (Bundtzen and Swanson, written commun., 1986) have argued that the Kuskokwim Mountains magmatism is the result of regional extension because (1) many of the rocks in the northern Kuskokwim Mountains plot in the alkalic field on total alkalis versus silica diagrams; (2) they believe that most of the rocks are a bimodal assemblage of basalt and rhyolite; and (3) the belt is associated with numerous normal/strike-slip faults. I distinguish between arc-related alkalic rocks (shoshonites) that are characterized by  $K_2O/Na_2O$  greater than, or equal to 1, low  $TiO_2$  and high field strength elements, and high alkalis and  $Al_2O_3$  (Morrison, 1980; JGR volume 91B). Most of the published analyses (Bundtzen and Laird, 1982, 1983a, b, c) have  $SiO_2$  greater than 53%, when the analyses are recalculated 100% anhydrous. Of 49 analyses having LOI less than 10%, 4 have less than 53%  $SiO_2$ , 36 have 53-63%  $SiO_2$ , 4 have 63-70%  $SiO_2$  (all plutonic), and 4 have more than 70%  $SiO_2$ . I classify rocks having more than 53%  $SiO_2$  as andesites even though many are olivine-bearing, and note that olivine is stable at a higher silica content in high-K suites than in low or moderate-K suites (Kushiro, 1975). Furthermore, the silica gap in the published data occurs in the dacite range (63 to 70%  $SiO_2$ ), which is not uncommon in modern arc volcanoes (Grove and Donnelly, 1986) and is distinct from the Miocene bimodal basalt-rhyolite suite associated with extension in the western United States (Christiansen and Lipman, 1972). Bundtzen and Swanson's argument for extension based on the abundance of faults is discussed further below.

The chemical data strongly support the interpretation that the Alaska Range-Talkeetna Mountains, Kuskokwim Mountains, and Yukon-Kanuti belts constituted an anomalously wide magmatic arc in the Late Cretaceous and early Tertiary, analogous to the western United States in the Oligocene. In their present configuration the three belts span a width of 550 km extending to about 900 km measured orthogonal to the Aleutian trench near the Alaska Peninsula. In the conterminous states, Oligocene volcanism as far east as the Rocky Mountains has been attributed to convergence along the western continental margin (Lipman and others, 1972; Snyder and others, 1976). The compositions of andesites in the northern Kuskokwim Mountains are remarkably similar to those in the Oligocene San Juan volcanic field in Colorado (Lipman and others, 1978; Table 3), which was located at least 900 km from the Oligocene continental margin.

Variation in  $K_2O$  within and among the three belts is complex because K and the other incompatible elements are affected by several factors, including the distance from the trench or depth to the subducting slab, the age and lithology of the underlying crust and mantle, the amount of crustal contamination, and the degree of partial melting of the source.  $K_2O$  contents are higher in the Kuskokwim Mountains

and Yukon-Kanuti belt (Fig. 5) than in the Alaska Range-Talkeetna Mountains belt, and the rocks in the northern Kuskokwim Mountains have more  $K_2O$  than either the southern Kuskokwim Mountains belt or the Yukon-Kanuti belt. As discussed earlier, high  $K_2O$  (and all incompatible elements) contents in the northern Kuskokwim Mountains belt may be partly related to interaction of the magmas with old, possibly thicker, continental crust or lithospheric mantle. Even moderate-K rocks in the northern Kuskokwim Mountains have higher K and incompatible element contents than rocks from any of the other areas or belts, suggesting that the magmas in the northern Kuskokwim Mountains have interacted with old continental crust or mantle, which has resulted in their enriched composition. The range in K within the northern Kuskokwim Mountains, however, is probably the result of some other factor, such as changing depth to the subducting slab.

The age data suggest that between 75 and 65 Ma the magmatic arc was narrower, consisting of only the Alaska Range-Talkeetna Mountains and Kuskokwim Mountains belts, and that the K gradient across the arc was steeper. During this period the K-content in the magmas increased across the arc, from moderate-K in the southern Alaska Range-Talkeetna Mountains belt, to high-K and shoshonitic in the northern Kuskokwim Mountains. Much of this steep gradient in K, across the Alaska Range-Talkeetna Mountains to the Kuskokwim Mountains belt, was probably tectonically controlled by increasing depth to the slab, but some K-enrichment in the northern Kuskokwim Mountains is probably due to lithospheric interaction.

During the period from 65 to 56 Ma the arc broadened to include the Yukon-Kanuti belt, and the K gradient across the arc was more gradual, probably due to a decrease in dip of the subducting slab. During this time period,  $K_2O$  at a given silica content was lower in the Yukon-Kanuti belt than in the northern Kuskokwim Mountains despite the fact that the Kuskokwim Mountains rocks were closer to the trench. This reversal in K trend across the arc is interpreted as the result of lithospheric interactions in the northern Kuskokwim Mountains. A change in crustal thickness, on the other hand, is not indicated by the gravity data and therefore was probably not a factor in the  $K_2O$  gradient. Gravity studies suggest that the crust under both the Yukon-Kanuti belt, in the study area, and the northern Kuskokwim Mountains is between 30 and 35 km thick (Barnes, 1977), and that it probably was about as thick in the early Tertiary. The  $K_2O$  content of the arc magmas in the southern Kuskokwim Mountains belt (Bristol Bay volcanic rocks) is about the same as in the Yukon-Kanuti belt, both of which are underlain by late Paleozoic and Mesozoic oceanic crust and island arc rocks of the Yukon-Koyukuk province and Togiak terrane (Fig. 5). The late Paleozoic and Mesozoic mafic and intermediate basement rocks would be expected to have much lower incompatible element contents, lower  $^{87}Sr/^{86}Sr$ , and higher  $^{143}Nd/^{144}Nd$  than the old sialic continental crustal rocks of the Ruby and Nixon Fork terranes, which underlie the northern Kuskokwim Mountains. The strong contrast between basement lithologies is evident in trace-element and isotope compositions of rhyolites from each area. Rhyolites from the Blackburn Hills, in the Yukon-Kanuti belt, have lower LREE, Rb, Th, U, Sn, F and initial  $^{87}Sr/^{86}Sr$  and higher  $^{143}Nd/^{144}Nd$  than rhyolites from the northern Kuskokwim Mountains (Moll and Patton, 1983; Moll and

Arth, 1985). Andesites from the Nowitna volcanic field show evidence of interaction with enriched continental crust, whereas those from the Blackburn Hills do not (Moll and Arth, 1985; Moll-Stalcup and Arth, 1989). I attribute the very high  $K_2O$  and incompatible-elements contents in the northern Kuskokwim Mountains magmas to the presence of old sialic crust under that area. Such old crust might also affect the  $K_2O$  gradient along the arc, from north to south, as well as cause the reversal in  $K_2O$  gradient across the arc from the northern Kuskokwim Mountains to the Yukon-Kanuti belt. Interaction with the crust seems especially common in subduction-related suites because these magmas rise slowly to the surface and have ample time to differentiate and assimilate country rock.

Bundtzen and Swanson (1984) note that many of the Late Cretaceous and early Tertiary volcanic and intrusive bodies are associated with normal or strike-slip faults. Little is known about the age and type of motion for most of these faults, but most appear to have significant right-lateral movement (Grantz, 1966). Some of the volcanic fields are cut by normal or strike-slip faults, but it is uncertain whether the faults entirely postdate the volcanism, and thus control only the exposure of the volcanic fields, or if some of the volcanism was contemporaneous with, or post-dates the faulting. Rhyolite domes dated at 58 Ma occur at Old Woman Mountain in the Kaltag fault zone just west of the Yukon River (W.W. Patton, Jr. and E.J. Moll, unpub. data, 1983) and similar domes dated at 61 m.y. occur in the Nixon-Fork Iditarod fault zone near McGrath (Bundtzen and Laird, 1983). Some of these rhyolite domes thus appear to have intruded into pre-existing fault zones. In addition, some felsic dikes or sills of Late Cretaceous and early Tertiary age strike parallel to major faults (Patton and others, 1980; Bundtzen and Laird, 1983b), which suggests some of the faulting had occurred by the time the dikes and sills were emplaced. Most of the Late Cretaceous and early Tertiary volcanic piles are gently folded, with maximum dips of 20 to 45 degrees, but most of the Middle Tertiary volcanic piles are flat lying and undeformed (Harris, 1985; W.W. Patton, Jr., oral comm., 1986). These data demonstrate that at least some of the deformation post-dates the Late Cretaceous and early Tertiary magmatism, and also suggests that some may have been contemporaneous with or predated it. I believe that these equivocal relations do not support the interpretation that the magmatism is the result of continental rifting. The chemical composition of rocks in both the Yukon-Kanuti and Kuskokwim Mountains belts was dominantly controlled by subduction, not continental rifting. Although the timing of movement along the faults is not well constrained, I find the idea that most of the movement along the faults took place in the Eocene and was related to the clockwise rotation or oroclinal bending of western Alaska (Globerman and Coe, 1984) very appealing. The implications of this idea are discussed further below.

In contrast to the 75 to 56 Ma rocks, many of the rocks in western Alaska that are 56 Ma or younger have chemistry and mineralogy that are not typical of subduction-related magmas. These rocks include the McKinley sequence granites and the quartz monzonite of Tired Pup in the western and northern Alaska Range, the rhyolites and basalts at the top of the Blackburn Hills, and the volcanic rocks in the Yukon River area in the Yukon-Kanuti belt. I believe subduction-related magmatism in western Alaska ceased at about 56 Ma, and the period from 56 to 50 Ma represents

a transition from subduction-related magmatism to post-subduction, possibly intraplate, magmatism, during which rocks typical of both environments erupted. This transitional period is marked by (1) a marked decrease in the volume of magma erupted over the entire province; (2) an increase in the proportion of basalt and the eruption of mildly alkalic basalt, latite, and alkali rhyolite in the Yukon-Kanuti belt and (3) the emplacement of peraluminous silicic granites of the McKinley sequence in the northern and western Alaska Range. Subduction-related volcanism younger than 56 Ma occurred in the Talkeetna Mountains at the eastern end of the Alaska Range-Talkeetna Mountains belt, but insufficient petrologic data is available on these rocks to determine whether a similar chemical or mineralogical transition marked the end of subduction-related magmatism there.

In summary, the age and compositional data suggest that the Alaska Range-Talkeetna Mountains, Kuskokwim Mountains, and Yukon-Kanuti belts constituted an anomalously wide magmatic arc related to north-directed subduction at a trench along the Paleocene continental margin of Alaska (Fig. 1) between 75 and 56 Ma. The data also suggest that only the southeastern two-thirds of the province was active between 75 and 66 Ma, broadening to the entire province between 66 and 56 Ma. A sharp decrease in magma volume and a mineralogical and compositional change at 56 Ma mark the end of the subduction event in most of the province except the easternmost end of the Talkeetna Mountains. Plate-motion models (Engebretson, 1982) show rapid north-northeast-directed subduction of the Kula plate under southern Alaska between 74 to 56 Ma. In early Eocene time the trench jumped out to its present position and the Aleutian arc began to form (Rea and Duncan, 1986; Scholl and others, 1986).

The location of the magmatic belts relative to each other and to their associated trench in Late Cretaceous and early Tertiary time is uncertain. The convergence angle between the present position of the three belts and the Paleocene plate-motion vectors is near the minimum (25 degrees) required for arc magmatism (Gill, 1981 p. 27; Wallace and Engebretson, 1984). Paleomagnetic data on the volcanic rocks indicate that western Alaska has rotated counter-clockwise 30 to 55 degrees but has not been latitudinally displaced since the Paleocene (Globerman and Coe, 1984; Hillhouse and Coe, in press). These data suggest that the magmatic belts may have had a convergence angle of 55 to 80 degrees prior to rotation in the Eocene. Paleomagnetic data from the Ghost Rocks Formation on Kodiak Island in southern Alaska indicate it has moved 2000 km to the north since the Paleocene (Plumley and others, 1982). These data suggest that a Tertiary suture associated with the accretion of southernmost Alaska should be located somewhere between southern Kodiak Island and the Alaska Range batholith (Thrupp and Coe, 1986). This hypothetical suture may represent a strike-slip fault (Moore and others, 1983) or the subduction zone associated with the wide Late Cretaceous and early Tertiary arc.

The location of the magmatic belts relative to the trench is further obscured by numerous right-lateral strike-slip faults which have apparently cut the wide magmatic arc since the Paleocene. The amount of offset since the Paleocene is not known. Most data suggest that the amount of offset along the strike-slip faults in western Alaska is considerably less than the amount of offset along the faults in eastern Alaska (Grantz, 1966). The Yukon-Kanuti belt appears to be offset along the Kaltag fault, which Patton

and Hoare (1968) suggest has had 60 to 130 km of right-lateral offset since the Late Cretaceous. My approximate reconstructions of fault movements and the approximate location of the trench lead me to estimate that the entire magmatic arc was within 750 km of the trench when it was active.

### MIDDLE TERTIARY MAGMATISM

The period from 55 to 43 Ma was characterized by a hiatus in magmatic activity in most of mainland Alaska except the Aleutian arc, the hinge line of the Alaska orocline (Talkeetna Mountains, Arkose Ridge, Prince William Sound, and Yakutat terrane), the Sanak-Baranof belt (Hudson, in press), and the Yukon River area. Small amounts of volcanic activity occurred in the Aleutian arc and Alaska Peninsula from 50 to 43 Ma (Scholl and others, 1986; Wilson, 1985; Vallier, and others, in press; Marsh, in press; Miller and Richter, in press). At about 40 Ma ( $\pm 3$ ), a brief magmatic pulse occurred in western interior Alaska and a major pulse of magmatic activity, which lasted 10 m.y., began in the Aleutian arc. The arc volcanism occurred in a narrow belt extending from the Aleutian Islands probably as far northeast as Sugar Loaf Mountain on the north side of the Denali fault in the central Alaska Range. Magmatism on the Aleutian Islands, on the west side of the Alaska Peninsula (the so-called Meshik arc of Wilson, 1985), on the north and west flank of the Alaska-Aleutian Range batholith (Merill Pass, Reed and Lanphere, 1973), at Mount Foraker and McGonagall (Reed and Lanphere, 1974b), at Mount Galen (Decker and Gilbert, 1978) and possibly as far east as Sugarloaf Mountain (Albanese, 1980; Albanese and Turner, 1980) was probably part of the middle Tertiary Alaska-Aleutian arc.

The Talkeetna Mountains and Yukon River areas were discussed in the previous section and are interpreted as the last remnants of a widespread Late Cretaceous and early Tertiary subduction event in western and southern Alaska. Eocene dikes in Prince William Sound are listed in Table 1) and are described in more detail by Hudson (in press), who considers them part of the Sanak-Baranof belt of anatectic granites. The Eocene volcanic rocks in south-central and western interior Alaska are described below.

### SOUTH-CENTRAL ALASKA

Many of the volcanic and plutonic rocks that give ages in the range 55 to 43 Ma. occur in the hinge of the Alaska orocline in an area between the Border Ranges and Denali fault or in the southern accretionary wedge south of the Border Ranges fault. Rocks in the hinge area include the Arkose Ridge Formation, the Talkeetna Mountains volcanic rocks, and many unnamed plutons in the northern Talkeetna Mountains.

Basalt dikes, sills, and altered tholeiitic flows (55-43 Ma) occur in the Arkose Ridge Formation of southern Alaska, just south of the Talkeetna Mountains volcanic rocks. A. Grantz (oral comm., 1986) considers these rocks to be late Paleocene in age on the basis of plant fossils in interbedded sandstones. The rocks have smaller Nb-Ta

depletions on multi-element diagrams than do contemporaneous arc rocks in the Talkeetna Mountains (Fig. 7). Also, their trace-element ratios do not unambiguously resolve whether they are arc tholeiites related to subduction or are intraplate rocks possibly related to the Castle Mountain fault. Therefore, both their tectonic environment and their precise age are uncertain.

Early to middle Eocene basaltic flows, hyaloclastites and flow breccias, along with interbedded clastic marine sedimentary rocks, occur in the Yakutat terrane south of the Border Ranges fault (Davis and Plafker, 1986). The basalts consist of LILE-depleted and LILE-enriched tholeiites, which Davis and Plafker (1986) interpret as normal mid-ocean ridge and oceanic island basalt. They suggest that these basalts are correlative with contemporaneous geochemically similar basalts, which occur in a linear belt extending from southern Vancouver Island to the southern Oregon Coast Range. They further suggest that the basalts originated as early to middle Eocene seamounts near the Kula-Farallon spreading center, and were accreted to the southern continental margin of Alaska at about 48 m.y., during subduction of the Kula-Farallon ridge and Kula plate.

#### WESTERN INTERIOR ALASKA

Volcanic rocks (37-43 Ma) occur on St. Lawrence Island, in the Melozitna area, in the Yukon River area and in the Sleetmute area (Fig. 8). Limited data suggest that these constitute a bimodal assemblage of felsic volcanic rocks and associated basalt that have calc-alkalic to mildly alkalic affinities.

Patton and Csejtey (1980) describe volcanic rocks on St. Lawrence Island (39.3 Ma) as rhyolite and dacite tuff, tuff breccia, and flows. The unit is not well exposed but is thought to be flat lying or gently dipping (Patton and Csejtey, 1980).

Three volcanic fields ( $40 \pm 3$  Ma) occur between the Melozitna and Koyukuk Rivers on the north side of the Kaltag fault at Indian Mountain, Takhakhdona Hills, and Dulbi River. All three consist chiefly of rhyolite tuff, flows, and breccia (Patton and others, 1978). Dark vesicular basalt flows also occur in the Takhakhdona Hills. Rhyolite obsidian (39.9-41.6 Ma) occurs at Indian Mountain (Miller and Lanphere, 1981) and is a possible source of the obsidian artifacts found in northwestern Alaska (Patton and Miller, 1970).

Harris (1985) reports K-Ar ages of  $40 \pm 3$  Ma at two localities on the Yukon River, one north of Morgan Island in the southernmost Nulato quadrangle (42.7 Ma) and one near the village of Grayling in the northernmost Holy Cross quadrangle (42.4 Ma). He also reports that the Oligocene volcanic rocks are flat-lying and undeformed. No lithologic data are given, but at least some are basalt because the K-Ar dates were run on basalt whole-rock samples. Additional volcanic rocks of this age may occur farther south in the Holy Cross quadrangle, where widespread undated volcanic rocks are exposed.

Little known volcanic rocks in this age range occur in the Sleetmute area. Olivine basalt (38.2 Ma) is reported from the Chuilnuk River area and a rhyolite sequence (43.8 Ma) is reported from Tang Mountain (Decker and others, 1986). The rhyolites at Tang Mountain are apparently interbedded(?) with trachyandesite near the base of the rhyolite section. Decker and others, (1986) describe a black glassy rhyolite unit, probably a basal vitrophyre, overlain by a gray rhyolite unit.

**PETROGENESIS:** Few petrologic data are available for the middle Tertiary volcanic rocks in western interior Alaska. Major-element data from St. Lawrence Island, Indian Mountain, the Takhakhdonga Hills, and Tang Mountain suggest that the rocks are dominantly felsic (66 to 77% SiO<sub>2</sub>), along with minor mildly alkalic basalt and andesite. Basalt occurs in the Takhakhdonga Hills and Yukon River area; andesite occurs at Tang Mountain. No chemical or mineralogical data are available for the basalt on the Yukon River. The basalt in the Takhakhdonga Hills has low Al<sub>2</sub>O<sub>3</sub> (14.0%) and TiO<sub>2</sub> (0.80%) and high total alkalis (5.5%; Patton and others, 1978); no trace-element data are available and their affinity (shoshonitic or mildly alkalic) is uncertain. Chemical analyses of the andesite at Tang Mountain (Decker and others, 1986) show that the rock has high TiO<sub>2</sub> (1.97%), K<sub>2</sub>O (3.00%), Nb (49 ppm), Ta (3.85 ppm), and La (51 ppm) and low Ba (445 ppm). The composition of this rock, along with its trace-element ratios (Ba/Ta= 116, La/Nb= 1.03), suggest it is not a subduction-related orogenic andesite but a trachyandesite of some other affinity. Normalized multi-element data on trachyandesites from Tang Mountain show very high LREE and virtually no Nb-Ta depletion (Fig. 9). The initial <sup>87</sup>Sr/<sup>86</sup>Sr on the same rock is 0.7033, much lower than initial <sup>87</sup>Sr/<sup>86</sup>Sr on the Late Cretaceous and early Tertiary rocks from the same area.

Mineralogical data are available only for rocks from a few localities and are summarized in Table 1). Most of the felsic rocks have minerals typical of calc-alkalic rocks, but some are more alkalic. Latite having anorthoclase, plagioclase, biotite, and oxides has been reported from the Takhakhdonga Hills, and trachyandesite having sanidine in addition to plagioclase and clinopyroxene occurs at Tang Mountain.

The tectonic affinity of these rocks is uncertain, but they do not appear to be related to subduction. The volcanic fields in the Takhakhdonga Hill, at Indian Mountain, the Dulbi River, and the Yukon River all occur within the Yukon-Koyukuk province along the southeast boundary where the sedimentary section is probably thinnest (W.W. Patton, Jr., pers. commun., 1984). Most of the rocks are flat lying and undeformed. The bimodal composition, high total alkali contents and lack of rocks having SiO<sub>2</sub> between 56 and 65% suggest that the volcanic rocks may have erupted in an extensional environment. The interior volcanic rocks erupted contemporaneously with the start of a major pulse of volcanism in the Aleutian-Alaska Range arc and the bend in the Hawaiian-Emperor seamount chain (Clague and others, 1975). Wallace and Engebretson (1984) also show a change from rapid northward-directed subduction of the Kula plate to slow northwest subduction of the Pacific plate at 43 Ma. The eruption of the volcanic rocks may be related to this change in the rate and angle of plate motion or to the switch from the Kula to the Pacific plate. Changes in the angle of subduction may have resulted in movement along the many strike-slip faults in the

area, however, the volcanic fields are not located on or near mapped faults, and their lack of deformation suggests that they have not experienced major movement. Also, paleomagnetic studies of Oligocene volcanic rocks in the Yukon River (Harris, 1985) and Lake Clark areas (Thrupp and Coe, 1986) suggest that the rocks have not rotated or moved latitudinally relative to North America since their formation.

Rocks of mid-Tertiary age may occur in at least two offshore basins. Eocene basalts having K-Ar ages of  $40.7 \pm 2$  and  $42.3 \pm 10$  Ma were found at the bottom of the Cape Espenberg and Nimiuk wells, drilled in the Kotzebue basin (Tolson, 1986). Poorly dated basalt flows or sills were also found near the bottom of wells drilled in Norton Sound basin and are thought to be middle to late Eocene (Kirschner, in press). Both basins are extensional basins with horsts and grabens (Norton Sound) or graben and half-graben (Kotzebue) structures (Kirschner, in press).

## LATE CENOZOIC VOLCANISM

Late Cenozoic volcanism in Alaska is dominated by calc-alkalic activity along the Aleutian arc and in the Wrangell Mountains and by contemporaneous alkalic and tholeiitic volcanism behind the arc in the Bering Sea region and easternmost interior Alaska (Fig. 10). The Aleutian arc has been active for at least the last 50 m.y. (Scholl and others, 1986), whereas most of the alkalic and tholeiitic Bering Sea basalts are restricted to the last 6 m.y.

## THE BERING SEA REGION

The Bering Sea volcanic province consists of a number of large late Cenozoic basalt fields that occur in a vast region extending from St. Lawrence and the Pribilof Islands on the submerged Bering Sea shelf landward to the Seward Peninsula and Togiak River valley in western Alaska (Fig. 10). Most fields consist of a broad plain or shield composed of voluminous basalt flows overlain by steep cones and maars of undersaturated highly alkalic magma. Basaltic volcanism began as an isolated event about 28 to 26 m.y. ago with the eruption of the Kugruk Volcanics on the Seward Peninsula (Hopkins and others, 1963; Swanson and others, 1981). However, most of the volcanism in the Bering Sea region is much younger. Volcanism on Nunivak Island began 6 m.y. ago (Hoare and others, 1968), about the same time it resumed on the Seward Peninsula (5.8 Ma; Swanson and others, 1981). Most of the other fields began erupting in the Pliocene and Pleistocene.

On Nunivak Island geologic mapping, paleomagnetic reversal stratigraphy, and K-Ar dating were used to determine the time and volume relationships of volcanism (Hoare and others, 1968). The volcanic rocks range in age from 6 Ma to Holocene; the older series (3 to 6 Ma) underlies the western third of the island and the younger series (0 to 1.7 Ma) covers the eastern two-thirds of the island.

Two suites were occur on Nunivak Island: 98% of the volcanic rocks form broad thin pahoehoe flows of alkali basalt and subordinate tholeiite that make up 30 to 50 small shield volcanoes. The remaining 2% are basanites and nephelinites that form small viscous flows, about 60 cinder cones, and ash deposits from four maar craters (J.M. Hoare, unpub. data, 1971). Subordinate eruptions of basanite or nephelinite commonly preceded and followed large eruptions of less alkalic basalt (J.M. Hoare, unpub. data, 1971). Analcime- or nepheline-bearing basanite or nephelinite overlie the Cretaceous sedimentary basement rocks on Nunivak Island, and in turn, are overlain by less alkalic basalts (J.M. Hoare, unpub. data, 1971). The youngest volcanism produced 40 to 50 basanite cones and four maar craters that erupted in the last 0.5 Ma in the south-central part of the island (Hoare and others, 1968). The cones and craters are aligned approximately east-west along several parallel fractures in a belt about 40 km long and 12 km wide. The basanite and nephelinite contain abundant inclusions of lherzolite, layered gabbro, chromite, conglomerate, sandstone, and basalt, and megacrysts up to 10 cm long of anorthoclase, augite, and kaersutite.

The late Cenozoic (0.24-1.5 Ma) volcanic shield in the Kookooligit Mountains on St. Lawrence Island covers 42 by 33 km, is at least 500 m thick, and overlies volcanic, plutonic, and sedimentary rocks of early Paleozoic to Tertiary age (Patton and Csejtey, 1980). About 95-97% of the volcanic rocks are alkali-olivine basalt and subordinate amounts of olivine tholeiite, and 3-5% are basanite and minor nephelinite (J.M. Hoare, unpub. data, 1981). Large fluid flows of alkali-olivine and olivine-tholeiite basalt erupted from many small craters (20 to 60 m in diameter) and from one or more larger craters (100 to 150 m in diameter) located in the central part of the field. Many small basanite flows and 60 to 80 cinder cones are aligned along an east-west belt across the north-central part of the field. Most of the basanite flows and cones contain xenoliths of deformed peridotite and/or gabbro. Sparse xenoliths of granitic rock and siliceous tuff in basanite cones and flows have been found at three localities i.

Widespread late Cenozoic volcanic fields cover about 10,000 km<sup>2</sup> of the Seward Peninsula (Till and Dumoulin, in press). The Imuruk Lake area was originally mapped by Hopkins (1963), who distinguished five volcanic formations on the basis of weathering, degree of frost brecciation and thickness of windblown silt. K-Ar studies in the Imuruk Lake region document the five eruptive episodes: (1) 28 to 26 Ma, (2) 5.8 to 2.2 Ma, (3) 0.9 to 0.8 Ma, (4) late Pleistocene, and (5) Holocene (Swanson and others, 1981). These data indicate that the earliest eruptive episode occurred 20 Ma before the onset of mafic alkalic volcanism elsewhere in the Bering Sea region. The Imuruk Lake volcanic field is composed dominantly of alkali-olivine basalt and subordinate olivine tholeiite, quartz tholeiite, and basanite erupted from small shield volcanoes, cinder cones, plugs, and maar craters. Xenolithic inclusions, which occur in alkalic basalt and basanite, consist dominantly of lherzolite and subordinate harzburgite, chromite, schist, and granite. Additional basalt fields on the Seward Peninsula include basanite, tephrite, and alkali olivine basalt from a 2.5-2.9 Ma field north of Teller on the margin of the Imuruk basin and one at Devil Mountain in the northernmost part of the Peninsula (S. Swanson and D. Turner, unpub. data, 1984).

The Pribilof Islands, located near the edge of the Bering Sea shelf, were also the site of late Cenozoic volcanism. Volcanism occurred on St. George, the northern island, between 2.2 to 1.6 Ma, and on St. Paul Island, located 70 km to the south, from 0.374 Ma to the present (Cox and others, 1966; Lee-Wong and others, 1979). Volcanism was active in the intervening time on a submarine ridge located between St. George and St. Paul Islands, where dredged whole-rock basalt samples yield ages of 0.774 and 0.836 Ma (Simpson and others, 1979). St. George Island consists chiefly of deeply dissected olivine tholeiite flows. Its topography is controlled by numerous east-west trending normal faults. St. Paul Island consists of coalescing small volcanoes, each composed of a central cinder cone and a surrounding shield of lava flows composed of alkali-olivine basalt and basanite. Dredged samples from the submarine ridge are chiefly olivine tholeiite, subordinate alkali-olivine basalt, minor basanite, and nephelinite (Lee-Wong and others, 1979).

The Togiak Basalt in southwestern Alaska (loc. 8 on Fig. 10) consists of about 9 km<sup>3</sup> of tholeiitic and alkali-olivine basalt flows (0.76 Ma) overlain by a tuya, or table mountain, formed by a subglacial basaltic eruption (Hoare and Conrad, 1978a, 1980). The tuya consists of palagonitized glassy tuff and pillow lava capped by glassy subaerial flows of alkali-olivine basalt that probably erupted during glacial advances more than 39,000 years ago.

The Yukon Delta contains numerous small volcanic fields that consist of alkali olivine basalt flows, erupted from low saucer-shaped cones, overlain by young steep-sided cones composed of basanite and nephelinite (Hoare and Condon, 1971a; J.M. Hoare, unpub. data, 1971). Olivine nephelinite ash, perhaps erupted from a maar crater in the Ingakslugwat Hills, contains abundant olivine gabbro and lherzolite nodules (Hoare and Condon, 1971a). Some of the young steep-sided cones are aligned west-northwest, indicating a probable fracture control (Hoare and Condon, 1971a).

A large volcanic field covering about 2,000 km<sup>2</sup> near St. Michael Island in southeastern Norton Sound consists chiefly of voluminous olivine tholeiite and alkali olivine basalt flows, and basanite tuffs, cones, and maar craters. Some of the young cones and short flows, such as those at Crater Mountain, consist of basanite with lherzolite nodules; others, such as a recent lava flow southeast of Crater Mountain, consist of olivine tholeiite. Flows from the base of the St. Michael volcanic field give whole-rock ages of 3.25 and 2.80 Ma (D.L. Turner, written comm., 1986). The approximate age of flows on St. Michael and the Yukon Delta were determined by Hoare and Condon (1966, 1968, 1971a, 1971b) and Hoare and Conrad (1959, 1978) using magnetic polarity and physiographic expression of the rocks. Their data indicates that the flows were erupted during both the Brunhes Normal- (0.7 Ma to the present) and Matuyama Reversed- (2.4 to 0.7 Ma) Polarity Chrons. A few highly dissected flows in these areas may be as old as Pliocene.

A small olivine basalt field at Flat Top Mountain east of the Kuskokwim delta (4.62-4.72 Ma; Decker and others, 1986) is presumed herein to be part of the Bering Sea province.

**PETROGENESIS:** Much confusion over the composition of the Bering Sea volcanic rocks exists because one of the few published papers describes the rocks on Nunivak Island as chiefly tholeiitic basalt and subordinate highly undersaturated alkalic basalt (Hoare and others, 1968), although alkali olivine basalt is the most common rock type (J.M. Hoare, unpub. data, 1971). More recent studies in the province have focused on the peridotite inclusions from Nunivak (Francis, 1976a and 1976b; 1978; Menzies and Murthy, 1980a; 1980b; Roden and Murthy, 1985; and Roden and others, 1984), the tectonic stress orientation of the province (Nakamura and others, 1977; 1980; Nakamura and Uyeda, 1980), and the Sr and Nd isotopic composition of the inclusions and volcanic rocks (Mark, 1971; Von Drach, and others, 1986). Most authors describe the rocks as "back-arc" basalt because of their position behind the Aleutian arc.

For this report I examined 200 analyses from St. Lawrence, Nunivak, St. Michael and Ingakslugwat (J.M. Hoare, unpub. data, 1967-1971), 36 from the Pribilof Islands (Lee-Wong and others, 1979), 3 of the Togiak Basalt (Hoare and Coonrad, 1980), and 19 from the Seward Peninsula (Swanson and Turner, unpub. man., 1981). Thin sections from the Hoare's collections show fresh phenocrysts and groundmass minerals, including olivine, and almost all of the analyses have less than 1% total H<sub>2</sub>O.

Volcanic rocks in most fields range from nephelinites having more than 25% normative nepheline to tholeiites having 15% normative hypersthene. Most of the fields are compositionally similar to the volcanic field on Nunivak. Alkaline olivine basalt and olivine tholeiite represent at least 95% of the volcanic rocks present in all the volcanic fields (J.M. Hoare, unpub. data, 1967-1972) and form flat-lying flows and shield volcanoes. About 2-3% of the rocks are highly alkalic undersaturated basanite and nephelinite which form short viscous flows, cones, and ash. Eruptions of small volumes of highly alkalic undersaturated magma generally preceded and postdated voluminous outpouring of alkali and tholeiitic basalt (J.M. Hoare, unpub. data, 1971).

Most of the volcanic rocks have a  $100\text{Mg}/(\text{Mg} + \text{Fe}^{2+})$  greater than 65, and many contain ultramafic mantle xenoliths, which suggest that they may be primary or near-primary melts of mantle peridotite, that have experienced little or no residence time in shallow magma chambers and relatively rapid rise to the surface. Data from both the basanite-nephelinite suite and the basalt suite cluster on AFM diagrams, further supporting the general lack of differentiation in the suite. Rare hawaiiite ( $100\text{Mg}/(\text{Mg} + \text{Fe}^{2+}) = 55$  to 64) has been reported on St. Lawrence Island, from the St. Michael volcanic field, and in dredgings from the ridge between St. Paul and St. George Islands, but has not been reported from the other areas.

Both the alkali-olivine and tholeiitic basalts have phenocrysts of olivine, plagioclase, clinopyroxene, magnetite, and ilmenite; they are characterized by diktytaxitic textures. Hawaiiites have less olivine; basanites lack plagioclase phenocrysts. Basanites have nepheline and analcime, and nephelinites have sodic pyroxene and nepheline in the groundmass mesotaxis (Hoare and others, 1968). None of the samples examined in thin section contained modal hypersthene, but it has been reported in the Imuruk Volcanics from the Imuruk Lake area (Hopkins, 1963).

The basanites and nephelinites commonly contain megacrysts and/or xenoliths of peridotite, gabbro, or bedrock. The megacrysts consist of unzoned anorthoclase, clinopyroxene, and kaersutite, as much as 10 cm long (Hoare, and others 1968). Most of the megacrysts are deformed, showing kink bands and undulatory extinction, and most show reaction relations to their host basalt. Aluminous clinopyroxene, kaersutite, and feldspar are common megacrysts in alkalic basalts around the world (Irving, 1974). Experimental studies of megacrysts and their host basalts suggest that megacrysts of clinopyroxene and kaersutite are near-liquidus phases of the host basalt at high pressures (10 to 20 kb). In contrast, experimental work suggests that anorthoclase is never a liquidus phase in a basalt or basanite at any pressure, and they therefore are generally thought to have precipitated from more evolved magmas which later mixed with the host basalt (Irving and Frey, 1984). The lack of differentiated magmas in the Bering Sea basalt province, however, makes this mechanism unlikely in these magmas. H. Wilshire (oral comm., 1985) believes that anorthoclase megacrysts in the Mojave Desert, Calif., formed in mantle veins due to several generations of partial melting. The anorthoclase megacrysts in the Bering Sea basalts may have formed in a similar manner and were disaggregated during their rise to the surface.

The most abundant xenoliths--75% of Nunivak's xenolith population (Francis, 1976a)-- are lherzolite nodules composed of olivine, enstatite, clinopyroxene, and spinel. Less than 1% of the lherzolite nodules on Nunivak contain chromian pargasitic amphibole, but about 50% contain zones of fine-grained diopside, olivine, spinel, and Al-rich glass, which Francis (1976a) interprets as relicts of melted amphibole. Red-brown chromian mica occurs between the amphibole and included spinel in some samples. Francis (1976a) believes that the spinel lherzolite xenoliths are accidental fragments of upper mantle and that amphibole formed during a mantle metasomatic event accompanied by infiltration of aqueous fluids enriched in alkalis and incompatible elements. He further suggests that the metasomatism predates entrainment of these nodules in the alkali basalt.

Other common xenoliths include corona-bearing pyroxene granulites (9% of Nunivak's xenolith population) which range from plagioclase to olivine dominated (Francis, 1976b). The reaction of olivine and plagioclase to clinopyroxene-spinel symplectite and aluminous orthopyroxene suggests that the xenoliths last equilibrated at 950 degrees C under at least 9 kbar pressure. Francis (1976b) interprets the xenoliths as fragments of the base of the crust, and proposes that the reaction took place in the corona structures in response to crustal thickening of the Bering Sea shelf from thin oceanic crust (10 km) in the early Mesozoic to thicker crust (about 30 km) in the Quaternary.

Other reported xenoliths include bedrock of obvious crustal origin and dunite, harzburgite, chromite, gabbro, and amphibole-bearing pyroxenite of less certain origin (Hoare and others, 1968; Francis, 1978). Lithologies of the bedrock xenoliths vary depending on the location of the volcanic field. At least some at each locality consist of fragments of underlying basalt flows.

The Bering Sea province is characterized by high alkali and low silica content and ranges in composition from nephelinite to basanite through alkali-olivine basalt to olivine tholeiite and tholeiite. Total alkalis decrease with increasing  $\text{SiO}_2$  (Fig. 11) as in the Hawaiian suites (Clague and Frey, 1982; Frey and Clague, 1983). But, Bering Sea basalts have lower CaO than Hawaiian lavas, which results in greater silica saturation at a given total alkali content. Figure 11 shows the dividing line for alkalic and tholeiitic rocks in the Bering Sea province and the line of silica saturation, as defined by MacDonald and Katsura (1964) for the Hawaiian Islands.

REE data from Nunivak (Roden, 1982), the Pribilof Islands (Kay, 1977; Florence Lee-Wong, unpub. data, 1986), and St. Michael volcanic field (E.J. Moll and W.W. Patton, Jr., unpub. data, 1983) show that lavas from these fields are LREE enriched. LREE are correlated with alkalinity: nephelinites have 40 ppm La and tholeiites have about 10 ppm La.

$\text{P}_2\text{O}_5$ , Rb, Sr, Ba,  $\text{K}_2\text{O}$ , and  $\text{Na}_2\text{O}$  all decrease with increasing silica and correlate positively with each other (J.M. Hoare, unpub. data, 1967-1971; Mark, 1971). These trends can not be due to crystal fractionation of any phenocryst commonly found in the basanites or basalts. Studies of suites of similar composition (Frey and others, 1978; Clague and Frey, 1982) have shown that basalt to nephelinite to melilite suites formed by decreasing degrees of partial melting of a garnet peridotite source. By analogy, this suggests that Bering Sea nephelinites, which have the highest Rb, Sr, Ba, K, Na, and P and the lowest  $\text{SiO}_2$ , originated by the smallest degree of partial melting; and that tholeiites, which have the lowest alkalis and highest  $\text{SiO}_2$ , originated by the greatest degree of partial melting. Experimental petrologists have suggested that strongly silica-undersaturated magmas form from a peridotite mantle rich in carbon (Wyllie and Huang, 1976; Egger and Holloway, 1977; Egger, 1978). I examined P and K data from over 200 major-element analyses of rocks from St. Lawrence Island to estimate the range of partial melting required this volcanic field using the method of Clague and Frey (1982). For the olivine tholeiite to nephelinite series,  $\text{K}_2\text{O}$  varies from 0.4 to 2.2% and  $\text{P}_2\text{O}_5$  varies from 0.22 to 0.71%. K content has the widest range, which suggests that it is the most incompatible major element. If K behaves as a totally incompatible element ( $D=0$ ) during partial melting, and no fractional crystallization occurs, its compositional range requires that the degree of partial melting vary by a factor of at least 5. Trace elements are probably more incompatible than major elements. Trace element data are not available for most of the Bering Sea basalt fields, but Rb data on a small suite of samples from Nunivak Island vary from 5.8 to 64.4 ppm (Mark, 1971), a factor of 11, over the range olivine tholeiite to nephelinite. A factor of 11 corresponds to a melting interval of at least 11% or 1 to 11%, 2 to 22%. However, as discussed below, Sr isotope data on the Bering Sea magmas suggest they have not originated in a mantle that has isotopically homogeneous  $^{87}\text{Sr}/^{86}\text{Sr}$ . The source for the nephelinites has lower  $^{87}\text{Sr}/^{86}\text{Sr}$ , and therefore probably lower Rb/Sr, than the source for the less alkalic basalts, which suggests that the melt interval is probably larger. Frey and others, (1978) suggest that 4 to 25% melting produced a compositional range from olivine melilite to quartz tholeiite in the Tertiary to Holocene basalts of Victoria, Australia, and Tasmania. Although I cannot

rigorously constrain the degree of partial melting for the Nunivak Island suite, the data suggest that the nephelinites form by small amounts of partial melting, probably on the order of 1 to 3%, and the tholeiites formed by larger amounts of melting, on the order of 11 to 33%.

Three isotopic studies have been done on the Bering Sea province and a summary of the reported data is shown in Figure 12. In the first study, Mark (1971) analyzed the Sr isotopic composition of more than 24 samples, ranging from nephelinite to tholeiite from Nunivak Island and of a few samples from St. Lawrence and St. Michael Islands, Ingakslugwat Hills and the Pribilof Islands. He found that the  $^{87}\text{Sr}/^{86}\text{Sr}$  = 0.7026 to 0.7033 of the Nunivak samples and decreased with increasing alkalinity and decreasing silica saturation. Nine basanites have  $^{87}\text{Sr}/^{86}\text{Sr}$  of  $0.70286 \pm 0.00018$  and 15 alkali olivine and tholeiitic basalts have  $^{87}\text{Sr}/^{86}\text{Sr}$  of  $0.70311 \pm 0.00013$ . Samples from the Pribilofs, Ingakslugwat, and St. Michael fall within the range of Sr isotope ratios given for Nunivak.  $^{87}\text{Sr}/^{86}\text{Sr}$  on samples from St. Lawrence Island, which overlies Paleozoic sedimentary rocks, are higher, ranging from 0.7036 to 0.7039 and also decrease with increasing alkalinity.

In the second isotopic study, Menzies and Murthy (1980a; 1980b) studied the Sr and Nd isotopic composition of basalts, kaersutite megacrysts, and paragasite from lherzolite nodules from Nunivak in the second isotopic study. They report Sr isotope data for volcanic rocks, ranging from nephelinites to olivine tholeiites, similar to values reported by Mark in 1971 ( $^{87}\text{Sr}/^{86}\text{Sr}$  = 0.70251 to 0.70330), and  $^{143}\text{Nd}/^{144}\text{Nd}$  ranging from 0.51289 to 0.51304. In contrast to Sr, the Nd isotopes do not correlate with alkalinity. The nodules and megacrysts have  $^{87}\text{Sr}/^{86}\text{Sr}$  ranging from 0.7027 to 0.7033, which suggests that the mantle under Nunivak is locally inhomogeneous in  $^{87}\text{Sr}/^{86}\text{Sr}$ . Coexisting paragasite and nodules have identical isotopic composition within analytical uncertainty. The Nunivak data plot on a  $^{87}\text{Sr}/^{86}\text{Sr}$  versus  $^{143}\text{Nd}/^{144}\text{Nd}$  diagram in the field where MORB (mid-ocean-ridge basalt) and OIB (oceanic-island basalt) overlap (Menzies and Murthy, 1980a; 1980b; Roden and others, 1984) but are LREE enriched (Roden, 1982). Thus the low Nd isotopic composition of the basalts and nodules requires a time-integrated LREE-depleted source for the basalts, but all the basalts, even the tholeiites, are LREE enriched (Fig. 13). Menzies and Murthy (1980a) suggest that the LREE and other incompatible elements were enriched in the source region by relatively recent (less than 200 Ma) mantle metasomatism and that the range in Sr isotopic composition can be explained by local inhomogeneities in Rb/Sr that developed during the metasomatic event and resulted in small variations in  $^{87}\text{Sr}/^{86}\text{Sr}$  over time.

The third isotope study (von Drach and others, 1986) focused on the Nd and Sr isotopic composition of the Aleutian Islands but also reported data from Nunivak, St. George, and St. Lawrence Islands. Data for Nunivak and St. George are identical to the previously published results on Nunivak (Menzies and Murthy, 1980) and plot on the  $^{87}\text{Sr}/^{86}\text{Sr}$ - $^{143}\text{Nd}/^{144}\text{Nd}$  diagram in the field where MORB and OIB overlap.  $^{143}\text{Nd}/^{144}\text{Nd}$  is  $0.5133 \pm 0.0002$  for six of the seven samples from both Nunivak and St. George.  $^{87}\text{Sr}/^{86}\text{Sr}$  for the same samples ranges from 0.7025 to 0.7033. Rocks

from St. Lawrence Island have significantly higher Sr and lower Nd and plot along the mantle array within the field for oceanic basalts (Fig. 12).

Data from all three studies suggest that volcanic rocks from Nunivak, the Pribilofs, Ingakslugwat, and St. Michael are isotopically similar and plot in the field where MORB and OIB overlap.  $^{143}\text{Nd}/^{144}\text{Nd}$  values are about 0.5132, and  $^{87}\text{Sr}/^{86}\text{Sr}$  ranges from 0.7025 to 0.7033 for all the analyzed fields except St. Lawrence Island, which has a basement composed of rocks at least as old as middle Paleozoic, and has higher  $^{87}\text{Sr}/^{86}\text{Sr}$  and lower  $^{143}\text{Nd}/^{144}\text{Nd}$  than the other volcanic fields (Fig. 12). In all the fields where data are sufficient,  $^{87}\text{Sr}/^{86}\text{Sr}$  appears to be negatively correlated with silica undersaturation and alkalinity. Hawaiian volcanic rocks on Oahu show a similar trend--the Honolulu Group, which are composed of undersaturated alkalic rocks have lower  $^{87}\text{Sr}/^{86}\text{Sr}$  (0.70331) than the underlying tholeiitic shield (0.70370) (Lanphere and others, 1980; Lanphere and Dalrymple, 1980; Clague and Frey, 1982). The source of the St. Lawrence Island magmas is either mantle that has higher  $^{87}\text{Sr}/^{86}\text{Sr}$  and lower  $^{143}\text{Nd}/^{144}\text{Nd}$  than the other volcanic fields (continental lithospheric mantle) or crustally contaminated isotopically similar mantle. Inclusions of sialic rock have been reported in highly alkalic magmas on St. Lawrence Island (Patton and Csejtey, 1980). However, the primitive composition of the magmas, which suggests little differentiation and thus little or no residence time in shallow magma chambers, argues against the crustal contamination hypothesis. Although, it is possible that crustal contamination may be responsible for the range in  $^{87}\text{Sr}/^{86}\text{Sr}$  within individual volcanic fields, it is not required because a similar range in Sr is found in the mantle, as evidenced from xenolith studies from Nunivak (Menzies and Murthy, 1980). The proposed metasomatic event, which enriched the mantle under the Bering Sea and western Alaska in K, LREE, and P probably occurred within the last 200 Ma, as suggested by Menzies and Murthy (1980), but was not synchronous with the alkalic volcanism, as suggested by Roden and others (1984). Consequent metasomatism as suggested by Roden and others (1984) does not account for the range in  $^{87}\text{Sr}/^{86}\text{Sr}$  in the basalts nor does it explain how large volumes of LREE-enriched magmas (including tholeiites) could form from a LREE-depleted mantle that had a small volume of LREE-enriched veins. It seems more likely that a separate metasomatic event, perhaps related to subduction in the Neocomian or Late Cretaceous and early Tertiary, was responsible for the enrichment.

Current theories on the geologic evolution of western Alaska center on controversy over the crustal structure of the Yukon-Koyukuk basin. Patton and Box (1989) think that the basement of the Yukon-Koyukuk is an accreted island arc that is Jurassic or younger in age, whereas other workers (Gemuts and others, 1983) argue that Precambrian and Paleozoic rocks of the Ruby, Seward, and Endicott terranes, which surround the basin and occur on St. Lawrence Island, extend under the basin beneath the arc rocks. The Ingakslugwat Hills, Nunivak Island, and St. Michael Islands lie within the Yukon-Koyukuk basin and the Pribilof Islands may lie within an offshore extension of this basin. Studies of crustal xenoliths included in the basaltic flows indicate that sialic rocks of pre-Cretaceous age do not occur in the St. Michael, Nunivak or Ingakslugwat volcanic fields (Hoare, unpub. data, 1981). The lack of pre-Cretaceous

sialic inclusions suggests that Paleozoic strata are not present beneath these areas and further suggests that the lithosphere under the basin might have lower  $^{87}\text{Sr}/^{86}\text{Sr}$  and higher  $^{143}\text{Nd}/^{144}\text{Nd}$  than the lithosphere under older, long-lived continental areas such as St. Lawrence Island.

In summary, the Bering Sea basalt suites of nephelinite to tholeiite probably originated by increasing degrees of partial melting of a peridotite mantle rich in carbon. Most of the magmas rose quickly to the surface and few, if any, were significantly differentiated. The basanites and nephelinites originated in a source having lower  $^{87}\text{Sr}/^{86}\text{Sr}$  than the source of the less alkalic basalts. The small range in  $^{87}\text{Sr}/^{86}\text{Sr}$  is probably due to mantle metasomatism that enriched the source area in K, LREE, Sr, Rb, and P within the last 200 m.y. (Menzies and Murthy, 1980). Furthermore, this metasomatism may be related to previous subduction events in western Alaska during the Neocomian or Late Cretaceous and early Tertiary. The volcanic rocks on St. Lawrence Island have more radiogenic  $^{87}\text{Sr}/^{86}\text{Sr}$  and less radiogenic  $^{143}\text{Nd}/^{144}\text{Nd}$  than volcanic rocks from the other fields and suggest the presence of more enriched mantle under St. Lawrence Island -- possibly continental lithospheric mantle. The correlation between exposed crustal type and Sr and Nd isotope composition can be explained by tectonic models for western Alaska that require younger and more mafic crust beneath the Yukon-Koyukuk province than the crust under the surrounding metamorphic borderlands and St. Lawrence Island.

#### EASTERN AND CENTRAL INTERIOR ALASKA

Several small isolated basaltic volcanoes occur in easternmost Alaska, between the Fortymile and Tanana Rivers. The cones and associated flows are undated but apparently are young, as evidenced by their well-preserved volcanic morphology. The best preserved and probably youngest cone is Prindle volcano, a small cinder cone that was the source of a narrow basanite lava flow more than 10 km long (Foster and others, 1966). The cone and adjacent flow contain abundant inclusions of harzburgite, wehrlite, lherzolite, pyroxenite, and granulite facies schist. Prindle volcano is undated but underlies the (informally named) White River ash bed, which was dated by  $\text{C}^{14}$  methods at approximately 1,900 yrs B.P. (Lerbekmo and Campbell, 1969).

Large volumes of olivine basalt flows, as much as 100 m thick, are exposed in fault blocks near the Porcupine and Black Rivers (Brosge' and Reiser, 1969). The flows are considered to be Tertiary or Quaternary because of their youthful appearance. One chemical analysis, of an alkali-olivine basalt, is available from flows along the Black River (Brabb and Hamachi, 1977).

Miscellaneous isolated volcanic rocks of late Cenozoic age also occur in central Alaska. A small isolated maar volcano composed of olivine basalt erupted 3000 yrs B.P. at Buzzard Creek on the north side of the Alaska Range (Albanese, 1980). Flat-lying olivine basalt flows cover approximately 100 km<sup>2</sup> of north-central Alaska between the Yukon and upper Koyukuk Rivers, about 125 km northeast of the town of

Tanana. These rocks are undated but their lack of deformation suggests they are late Tertiary or Quaternary in age (Patton and Miller, 1973).

The few available analyses of volcanic rocks in central and eastern Alaska indicate that the rocks are compositionally similar to Bering Sea basalts. Alkalic basalts in the Porcupine and Yukon-Tanana Upland area are located north of arc volcanoes in the Wrangell Mountains (Fig. 10). By analogy with Bering Sea basalts, these magmas probably represent regional extension behind the present arc. Cones and flows of olivine basalt, some of which contain ultramafic inclusions, also occur in a south-trending regional belt that extends from eastern Alaska into the Yukon Territory and continues down along the western North America continental margin through British Columbia (Foster and others, 1966; Sinclair and others, 1978).

Jumbo Dome, the only occurrence of Quaternary (0.80-2.8 Ma) orogenic andesite in interior Alaska, occurs about 10 km southwest of the maar volcano at Buzzard Creek (Fig. 10) (Albanese, 1980). The hornblende andesite dome is probably related to subduction under this area, making it the only Quaternary occurrence of arc volcanism in the 300-km-wide magmatic gap between Mount Spurr, at the northeast end of the Aleutian arc, and the Wrangell Mountains.

#### LATE CENOZOIC TECTONIC IMPLICATIONS

Bering Sea basalts have compositions similar to suites found in a variety of tectonic environments including oceanic islands (Hawaii: Clague and Frey, 1982; Frey and Clague, 1983), stable continents associated with regional faulting (southern and eastern Australia: Irving, 1974; Frey and others, 1978, and the western U.S.: Menzies and others, in press), continental rifts (east Africa: King, 1970), and behind volcanic arcs a great distance from the arc (China and Korea: Nakamura et al, 1985). The locations of the various volcanic fields do not define a narrow volcanic belt, a rift axis or a "hot-spot" trend. Most recent authors (Nakamura and others, 1977; von Drach and others, 1986) have labeled the Bering Sea basalts as "back-arc" basalts, although they do not constitute a classic back arc characterized by a spreading rift axis or by high heat flow (Marshall, 1978; Smirnov and Sugrobov, 1979), nor do they have typical "back-arc" compositions which usually range from N-MORB to arc tholeiite (Saunders and Tarney, 1984; Hawkins and Melchoir, 1985). Major and trace-element data suggest that the Bering Sea basalts came from a source similar to oceanic island basalts. Trace-element ratios and isotopic compositions of the Bering Sea basalts are similar to Hawaiian volcanic rocks and different from N-MORB (Table 4). The Bering Sea volcanic fields, however, are not aligned along the trend of a hot-spot. Chondrite normalized multi-element diagrams show that, unlike arc basalts, Bering Sea basalts are not depleted in Nb or Ta (Fig. 13) and, therefore, have a different source than the Aleutian arc.

Many of the voluminous basalt fields are located on or near strike-slip or normal faults, and fault displacements suggest that at least some of the faulting began while volcanism was still active. The Togiak Basalt is located in a north-northeast-trending

graben (Hoare and Coonrad, 1980; Globberman, 1985). Late Pliocene volcanic rocks on St. George Island in the Pribilofs are cut by numerous normal faults, most of which trend approximately east-northeast (Hopkins, 1976). Several of the Yukon delta volcanic fields are located along a trace of the Anvik fault, and the volcanic field near St. Michael Island is probably intersected by a trace of a Kaltag fault splay (W.W. Patton, Jr., oral comm., 1984). Young volcanic cones are aligned approximately east-west, apparently defining a fracture or fault in the St. Lawrence, Nunivak, and St. Michael volcanic fields, and in a small field north of Aropuk Lake on the Yukon delta. Late Cenozoic volcanism in the Seward Peninsula is associated with transform faulting, geothermal anomalies, large late Tertiary grabens, and high levels of seismicity in the central Seward Peninsula (Turner and Forbes, 1980).

The distribution of active faults and monogenetic cones in late Quaternary volcanic fields on St. Lawrence, St. Michael, Ingakslugwat, Nunivak, the Pribilofs, and the Seward Peninsula were used by Nakamura (Nakamura and others, 1977; Nakamura and Uyeda, 1980; and Nakamura and others, 1980) to define the tectonic stress field in the Bering Sea region in the late Quaternary. Cones on most of the volcanic fields are aligned east-west, corresponding to east-west maximum horizontal compression. Maximum horizontal compression in the Aleutian arc is oriented north-south perpendicular to the trench. The axis of maximum horizontal compression (MHC) can represent either the intermediate stress axis or the maximum stress axis. Nakamura uses a presumed tectonic environment of the volcanism to interpret the MHC in the Bering Sea region as representing north-south extension and the MHC in the arc as representing north-south compression.

Holocene surface faults (Hudson and Plafker, 1978; Plafker and Gilpin, in press; Plafker and others, in press) and focal-mechanism solutions for earthquakes on the Seward Peninsula and adjacent northwestern Alaska show dominantly normal fault movement with extension in the northwest or northeast directions (Biswas and others, 1986). Biswas and others (1986) classify all of western Alaska and the Bering Sea shelf as an area of "tensional stress regime". However, they provide no mechanisms for western Alaska south of the Seward Peninsula area or for the Bering Sea shelf.

In late Cenozoic time the Bering Sea shelf was located in the vicinity of the Eurasian, North American, and Pacific plates. The shelf was probably part of the North American plate and the plate boundary between Eurasia and North America was located to the west of the Bering Sea shelf in eastern Siberia (Zonhshain and others, 1985). Harbert and others (in press) suggest slight convergence between Alaska and Eurasia in a north to northeast direction from 37-0 Ma in the Bering Sea region based on spreading patterns in the North Atlantic. North to northeast convergence is in contradiction with Nakamura and Biswas's studies and suggests that compressive stress from convergence between the North American and Eurasian plates was localized along the plate boundary in Siberia and did not affect the Bering Sea shelf. Therefore, plate motion between the North American and Eurasian plates does not appear to be responsible for volcanism on the Bering Sea shelf.

In the Bering Sea region, motion between the North American and Pacific plates was dominated by north-directed subduction of the Pacific plate along the Aleutian

trench for at least the last 50 Ma. None of the Bering Sea basalts, not even those from the Pribilof Islands only 550 km from the trench, have the Nb-Ta depletions characteristic of arc magmas. Neither do they occur along a rift axis, nor do they have typical back-arc compositions. However, they were erupted in a broad extensional environment located behind the Aleutian arc. Thus, although the rocks do not constitute a classic back arc, the occurrence of the tholeiitic and alkalic basalt in the Bering Sea region behind the Aleutian arc and in east-central Alaska behind the Wrangell Mountains volcanoes suggests that they represent a broad zone of regional extension behind the arc. Alkaline basalts that lack Nb-Ta anomalies occur behind the Japanese arc in Korea and China (Nakamura and others, 1985) and this occurrence may be similar to that of the Bering Sea basalts.

Voluminous eruptions of Bering Sea basalts began at 6 Ma, contemporaneous with small changes in Pacific plate motion (Barron, 1986; Cox and Engebretson, 1985) and the start of a major pulse in volcanic activity in the Aleutian arc that continues to the present. The timing of eruptions in the Bering Sea region may be related to the change in the angle of Pacific plate motion at 6 Ma, or it may possibly represent the time necessary to heat the back-arc region before volcanism began.

## SUMMARY AND CONCLUSIONS

The Alaska Range-Talkeetna Mountains, Kuskokwim Mountains, and Yukon-Kanuti belts constitute an anomalously wide volcanic arc that was active during the Late Cretaceous and early Tertiary. The arc was narrower, consisting of the Alaska Range-Talkeetna Mountains and Kuskokwim Mountains belts from 75 to 66 Ma and broadened considerably to include the Yukon-Kanuti belt from 65 to 56 Ma. Plate motion models predict rapid north-northeast-directed subduction of the Kula plate under southern Alaska between 75 and 56 Ma (Engebretson, 1982). The angle of convergence between Paleocene plate motions and the present continental margin and three parallel magmatic belts is too small to generate arc magmatism (Wallace and Engebretson 1984; Gill, 1981). This enigma is resolved by paleomagnetic models that suggest that western Alaska has been rotated 30 to 55 degrees counterclockwise since the Paleocene (Globerman and Coe, 1984; Hillhouse and Coe, in press).

Assuming models for counterclockwise rotation of western Alaska are correct, the continental margin of southern Alaska, which now has a tightly curved S-shape, may have had a more open S-shape in the Late Cretaceous and early Tertiary. This configuration places St. Matthew Island close to the trench in Paleocene time, which is consistent with its low K-contents and tentative correlation with the Alaska Range-Talkeetna Mountains belt. Unrotating western Alaska also places the continental margin and three magmatic belts approximately east-west in Paleocene time--orthogonal to the direction of subduction. Compression between Alaska and Eurasia related to opening in the North Atlantic was probably responsible for flexure of the southern continental margin into its present tight S-curve. This bending is probably responsible for the post-Paleocene counterclockwise rotation of western Alaska.

At about 56 Ma, the trench jumped away from the continental margin to its present position, and formation of the Aleutian ridge began (Scholl and others, in press). Paleomagnetic data on Paleocene and Oligocene volcanic rocks suggest that rotation of western Alaska occurred between 56 and 43 Ma (Thrupp and Coe, 1986; Harris, 1985). Between 56 and 43 Ma, Engebretson (1982) show rapid north-directed subduction under southern Alaska. Subduction-related volcanic rocks between 56 and 48 Ma are restricted to the hinge line of the oroclinal bend (Arkose Ridge Formation and Talkeetna Mountains volcanic rocks), which may have been more orthogonal to the plate motion than the rotated (or rotating?) southwestern continental margin in the Eocene.

Bimodal mildly alkalic volcanism in interior Alaska is of the intraplate type and may be related to regional extension or movement along strike-slip faults. This volcanism occurred at  $40 \pm 3$  Ma, coincident with a change in the angle of Pacific plate motion at 43 Ma and the start of a peak in magmatic activity in the Aleutian arc that occurred between 40 and 30 Ma.

Bering Sea basalts were erupted in a broad extensional environment behind the Aleutian arc, but are not a classic back arc. Eruptions of the basalts, which started at about 6 Ma, are contemporaneous with changes in Pacific plate motion and the beginning of a major eruptive pulse in the Aleutian arc. Bering Sea basalts originated in a mantle source similar to that for oceanic island basalts. The source of the magmas had been previously metasomatized by the addition of K, P, REE, and Ti. This metasomatic event occurred within the last 200 m.y. (Menzies and Murthy, 1980), probably during the widespread early or Late Cretaceous and early Tertiary subduction events.

## ACKNOWLEDGMENTS

Many of the ideas contained in this report came from stimulating discussions with my colleagues at the U.S. Geological Survey, Stanford University, and the Alaska Division of Geological and Geophysical Surveys. I would especially like to thank Bill Patton, Steve Box, Gail Mahood, Dave Clague, and Howard Wilshire. I thank Sam Swanson, Don Turner, Michael Roden, Florence Lee-Wong, Alicia Davis, Gordon Thrupp, Brian Globberman, Art Grantz, Mark Robinson, John Decker, and Rocky Reifensuhl for contributing unpublished data. Much of the section on the Bering Sea basalts is based on unpublished manuscripts and data from the late Joe Hoare. I thank Warren Coonrad for organizing Joe's field notes, chemical analyses, thin sections, and maps for my study. Comments by Steve Box, Gail Mahood, Allan Cox, Bill Patton, Elizabeth Miller and Bob Coleman improved an early draft of the manuscript. The paper received helpful reviews from Tracy Vallier and Joe Arth.

## REFERENCES CITED

- Albanese, M.. D. 1980, The geology map of three extrusive bodies in the central Alaska Range: unpublished M.S. thesis, University of Alaska, Fairbanks, 104 p.
- Albanese, M.D. and Turner, D.L.,1980,  $^{40}\text{K}$ - $^{40}\text{Ar}$  ages from rhyolite of Sugar Loaf Mountain, central Alaska Range, Implications for offset along the Hines Creek Strand of the Denali Fault system: Short notes on Alaska Geology 1979-1980, Alaska State Division of Geological and Geophysical Surveys, 1980, p. 7-10.
- Arculus, R.J. and Powell, 1986, Source component mixing in the regions of arc magma generation, *Journal of Geophysical Research*: v. 91, p. 5913-5926.
- Arth, J.G., Barker, F., Peterman, Z.E., and Friedman, I.,1978, Geochemistry of the Gabbro-Diorite-Tonalite-Trondhjemite suite of southwest Finland and its implications for the origin of tonalitic and trondhjemitic magmas: *Journal of Petrology*, v. 19, p. 289-316.
- Barnes, D.F., Bouguer gravity map of Alaska: U.S. Geological Survey Geophysical Investigations Map GP-913, 1:2,500,000 scale.
- Barron, J.A., 198\_, Paleooceanographic and tectonic controls on deposition of the Monterey Formation and related siliceous rocks in California: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 53, p. 27-45.
- Beikman, H.M., 1974, Preliminary geologic map of the southwest quadrant of Alaska, U.S. Geological Survey Miscellaneous Field Studies Map MF-611, 2 sheets, 1:1,000,000 scale.
- Biswas, N.N., Akim K., Pulpan, H., and Tytgat, G., 1986, Characteristics of regional stresses in Alaska and neighboring areas: *Geophysical Research Letters*, v. 13, p. 177-180.
- Box, S.E., 1985:, Mesozoic tectonic evolution of the northern Bristol Bay region, southwestern Alaska: unpublished Ph.D. thesis, Santa Cruz, University of California, 125 p.
- Brabb, E.E. and Hamachi, B.R., 1977, Chemical composition of Precambrian, Paleozoic, Mesozoic, and Tertiary rocks from east-central Alaska: U.S. Geological Survey Open-File Report 77-631, p. 87.
- Brew, D. A. , in press, Latest Mesozoic and Cenozoic magmatism in southeastern Alaska: *in* Plafker, G., Jones, D.L., and Berg, H.C., eds., *Geology of Alaska*: Boulder, Colorado, Geological Society of America, *The Geology of North America*, Chapter 5F, p.--.

- Brosgé, W.P., and Reiser, H.N., 1969, Preliminary geologic map of the Coleen quadrangle, Alaska: U.S. Geological Survey Open-File Report 263, scale 1:250,000.
- Bundtzen, T.K. and Gilbert, W.G., 1983, Outline of geology and mineral resources of Upper Kuskokwim region, Alaska: Journal of Alaska Geological Society, v.3, p. 101-117.
- Bundtzen, T.K. and Laird, G.M., 1982, Geologic map of the Iditarod D-2 and Eastern D-3 Quadrangles, Alaska: Alaska Division of Geological and Geophysical Surveys Geologic Report 72, scale 1:63,360.
- Bundtzen, T.K. and Laird, G.M., 1983a, Geologic Map of the McGrath D-6 Quadrangle, Alaska: Alaska Geological and Geophysical Surveys Professional Report 79, scale 1:63,360.
- Bundtzen, T.K. and Laird, G.M., 1983b, Geologic Map of the Iditarod D-1 Quadrangle, Alaska: Alaska Geological and Geophysical Surveys Professional Report 78, scale 1:63,360.
- Bundtzen, T.K. and Laird, G.M., 1983c, Preliminary geologic map of northeastern Iditarod C-3 Quadrangle, Alaska, Alaska Geological and Geophysical Surveys Report of Investigations 83-13, scale 1:63,360.
- Bundtzen, T.K. and Swanson, S.E., 1984, Geology and petrology of igneous rocks in Innoko River area, Alaska, Geological Society of America Abstracts with Programs, v. 16, no. 5, p. 273.
- Cady, W.M., Wallace, R.E., Hoare, J.M., and Webber, E.J., 1955, The central Kuskokwim Region, Alaska: U.S. Geological Survey Professional Paper 268, 132 p.
- Chapman, R.M., Patton, W.W., Jr., and Moll, E.J., 1985, Reconnaissance geologic map of the Ophir quadrangle, Alaska: U.S. Geological Survey Open-File Report 85-302, 17 pages.
- Chapman, R.M., Weber, F.R., and Taber, B., 1971, Preliminary geologic map of the Livengood quadrangle, Alaska: U.S. Geological Survey Open-File Report OF-483, scale 1:250,000.
- Christiansen, R.L. and Lipman, P.W., 1972, Cenozoic volcanism and plate-tectonic evolution of the western United States-Part II, Late Cenozoic: Phil Trans. R. Soc. London A, v. 271, p. 249-284.
- Clague, D.A., Dalrymple, G.B., and Moberly, R., 1975, Petrography and K-Ar ages of dredged volcanic rocks of the western Hawaiian Ridge and southern Emperor Seamountchain: Geological Society of America Bulletin, v.86, p. 991-998.

- Clague, D.A. and Frey, F.A., 1982, Petrology and trace element geochemistry of the Honolulu Volcanics, Oahu: Implications for the oceanic mantle below Hawaii, *Journal of Petrology*, v. 23, p. 447-504.
- Coe, R.S., Globberman, B.R., Plumley, P.W., and Thrupp, G.A., 1985, Paleomagnetic results from Alaska and their tectonic implications, *in* Howell, D.G., ed., *Tectonostratigraphic terranes of the Circum-Pacific region: Houston, Circum-Pacific Council for Energy and Mineral Resources*, v.1, p.85-108.
- Cooper, A.K., Marlow, M.S., and Scholl, D.W., 1986, Geologic framework of the Bering Sea crust, *in* D.W. Scholl, ed., *Geology and Resource Potential of the Continental Margin of Western North America and Adjacent Ocean Basins-- Beaufort Sea to Baja California: Circum-Pacific Council for Energy and Mineral Resources Earth Science Series*, v. 6, Houston, Texas, p. ????
- Cox, A. and Engebretson, D., 1985, Change in motion of Pacific plate at 5 M.y. BP: *Nature*, v. 313, p. 472-474.
- Cox, A., Hopkins, D.M., and Dalrymple, G.B., 1966, Geomagnetic polarity epochs: Pribilof Islands, Alaska: *Geological Society of America Bulletin*, v. 77, p. 883-910.
- Csejtey, B., Jr., 1974, Reconnaissance geologic investigations in the Talkeetna Mountains, Alaska: U.S. Geological Survey Open-File Report 74-147, 53 p, 1:63,360 scale.
- Csejtey, B., Jr., Mullen, M.W., Cox, D.P., Gilbert, W.G., Yeend, W.E., Smith, T.E., Wahrhaftig, C., Craddock, C., Brewer, W.M., Sherwood, K.W., Hickman, R.G., Stricker, G.D., St. Aubin, D.R., and Goertz, D.J., III, 1986, Geology and geochronology of the Healy quadrangle: U.S. Geological Survey Open-File Report 86-396, 90 p., 4 sheets, 1:250,000 scale.
- Csejtey, B., Jr., Nelson, W.H., Jones, D.L., Silberling, N.J., Dean, R.M., Morris, M.S., Lanphere, M.A., Smith, J.G., and Silberman, M.L., 1978, Reconnaissance geologic map and geochronology, Talkeetna Mountains quadrangle, northern part of Anchorage quadrangle, and southwest corner of Healy quadrangle, Alaska, U.S. Geological Survey Open-File Report 78-558-A, 60 p., 1 sheet, 1:250,000 scale.
- Davis, A.S. and Plafker, G., 1986, Eocene basalts from the Yakutat terrane: Evidence for the origin of an accreting terrane in southern Alaska: *Geology*, v.14, p. 963-966.
- Decker, J.E. and Gilbert, W.G., 1978, The Mount Galen volcanics - A new middle Tertiary volcanic formation in the central Alaska Range: Alaska Division of Geological and Geophysical Surveys Geologic Report 59, 11 p., 1:63,360 scale.
- Decker, J., Reifentstahl, R.R., and Coonrad, W.L., 1984, Compilation of geologic data from the Russian Mission A-3 quadrangle Alaska: Alaska Division of Geological and Geophysical Surveys Report of Investigations, 84-19, 1:63,360 scale.

- Decker, J., Reifentstahl, R.R., and Coonrad, W.L., 1985, Compilation of geologic data from the Sleetmute A-7 quadrangle, southwestern Alaska: Alaska Division of Geological and Geophysical Surveys report of Investigations 85-1, 1:63,360 scale.
- Decker, J., Reifentstahl, R.R., Robinson, M.F., and Waythomas, C.F., 1986, Geologic map of the Sleetmute A-5, A-6, B-5, and B-6 Quadrangles, Alaska: Alaska Division of Geological and Geophysical Surveys Professional Report 93, 22 p., 1 sheet, 1:250,000 scale.
- Eakin, G.R., Gilbert, W.G., and Bundtzen, T.K., 1978, Preliminary bedrock geology and mineral resource potential of west-central Lake Clark quadrangle, Alaska, Alaska Division of Geological and Geophysical Surveys Open-File Report AOF 118, 15 p., scale 1:125,000.
- Eggler, D.H., 1978, The effect of CO<sub>2</sub> upon partial melting of peridotite in the system Na<sub>2</sub>O-CaO-Al<sub>2</sub>O<sub>3</sub>-MgO-SiO<sub>2</sub>-CO<sub>2</sub> to 35 kb, with an analysis of melting in a peridotite-H<sub>2</sub>O-CO<sub>2</sub> system: American Journal of Science, v. 278, p. 305-343.
- Eggler D.H. and Holloway, J.R., 1977, Partial melting of peridotite in the presence of H<sub>2</sub>O and CO<sub>2</sub>: principles and review, in Dick, H.J.B., ed., Magma Genesis: Bulletin Oregon Department of Mineral Industries, v. 96, 169-83.
- Engelbretson, D.C., Cox, A., and Gordon, R.G., 1982, Relative motions between oceanic and continental plates in the Pacific Basin: Geological Society of America Special Paper 206, 59 p.
- Foster, H.L., 1976, Geologic Map of the Eagle Quadrangle, Alaska: U.S. Geological Survey Map I-922, scale 1:250,000.
- \_\_\_\_\_, 1970, Reconnaissance geologic map of the Tanacross quadrangle, Alaska: U.S. Geological Survey Map I-593, scale 1:250,000.
- \_\_\_\_\_, 1967, Geology of the Mount Fairplay area, Alaska: U.S. Geological Survey Bulletin 1241-B.
- Foster, H.L., Forbes, R.B., and Ragan, D.M., 1966, Granulite and peridotite inclusions from Prindle Volcano, Yukon-Tanana Upland, Alaska: U.S. Geological Survey Professional Paper 550-B, p. B115-B119.
- Foster, H.L., Laird, J., Keith, T.E.C., Cushing, G.W. and Menzies, W.D., 1983, Preliminary geologic map of the Circle quadrangle, Alaska: U.S. Geological Survey Open-File Report 83-170A, 29 p., scale 1:250,000.
- Francis, D.M., 1976a, The origin of amphibole in lherzolite xenoliths from Nunivak Island, Alaska: Journal of Petrology, v. 17, pp. 357-378.

- \_\_\_\_\_, 1976b, Corona-bearing pyroxene granulite xenoliths and the lower crust beneath Nunivak Island, Alaska: *Canadian Mineralogist*, v. 14, p. 291-298.
- \_\_\_\_\_, 1978, The implications of the compositional dependence of texture in spinel lherzolite xenoliths: *Journal of Geology*, v. 86, p. 473-486.
- Frey, F.A. and Clague, D.A., 1983, Geochemistry of diverse basalt types from Loihi Seamount Hawaii: Petrogenic implications: *Earth and Planetary Science Letters*, v. 66, p. 337-355.
- Frey, F.A., Green, D.H., and Roy, S.D., 1978, Integrated models of basalt petrogenesis: A study of quartz tholeiites to olivine melilitites from south eastern Australia utilizing geochemical and experimental petrological data: *Journal of Petrology*, v. 19, p. 463-513.
- Gemuts, I., Puchner, C.C., and Steefel, C.I., 1983, Regional geology and tectonic history of western Alaska: *Journal of Alaska Geological Society*, v. 3, p. 67-85.
- Gilbert, W.G., Ferrell, V.M., and Turner, D.L., 1976, The Teklanika Formation - a new Paleocene volcanic formation in the central Alaska Range, Alaska State Division of Geological and Geophysical Surveys Geologic Report 47, 16 pages.
- Gill, J.B., 1981, *Orogenic Andesites and Plate Tectonics*: Springer-Verlag, New York, 390 p.
- Globerman, B.R. and Coe, R.S., 1984, Paleomagnetic results from Upper Cretaceous volcanic rocks in northern Bristol Bay, SW Alaska, and tectonic implications, In Howell, D.G., Jones, D.L., Cox, A. and Nur, A. eds., *Proceedings of the Circum-Pacific terrane conference*: Stanford University Publications in the Geological Sciences, v. 18, p. 98-102.
- Globerman, B.R., 1985, A paleomagnetic and geochemical study of upper Cretaceous to lower Tertiary volcanic rocks from the Bristol Bay region, southwestern Alaska, unpublished Ph.D. thesis, University of California Santa Cruz, 292 p.
- Godson, R.H., 1984, Composite Magnetic Anomaly Map of the United States, Part B: Alaska and Hawaii: U.S. Geological Survey Geophysical Investigations Map GP-954-B, scale 1:2.5 million.
- Grantz, A., 1966, Strike-slip faults in Alaska: U.S. Geological Survey Open-File Report 267, 82p.
- Grove, T.L. and Donnelly-Nolan, J.M., 1986, The evolution of young silicic lavas at Medicine Lake Volcano, California: implications for the origin of compositional gaps in calc-alkaline series lavas: *Contributions to Mineralogy and Petrology*, v. 92, p. 281-302.

- Harbert, W.P., Frei, L.S., Cox, A., and Engebretson, D.C., in press Relative motions between Eurasia and north America in the Bering Sea Region: Tectonophysics.
- Harris, R.A., 1985, Paleomagnetism, geochronology and paleotemperature of the Yukon-Koyukuk province, Alaska, unpublished M.S. thesis, University of Alaska, Fairbanks, 143 p.
- Hawkins, J.W. and Melchior, J.T., 1985, Petrology of Mariana Trough and Lau Basin Basalts: *Journal of Geophysical Research*, v. 90, p. 11, 431-11, 468.
- Hildreth, W.E., 1981, Gradients in silicic magma chambers: Implications for lithospheric magmatism: *Journal of Geophysical Research*, v. 86, p. 10153-10192.
- Hildreth, W.E. and Moorbath, S., 1986, Continental influence on Chilean arc magmatism, in Sixth International Conference, Geochronology, Cosmochronology and Isotope Geology, Abstracts: *Terra Cognita*, v. 6, no. 2, p. 196.
- Hillhouse, J.W. and Gromme', C.S., 1982, Limits to northward drift of the Paleocene Cantwell Formation, central Alaska: *Geology*, v. 10, p. 552-556.
- Hillhouse, J.W., Gromme', C.S., and Csejtey, B., Jr., 1985, Tectonic implications of paleomagnetic poles from early Tertiary volcanic rocks, south-central Alaska: *Journal of Geophysical Research*, v. 90.
- Hillhouse, J.W. and Coe, R.S., in press, Paleomagnetic data from Alaska: in Plafker, G., Jones, D.L., and Berg, H.C., eds., *Geology of Alaska: Boulder, Colorado, Geological Society of America, The Geology of North America, Chapter 5F, p.--*.
- Hoare, J.M. and Condon, W.H., 1966, Geologic Map of the Kwiguk and Black Quadrangles, western Alaska, U.S. Geological Survey Miscellaneous Geological Investigation Map I-469, scale 1:250,000.
- Hoare, J.M. and Condon, W.H., 1968, Geologic map of the Hooper Bay quadrangle, Alaska: U.S. Geological Survey Miscellaneous Geological Investigation Map I-523, scale 1:250,000.
- Hoare, J.M. and Condon, W.H., 1971a, Geologic Map of the Marshall Quadrangle, Western Alaska, U.S. Geological Survey Miscellaneous Geological Investigation Map I-668, scale 1:250,000.
- Hoare, J.M. and Condon, W.H., 1971b, Geologic Map of the St. Michael quadrangle, Alaska, U.S. Geological Survey Miscellaneous Geologic Investigations Map I-682, scale 1:250,000.
- Hoare, J.M., Condon, W.H., Cox, A., and Dalrymple, G.B., 1968, Geology, paleomagnetism, and potassium-argon ages of basalts from Nunivak Island, Alaska, in Coats, R.R.,

- Hay, R.L., and Anderson, C.A., eds., *Studies in Volcanology: A memoir in honor of Howell Williams*, Geological Society of America Memoir 116, p. 377-414.
- Hoare, J.M. and Coonrad, W.L., 1959b, *Geology of the Russian Mission quadrangle, Alaska*, U.S. Geological Survey Miscellaneous Geologic Investigations Map I-292, scale 1:250,000.
- Hoare, J.M. and Coonrad, W.L., 1978a, *A Tuya in Togiak Valley, southwest Alaska*: U.S. Geological Survey *Journal of Research*, v. 6, p. 193-201.
- Hoare, J.M. and Coonrad, W.L., 1978b, *Geologic Map of the Goodnews and Hagemeister Island Quadrangles Region, Southwestern Alaska*, U.S. Geological Survey Open-File Report 78-9B, scale 1:250,000.
- Hoare, J.M. and Coonrad, W.L., 1980, *The Togiak Basalt: A new formation in southwestern Alaska*: U.S. Geological Survey Bulletin 1482-C, 11 p.
- Hopkins, D.M., 1976, *Fault history of Pribilof Islands and its relevance to bottom stability in St. George Basin* *in* *Environment Assessment of the Alaskan Continental Shelf*: Boulder, Colorado, Environmental Research Laboratories, v. 13, p. 41-67.
- Hopkins, D.M., 1963, *Geology of the Imuruk Lake Area, Seward Peninsula, Alaska*, U.S. Geological Survey Bulletin 1141-C, 101 p.
- Hudson, Travis, 1979, *Mesozoic plutonic belts of southern Alaska*: *Geology*, v. 7, p. 230-234.
- Hudson, Travis, in press, *Crustal melting events in Alaska*, *in* Plafker, G., Jones, D.L., and Berg, H.C., eds., *Geology of Alaska*: Boulder, Colorado, Geological Society of America, *The Geology of North America*, Chapter 5F, p.--.
- Hudson, Travis and Plafker, George, 1978, *Kigluaik and Bendeleben faults, Seward Peninsula*: U.S. Geological Survey Circular 772-B, p. B47-B50.
- Irvine, T.N. and Baragar, W.R.A., 1971, *A guide to the chemical classification of the common volcanic rocks*: *Canadian Journal of Earth Sciences*, v. 8, p. 523-548.
- Irving, A.J., 1974, *Megacrysts from the Newer Basalts and other basaltic rocks of southeastern Australia*: *Geological Society of America Bulletin*, 1974, v. 85, p. 1503-1514.
- Irving, A.J. and Frey, F.A., 1984, *Trace element abundances in megacrysts and their host basalts: Constraints on partition coefficients and megacryst genesis*: *Geochimica et Cosmochimica Acta*, v. 48, p. 1201-1221.
- Jones, D.L., Silberling, N.J., Coney, P. and Plafker, G., 1987, *Lithotectonic terrane maps of Alaska (west of the 141st meridian)*, U.S. Geological Survey Miscellaneous Field Studies Map MF-1874-A, scale 1:2,500,000.

- Kay, R.W., 1977, Geochemical constraints on the origin of Aleutian magmas, *in* Talwani, M. and Pittman, eds., *Island Arcs, Seep Sea Trenches and Back-Arc Basins*, Maurice Ewing Series 1, AGU, p.229-242.
- Kay, R.W., Sun, S.S., and, Hu Lee, C.N.,1978, Pb and Sr isotopes in volcanic rocks from the Aleutian Islands and Pribilof Islands, Alaska: *Geochimica et Cosmochimica Acta*, v.42, p.263-273.
- Kerin, L.J., 1976, The reconnaissance petrology of the Mt. Fairplay igneous complex, unpublished M.S. thesis, University of Alaska, Fairbanks, 95 p.
- King, B.C., 1970, Volcanicity and rift tectonics in East Africa, *in* Clifford, T.N. and Gass, I.G., eds., *African Magmatism and Tectonics*, Hofner, New York, P.263-283.
- Kirschner, C.E., in press a, Interior Basins of Alaska, *in* Plafker, G., Jones, D.L., and Berg, H.C., eds., *Geology of Alaska: Boulder, Colorado, Geological Society of America, The Geology of North America*, Chapter 5F, p.--.
- Kushiro, I., 1975, On the nature of silicate melt and its significance in magma genesis: regularities in the shift of the liquidus boundaries involving olivine, pyroxene, and silica minerals: *American Journal of Science*, v. 275, p. 411-43.
- Lanphere, M.A. and Dalrymple, G.B., 1980, Age and strontium isotopic composition of the Honolulu Volcanic Series, Oahu, Hawaii: *American Journal of Science*, V. 280-A, p. 736-751.
- Lanphere, M.A., Dalrymple, G.B., and Clague, D.A., 1980, Rb-Sr systematics of basalts from the Hawaiian-Emperor volcanic chain: *Initial Report Deep Sea Drilling Project*, v. 55, p. 695-706.
- Lanphere, M.A. and Reed, B.L., 1985, The McKinley Sequence of granitic rocks: A key element in the accretionary history of southern Alaska: *Journal of Geophysical Research*, v. 90, p. 11,413-11,430.
- Lee-Wong, Florence, Vallier, T.L., Hopkins, D.M., and Silberman, M.L., 1979, Preliminary report on the petrography and geochemistry of basalts from the Pribilof Islands and vicinity, southern Bering Sea: *U.S. Geological Survey Open-File Report 79-1556*, 51 p.
- Lerbekmo, T.F. and Campbell, F.A., 1969, Distribution, composition, and source of the White River ash, Yukon Territory: *Canadian Journal of Earth Science*, v. 6, p. 109-116.
- Lipman, P.W. Doe, B.R., Hedge, C.E., Steven, T.A., 1978, Petrologic evolution of the San Juan volcanic field, southwestern Colorado: Lead and strontium isotopic evidence: *Geological Society of America Bulletin*, v. 89, p. 59-82.

- Lipman, P., Prostka, H.J., and Christiansen, R.L., 1972, Cenozoic volcanism and plate tectonic evolution of western United States, part I, Early and middle Cenozoic: *Philosophical Transactions, Royal Society of London A*, v. 271, p. 249-284.
- Luedke, R.G. and Smith, R.L., 1986, Map showing distribution, composition, and age of Late Cenozoic volcanic centers in Alaska: U.S. Geological Survey Miscellaneous Investigations Series Map I-1091-F, scale 1:1,000,000.
- MacDonald, G.A. and Katsura, T., 1964, Chemical composition of Hawaiian lavas: *Journal of Petrology*, v. 6, p. 82-133.
- Magoon, L.B., Adkison, W.L., and Egbert, R.M., 1976, K-Ar age dates, and petroleum operations, Cook Inlet area, Alaska: U.S. Geological Miscellaneous Investigations Series Map I-1019, scale 1:250,000.
- Mahood, G. and Hildreth, W., 1983, Large partition coefficients for trace elements in high-silica rhyolites: *Geochimica et Cosmochimica Acta*, v. 47, p. 11-30.
- Mark, Robert Kent, 1971, Strontium isotopic study of basalts from Nunivak Island, Alaska, unpublished Ph.D. thesis Stanford University, 50 p.
- Marlow, M.S. and Cooper, A.K., 1986, Regional geology of the Beringian margin, *in* Proceedings of O.J.I., International Seminar on the formation of ocean margins: University of Tokyo, Tokyo, Japan, p.
- Marsh, B.D., in press, Age, Character, and significance of Aleutian arc volcanic rocks, *in* Plafker, G., Jones, D.L., and Berg, H.C., eds., *Geology of Alaska: Boulder, Colorado, Geological Society of America, The Geology of North America, Chapter 5F, p.--*.
- Marshall, M., 1978, The magnetic properties of some DSDP basalts from the North Pacific and inferences for Pacific plate tectonics: *Journal of Geophysical Research*, V.83, no. B1, p. 289-308.
- Miller, T.P. and Richter, D.H., in press, Quaternary volcanism in the Alaska Peninsula and Wrangell Mountains, Alaska, *in* Plafker, G., Jones, D.L., and Berg, H.C., eds., *Geology of Alaska: Boulder, Colorado, Geological Society of America, The Geology of North America, Chapter 5F, p.--*.
- Menzies, M.A., Arculus, R.J., Best, M.G., Bergman, S.C., Ehrenberg, S.N., Irving, A.J., Roden, M.F., and Schulze, D.J., 1987, A record of subduction processes and within-plate volcanism in lithospheric xenoliths of the southwestern USA *in* Nixon, P.H., *ed.*, *Mantle Xenoliths*, J.Wiley and Sons, LTD, p. 59-74.
- Menzies, Martin and Murthy, V. Rama, 1980a, Mantle metasomatism as a precursor to the genesis of alkaline magmas - isotopic evidence: *American Journal of Science*, v. 280-A, p. 622-638.

- Menzies, Martin and Murthy, V. Rama, 1980b, Nd and Sr isotope geochemistry of hydro mantle nodules and their host alkali basalts. Implications for local heterogeneities in metasomatically veined mantle: *Earth and Planetary Science Letters*, 46, p. 323-334.
- Miller, T.P. and Lanphere, M.A., 1981, K-Ar age measurements on obsidian from the Little Indian River locality in interior Alaska, *in* Albert, N.R.D., and Hudson, T., *The United States Geological Survey in Alaska: Accomplishments during 1979*, U.S. Geological Survey Circular 823-B, p. B39-B42.
- Moll, E.J. and Arth, J.G., 1985, Sr and Nd isotopes from Late Cretaceous-early Tertiary volcanic fields in western Alaska: Evidence against old radiogenic continental crust under the Yukon-Koyukuk basin, *EOS*, v. 66, no. 46, p. 1102.
- Moll, E.J. and Patton, W.W., Jr., 1982, Preliminary report on the Late Cretaceous-early Tertiary volcanic and related plutonic rocks in western Alaska, *in* Coonrad, W.L., ed., *The United States Geological Survey in Alaska: Accomplishments during 1980*, Geological Survey Circular 844, p. 73-76.
- Moll, E.J. and Patton, W.W., Jr., 1983, Late-Cretaceous-early Tertiary calc-alkalic volcanic rocks of western Alaska: *Geological Society of America Abstracts with Programs*, 1983, v. 15, no. 5, p. 406.
- Moll, E.J., Silberman, M.L., and Patton, W.W., Jr., 1981, Chemistry, mineralogy, and K-Ar ages of igneous and metamorphic rocks of the Medfra Quadrangle, Alaska: U.S. Geological Survey Open-File Report 80-81C, 2 sheets, scale 1:250,000.
- Moll-Stalcup, E.J., 1987, The petrology and Sr and Nd isotopic characteristic of five Late Cretaceous-early Tertiary volcanic fields in western Alaska: unpublished Ph.D. thesis, Stanford University, 310 p.
- Moll-Stalcup, E.J. and Arth, J.G., 1989, The nature of the crust in the Yukon-Koyukuk province as inferred from the chemical and isotopic composition of five Late Cretaceous and early Tertiary volcanic fields in western Alaska, *Journal of Geophysical Research*, v.94, p.15,989-16,020.
- Moll-Stalcup, E.J. and Arth, J.G., in prep, The petrology and Sr and Nd isotopic composition of the Blackburn Hills volcanic field, western Alaska: submitted to *Journal of Petrology*.
- Moll-Stalcup, E.J., Brew, D.A., and Vallier, T.L., in press, Latest Cretaceous and Cenozoic magmatic rocks in Alaska, *in* Plafker, G., Jones, D.L., and Berg, H.C., eds., *Geology of Alaska: Boulder, Colorado, Geological Society of America, The Geology of North America*, Chapter 5F, p.--.
- Moore, J.C., Byrne, T., Plumley, P.W., Reid, M., Gibbons, H., and Coe, R.S., 1983, Paleogene evolution of the Kodiak Islands, Alaska: Consequences of Ridge-trench interaction in a more southerly latitude: *Tectonics*, v. 2, p. 265-293.

- Morrison, G.W., 1980:, Characteristics and Tectonic setting of the shoshonite rock association: *Lithos*, v.13, p. 97-108.
- Nakamura, E., Campbell, I.H., and Sun, S., 1985, The influence of subduction processes on the geochemistry of Japanese alkaline basalts: *Nature*, v. 316, p. 55-58
- Nakamura, K., Jacob, K.H., and Davies, J.H., 1977, Volcanoes as possible indicators of tectonic stress orientation: *Aleutians and Alaska: Pageoph*, Birkhauser Verlag, Basel v. 115, p. 87-112.
- Nakamura, K., and Uyeda, S., 1980, Stress gradient in arc-back arc regions and plate subduction: *Journal of Geophysical Research*, v. 85, p. 6419-6428.
- Nakamura, K., Plafker, G., Jacob, K.H., and Davies, J.N., 1980, A tectonic stress trajectory map of Alaska using information from volcanoes and faults: *Bulletin of the Earthquake Research Institute*, v. 55, p. 89-100.
- Nelson, S.W., Dumoulin, J.A., and Miller, M.L., 1985, Geologic map of the Chugach National Forest, Alaska, U.S. Geological Survey Map-1645-B, 16 p. , one sheet, scale 1:250,000
- Nelson, W.H., Carlson, C., and Case, J.E., 1983, Geologic Map of the Lake Clark Quadrangle, Alaska, U.S. Geological Survey Miscellaneous Field Studies Map, MF1114-A, scale 1:250,000.
- Nokleberg, W.J., Albert, N.R.D., Bond, G.C., Herzon, P.L., Miyaoka, R.T., Nelson, W.H., Richter, D.H., Smith, T.E., Stout, J.H., Yeend, W., and Zehner, R.E., 1982, Geologic map of the southern Mount Hayes quadrangle, Alaska: U.S. Geological Survey Open-File Report 82-52, 27 p., 1 sheet, scale 1:250,000.
- Nokleberg, W.J., Jones, D.L., and Silberling, N.J., 1985, Origin and tectonic evolution of the MacLaren and Wrangellia terranes, eastern Alaska Range, Alaska: *Geological Society of America Bulletin*, v. 96, p. 1251-1270.
- Patton, W.W., Jr., 1966, Regional geology of the Kateel River Quadrangle, Alaska, U.S. Geological Survey Miscellaneous Geologic Investigations Map I-437, scale 1:250,000.
- Patton, W.W., Jr., Box, S.E., and Moll-Stalcup, E.J., and Miller, T.P., in press b, Geology of west-central Alaska, *in* Plafker, G., Jones, D.L., and Berg, H.C., eds., *Geology of Alaska*: Boulder, Colorado, Geological Society of America, *The Geology of North America*, Chapter 5F, p.--.
- Patton, W.W., Jr. and Csejtey, Be'la Jr., 1980, Geologic Map of St. Lawrence Island, Alaska: U.S. Geological Survey Miscellaneous Geologic Investigations Map I-1203, scale 1:250,000.

- Patton, W.W., Jr. and Hoare, J.M., 1968, The Kaltag fault, west-central Alaska, in Geological Survey Research in 1968: U.S. Geological Survey Professional Paper 600D, p. D147-D153.
- Patton, W.W., Jr., Lanphere, M.A., Miller, T.P., and Scott, R.A., 1976, Age and tectonic significance of volcanic rocks on St. Matthew Island, Bering Sea, Alaska: U.S. Geological Survey Journal of Research, v. 4, p. 67-73.
- Patton, W.W., Jr. and Miller, T.P., 1970, A possible source for obsidian found in archeological sites in northwestern Alaska: Science, V. 169, p.760-761.
- Patton, W.W., Jr. and Miller, T.P., 1973, Bedrock geologic map of Beetles and southern part of Wiseman Quadrangles, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-492, scale 1:250,000.
- Patton, W.W., Jr., Miller, T.P., Berg, H.C., Gryc, G., Hoare, J.M., and Ovenshine, A.T., 1975, Reconnaissance geologic map of St. Matthew Island, Bering Sea, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map, MF-642, scale 1:125,000.
- Patton, W.W., Jr., Miller, T.P., Chapman, R.M., and Yeend, W., 1978, Geologic Map of the Melozitna Quadrangle: U.S. Geological Survey Miscellaneous Investigation Series Map I-1071, scale 1:250,000.
- Patton, W.W., Jr. and Moll, E.J., 1985, Geologic map of northern and central parts of Unalakleet quadrangle, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1749, scale 1:250,000.
- Patton, W.W., Jr., Moll, E.J., Dutro, J.T., Jr., Silberman, M.L., and Chapman, R.M., 1980, Preliminary geologic map of the Medfra quadrangle: U.S. Geological Survey Open-File Report 80-811A, scale 1:250,000.
- Perfit, M.R., Gust, D.A., Bence, A.E., Arculus, R.J. and Taylor, S.R., 1980, Chemical characteristics of island-arc basalts: Implications for mantle sources: Chemical Geology, v.30, p.227-256.
- Plumley, P.W., Coe, R.S., Byrne, T., Reed, M.R., and Moore, J.C., 1982, Paleomagnetism of volcanic rocks of the Kodiak Islands indicates northward latitudinal displacement: Nature, v. 300, p. 50-52.
- Rea, D.K., and Duncan, R.A., 1986, North Pacific plate convergence: A quantitative record of the past 140 m.y.: Geology, V.14, p.373-376.
- Reed, B.L. and Lanphere, M.A., 1972, Generalized geologic map of the Alaska-Aleutian Range batholith showing potassium-argon ages of the plutonic rocks: U.S. Geological Survey Miscellaneous Field Studies Map MF-372. 2 sheets, scale 1:1,000,000.

- Reed, B.L. and Lanphere, M.A., 1973, The Alaska-Aleutian Range batholith-Geochronology, chemistry, and relation to circum-Pacific plutonism: *Geological Society of America Bulletin*, v. 84, p. 2583-2610.
- Reed, B.L. and Lanphere, M.A., 1974a, Chemical variations across the Alaska-Aleutian Range batholith: *Journal of Research, U.S. Geological Survey*, v. 2, p. 343-352.
- Reed, B.L. and Lanphere, M.A., 1974b, Offset plutons and history of movement along the McKinley segment of the Denali fault system, Alaska: *Geological Society of America Bulletin*, v. 85, p. 1883-1892.
- Reed, B.L. and Nelson, S.W., 1980, Geologic Map of the Talkeetna Quadrangle, Alaska, U.S. Geological Survey Miscellaneous Investigations Series Map MI-1174, scale 1:250,000.
- Reifenstuhl, R.R., Decker, J., and Coonrad, W.L., 1985, Compilation of geologic data from the Taylor Mountains D-8 quadrangle, southwestern Alaska, Alaska Division of Geological and Geophysical Surveys Report of Investigations 85-4, scale 1:63,360.
- Reifenstuhl, R.R., Robinson, M.S., Smith, T.E., Albanese, M.D., and Allegro, G.A., 1984, Geologic Map of the Sleetemute B-6 Quadrangle, Alaska: Alaska Division of Geological and Geophysical Surveys Report of Investigations 84-12, scale 1:63,360.
- Robinson, M.S., Decker, J., Reifenstuhl, R.R., Murphy, J.M., and Box, S.E., 1984, Geologic Map of the Sleetemute B-5 quadrangle, Alaska, Alaska Division of Geological and Geophysical Surveys Report of Investigations 84-10, scale 1:63,360.
- Roden, M.F., 1982, Geochemistry of the earth's mantle, Nunivak Island, Alaska and other areas: evidence from xenolith studies: unpublished Ph.D. thesis, Massachusetts Institute of Technology, p. .
- Roden, M.F., Frey, F.A. and Francis, D.M., 1984, An example of consequent mantle metasomatism in peridotite inclusions from Nunivak Island, Alaska: *Journal of Petrology*, v. 25, p. 546-577.
- Roden, M.F. and Murthy, V. Rama, 1985, Mantle metasomatism: *Annual Review of Earth and Planetary Sciences*, v. 13, p. 269-96.
- Saunders, A.D. and Tarney, J., 1984, Geochemical characteristics of basaltic volcanism within backarc basins, *in* Kolelaar, B.P., and Howells, M.F., *Marginal Basin Geology*, Oxford, p. 59-76
- Savostin, L.A. and Karasik, A.M., 1981, Recent plate tectonics of the Arctic Basin and of northeastern Asia: *Tectonophysics*, v.74, p.11-145.

- Scholl, D.W., Vallier, T.L., and Stevenson, A.J., 1986, Geologic evolution and petroleum geology of the Aleutian ridge, *in* D.W. Scholl, ed., *Geology and Resource Potential of the Continental Margin of Western North America and Adjacent Ocean Basins - Beaufort Sea to Baja California: Circum-Pacific Council for Energy and Mineral Resources Earth Science Series, v. 6, Houston, Texas, p. 59-72.*
- Silberman, M.L. and Grantz, A., 1984, Paleogene volcanic rocks of the Matanuska Valley area and the displacement history of the Castle Mountain fault, *in* Coonrad, W.L. and Elliott, R.L., eds. *The U.S. Geological Survey in Alaska: Accomplishments During 1981: U.S. Geological Survey Circular 868, p. 82-86.*
- Silberman, M.L., Moll, E.J., Chapman, R.M., Patton, W.W., Jr., and Connor, C.L., 1979, Potassium-argon age of granitic and volcanic rocks from the Ruby, Medfra, and surrounding quadrangles, west-central Alaska, *in* Johnson, K.M., and Williams, J.R., eds., *The United States Geological Survey in Alaska: Accomplishments during 1978: U.S. Geological Survey Circular 804-B, p. B863-B866.*
- Simpson, G.L., Vallier, T.L., Pearl, J.E., and Lee-Wong, Florence, 1979, Potassium-argon ages and geochemistry of basalt dredged near Saint George Island, southern Bering Sea, *in* Johnson, K.M., and Williams, J.R., eds, *The United Geological Survey in Alaska: Accomplishments during 1978: U.S. Geological Survey Circular 804-B, p. B134-B136.*
- Sinclair, P.D., Templeman-Kluit, D.J., and Medaris, L.G., Jr., 1978, Lherzolite nodules from a Pleistocene cinder cone in central Yukon, *Canadian Journal of Earth Sciences, v. 15, p. 220-226.*
- Smirnov, Ya.B., and Sugrobov, V.M., 1979, Heat flow in the Northwest Pacific Ocean, *Priroda, v. 8, p.94-101.*
- Smirnov, Ya.B., and Sugrobov, V.M., 1979, Terrestrial heat flow in the Kuril-Kamchatda and Aleutian provinces:1, Heat flow and tectonics: *Vulkanologiya i Seismologiya Akademiya Nauk SSSR, Moscow, no. 1, p. 59-73.*
- Snyder, W.S., Dickinson, W.R., and Silberman, M.L., 1976, Tectonic implications of space-time patterns of Cenozoic magmatism in the western United States: *Earth and Planetary Science Letters, v. 32, p. 91-106.*
- Streckeisen, A., 1980, Classification and nomenclature of volcanic rocks, lamprophyres, carbonatites, and melilitic rocks: IUGS subcommission on the systematics of igneous rocks, *Geology, v. 7, p. 331-335.*
- Swanson, S.E., Turner, D.L., and Fores, R.B., 1981, Petrology and geochemistry of Tertiary and Quaternary basalts from the Seward Peninsula, western Alaska: *Geological Society of America Abstract with Programs, v. 13, no. 7, p. 563.*

- Thompson, R.N., Morrison, M.A., Dickin, A.P., and Hendry, G.L., 1982, Continental flood basalts .... Arachnids rule OK?, *in* Hawksworth, C.J. and Norry, eds., *Continental Basalts and Mantle Xenoliths*, Shiva, p. 158-185.
- Thompson, R.N., Morrison, M.A., Hendry, G.L., and Parry, S.J., 1984, An assessment of the relative roles of crust and mantle in magma genesis: an elemental approach: *Phil. Trans. Royal Soc. London A*, v. 310, p. 549-590.
- Throckmorton, M.L. and Patton, W.W., Jr., 1978, Contact-metasomatic magnetite deposit, Medfra quadrangle, Alaska: U.S. Geological Survey Open-File Report 78-26, 13 p.
- Thrupp, G.A. and Coe, R.S., 1986, Early Tertiary paleomagnetic evidence and the displacement of southern Alaska: *Geology*, v. 14, p. 213-217.
- Till, A.B. and Dumoulin, J., in press, Geology of Seward Peninsula and St. Lawrence Island, Alaska *in* Plafker, G., Jones, D.L., and Berg, H.C., eds., *Geology of Alaska: Boulder, Colorado, Geological Society of America, The Geology of North America, Chapter 5F*, p.--.
- Tolson, R.B., 1986, Structure and stratigraphy of the Hope Basin, *in* D.W. Scholl, ed., *Geology and Resource Potential of the Continental Margin of Western North America and Adjacent Ocean Basins-- Beaufort Sea to Baja California: Circum-Pacific Council for Energy and Mineral Resources Earth Science Series*, v. 6, Houston, Texas, p. 59-72.
- Turner, D.L. and Forbes, R.B., 1980, A geological and geophysical study of the geothermal energy potential of Pilgrim Springs, Alaska: University of Alaska, Geophysical Institute Report UAG R-271, 165 p.
- Turner, D.L., Swanson, S.E., and Wescott, E., 1981, Continental rifting--A new tectonic model for geothermal exploration of the central Seward Peninsula, Alaska: Geothermal Resources Council, *TRANSACTIONS*, v.5, p.213-216.
- Tysdal, R.G. and Case, J.E., 1979, Geologic map of the Seward and Blying Sound quadrangle, U.S. Geological Survey Miscellaneous Investigations Series Map I-1150, scale 1:250,000.
- Vallier, T.L., Scholl, D.W., Fisher, M.A., Bruns, T.R., von Huene, R., and Stevenson, A.J., in press, Geology of the Aleutian structural arc, *in* Plafker, G., Jones, D.L., and Berg, H.C., eds., *Geology of Alaska: Boulder, Colorado, Geological Society of America, The Geology of North America, Chapter 5F*, p.--.
- Von Drach, V., Marsh, B.D., Wasserburg, G.J., 1986, Nd and Sr isotopes in the Aleutians: multicomponent parenthood of island-arc magmas: *Contributions to Mineralogy and Petrology*, v. 92, p. 13-34.

- Wallace, W.K. and Engebretson, D.C., 1984, Relationship between plate motions and Late Cretaceous to Paleogene magmatism in southwestern Alaska: *Tectonics*, v. 3, p. 295-315.
- Weber, F.R., Foster, H.L., Keith, T.E.C., and Dusel-Bacon, C., 1978, Preliminary geologic map of the Big Delta quadrangle, Alaska: U.S. Geological Survey Open-File Report 78-529A, 1 pl., scale:1:250,000.
- Wilson, F.H., 1977, Some plutonic rocks of southwestern Alaska, a data compilation: U.S. Geological Survey Open-File Report 77-501, 7 p., 4 sheets.
- Wilson, F.H., 1985, The Meshik arc - an Eocene to earliest Miocene magmatic arc on the Alaska Peninsula: Alaska Division of Geological and Geophysical Surveys Professional Report 88, 14 p.
- Wood, D.A., 1979, A variably veined suboceanic upper mantle---genetic significance for mid-ocean ridge basalts from geochemical evidence: *Geology*, v. 7, p. 499-503.
- Wyllie, P.J. and Huang, W-L., 1976, Carbonation and melting reactions in the system CaO-MgO-SiO<sub>2</sub>-CO<sub>2</sub> at mantle pressures with geophysical and petrological applications: *Contributions to Mineralogy and Petrology*, v. 54, 140-173.
- Zielinski, R.A. and Lipman, P.W., 1976, Trace-element variations at Summer Coon volcano, San Juan Mountains, Colorado, and the origin of continental-interior andesite: *Geological Society of America Bulletin*, v. 87, p. 1477-1485.
- Zonenshain, L.P., Savostin, L.A., and Sedov, A.P., 1984, Global paleogeodynamic reconstructions for the last 160 million years: *Geotectonics*, v. 18, p. 181-195.

Table 1: Tables summarizing age, petrologic, and chemical data on Late Cretaceous and Cenozoic volcanic and plutonic rocks

Late Cretaceous and early Tertiary  
Alaska Range-Talkeetna Mountains belt

Unit/Name	Age (Ma)	Classification <sup>1</sup>	K <sub>2</sub> O type <sup>2</sup>	Rock types	SiO <sub>2</sub> range (wt. %)	Mineralogy	Other comments	References
Summit Lake	73-58	Calc-alkalic (61)	Moderate-K	Granodiorite, quartz diorite	54.5 - 70	Biot, hmbld reported	Rocks having more than 65 % SiO <sub>2</sub> divided into two suites: normal calc-alkalic and trondhjemite-tonalite.	Reed and Lamphere, 1972; 1974a; M.A. Lamphere, oral comm., 1984
Mt. Susitna	72.0	Presumed calc-alkalic	No data	Granodiorite, quartz diorite, diorite	No data	No data		Magoon and others, 1976; Reed and Lamphere, 1972
Tonalite of Talkeetna Mountains	74 - 61	Calcic 62	Moderate-K	Tonalite, quartz diorite	58 - 63	Andesine, qtz, ortho, hmbld, biot, mt	Granite and granodiorite body to the west intrudes the tonalite body and is 65-67 Ma (has biot+ms)	Caejey, 1976; 1974; Caejey and others, 1978
Quartz monzonite of Tired Pup	58-55	Calc-alkalic (does not cross)	High-K	Granodiorite, granite	69.8 - 70.1	Biot; biot+ms reported	Similar to McKinley sequence (Lamphere oral comm., 1984)	Reed and Lamphere, 1972; Reed and Lamphere, 1974
McKinley sequence	57.9-56.6	Calc-alkalic (does not cross)	High-K	Granite, granodiorite	65.9 - 77.6	biot; biot+ms reported	SIR = 0.7054 - 0.7085 Rocks have normative corundum. Limited trace element data—Low Ba (150-472), low Sr (6-47), and high Nb (24-44)—suggests rocks are not arc-related	Lamphere and Reed, 1985
Crystal Creek sequence	60.5-58	Calc-alkalic	High-K	Granite, granodiorite	70.5 and 75.2	Biot and biot+hmbld reported		Reed and Lamphere, 1972; 1974a
Middle Fork pluton	58-55	Alkali-calcic		Gabbro, diorite, qtz monzdr., syenite, granite, Alkali feldspar granite	42 - 53 and 60 - 75	Gb and dior: plag ±olt:opx+cpx+bt ±K-spar; syenites: K-spar: qtz+cpx+fsyal:to; granite: plag+K-spar +qtz+biot+hmbld; alkali feldspar granite: plag+K-spar+biot +Na-amphiboles	An orogenic type bimodal suite of multiple coeval magmas. Alkali feldspar granite is peralkaline	Soie, D.N., 1988
Granite intrusions of the northern Talkeetna Mountains	58.6-50.4	Probably calc-alkalic	No data	Granite, granodiorite	No data	Qtz+k-spar+plag+biot, rare hmbld	Includes Tertiary granite intrusions in the Healy and Talkeetna Mountains Quadrangles. Several plutons are associated with tin mineralization. Caejey considers these plutons to be the subvolcanic equivalents of nearby volcanic rocks	Caejey, 1974; Caejey and others, 1978; Caejey and others, 1986
Volcanic and related plutonic rocks of Lake Clark region	71-58	Presumed calc-alkalic	No data	Rhyodacite, intermediate to mafic volcanic rocks	No data	Rhyodacite: plag: qtz:biot:hmbld intermediate-mafic: plag+hmbld:pyr	Age uncertain. Description includes undivided rocks of Paleocene and Eocene age. Rhyodacites include breccia flows, ash-flow tuffs and intrusive rocks	Nelson and others, 1983; G.A. Thrupp, written comm., 1984; Eakin and others, 1978; Thrupp and Coe, 1986
Talkeetna Mountains volcanic field	56-51	Calc-alkaline	Moderate-K	Basalt, andesite, dacite, rhyolite	52 - 60 and 68 - 71	Basalt: ol+plag+cpx+mt+il andesite: plag+cpx:opx+mt dacite: plag:biot: qtz:biot: rhyolitic: san+plag in perlitic glass.	Volcanic field consists of a lower sequence composed of quartz latite, rhyolite, latite, and minor andesite, and an upper sequence of andesite and basalt. Description may include rocks of the upper sequence that may be as young as Miocene. Rocks have high Ba (300-700 ppm), moderate Sr (200-500), moderate Y(10-30), moderate, Yb(1.5-3) and low Zr(70-150)	Caejey and others, 1978; Gilbert and others, 1976; Hillhouse, Grommé, and Caejey, 1985
Volcanic rocks of Cantwell Formation	61-42 (Gilbert and others, 1976) consider the suite Paleocene age)	Calc-alkalic or calcic	Moderate to high-K	Andesite, rhyolite, dacite, minor basalt, quartz diorite	52 - 62 and 71 - 77	Rhy: san:dt:plag: qtz:biot; Mafic-intermediate: labrad-andes+cpx+mt:ol; Qtz-diorite: plag: andesine rimmed by K-spar+cpx+qtz:ol: garnet	Data show considerable scatter on major element Harker diagrams, probably due to post-depositional alteration. Suite has more TiO <sub>2</sub> - about 2 weight percent, than other contemporaneous suites. Equivalent to Teklanika Formation of Gilbert and others (1976).	Gilbert and others, 1976; Caejey and others, 1986

<sup>1</sup>Classification for Late Cretaceous to early Tertiary suites based on Peacock diagram. Numbers refer to weight percent SiO<sub>2</sub> where CaO and Na<sub>2</sub>O<sub>3</sub> + K<sub>2</sub>O cross. Late Cenozoic rocks are classified on the basis of normative mineralogy. Basalts having normative nepheline are alkali basalt or alkali olivine basalt if they contain more than 10% normative olivine. Basalt having normative hypersthene is called tholeiitic or olivine tholeiitic if it contains more than 10% normative olivine. Basanites have 10 to 20% normative nepheline; nephelinites have more than 20% normative nepheline. Hailites have more than 5% total alkalis and less than 5% MgO. Fe<sub>2</sub>O<sub>3</sub>/FeO set to 0.15 before norms are calculated.

<sup>2</sup>K<sub>2</sub>O-type defined by Gill, (1981). Shoshonites have higher K<sub>2</sub>O than orogenic andesites but are similar to orogenic andesites in other major elements.

Late Cretaceous and early Tertiary  
Kuskokwim Mountains belt  
northern volcanic fields

Unit/Name	Age (Ma)	Classification	K <sub>2</sub> O type	Rock types	SiO <sub>2</sub> range (wt. %)	Mineralogy	Other comments	References
East Susitna batholith	57.0-54.5 70 (U-Pb)			Diorite, gneissodiorite; subordinate quartz monzonite		Biot, hornbl	Nokleberg and others believe this pluton was 400 km to the south during emplacement so it may not be part of this belt. Pluton is regionally metamorphosed	Nokleberg and others, 1982
Sisach volcanic field	71.0-66.3	Calc-alkalic (do not cross but project to cross at 60.5)	High K <sub>2</sub> O	Rhyolite, dacite, minor andesite	62 and 69 - 77.5	Rhyolite: san+qtz+plag+ altered mafics; olig+san+biot ±fayalite dacite: andesine+qtz+ biot+hornbl andesite: plag+ biot+mt	Volcanic sequence at least 500 m. thick. May be borderline alkalic. K <sub>2</sub> O and Na <sub>2</sub> O appear to be strongly modified by post depositional alteration.  SIR greater than 0.7080	Moll and others, 1981; E.J. Moll-Stalcup, and R.M. Chapman, unpub. data, 1985
Nowina volcanic field	64.2-63.8	calc-alkalic (60.5)	Moderate to high K  (K <sub>2</sub> O/Na <sub>2</sub> O=0.66 for andesite)	andesite, rhyolite	55 - 63.5 and 73	Andesite: plag+cpx:opx+mt rhyolite: plag+san: biot:mt.	Andesite occurs as thick sequence of flows. Rhyolite as small domes overlying andesite sequence	Moll and others, 1981
Dishna volcanic field	Undated	Calc-alkalic	High K <sub>2</sub> O  (K <sub>2</sub> O/Na <sub>2</sub> O=1)	Rhyolite dacite	65 - 78	Rhyolite: qtz+san+biot dacite: qtz+plag+opx+ biot	Information based on only 7 analyses	Chapman and others, 1985; E.J. Moll and R.M. Chapman, unpub. data, 1985
Yema volcanic field	69-55	Calc-alkalic	Unknown	Andesite rhyolite basalt	Unknown	Rhyo: K-spar+ qtz+biot dacite: plag+ cpx+biot:hornbl andesite: plag+ pyri:biot:hornbl basalt: ol+plag +pyr	Probably moderate to high-K	M. L. Miller and T.K. Bundtzen, unpub. data, 1986

Kuskokwim Mountains belt  
volcanoplutonic complexes

Page Mtn., Lonesome Hills, and Cloudy Mountain	71-65	Borderline alkali-calcic--calc-alkalic (60)	Shoshonitic (K <sub>2</sub> O/Na <sub>2</sub> O =1.21)	Andesite, dacite, basalt, monzonite, quartz-monzonite, gneissodiorite	56 - 65.5	Basalt: ol+plag :cpx+opx andesite: plag+ cpx:biot dacite: plag+ biot+qtz+ altered mafics intrusive rocks: plag+K-spar+ cpx+biot:amph+ opaque oxides	Data include intrusive and volcanic rocks. Volcanic sequences generally hydrothermally altered by intrusive rocks. REE data suggest the volcanic rocks may be compositionally similar to less altered Nowina volcanic rocks. Lonesome Hills formerly Alone benchmark on 1:250,000 map	Moll and others, 1981
Beaver Mtns, Candle Hills, Takotna Mtn.	73-70	Borderline alkali-calcic--calc-alkalic  (data show much scatter cross near 55-58)	Shoshonitic (K <sub>2</sub> O/Na <sub>2</sub> O=1.5)	Basalt, andesite, dacite, quartz monzonite, monzonite, monzodiorite; minor syenite	51.7 - 64	Bas: plag+cpx+ol; Andes and dacite: plag+pyr; Syenite: K-spar+ qtz+plag+biot: cpx; monz and monzodiorite: K-spar+qtz+plag+ biot+cpx; qtz monzo: ortho+ qtz+plag+biot	Data include intrusive and volcanic rocks. Whole rock chemical analyses having sums less than 98 wt % or more than 7% volatiles were not used. Candle Hills are located in the northwest McGrath quadrangle; Takotna Mtn. is in the northeast Iditarod quadrangle	Bundtzen and Laird, 1982, 1983a, 1983b; 1983c;

Kuskokwim Mountains belt

intrusive rocks of the northern Kuskokwim Mountains

VonFrank Mountain, Whirlwind Ridge, Nixon Fork, Cripple Creek, and Appel Mtn. plutons	70.4-66.4	Alkali-calcic (57)	High-K to shoshonitic K <sub>2</sub> O/Na <sub>2</sub> O=1	Monzonite, quartz monzodiorite, monzonite, monzogabbro. Gabbro at Von Frank Mountain only	53.3 - 64.4	Qtz-monz: plag+ ortho+qtz+ gm amph+biot; Monz: plag+ ortho+cpx+opx+ biot+gm amph: qtz; Monzogh and gb: plag:ortho:qtz+ biot+cpx+opx+ ol:gm amph	Nixon Fork pluton intrudes limestone and is associated with a gold copper skarn. Von Frank Mountain pluton is compositionally zoned from quartz-rich monzonite in the core to layered monzogabbro and gabbro at the rim. Plutons located in the central Medfra quadrangle; see Moll and others (1981) for location of specific plutons	Moll and others, 1981
---	-----------	--------------------	--	---	-------------	---	--	-----------------------

Late Cretaceous and early Tertiary  
Kuskokwim Mountains belt  
Intrusive rocks of the northern Kuskokwim Mountains, cont'd

Unit/Name	Age (Ma)	Classification	K <sub>2</sub> O type	Rock types	SiO <sub>2</sub> range (wt. %)	Mineralogy	Other comments	References
Telida Mtn. and upper Sulukna River plutons	70.5 and 69.2	Alkali-calcic  (do not cross)	Shoshonitic (K <sub>2</sub> O/Na <sub>2</sub> O=2.25)	Granite and qtz monz; (very minor syenite at upper Sulukna River pluton)	71 and 76  (two analyses only)	Telida: plag+k-spar+qtz+biot: hmbld:opaque oxides; Upper Sulukna River pluton: granite and qtz monz: plag+ortho+qtz+biot+hmbld+cpx Syenite: ortho+qtz+sphene+diopsid+plag	Upper Sulukna River pluton intrudes limestone and is associated with a magnetite skarn. Syenite occurs as a border phase at upper Sulukna River pluton. Telida pluton is associated with abundant tourmaline. Telida pluton is located in the eastern part of the Medfra quadrangle	Moll and others, 1981; Throckmorton, and Patton, 1978
Sunshine Mountain, Meadow Creek, and West Fork plutons	63.4 and 62.5	Calc-alkalic  (do not cross, project to cross around 60.5)	High-K  (K <sub>2</sub> O/Na <sub>2</sub> O=1.14)	Granite	69 - 75	Plag+qtz+ortho or san+biot: hmbld	Meadow Creek pluton is dominantly a porphyritic hypabssal intrusion having andesine instead of orthoclase. Plutons located in the west-central part of the Medfra	Moll and others, 1981
Intrusions of Shepard Creek and intrusions of Mystery Mountains	69.9 - 62.2	Borderline calc-alkalic (61.5 Shepard Creek only)	High-K (K <sub>2</sub> O/Na <sub>2</sub> O=0.93)	Rhyolite, dacite, high silica andesite, granodiorite, qtz monz, qtz-rich monz and qtz monzodiorite	61 - 75	Rhy: plag+qtz+san+biot dacite: plag+qtz+biot andesite: plag hmbld+biot plutons: qtz+ortho+plag+biot hmbld:cpx:opx	Shepard Creek rocks consist of shallow hypabyssal rocks. Mystery Mountains rocks consist of a number of long parallel sills, small plugs, and flows. Sills are highly altered and contain abundant tourmaline. Intrusions are located in the central Medfra quadrangle	Moll and others, 1981
Stone Mtn pluton	68.3 and 66.4	Calc-alkalic (60-61)	Moderate-K (K <sub>2</sub> O/Na <sub>2</sub> O=0.47)	Monzonite, monzodiorite, tonalite, gabbro, monzogabbro	55.7 - 63	Monzogabbro and gabbro: ol+cpx+opx+plag: K-spar Monzonitic rocks: plag+ortho: qtz+biot+green hmbld:opx: cpx+opaque oxides	Layered gabbro having less than 48 wt. percent Si <sub>2</sub> O appears cumulate and was not included. Small pluton in the central Medfra quadrangle	Moll and others, 1981
Rhyolites of Nixon Fork-Iditarod area	65.8	Presumed calc-alkalic (does not cross)	High-K K <sub>2</sub> O/Na <sub>2</sub> O=1.2	Rhyolite	73 - 75	Biot+qtz are listed. Presumed to also have plag and andesine	These are small intrusive rhyolites bodies intruded along the Nixon Fork-Iditarod fault in the central Iditarod quadrangle. Most are peraluminous; some are garnet-bearing	Bundson, and Laird, 1982; M.L. Miller, oral comm., 1986
Kuskokwim Mountains belt Sleetmute-Nyac area								
Holokuk Basalt	74.5 - 64.3	Calc-alkalic (59)	Chiefly moderate-K, some high-K rhyolite	Basaltic andesite, andesite, rhyolite	52 - 60 (no chemical analyses on rhyolite)	Andesite: plag+hmbld(?) basaltic andesite: plag+cpx+opx+ol rhyolite: san+qtz	Holokuk Basalt is over 1000m thick, and actually consists of chiefly andesite lahars and flows, felsic breccia, tuff breccia, and agglomerate. Andesitic rocks have SIR 0.7040-0.7050	M.F. Robinson, oral comm.; 1985; Decker and others, 1984; Decker and others, 1986
Rhyolite and rhyolite porphyry	Chiefly 64 to 61.5 maybe as old as 70.5 at Barometer Mtn	Calc-alkalic (does not cross, plots along trend with Holokuk Basalt)		Rhyolite porphyry and rhyolite	62 and 68 - 76	Qtz+plag+san+biot:white mica:tourmaline	Contains tourmaline and quartz tourmaline breccia	Decker and others, 1986; M.F. Robinson, oral comm., 1985
Chailuk and Kialuk plutons	68.7-67.5	Calc-alkalic (does not cross)		Granodiorite (granite border-phase on Chailuk pluton)	64 - 73	Granodiorite: ande+K-spar+qtz+biot+hmbld:ms	Contains cross-cutting qtz-tourmaline veins	Decker and others, 1986
Intermediate stocks and dikes	69.8	Calc-alkalic		Diorite to granodiorite	58 - 63	Plag:qtz:K-spar: cpx:biot: hmbld	Dikes associated with lode occurrences of cinnabar and antimony	Decker and others, 1986; M.F. Robinson, oral comm., 1986

Late Cretaceous and early Tertiary  
Kuskokwim Mountains belt  
Titchik Lake plutons

Unit/Name	Age (Ma)	Classification	K <sub>2</sub> O type	Rock types	SiO <sub>2</sub> range (wt. %)	Mineralogy	Other comments	References
Kernak, Otstakuk Hills, and Ualik Lake plutons	Undated	Alkali-calcic (does not cross, projects to cross at 52)	Shoshonite	Granite, monzonite	57 - 68	Perthitic fsp: olig:qtz+pyr ±amph:biot	Data based on only three samples having higher K <sub>2</sub> O than other Titchik Lake region plutons.	Wilson, 1977 Hoare and Conrad, 1978b
Other plutons in Titchik Lake region	72.5-60.7	Calc-alkalic (does not cross but projects to cross at about 59)	High-K	Granite, qtz monzonite, grano-diorite, qtz monodiorite, monzonite, qtz diorite, tonalite, diorite	56 - 76	Granite: plag+qtz+K-spar+biot ±hbl:dt:cpx monzonite: plag+K-spar+biot+pyr+±qtz diorite: plag+opx+cpx+K-spar:qtz tonalite: plag+biot+qtz	Includes plutons at Nayoruran, Shotgun Hills, Titchik Lake, Akulutuk, Togiak Lake, Sunday Creek, Mt. Oratia, Mt. Waskey, Kashaik, Zomo Creek, Wattamuse, Salutak, Hobolima, Crooked Mtn, Kulik Lake, Hook, Muklung Hills, Kulukak, Mt. Chamchivak, Kasigluk River, Kasigluk Dome, Bek River Sill. Exact location of plutons given in Wilson, 1977.	Wilson, 1977; Hoare and Conrad, 1978b

Kuskowim Mountains Belt

Bristol Bay volcanic rocks

Bristol Bay volcanic rocks	68.7-64.5	Calc-alkaline	Moderate-K (K <sub>2</sub> O/Na <sub>2</sub> O=0.32)	Andesite, basalt, dacite	51.6 - 66.4	Andesites: plag+cpx+opx Basalt: plag+cpx+opx:ol	SIR = 0.7037- 0.7041	Hoare, unpub. data Globerman, 1985 Box, 1985
----------------------------	-----------	---------------	--	--------------------------	-------------	--	----------------------	--

Berling Sea shelf

St. Matthew Island	Volcanic rocks 77-66; plutonic sample 61	Calc-alkalic (60)	Low to moderate-K (K <sub>2</sub> O/Na <sub>2</sub> O=0.5)	Basalt, andesite, dacite, rhyolite, granodiorite	Volcanic rocks 52 - 78 plutonic rocks 63 - 65	Basalt: plag+aug+cpx+oxides andesite: plag+cpx+opx+oxides dacite: plag:qtz:cpx rhyolite: plag+qtz+biot granodiorite: plag+ortho+qtz+hbl:dt	Entire island (350 km <sup>2</sup> ) is Late Cretaceous to early Tertiary volcanic and plutonic rocks. Lower sequence is pyroclastic andesite, dacite, and rhyolite, overlain by andesite and basalt flows at least 250 m. thick. Youngest sequence is rhyolite and dacite welded tuff and hypabyssal rocks.	Patton and others, 1975; Patton and others, 1976; W.W. Patton, Jr., oral comm., 1986
St. Lawrence Island	64.4, 64.0, and 62.1	Probably calc-alkalic	Probably high-K	Basalt, soda rhyolite, trachyandesite, andesite	48.4 and 68.3-70.8	Soda rhyolite: san+hbl:dt no data on other rock types	Insufficient data to classify suite.	Patton and Coejoy, 1980

Yukon-Kanuti belt

Kanuti volcanic field	60 - 56	Calc-alkalic (does not cross; projects to cross at 60.5)	High-K; (K <sub>2</sub> O/Na <sub>2</sub> O=0.8)	Rhyodacite, dacite, hi-silica andesite	62 - 73	Rhyodacites: plag+biot+qtz; Dacites: plag+hbl:dt:qtz+mt Andesite: plag+hbl:dt:qtz+mt+opx+cpx	Rocks have high LREE (La=54-77) and low HREE (La=0.15-0.23) contents. Volcanic field located near the margin of the Yukon-Koyukuk depression	Patton and Miller, 1973 Moll-Stalcup and Arth, 1989; M.A. Lauphere, oral comm., 1986
Blackburn Hill volcanic field	65 - 56	Calc-alkalic 60.0	Moderate to high-K	Andesite, minor basalt rhyodacite, alkali rhyolite granodiorite	48 - 77	Andesite: plag+cpx:opx+mt or +plag+cpx+hbl:dt+mt basalt: ol+plag+cpx+mt rhyolite: olig:dt:biot:hbl:dt alkali rhyolite: sm:orth:ol:igo+hedenbergite or opx	Best exposed volcanic field in the Yukon-Kanuti or Kuskokwim Mountains belt. Alkali rhyolite occurs at top of section and is dated at 56 Ma. Volcanic suite has a silica gap in the dacite range. Samples from granodiorite pluton plot in the gap.	Moll-Stalcup and Arth, in review

Late Cretaceous and early Tertiary  
Yukon-Kanuti belt, cont'd

Unit/Name	Age (Ma)	Classification	K <sub>2</sub> O type	Rock types	SiO <sub>2</sub> range (wt. %)	Mineralogy	Other comments	References
Holy Cross quadrangle	Undated	Presumed calc-alkalic	No data	Basalt to rhyolite	No data	No published data. Known to include rhyolite: qtz+plag+san+biot	Widespread but little known volcanic and plutonic rocks occur in the region. Rhyolite ash-flow tuffs identified in thin sections from Hoare's 1962 collections	Beikman, 1974; J.M. Hoare, unpub. field data, 1960-1963
Yukon River volcanic rocks	53 - 48 and 44 - 43	Calc-alkalic (60.8)	Moderate to high-K	Basalt, andesite, dacite, rhyolite	48 - 76.5	Basalt: ol+cpx+plag+biot Andesite: plag+cpx+opx or plag+hmbld Dacite: qtz+plag+hmbld;cpx Rhyolite: qtz+plag+amorth;san Alkali rhy: smorth+clig+san+biot+hmbld	The mineralogy of some of the younger rocks suggests they are alkalic. Harris, (1985) reports ages of 43 to 44 Ma on basalts from this area	Patton and Mell, 1985; Moll-Stalcup and Arth, 1989; Harris, 1985
Beringian shelf edge	54.4-50.2	Calc-alkalic	Moderate-K (K <sub>2</sub> O/Na <sub>2</sub> O=0.4)	Andesite, basalt, dacite	50.9-65.1	Basalt: ol+plag+cpx; Andesite and dacite: plag+cpx+opx	Samples dredged from the Bering Sea shelf at about 58°30' lat., 176-177° long.	Devis and others, 1987a; Marlow and Cooper, 1986
Tokatjikh volcanic field	Unknown probably 59-56	Calc-alkalic (do not cross, project to cross at 62)	Transitional between moderate and high-K	Dacite	65.3-69.8	Dacite: plag+cpx+opx;qtz;biot or plag+hmbld+cpx+opaque oxides	Located on strike with the Kanuti volcanic field	Patton and others, 1978; W.W. Patton, Jr., and E.J. Moll-Stalcup, unpub. data, 1986
Roundabout Mtn. volcanic field	Unknown probably 59-38	Probably calc-alkalic	No data	Andesite	No data	Andesite: plag+cpx+mt	Occurrence of little known, poorly exposed, very fresh volcanic rocks that discontinue usually crop out over a 600 km <sup>2</sup> area along the Koyukuk River south of Hualia. Shown on map as unit KJv.	W.W. Patton, Jr., pers. comm., 1985; Patton, 1966

Yukon-Tanana Upland

Granitic plutons	69.5-56.7	Calc-alkalic, minor shoshonitic	Chiefly high-K some shoshonitic and Mod-K	Granite, granodiorite, qtz diorite, qtz monzonite, monzonite	65 - 77	Plag+K-spar+qtz+biot+hmbld+trms	Includes numerous plutons in the Circle and Livengood quadrangles and northern and northeastern Big Delta quadrangle.	Foster and others, 1983; Chapman and others, 1971; Weber and others, 1978
Mafic and intermediate rocks	67.4, 65.0 64.5 (on basalt)	Probably calc-alkalic	No data	Andesite Some gabbro	No data	No data	Intermediate lavas, breccia tuff, and plugs occur in the eastern Tanacross quadrangle; Gabbro occurs in the west-central Tanacross quadrangle.	Foster, 1970
Siliceous volcanic rocks	64.1-57.8	Probably calc-alkalic	No data	Felsic porphyries, welded tuff, and breccias	No data	No data	Includes felsic volcanic and hypabyssal rocks in the eastern Big Delta, west-central Eagle, and eastern Tanacross quadrangles. Lithologically similar rocks give ages of 90 Ma and the two events are difficult to distinguish.	Foster, 1976; H.L. Foster, oral comm., 1985; Foster, 1970; Weber and others, 1978
Mount Fairplay pluton	69 to 58	Alkalic	(K <sub>2</sub> O/Na <sub>2</sub> O=1.2)	Diorite, monzonite, some syenite	55.4	Diorite: cpx+hmbld+biot+plag; Monzonite: plag+K-spar+hmbld;biot;cpx; Syenite: plag+K-spar+hmbld;biot;cpx; Qtz Monz: plag+K-spar+biot+hmbld;px;opx	One chemical analyses only. At least some of the rocks are nepheline normative	Foster, 1967; Kerin, 1976

Early Tertiary  
Snak-Baranof belt

Unit/Name	Age (Ma)	Classification	K <sub>2</sub> O type	Rock types	SiO <sub>2</sub> O range (wt. %)	Mineralogy	Other comments	References
Boring Glacier, Valdes quadrangle	53-46 (biot) 64-48 (hornbl)	Calcic (64)	Med to high K (granite mostly high-K)  (K <sub>2</sub> O/Na <sub>2</sub> O varies, mostly 0.3-1.1 (higher values for granite))	Tonalite, granodiorite, granite	63-76	Tonalite: plag+qtz+biot+hornbl+K-spar+mu; mus+garn; ap+rt+tr+cl+ph; Granite and gabbro: plag+qtz+K-spar+biot+mu; garnet	Only one sample of granite (Boring Glacier quadrangle)	Hudson and others, 1977
Stagway quadrangle	41-26 (biot) 51.4-45.5 (hornbl) Biotite ages not considered reliable	Calcic (64)	Low to med-K (K <sub>2</sub> O/Na <sub>2</sub> O=0.2-0.8)	Tonalite, granodiorite, qtz diorite	61-79	Plag+qtz+biot+hornbl+K-spar+mu; garn+ap+tr+mt+rt+ph	Tonalite from one locality contain magmatic(?) red epidote (thulite) and one sample also contains garnet. These rocks have 79% SiO <sub>2</sub> , highly enriched LREE (La=400-500x chond) and variable HREE (La=4-40x chond). One granodiorite sample contains garnet	T. Hudson and G. Pfaffner, unpub. data, 1987; J. Lull and G. Pfaffner, oral comm., 1978
Cardova quadrangle	53-51 (biot) 54-50 (hornbl)	Calcic (64)	Low to high-K (K <sub>2</sub> O/Na <sub>2</sub> O=0.1-1.4; higher values for granite)	Chiefly granodiorite; minor granite, gabbro, orthopyroxenite, and peridotite.	47-79	Granitic: plag+qtz+K-spar+biot+hornbl+mu; garn+ap+tr+mt; Gabbro: plag+hornbl+ep+rt+opx+qtz+biot+ap+mt; Orthopyx: Opx+amph+scap; Peridotite: ol+opx+opx+plag+hornbl	REE trends for granodiorite are very consistent: La 70-120x; La 12-20x chondrites; moderate Eu anomaly. Orthopyroxenite occurs only as stopped block(?) or xenolith(?) in one granodiorite sample from the Rude River pluton. Peridotite occurs at one locality	J.S. Lull and George Pfaffner, oral comm., 1988 Winkler and Pfaffner, 1981
Yakutat and Mt. St. Elias quadrangle	48-43 (biot) 54-51 (hornbl) 48-47 (mu)	Calcic (64)	Low to high-K (K <sub>2</sub> O/Na <sub>2</sub> O=0.1-2.0)	Granite, granodiorite, diorite	52-76	Granitic: plag+qtz+K-spar+biot+hornbl+mu; garn+ap+tr+mt; Diorite: plag+opx+hornbl+biot+qtz+ap+mt	One pluton gives a U-Pb zircon age of 36 Ma and biotite K-Ar ages of 21-23 Ma	Hudson and others, 1977; M.A. Lamphere, unpub. K-Ar age, 1979
Snak pluton	60	Calc-alkalic (60)	High-K (K <sub>2</sub> O/Na <sub>2</sub> O=1.0-1.4)	Granodiorite, granite	66-73	Qtz+plag+K-spar+biot+mu (up to 2%) +Fe-Ti oxides		Moore, 1974a; Hill, 1979
Kodiak batholith	60-57	Calcic (61.5)	Med-K (K <sub>2</sub> O/Na <sub>2</sub> O=0.9-1.0)	Tonalite, granodiorite	63-70	Qtz+plag+K-spar+biot+Fe-Ti oxides	Some garnet-bearing two-mica granite	Hill, 1979; Shew and Wilson, 1981
Shumagin batholith	64-56	Calcic (61.5)	High-K (K <sub>2</sub> O/Na <sub>2</sub> O=0.7-1.1)	Granodiorite, granite	63-76	Qtz+plag+K-spar+mu+Fe-Ti oxide		Moore, 1974b; Hill, 1979; Burk, 1965

Middle Tertiary  
Interior west-central Alaska

Takhakhdona and Toksojikh volcanic fields	Unknown; probably 59-39	Calc-alkalic (61)	Transitional between moderate and high-K	Trachybasalt, andesite, rhyolite, latite	52.7 and 64.5 - 77	Trachybasalt: no data; andesite: plag+opx+tr+hornbl+oxides; rhyolite: plag+K-spar+qtz+biot+hornbl; latite: amorph+plag+biot+opaques	One analysis having SiO <sub>2</sub> < 64 is a trachybasalt having aluminous affinities. Sparse data from these poorly studied fields were combined, but the fields may be of different ages	Patton and others, 1978; W.W. Patton, Jr. and E.J. Moll-Stalcup, unpub. data, 1986
Roundabout Mtn. volcanic field	Unknown; probably 59-38	Probably calc-alkalic	No data	Andesite	No data	Andesite: plag+opx+mt	Little known, poorly exposed very fresh volcanic rocks that discontinuously crop out over a 600 km <sup>2</sup> area along the Koyukuk River south of Huslia.	W.W. Patton, Jr., pers. comm., 1985; Patton, 1966
Indian Mtn. volcanic field	41.6 and 39.9	Calc-alkalic	High-K (K <sub>2</sub> O/Na <sub>2</sub> O=1)	Rhyolite and dacite	72.5 - 77.5	Glass, no micro-lites	Consists of tuff, breccia and flows. Oldest known obsidian locality in the U.S. Covers an area of about 40 km <sup>2</sup>	Miler and Lamphere, 1981
Dulbi River volcanic field	43.2	Calc-alkalic	High-K (K <sub>2</sub> O/Na <sub>2</sub> O=1)	Rhyolite	76.8	Qtz+plag+K-spar in perlitic	Rhyolite in this area is highly altered	W.W. Patton, Jr. and Moll-Stalcup, unpub. data, 1985

Middle Tertiary,  
Interior west-central Alaska

Unit/Name	Age (Ma)	Classification	K <sub>2</sub> O type	Rock types	SiO <sub>2</sub> range (wt. %)	Mineralogy	Other comments	References
Yukon River volcanic rocks	42.7	No data	No data	Basalt	No data	No data	No lithologic data were reported but we know basalt occurs because basalt whole-rock samples were dated. Other lithologies may be more common.	Harris, 1985
Western Alaska wells	42.3-39.7	Tholeiitic(?)		Basalt	52.4-53.9	Ba: plag+cpx+ce, chal+clay	Middle Tertiary basalts have been found in the Norton Cost well #1, the Cape Espenberg well #1, Social Nimiak well and the Bethel Basin (Napatak #1) well. Chemical data is on basalt from the Norton Cost well only. The rocks there are highly altered and have at least 10% CaO	Turner and others, 1983; Tolson, 1986; and unpublished oil company data
Basalt in Chukmuk River area	38.2	No data	No data	No data	No data	Ol+cpx+plag+opx?	Small isolated occurrence of fresh olivine basalt located at lat. 61°07', long. 157°57'	Decker and others, 1986; M.F. Robinson, oral comm., 1986
Rhyolites at Tang Mtn. (equivalent to the black and gray rhyolite unit of Tang Mtn. of Decker) and others, 1986)	43.8	Rhyolites are probably calc-alkalic; tachyandesite is alkalic	Probably high-K rhyolites: K <sub>2</sub> O/Na <sub>2</sub> O=1.57 trachyte: K <sub>2</sub> O/Na <sub>2</sub> O=0.8	Rhyolite tuff and breccia, tachyandesite	75-76 and 55	Rhyolite: san+qz in glass tachyandesite: plag+san+cpx	Major and trace element data suggest the tachyandesite is not orogenic. It has high TiO <sub>2</sub> (1.97%), high K <sub>2</sub> O and La and lacks a Nb-Ta anomaly. SIR on the tachyandesite is 0.7033 assuming an age of 43 Ma. Tang Mtn. is an informal name given to orange-colored hills located at lat. 61°27', long. 157°50'	Decker and others, 1986; M.F. Robinson, oral comm., 1986
Middle Tertiary								
Bering Sea region								
St Lawrence Island volcanic rocks	39.3	Probably calc-alkalic	Probably high-K (K <sub>2</sub> O/Na <sub>2</sub> O=1)	Rhyolite and dacite	67 and 76.5	No data	Consists of tuff, breccia and flows. Only two chemical analyses are available for these rocks which makes them difficult to classify	Patten and Coejuay, 1980
Middle Tertiary Southern Alaska								
Volcanic rocks of Matanuska Valley	45.5-37	Tholeiitic and calc-alkalic	No data	Basalt, dacite, rhyolite	45 - 53 and 69-78	No data	Matanuska Valley rocks are hypabyssal intrusive rocks having SIR=.7036-.7042	Silberman and Grantz, 1984
Arkos Ridge Formation	56-43	Tholeiitic	Low to moderate-K	Basalt, andesite	48 - 62	Plag+ol+pyr+oxides	Rocks are highly altered. K-Ar ages considered minimum ages. Grantz and Silberman consider its age as 50-60 Ma. Grantz reports that these basalts are similar in appearance to the basalts exposed on the western flank of the Talkeetna Mountains.	Silberman and Grantz, 1984; A. Grantz, oral and written comm., 1986; M.L. Silberman, written comm., 1986
Plutons of Prince William Sound region	36.6-34.2	Calcic (62.5)	Moderate-K	Chiefly granite; subordinate granodiorite, tonalite, diorite, gabbro	52 - 54 and 67 - 77	Granite and granodiorite: oligo to and+biot; hornbl (rare biot: mus) diorite: and+hornbl:biot gabbro: cpx+opx+labradiorite:qtz	Includes plutons on Esther Perry and Culross Islands and and at Billings Glacier, Eshamy Lagoon, Granite Mine, Nellie Juan, Passage Canal, Minors Lake and Yale Arm. Billings Glacier is at lat. 60°50', long. 148°35' and Yale Arm is at 61°12' 147°46'.	Tydal and Case, 1979; Nelson, and others, 1985; S.W. Nelson, written comm., 1986
Sills and dikes Prince William Sound region	54.8-43.6	No data	No data	Hypabyssal dacite and rhyolite	No data	Plag+hornbl (frequently altered to sericite calcite and chlorite)	These hypabyssal intrusive bodies cut the Valdez Group	Nelson, and others, 1985

## Middle Tertiary

## Southern Alaska, cont'd

Unit/Name	Age (Ma)	Classification	K <sub>2</sub> O type	Rock types	SiO <sub>2</sub> range (wt. %)	Mineralogy	Other Comments	References
Plutons of Yakutat-Mt. St. Elias area	30.6-18.5	Not a coherent suite	Low to high-K (K <sub>2</sub> O/Na <sub>2</sub> O)= 0.1-0.5	Granite, granodiorite, tonalite, diorite	52-73	Granitic: plag+qtz+K-spar: hornbl: biot+Fe-Ti oxides: cpx: epidote: apatite	Includes samples from Mt. Owen pluton, Brabason pluton, Mt. Forseta pluton, and Valerie Glacier pluton. Tonalite at Mt. Forseta is foliated. Two tonalites contain muscovite	Hudson and others, 1977
Dikes and plugs in the Yakutat-Mt. St. Elias area	Probably Tertiary	Not a coherent suite	Low to high-K (K <sub>2</sub> O/Na <sub>2</sub> O)= 0.1-0.51	Dacite porphyry, basalt, diabase	47-68	Dacite: plag+hornbl+Fe-Ti oxides; BS: plag+oxides	Basalt has amygdale filled with calcite, quartz, and chlorite Diabase has secondary chlorite epidote, and calcite	Hudson and others, 1977
Volcanic rocks in Yakutat-Mt. St. Elias area			High-K (K <sub>2</sub> O/Na <sub>2</sub> O= 0.1-0.5)	Rhyolite porphyry	76-77	Plag+qtz+san: biot: trma: sphene: apatite	Related to Mt. Owen's pluton. data base on only two samples	Hudson and others, 1977

## Late Tertiary and Quaternary

## Bering Sea basalt province, Seward Peninsula

Imuruk Lake area: Five volcanic formations:							Hopkins, 1963, reports inclusions of granite rock, dunite, chromite and carbonate and xenocrysts of orthoclase and oligoclase. Swanson and Turner report inclusions of hornblite and hornburgite and xenocrysts of pyroxene and olivine.	Hopkins, 1963; S.E. Swanson and D.L. Turner, unpub. report, 1984; Swanson, and others, 1981
Kagrak Volcanics	29-26	Alkalic and tholeiitic		Basanite, ol tholeiite, AOB	44 - 48	Ol+cpx+plag+mt		
Imuruk Volcanics	5.8-2.2	Tholeiitic		Tholeiite, ol tholeiite	49 - 52	Plag+ol+cpx+opx+opaque oxides	No inclusions reported	
Gooling Volcanics	0.9-0.8	Alkalic and tholeiitic		Basanite, ol tholeiite	44 - 51	Ol+cpx+plag+opaque oxides	Inclusion bearing	
Camille Basalt	middle and/or late Pleistocene	No data		No data	No data	Presumed tholeiitic	No inclusions reported.	
Lost Jim Basalt	Holocene	No data		No data		Plag+ol+cpx+opaque oxides	No inclusions reported. Presumed tholeiitic	
Unnamed volcanic field north of Teller	2.9 - 2.5	Alkalic		Basanite, tephrite, AOB	44.8 - 47.7	No data	Includes Makacherni Mtn., Eva Mtn., and "hill 1220" Locations shown on 1:63,300-scale Teller B-3 quadrangle map	S.E. Swanson and D.L. Turner, unpub. report, 1984

## Late Tertiary and Quaternary

## Bering Sea basalt province western Alaska

St Michael volcanic field	3.25 - Holocene	Chiefly alkalic, some tholeiitic		Ol tholeiite, AOB, basanite hawaiiite	44 - 51	Basalt: ol+cpx+plag+mt; Basanite: ol+cpx+opaque oxides; plagiocl: analcime; nepheline	Highly alkalic rocks have ilmenite nodules. SIR= 0.7027	J.M. Hoare, unpub. data, 1968; Patton and Moll, 1965; Mark, 1971
Yukon Delta area: Ingalukwat volcanic field	Presumed to be Pliocene and younger	Alkalic		AOB, basanite, nepheline	41 - 51	Ol+cpx+plag+mt;	Field covers 200 square miles. Some of the undersaturated rocks have inclusions of olivine gabbro and ilmenite. SIR = 0.7028	J.M. Hoare, 22 unpub. chem. analyses, 1968; Hoare and Condon, 1971a
Nunivak Island	Five episodes: 6-4, 3.4-3.1, 1.7-1.5, 0.9-0.3 and Holocene	Alkalic and tholeiitic		Ol tholeiite, basanite, AOB, nepheline	44 - 51	Basaltic: ol+cpx+plag+mt Basanite: ol+cpx+plag+neph: analcime +nepheline; augite and kaersutite	Undersaturated alkalic rock contain inclusions of layered gabbro, ilmenite, spinel-pyroxene, sandstone and conglomerate and xenocrysts of anorthoclase, augite and kaersutite SIR=0.7026-0.7033, (Mark); 0.70251-0.70330, (Mezianis); 0.70279-0.70327, (Roden)	Mark, 1971; Mezianis and Murthy, 1980a; and 1980b; Roden and others, 1984; Hoare and others, 1968; J.M. Hoare, unpub. data, 1971-1981; Roden, 1982

Bering Sea basalt province  
Western Alaska, cont'd.

Unit/Name	Age (Ma)	Classification	K <sub>2</sub> O type	Rock types	SiO <sub>2</sub> range (wt. %)	Mineralogy	Other Comment	References
Devil Mountain and volcanic fields to the east	Undated	Presumed alkalic and tholeiitic		No data	No data	No data	Undated but presumed to be similar to the other Bering Sea basalt fields	S.E. Swanson and D.L. Turner unpub. report 1984; Lundke and Smith, 1986
Basalt of Flat Top Mtn.	4.72 and 4.62	alkalic		Basalt	49.4	Ol+cpx+plag	Located in the valley of the Kwetluk, Kisaralik, and Aniak Rivers. One chemical analysis	M.F. Robinson, oral comm., 1986; Decker and others, 1986; Box and Moll-Stalcup, unpublished data, 1989
Togjak Basalt	0.758 and younger	Tholeiitic and alkalic		Tholeiite and AOB	48.6 - 51.7	Plag+cpx+ol+mt	The youngest volcanic rocks in this field form a tuya from a subglacial eruption	Hoare and Coonrad, 1978a; Hoare and Coonrad, 1980

Bering Sea basalt province  
Bering Sea shelf

Pribilof Islands: St. George Island	2.2 - 1.6	Tholeiitic		Ol tholeiite	48.5 - 49.1	Plag+cpx+ol+cl+mt	SIR St. George = 0.7027; St. Paul = 0.7031. Nepheline found in two samples.	Lao-Wong and others, 1979; Simpson and others, 1979;
Dredged rocks from submarine ridge	0.836 - 0.774	Tholeiitic and alkalic		Ol tholeiite, AOB, hawaiite	49 - 50.7		Submarine ridge located between St. George and St. Paul. Inclusions of dunite or peridotite and of aplite or fine-grained granite are reported in the highly alkalic rocks	Mark, 1971; Cox and others, 1966
St. Paul Island	0.374 to Holocene	Alkalic		Basaltic, tepehite, AOB, hawaiite	44.7 - 46.8			
Dredged rocks from Bering Shelf edge	0.35-1.1	Alkalic		Basaltic	43.1-47.3	Ol+cpx phenoc+plag microlites	Samples dredged from Bering Sea shelf near 58°30' lat., 177°20' long.	Davis and others, 1987b
Kookooligit volcanic field	1.46-238	Chiefly alkalic, some tholeiitic		Basaltic, ol tholeiite, AOB, hawaiite, nephelinitic	44 - 55	Basalt: ol+cpx+plag+mt; Basaltic: ol+cpx+plag+nephelinitic	Field occurs on St. Lawrence Island. SIR = 0.7036-0.7039	J.M. Hoare, unpub. data, 1971; Mark, 1971; Patton and Coejoy, 1980

Eastern Alaska

Prindle Volcano	Post-early Pleistocene	Alkalic		Basaltic	42.8	Cpx+ol+opaque oxides (nepheline + K-spar in the groundmass)	Prindle Volcano has peridotite and granulite inclusions. Prindle Volcano and the older alkali-olivine basalts are part of a south trending belt of basalt and ultrabasic inclusion-bearing lavas extending from eastern Alaska down through British Columbia	Foster, 1981; Foster and others, 1966; H.L. Foster, oral comm., 1984; Foster, 1970
Other volcanoes in the vicinity of Prindle Volcano	Undated presumed older than Prindle Volcano	Alkalic		AOB	Unknown	Basalt: ol+plag+cpx+opaque oxides		
Basalt of Porcupine-Blue River area	16.5	Alkalic and Tholeiitic		Ol tholeiite, AOB, basaltic	43-46	Plag+cpx+ol	SIR=0.7041-0.7042; NIR=0.51287-0.51291	Flumley and Vance, 1988; Wirth and Bird, 1988; Brabb and Hamachi, 1977; Brabb, 1970; Brabb and Clarkin, 1969; Brogé and Reiser, 1969; Wirth, 1988; Wirth, K.R., pers. comm., 1988

Late Tertiary and Quaternary  
Eastern Alaska, cont'd

Unit/Name	Age (Ma)	Classification	K <sub>2</sub> O type	Rock types	SiO <sub>2</sub> range (wt. %)	Mineralogy	Other Comment	References
Wrangell volcanic field	26-0	Calc-alkalic. Locally transitional to tholeiitic	Medium-K	Chiefly andesite but range from basalt to rhyolite	74.0-49.5	Predominantly mineralogy Andesite: plag+cpx+hyp; plag+hyp+ol; plag+hyp; plag+ol; Dacite: plag+hmbld; plag+hmbld+biot; plag+hmbld+hyp; Basalts: plag+ol; plag+ol+hyp; Rhyolite: plag+biot plag+biot+hmbld	Large, complex field covers more than 10,000 km <sup>2</sup> in north-central Alaska and consists chiefly of a number of large coalescent shield volcanoes. Available K-Ar ages suggest, with exceptions that field youngs to west. Mt. Wrangell in west end of field is geothermally active; K-A ages of 20-26 Ma have been obtained from volcanic centers at east end of field along the Yukon Territory border	Richter and others, 1979; Lowe and others, 1982; Nye, 1983; Richter and others, 1984; Miller and Richter, this vol.
<b>Central Alaska</b>								
Ray Mtns. basalt flows	32.3 (Patton believes it is younger than the K-Ar age)	Tholeiitic		Tholeiite	50-52	Plag+cpx, minor ol	Little known volcanic field covering about 50 km <sup>2</sup> between the Yukon River and the Kamuti volcanic field. Shown on 1:250,000-scale reconnaissance map of the Bettles quadrangle. SIR=7046; NIR=0.51282	Patton and Miller, 1973; Wirth, K.R., 1988; Wirth, K.R., pers. comm., 1988; Patton, W.W., Jr., pers. comm., 1984 (K-Ar age)
Buzzard Creek	0.003	Tholeiitic transitional		Ol basalt	47.4-49.9	Ol+plag+cpx+opx	Ol basalt on the Maar volcano north side of the Alaska Range near Buzzard Creek. Chemically similar to Bering Sea basalts	Fe'we' and others, 1966; Albanese, 1980
Jumbo Dome	2.72 and younger	Calc-alkalic	Moderate to high-K	Hmbld andesite	56.2-59.0	Hmbld+plag+cpx opaque oxides	This dome is related to subduction of the Pacific plate.	Wahrhaftig, 1970; Albanese, 1980

Abbreviations are as follows: amph=amphibole, andes=andesite, andst=andesite, north=northoclase, AOB=alkali-olivine basalt, aug=augite, biot=biotite, bs=basalt, chl=chlorite, dior=diorite, fsp=feldspar, gar=garnet, gb=gabbro, gr=green, hmbld=hmbld, hyp=hypersthene, K-spar=K-feldspar, labrad=labradorite, monz=monzonite, monzgb=monzogabbro, monzdr=monzodiorite, mt=magnetite, mus=muiscovite, neph=nepheline, ol=olivine, olig=oligoclase, opx=orthopyroxene, orth=orthoclase, oxides=Fe-Ti oxides, plag=plagioclase, pyr=pyroxene, qtz=quartz, san=sanidine, sph=sphene, zr=zircon, SIR=initial <sup>87</sup>Si/<sup>86</sup>Si, NIR=initial <sup>143</sup>Nd/<sup>144</sup>Nd.

Table 2. Incompatible elements in andesites from the Kuskokwim Mountains belt

	Von Frank	Nowitna	Page Mtn.	Sleetemute	Bristol Bay	St. Matthew Island
	(shoshonitic)	(moderate to high-K)	(altered probably high-K to shoshonit)	(moderate-K)	(moderate-K)	(low to moderate K)
	n=5	n=7	n=7	n=12	n=6	n=6
SiO <sub>2</sub>	53.8 - 59.1	55.0-62.0	57.8-61.9	54.0-59.5	53.3-56.9	54.8-64.0
Nb	8-16	9-18	9-14	7.7-10.7	4-6	5-11
Zr	92-239	174-253	145-183	109-168	97-127	117-158
Y	17-25	22-33	14-25	17-26	21-32	23-26
Sr	911-1420	475-525	466-670	307-547	511-760	350-544
Rb	59-155	48-90	79-132	38-64	4-32	14-25
Ba	1792-2210	846-1350	1097-1870	349-910	376-786	318-470
La	49-57	30-44	25-37	12.8-28.8	12-21	1-16
Th	16.6-30.5	8.2-13.1	9.74-12.2	3.0-6.5	n.d.	1.0-2.0
Ta	0.902-1.18	0.79-1.13	0.664-0.85	0.43-1.06	n.d.	0.38-0.71
Hf	5.0-6.6	4.1-5.8	3.4-4.3	2.42-4.24	n.d.	2.5-4.1
<sup>87</sup> Sr/ <sup>86</sup> Sr	0.7047- 0.7051	0.70434- 0.70508	0.7049- 0.7059	0.70403- 0.70601	0.70370- 0.70414	No data

Note: Von Frank, Nowitna, and Page Mountain are in the northern Kuskokwim Mountains, Sleetemute is in the central Kuskokwim Mountains, and Bristol Bay is at the southern end of the Kuskokwim Mountains. Incompatible elements increase from north to south, left to right. SiO<sub>2</sub> is in weight percent, all other elements are in parts per million.

Table 3. Trace element abundances (ppm) and ratios in representative andesite samples from the San Juan volcanic field (Summer Coon), Colorado, and the northern Kuskokwim Mountains belt (Nowitna) and the Yukon-Kanuti belt (Blackburn Hills), Alaska

---

	Summer Coon <sup>1</sup>	Nowitna <sup>2</sup>	Blackburn Hills <sup>2</sup>
SiO <sub>2</sub>	56	57	56.5
Rb	65	52	35
Ba	1,160	870	891
Sr	930	505	493
K/Rb	344	298	299
La	30	32	30
Ce	79	66	52
Yb	1.5	2.4	2.2
La/Yb	20	13.3	13.6
Th	3.6	8.3	6.1
U	1.1	3.1	2.2
Th/U	3.2	2.7	2.8

---

1) Zielinski and Lipman (1976)

2) Moll-Stalcup (1987)

TABLE 4. Comparison of selected trace-element ratios for a Bering Sea tholeiite with a Hawaiian tholeiite and N-MORB.

	Bering Sea basalts St. Michael volcanic field (80ML57B)	Hawaii (Clague and Frey, 1982)	Average N-MORB (Wood, 1979)
P <sub>2</sub> O <sub>5</sub> /Ce	87.0	81.3	0.02
Rb/Sr	0.044	0.031	0.008
K/Rb	409.	432.	1,060.
Zr/Hf	46.4	45 ± 4	33.5
Hf/Ta	2.44	1.35	15.45
Th/La	0.125	0.091	0.065
Th/Ce	0.065	0.047	0.021
Th/Sm	0.47	0.39	0.063
Th/Nd	0.088	0.094	0.023
Sr/Th	214.	246.	660.
Ba/Th	88.7	165.0	60.0
Sr/Ba	2.4	1.52	11.0
Ba/La	11.1	14.9	3.9
Ba/Ce	5.8	7.6	1.3
Zr/Ta	113.0	60.0	518.0
Sr/Ce	14.0	11.3	13.9

## FIGURE CAPTIONS

- Figure 1:** Location of Late Cretaceous and early Tertiary volcanic and plutonic belts of mainland Alaska. Proposed Paleocene continental margin shown in red. A few locations are: 1, Blackburn Hills volcanic field; 2, Yukon River area; 3, Kanuti volcanic field; 4, Sischu volcanic field; 5, Nowitna volcanic field; 6, Sleetmute area; 7, Bristol Bay; and 8, Talkeetna Mountains. Dashed line marking 180-m water depth delineates edge of the Bering Sea shelf.
- Figure 2:** Chondrite-normalized spidergrams for rocks from the McKinley sequence. Data from Lanphere and Reed (1985); normalization factors from Thompson and others (1984). The rocks have deep Nb-Ta troughs characteristic of subduction-related magmas. Sharp spikes at Sr, P and Ti suggest the rocks are highly differentiated and have fractionated plagioclase, Fe-Ti oxides and apatite.
- Figure 3:** Chondrite-normalized spidergrams for andesites from the Kuskokwim Mountains belt and St. Matthew Island. Data for northern Kuskokwim Mountains (Nowitna) and St. Matthew Island from E.J. Moll-Stalcup and W.W. Patton, Jr., (unpub. data, 1985). Data for Sleetmute from Decker and others (1986); for Bristol Bay from Globberman (1985).
- Figure 4:** Eight chondrite-normalized rare-earth-element diagrams for rocks from five areas. A, andesite and rhyolites from the Sischu volcanic field having low HREE that probably indicates hornblende or garnet fractionation. B, patterns for highly fractionated (high silica?) rhyolites from the Sischu volcanic field. C, rhyolite and granites from the Sleetmute area. D, REE data for andesites from the northern Kuskokwim Mountains: Von Frank = high K to shoshonitic; Nowitna = moderate to high K; Stone Mountain = moderate K; Cloudy Mountain = highly altered, but are thought to be shoshonitic on the basis of  $K_2O$  content. However, REE data suggest it may actually be moderate K. E, REE data for Sleetmute andesites. Chondrite normalized La values for andesites of the Bristol Bay volcanic unit shown by bar at La. F, REE data for andesites from St. Matthew Island. These rocks have very low K, Rb, Th, and LREE and may be correlative with the Alaska Range-Talkeetna Mountains belt for which we have no REE data. G and H, REE data for the Blackburn Hills volcanic field, showing the two andesite types that are distinguished on the basis of HREE: G, group 1 pyroxene andesites; H, group 2 pyroxene and hornblende andesites. Data for Sleetmute from Decker and others (1986); for Bristol Bay from Globberman (1985). All other data from E.J. Moll-Stalcup and W.W. Patton, Jr., (unpub. data, 1979-1986).
- Figure 5:**  $K_2O$  versus  $SiO_2$  for Late Cretaceous and early Tertiary volcanic and plutonic suites. Alaska Range-Talkeetna Mountains belt (Summit Lake plutonic rocks; Reed and Lanphere, 1974); St. Matthew Island; Bristol Bay volcanic unit,

southern Kuskokwim Mountains; Nowitna volcanic field and Von Frank Mountain pluton (shoshonitic suite), northern Kuskokwim Mountains; Blackburn Hills and Kanuti volcanic field, Yukon-Kanuti belt.

- Figure 6: Chondrite-normalized spidergrams. A, samples from the Blackburn Hills in the Yukon-Kanuti belt. Samples 52 and 30d are from the older andesite section; samples 9f and 65 are basalts that overlie rhyolite domes or flows and are younger than 56 Ma. The younger rocks are more mafic, have lower alkalis relative to La and smaller Nb-Ta depletions. B, samples from Yukon River Eocene volcanic rocks. All data from E.J. Moll-Stalcup and W.W. Patton, Jr., (unpub. data, 1981). All the rocks except 332, a basalt dated at 53 Ma., have large Nb-Ta anomalies. The more siliceous rocks have negative spikes at Sr, P, and Ti due to fractionation. C, samples from the Kanuti volcanic field. All data from E.J. Moll-Stalcup and W.W. Patton, Jr. (unpub. data, 1980-1981).
- Figure 7: Chondrite-normalized spidergrams for the Arkose Ridge Formation and Talkeetna Mountains volcanic rocks. Data from A. Grantz and M.L. Silberman (unpub. data, 1981). The variation in the shape of the pattern for the Arkose Ridge samples for Ba, Rb, Th and K suggests that these rocks are altered.
- Figure 8: Distribution of middle Tertiary igneous rocks in Alaska. Dashed line marks water depth of 180 m. Middle Tertiary Alaska-Aleutian range arc shown in red. Middle Tertiary igneous localities shown in blue. Locations are: 1, Indian Mountain; 2, Dulbi River; 3, Takhakhadona Hills; 4, Kateel River; 5, Yukon River; 6, Sleetmute area; 7, Matanuska Valley and Arkose Ridge; 8, Prince William Sound; and 9, St. Lawrence Island.
- Figure 9: Chondrite-normalized spidergrams of a sample of trachyandesite from Tang Mountain in the Sleetmute area. Rock has higher Ta relative to La and only slightly less Nb relative to La, and is not considered an orogenic andesite. Data from Decker and others (1986).
- Figure 10: Location of late Cenozoic (0-6 Ma) volcanic fields in Alaska. 1, Imuruk Lake area; 2, Devil Mountain; 3, St. Michael volcanic field; 4, St. Lawrence Island; 5, Ingakslugwat volcanic field; 6, Nunivak Island; 7, Flat Top Mountain; 8, Togiak Basalt; 9, Pribilof Islands; 10, Prindle Volcano; 11, Porcupine-Black Rivers; 12, Ray Mountains; 13, Jumbo Dome (west) and Buzzard Creek (east). Blue, chiefly basalt and basanite; red, chiefly arc volcanic rocks. Dashed line marking 180-m water depth delineates the edge of the Bering Sea shelf.
- Figure 11: Plot of total alkalis versus SiO<sub>2</sub> for Bering Sea basalts. Lines divide silica-saturated and silica-undersaturated rocks on Hawaii and in Bering Sea basalts. Data from J.M. Hoare (unpub. data); S. Swanson and D. Turner (unpub. data, 1985) and Lee-Wong and others (19??).
- Figure 12: <sup>87</sup>Sr/<sup>86</sup>Sr and <sup>143</sup>Nd/<sup>144</sup>Nd data for Bering Sea basalts. Data from Menzies and Murthy (1980), Von Drach and others (1986) and Roden (1982). Rocks from St. George in the Pribilof Islands and Nunivak Island plot in the field

where values for MORB (mid-ocean-ridge basalt) and oceanic island basalt overlap. Analyses of samples from St. Lawrence Island plot closer to bulk-earth compositions. Bulk-earth values from Allegre, Hart, and Minster (1983).

Figure 13: Chondrite-normalized spidergrams for volcanic rocks from the St. Michael volcanic field (57B) and the Pribilof Islands (26B, G142A, 27B, HP100, and G115B). Data from E.J. Moll-Stalcup and W.W. Patton, Jr., (unpub. data, 1980-1981) and F. Lee-Wong (unpub. data, 1983). The rocks having the highest alkalis and LREE are nephelinites; those having the lowest are tholeiites. Note the positive Nb-Ta anomaly.




Figure 1




Figure 2




Figure 3




Figure 4





Figure 4 cont'd.



**EXPLANATION**

- ▣ Von Frank Mountain,  
Whitwind Ridge plutonic rocks
- Nowlina volcanic field
- △ Blackburn Hills volcanic field
- ▴ Kanuit volcanic field
- ▾ St. Matthew Island
- ◇ Bristol Bay volcanic unit
- Alaska Range—Talketna Mountains  
belt (Summit Lake plutonic rocks)

Figure 5






Figure 6




Figure 7




Figure 8




Figure 9




Figure 10





Figure 11



**EXPLANATION**

- Nunivak Island**  
 ○ Basanite  
 ◇ Olivine tholeiite  
 ● Inclusions and megacrysts
- Pribilof Islands (St. George)**  
 △ Alkali olivine basalt
- St. Lawrence Island**  
 ● Basanite  
 ◆ Olivine tholeiite

Figure 12




Figure 13