

**RECONNAISSANCE DATA FOR SELECTED HERBICIDES,  
TWO ATRAZINE METABOLITES, AND NITRATE IN SURFACE  
WATER OF THE MIDWESTERN UNITED STATES, 1989-90**

**By Elisabeth A. Scribner, E. Michael Thurman, Donald A. Goolsby,  
Michael T. Meyer, Margaret S. Mills, and Michael L. Pomes**

---

**U.S. GEOLOGICAL SURVEY**

**Open-File Report 93-457**



**Lawrence, Kansas  
1993**

**U.S. DEPARTMENT OF THE INTERIOR**

**BRUCE BABBITT, Secretary**

**U.S. GEOLOGICAL SURVEY**

**Robert M. Hirsch, Acting Director**

---

**For additional information write to:**

District Chief  
U.S. Geological Survey  
4821 Quail Crest Place  
Lawrence, Kansas 66049-3839

**Copies of this report can be purchased from:**

U.S. Geological Survey  
Books and Open-File Reports Section  
Federal Center, Box 25425  
Denver, Colorado 80225

## CONTENTS

	<b>Page</b>
Abstract .....	1
Introduction.....	1
Methods .....	3
Sampling design .....	3
Data-collection techniques.....	3
Analytical procedures .....	3
Quality assurance .....	10
Discussion.....	11
References cited.....	17

## ILLUSTRATIONS

Figures 1-5. -- Map showing:

1. Location of study area, hydrologic units, and sites sampled during 1989 and 1990 in the midwestern United States.....	2
2. Location of sites sampled in South Dakota, Nebraska, and Kansas, map numbers 1 through 30 .....	8
3. Location of sites sampled in Minnesota, Iowa, and Missouri, map numbers 31 through 78.....	9
4. Location of sites sampled in Wisconsin and Illinois, map numbers 79 through 115.....	10
5. Location of sites sampled in Indiana and Ohio, map numbers 116 through 147.....	11

Figures 6-9. -- Graph showing:

6. Concentrations of alachlor during 1989-90 reconnaissance study .....	18
7. Concentrations of atrazine during 1989-90 reconnaissance study .....	19
8. Concentrations of cyanazine during 1989-90 reconnaissance study .....	20
9. Concentrations of metolachlor during 1989-90 reconnaissance study .....	21

## TABLES

Table 1. Quantities of four major herbicides applied in 10 agricultural Midwestern States 1987-89 .....	2
Table 2. Sampling-site names, locations, and drainage areas .....	4
Table 3. Concentrations of selected herbicides and two atrazine metabolites in quality-assurance samples, 1989-90.....	12

## **TABLES--Continued**

	<b>Page</b>
Table 4. Herbicide compounds analyzed and percentage of detections in pre-application, post-application, and fall low-flow samples during 1989-90.....	16
5. Comparison of herbicide samples collected during 1989-90 with Maximum Contaminant Levels and Health Advisory Levels for drinking water in effect at the time of this study .....	17
6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90.....	23
7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90.....	51

## **CONVERSION FACTORS AND ABBREVIATED WATER-QUALITY UNITS**

<i>Multiply</i>	<i>By</i>	<i>To Obtain</i>
cubic foot per second (ft <sup>3</sup> /s)	0.02832	cubic meter per second
mile (mi)	1.609	kilometer
pound (lb)	453.6	gram
square mile (mi <sup>2</sup> )	2.590	square kilometer

Temperature can be converted to degrees Celsius ( $^{\circ}\text{C}$ ) or degrees Fahrenheit ( $^{\circ}\text{F}$ ) by the equations:

$$\begin{aligned}^{\circ}\text{C} &= \frac{5}{9} (^{\circ}\text{F} - 32) \\ ^{\circ}\text{F} &= \frac{9}{5} (^{\circ}\text{C}) + 32.\end{aligned}$$

### Abbreviated Water-Quality Units

- microgram per liter ( $\mu\text{g/L}$ )
- microsiemens per centimeter at 25 degrees Celsius ( $\mu\text{S/cm}$ )
- milligram (mg)
- milligram per liter (mg/L)
- milliliter (mL)
- milliliter per minute (mL/min)
- nanogram per microliter (ng/ $\mu\text{L}$ )



# RECONNAISSANCE DATA FOR SELECTED HERBICIDES, TWO ATRAZINE METABOLITES, AND NITRATE IN SURFACE WATER OF THE MIDWESTERN UNITED STATES, 1989-90

By Elisabeth A. Scribner, E. Michael Thurman, Donald A. Goolsby,  
Michael T. Meyer, Margaret S. Mills, and Michael L. Pomes

## ABSTRACT

Water-quality data were collected from 147 rivers and streams during 1989-90 to determine the temporal and geographic distribution of selected preemergent herbicides, two atrazine metabolites, and nitrate in 10 Midwestern States. This report includes a description of the sampling design, data-collection techniques, laboratory and analytical methods, and a compilation of constituent concentrations and quality-assurance data. All water samples were collected by depth-integrating techniques at three to five locations across the wetted perimeter of each stream. Sites were sampled three times in 1989--before application of herbicides, during the first major runoff after application of herbicides, and in the fall during a low-flow period when most of the streamflow was derived from ground water. About 50 sites were selected by a stratified random procedure and resampled for both pre- and post-application herbicide concentrations in 1990 to verify the 1989 results. Laboratory analyses consisted of both enzyme-linked immunosorbent assay (ELISA) and confirmation by gas chromatography/mass spectrometry (GC/MS). The data have been useful in studying herbicide transport, in comparison of the spatial distribution of the post-application concentrations of 11 herbicides and 2 atrazine metabolites (deethylatrazine and deisopropylatrazine) in streams and rivers at a regional scale, in examination of the annual persistence of herbicides and two atrazine metabolites in surface water, and in assessment of atrazine metabolites as indicators of surface- and ground-water interaction.


## INTRODUCTION

This is one of several water-quality reports intended to present the results of a reconnaissance study of preemergent herbicides, two atrazine metabolites, and nitrite plus nitrate in surface water of a 10-state region in the midwestern United States (fig. 1). The reconnaissance study was conducted during

1989-90 by the U.S. Geological Survey as part of the Toxic Substances Hydrology Program.

During 1987-89, about 136 million pounds per year of four major herbicides were applied in the 10-state region (table 1). The herbicides alachlor, atrazine, cyanazine, and metolachlor accounted for about 73 percent of the pesticides applied (Gianessi and Puffer, 1990). The intense use of herbicides, their moderate water solubility, and mobility can cause them to leach into ground water, to run off in surface water, as well as to be transported in air and in precipitation (Goolsby and others, 1990). Goolsby and others (1991a) identified substantial increases in herbicide concentrations in streams and rivers and in runoff from fields immediately after herbicide application. Nonpoint-source contamination of surface- and ground-water supplies may be associated with the intense use of agricultural chemicals. Drinking-water quality also may be affected because conventional water-treatment practices do not remove these moderately soluble herbicides (Goolsby and others, 1990).

This report presents the water-quality data collected during a reconnaissance study of surface water in the midwestern United States. The data are being used in subsequent reports to document the occurrence, distribution, and concentrations of selected preemergent herbicides, two atrazine metabolites, and nitrite plus nitrate, to understand the geographic and seasonal distribution of nitrite plus nitrate and commonly used herbicides in streams of different size throughout the 10-state area, and to examine the usefulness of a low-cost immunoassay analysis technique for determining atrazine concentrations in a regional-scale reconnaissance (Goolsby and others, 1990). The scope of this report includes a discussion of the sampling design, data-collection techniques, analytical procedures, and a compilation of surface-water quality and quality-assurance data.



**Figure 1.** Location of study area, hydrologic units, and sites sampled during 1989 and 1990 in the midwestern United States.

**Table 1.** Quantities of four major herbicides applied in 10 agricultural Midwestern States, 1987-89  
 [From Gianessi and Puffer, 1990; values shown are in millions of pounds of active ingredient per year]

State	Alachlor	Atrazine	Cyanazine	Metolachlor
Illinois	8.0	8.5	3.1	8.1
Indiana	6.9	5.7	1.7	3.5
Iowa	6.4	5.6	3.7	8.5
Kansas	1.9	4.7	.2	2.2
Minnesota	4.0	1.5	2.8	2.4
Missouri	1.8	3.1	1.2	1.6
Nebraska	3.8	7.1	1.8	1.9
Ohio	3.7	3.8	1.6	4.0
South Dakota	1.9	.5	.3	1.0
Wisconsin	1.3	2.7	1.7	1.4
<b>Total</b>	<b>39.7</b>	<b>43.2</b>	<b>18.1</b>	<b>34.6</b>
Total of four major herbicides: 135.6				

The study area includes the following states: Illinois, Indiana, Iowa, Kansas, Minnesota, Missouri, Nebraska, Ohio, South Dakota, and Wisconsin. Sampling-site names and locations are given in table 2, and the corresponding map locations are shown in figures 2 through 5. Also, included in table 2 is the drainage area of each sampling site.

## METHODS

Sampling sites were geographically distributed throughout the study area and represented streams and rivers of all sizes. A sampling schedule was established for each calendar year. Sampling techniques, including data-collection methods, processing, packaging, and shipping, were established. Laboratory methods were developed for enzyme-linked immunosorbent assay (ELISA) and gas chromatography/mass spectrometry (GC/MS).

### Sampling Design

Sampling sites were selected at 147 U.S. Geological Survey streamflow-gaging stations by a stratified random-sampling procedure designed to ensure geographic distribution and regional-scale interpretation of the data. The number of sites per state was proportional to the amount of corn and soybean production in each state, and sites were chosen randomly by county. The drainage area of the basins sampled ranged from 66 to more than 700,000 mi<sup>2</sup>, with a median drainage area of 770 mi<sup>2</sup>. Most of the sites were sampled three times in 1989--(1) before application of herbicides (March or April); (2) after application and during the first major runoff (May or June); and (3) in the fall during a low-flow period (October or November) when most of the streamflow was derived from ground water. A few sites were not sampled after herbicide application because of drought conditions or no runoff.

About one-third of the sites sampled in 1989 were resampled during 1990 before (March or April) and after (May or June) herbicide application. Based on a ranking of 1989 herbicide concentrations from highest to lowest values, sampling sites in 1990 were selected as follows: 50 percent of the sites to be sampled in 1990 were randomly selected from the upper one-third of 1989 herbicide concentrations, 25 percent from the middle one-third, and 25 percent from the lower one-third.

### Data-Collection Techniques

All water samples were collected by depth-integrating techniques at three to five locations across each stream (Thurman and others, 1992). The herbicide samples were collected in glass or Teflon<sup>1</sup> containers, composited in large glass containers, and filtered through 1-μm (micrometer) pore diameter, glass-fiber filters into baked glass bottles prior to shipment to the laboratory.

### Analytical Procedures

Specific conductance and pH measurements were made onsite. Samples, preserved with mercuric chloride, were analyzed for nitrate plus nitrite at the National Water Quality Laboratory, Arvada, Colorado (Fishman and Friedman, 1989). Herbicides were analyzed by enzyme-linked immunosorbent assay at the U.S. Geological Survey laboratory in Iowa City, Iowa (Goolsby and others, 1991b; table 6 at the end of this report). Most samples were also analyzed by gas chromatography/mass spectrometry (Thurman and others, 1990) at the U.S. Geological Survey laboratory in Lawrence, Kansas (table 7 at the end of this report). A comparison of the immunoassay method to the GC/MS analysis is presented in Goolsby and others (1991b).

Solvents used for analyzing samples included pesticide-grade methanol, ethyl acetate, and isoctane. Deionized water was charcoal filtered and glass distilled prior to use. Internal quantitative standard solutions were prepared in methanol and phenanthrene-d<sub>10</sub>. Assay kits were used for the immunoassay analysis. The kits used polyclonal antibodies coated on the walls of polystyrene test tubes and an atrazine-enzyme conjugate prepared by covalently binding atrazine to horseradish peroxidase by a modified carbodiimide technique (Bushway and others, 1988). The immunoassay procedure is described by Goolsby and others (1991b).

An automated workstation was used for solid-phase extraction of the herbicide compounds. Cartridges were preconditioned

<sup>1</sup>The use of brand names in this report is for identification purposes only and does not constitute endorsement by the U.S. Geological Survey.

**Table 2.** *Sampling-site names, locations, and drainage areas*

Map no. (figs. 2-5)	State	Site identifier	Site name	Latitude (degrees, minutes, seconds)	Longitude (degrees, minutes, seconds)	Drainage area (square miles)
1	S. Dak.	06479438	Big Sioux River near Watertown	45°00'22"	97°09'53"	1,007
		06479525	Big Sioux River near Castlewood	44°43'54"	97°02'39"	1,997
		06480000	Big Sioux River near Brookings	44°13'27"	96°46'06"	3,898
		06482020	Big Sioux River at N. Cliff Avenue	43°34'01"	96°42'39"	5,216
		06478500	James River near Scotland	43°11'09"	97°38'07"	20,653
		06479010	Vermillion River near Vermillion	42°49'02"	96°55'26"	1,779
		06485696	Brule Creek near Elk Point	42°48'32"	96°41'11"	204
		06485500	Big Sioux River at Akron	42°50'14"	96°33'41"	8,424
9	Nebr.	06478518	Bow Creek near St. James	42°43'48"	97°08'53"	304
		06796973	Elkhorn River near Atkinson	42°29'12"	98°54'42"	586
		06798300	Clearwater Creek near Clearwater	42°08'20"	98°12'10"	210
		06788988	Mira Creek near Northloup	41°30'09"	98°47'47"	66
		06795500	Shell Creek near Columbus	41°34'33"	97°46'55"	270
		06800000	Maple Creek near Nickerson	41°32'44"	96°30'09"	450
		06783500	Mud Creek at Sweetwater	41°02'15"	98°59'35"	707
		06879900	Big Blue River at Surprise	41°06'05"	97°18'35"	345
		06804000	Wahoo Creek at Itica	41°08'40"	96°32'10"	271
		06880800	W. Fork Big Blue River near Dorchester	40°43'52"	97°10'38"	1,206
		06803000	Salt Creek at Roca	40°39'29"	96°39'55"	167
		06844000	Muddy Creek near Arapahoe	40°18'20"	99°54'40"	246
		06884000	Little Blue River near Fairbury	40°06'54"	97°10'13"	2,350
		06811500	Little Nemaha River at Auburn	40°23'33"	95°48'46"	793
		06882000	Big Blue River at Barneston	40°02'40"	96°35'12"	4,447
		06815000	Big Nemaha River at Fall City	40°02'08"	95°35'45"	1,340
25	Kans.	06814000	Turkey Creek near Seneca	39°56'52"	96°06'30"	276
		06885500	Black Vermillion River near Frankfort	39°41'03"	96°26'15"	410
		06890100	Delaware River near Muscotah	39°31'17"	95°31'57"	431
		06876700	Salt Creek near Ada	39°08'30"	97°50'10"	384
		06888500	Mill Creek near Paxico	39°03'44"	96°10'52"	316
		06889000	Kansas River at Topeka	39°04'00"	95°38'58"	56,720
31	Minn.	05337400	Knife River near Mora	45°55'12"	93°18'26"	102
		05270500	Sauk River near St. Cloud	45°33'35"	94°14'00"	925
		05294000	Pomme de Terre River at Appleton	45°12'10"	96°01'20"	905
		05286000	Rum River near St. Francis	45°19'40"	93°22'20"	1,360
35	Wisc.	05340500	St. Croix River at St. Croix Falls	45°24'25"	92°38'49"	6,240

**Table 2.** Sampling-site names, locations, and drainage areas--Continued


Map no. (figs. 2-5)	State	Site identifier	Site name	Latitude (degrees, minutes, seconds)	Longitude (degrees, minutes, seconds)	Drainage area (square miles)
36	Minn.	05280000	Crow River at Rockford	45°05'12"	93°44'02"	2,520
37		05330000	Minnesota River near Jordan	44°41'35"	93°38'30"	16,200
38		05316500	Redwood River near Redwood Falls	44°31'25"	95°10'20"	697
39		05317000	Cottonwood River near New Ulm	44°17'29"	94°26'24"	1,280
40		05320500	Le Sueur River near Rapidan	44°06'40"	94°02'28"	1,100
41		05353800	Straight River near Faribault	44°15'29"	93°13'51"	442
42		05376000	N. Fork Whitewater River near Elba	44°05'30"	92°03'57"	101
43		06483000	Rock River at Luverne	43°39'15"	96°12'03"	425
44		05476000	Des Moines River at Jackson	43°37'10"	94°59'10"	1,220
45		05384500	Rush Creek near Rushford	43°50'00"	91°46'40"	129
46	Iowa	06605000	Ocheyedan River near Spencer	43°07'44"	95°12'37"	426
47		05476500	Des Moines River at Estherville	43°23'51"	94°50'38"	1,372
48		05459500	Winnebago River at Mason City	43°09'54"	93°11'33"	526
49		05458000	Little Cedar River near Ionia	43°02'05"	92°30'05"	306
50		05411600	Turkey River at Spillville	43°12'28"	91°56'56"	177
51		05388250	Upper Iowa River near Dorchester	43°25'16"	91°30'31"	770
52		06606600	Little Sioux River at Correctionville	42°28'20"	95°47'49"	2,500
53		05482300	North Raccoon River near Sac City	42°20'28"	94°59'05"	713
54		05480500	Des Moines River at Fort Dodge	42°30'22"	94°12'04"	4,190
55		05449500	Iowa River near Rowan	42°45'36"	93°37'23"	429
56		05463050	Cedar River at Cedar Falls	42°30'50"	92°37'55"	4,734
57		05463500	Black Hawk Creek at Hudson	42°24'28"	92°27'47"	303
58		05421000	Wapsipinicon River at Independence	42°27'49"	91°53'42"	1,048
59		06607200	Maple River at Mapleton	42°09'28"	95°48'27"	669
60		06609500	Boyer River at Logan	41°38'33"	95°46'57"	871
61		05484500	Raccoon River at Van Meter	41°32'02"	93°56'59"	3,441
62		05471200	Indian Creek near Mingo	41°48'17"	93°18'26"	276
63		05453100	Iowa River near Marengo	41°48'41"	92°03'42"	2,794
64		05455100	Old Mans Creek near Iowa City	41°36'25"	91°36'40"	201
65		05418450	N. Fork Maquoketa River at Fulton	42°08'42"	90°40'55"	516
66		05471500	S. Skunk River near Oskaloosa	41°21'19"	92°39'31"	1,635
67		05472500	N. Skunk River near Sigourney	41°18'03"	92°12'16"	730
68		06810000	Nishnabotna River above Hamburg	40°37'57"	95°37'32"	2,806
69		06903400	Chariton River near Chariton	40°57'12"	93°15'37"	182
70		06903700	S. Fork Chariton River near Promise City	40°48'02"	93°11'32"	168
71		05474000	Skunk River at Augusta	40°45'13"	91°16'40"	4,303
72	Mo.	06817700	Nodaway River near Graham	40°12'08"	95°04'07"	1,320
73		05500000	South Fabius River near Taylor	39°53'49"	91°34'49"	620

**Table 2. Sampling-site names, locations, and drainage areas--Continued**

Map no. (figs. 2-5)	State	Site identifier	Site name	Latitude (degrees, minutes, seconds)	Longitude (degrees, minutes, seconds)	Drainage area (square miles)
74 Mo.		06902000	Grand River near Sumner	39°38'25"	93°16'25"	6,880
75		05506500	Middle Fork Sal River at Paris	39°29'05"	91°59'50"	356
76		06908000	Blackwater River at Blue Lick	38°59'32"	93°11'48"	1,120
77		06934500	Missouri River at Hermann	38°42'36"	91°26'21"	524,200
78		05587450	Mississippi River at Grafton	38°48'46"	90°40'38"	141,000
79 Wisc.		05394500	Prairie River near Merrill	45°14'09"	89°38'59"	184
80		05379500	Trempealeau River at Dodge	44°07'55"	91°33'14"	643
81		05382000	Black River near Galesville	44°04'22"	91°17'41"	2,080
82		05407000	Wisconsin River at Muscada	43°11'54"	90°26'26"	10,400
83		05413500	Grant River near Burton	42°43'13"	90°49'09"	269
84		05434500	Pecatonic River at Martintown	42°30'34"	89°47'58"	1,034
85		05430500	Rock River at Afton	42°36'33"	89°04'14"	3,340
86		05543830	Fox River at Wakesha	43°00'17"	88°14'37"	126
87		04087240	Root River at Racine	42°45'05"	87°49'25"	190
88 Ill.		05527800	Des Plains River at Russell	42°29'22"	87°55'32"	123
89		05548280	Nippersink Creek at Spring Cove	42°26'37"	88°14'51"	192
90		05439500	S. Br. Kishwaukee River at Fairdale	42°06'40"	88°54'00"	387
91		05440000	Kishwaukee River near Perryville	42°11'45"	88°59'55"	1,099
92		05444000	Elkhorn Creek near Penrose	41°54'10"	89°41'40"	146
93		05543500	Illinois River near Marseilles	41°19'40"	88°43'10"	8,259
94		05552500	Fox River near Dayton	41°23'12"	88°47'26"	2,642
95		05540500	Dupage River near Shorwood	41°31'20"	88°11'35"	324
96		05526000	Iroquois River near Chebanse	41°00'32"	87°49'27"	2,091
97		05466500	Edwards River near New Boston	41°11'15"	90°58'05"	445
98		05569500	Spoon River at London Mills	40°42'32"	90°16'53"	1,072
99		05584500	La Moine River at Colmar	40°19'45"	90°53'55"	655
100		05583000	Sangamon River at Oaksford	40°07'25"	89°59'05"	5,093
101		05579500	Lake Fork near Cornland	39°57'00"	89°23'10"	214
102		05576500	Sangamon River at Riverton	39°50'34"	89°32'52"	2,618
103		05587060	Illinois River at Hardin	39°09'37"	90°36'55"	28,690
104		05587000	Macoupin Creek near Kane	39°14'03"	90°23'40"	868
105		05592100	Kaskaskia River near Cowden	39°13'50"	88°50'33"	1,330
106		05592500	Kaskaskia River at Vandalia	38°57'35"	89°05'20"	1,940
107		03345500	Embarрас River at Ste. Marie	38°56'10"	88°01'10"	1,516
108		05594000	Shoal Creek near Breese	38°36'35"	89°29'40"	735
109		05594800	Silver Creek near Freeburg	38°24'22"	89°52'26"	464
110		03378000	Bonpas Creek at Browns	38°23'11"	87°58'32"	228

**Table 2.** Sampling-site names, locations, and drainage areas--Continued

Map no. (figs. 2-5)	State	Site identifier	Site name	Latitude (degrees, minutes, seconds)	Longitude (degrees, minutes, seconds)	Drainage area (square miles)
111	Ind.	03378500	Wabash River near New Harmony	38°13'42"	87°59'04"	29,080
112	Ill.	03381495	Little Wabash River at Carmi	38°05'32"	88°09'22"	3,088
113		05599500	Big Muddy River near Murphysboro	37°44'55"	89°20'45"	2,169
114	Mo.	07022000	Mississippi River at Thebes	37°13'00"	89°27'50"	713,200
115	Ill.	03612500	Ohio River near Grand Chain	37°10'34"	89°04'02"	203,100
116	Ind.	05518000	Kankakee River at Shelby	41°10'58"	87°20'33"	1,779
117		03331500	Tippecanoe River near Ora	41°09'26"	86°33'49"	856
118		05524500	Iroquois River near Foresman	40°52'14"	87°18'24"	449
119		03328500	Eel River near Logansport	40°46'55"	86°15'50"	789
120		03322900	Wabash River at Linn Grove	40°39'22"	85°01'58"	453
121		03335000	Wildcat Creek near Lafayette	40°26'26"	86°49'45"	794
122		03333450	Wildcat Creek near Jerome	40°26'29"	85°55'08"	146
123		03336000	Wabash River at Covington	40°08'24"	87°24'24"	8,218
124		03351000	White River near Nora	39°54'35"	86°06'20"	1,219
125		03361000	Big Blue River at Carthage	39°44'38"	85°34'33"	184
126		03275000	Whitewater River near Alpine	39°34'46"	85°09'29"	522
127		03357500	Big Walnut Creek near Reelsville	39°32'11"	86°58'35"	326
128		03354000	White River near Centerton	39°29'51"	86°24'02"	2,444
129		03362500	Sugar Creek near Edinburgh	39°21'39"	85°59'51"	474
130		03363900	Flatrock River at Columbus	39°14'06"	85°55'36"	534
131		03342500	Busseron Creek near Carlisle	38°58'26"	87°25'33"	228
133		03366500	Muscatatuck River near Deputy	38°48'15"	85°40'26"	293
134		03302800	Blue River at Fredricksburg	38°26'02"	86°11'31"	283
135	Ohio	04185000	Tiffin River at Stryker	41°30'16"	84°25'47"	410
136		04193500	Maumee River at Waterville	41°28'34"	83°44'20"	6,330
137		04198000	Sandusky River near Fremont	41°18'28"	83°09'32"	1,251
138		04186500	Auglaize River near Fort Jennings	40°56'55"	84°15'58"	332
139		03223000	Olentangy River at Claridon	40°34'58"	82°59'20"	157
140		03219500	Scioto River near Prospect	40°25'10"	83°11'50"	567
141		03136500	Kokosing River at Mount Vernon	40°24'20"	82°30'00"	202
142		03267900	Mad River at Eagle City	39°57'51"	83°49'54"	310
143		03240000	Little Miami River near Oldtown	39°44'54"	83°55'53"	129
144		03230500	Big Darby Creek at Darbyville	39°42'02"	83°06'37"	534
145		03157000	Clear Creek near Rockbridge	39°35'18"	82°34'43"	89
146		03234500	Scioto River at Higby	39°12'44"	82°51'50"	5,131
147		03245500	Little Miami River at Milford	39°10'17"	84°17'53"	1,203




**Figure 2.** Location of sites sampled in South Dakota, Nebraska, and Kansas, map numbers 1 through 30.


sequentially with 2 mL methanol, 6 mL ethyl acetate, 2 mL methanol, and 2 mL distilled water. Each 123-mL water sample was spiked with a surrogate standard, 2.4 ng/ $\mu$ L terbutylazine, and pumped through the cartridge at a rate of 20 mL/min by the robotic probe. Herbicide compounds were eluted with ethyl acetate and spiked robotically with phenanthrene-d<sub>10</sub>. The ethyl-acetate layer was transferred by probe to a clean test tube. Finally, the extract was evaporated automatically by a turbovap at 45 °C under a nitrogen stream to approximately 100  $\mu$ L. The robotic probe was washed between samples by immersing in 4 mL of ethyl acetate and bubbling filtered compressed air through

the probe to ensure thorough removal of any herbicide or spike residues adhering to the outside of the probe.

Automated GC/MS analyses of the eluates were performed on a mass selective detector. Operating conditions were as follows: ionization voltage, 70 electronvolts; ion source temperature, 250 °C; electron multiplier, 2,200 volts; direct capillary interface at 280 °C, tuned daily with perfluorotributylamine; dwell time, 50 milliseconds. Separation of the herbicides was carried out using a 12-meter fused-silica capillary column, 0.2 millimeter in diameter with a methyl silicone stationary phase,



**Figure 3.** Location of sites sampled in Minnesota, Iowa, and Missouri, map numbers 31 through 78 (site 35 sampled in Wisconsin).




**Figure 4.** Location of sites sampled in Wisconsin and Illinois, map numbers 79 through 115 (site 111 sampled in Indiana and site 114 sampled in Missouri).

0.33  $\mu\text{m}$  thick. Helium was used as the carrier gas at a flow rate of 1 mL/min and a head pressure of 35 kilopascals. The column temperature was held at 50 °C for 1 minute, then ramped at 6 °C per minute to 250 °C. Injector temperature was 280 °C. The filament and multiplier were not turned on until 5 minutes into the analysis. Quantification of the base peak of each compound was based on the response of the 188 ion of the internal standard, phenanthrene-d<sub>10</sub>. Confirmation of the compound was based on the presence of the

molecular ion and one to two confirming ions with a retention-time match of + 0.2 percent relative to phenanthrene-d<sub>10</sub>. This procedure is described by Thurman and others (1990) and Meyer and others (1993).

## Quality Assurance

All water-quality measurements and water samples for this study were collected by U.S. Geological Survey personnel. All samples for GC/MS analysis were collected in duplicate to



**Figure 5.** Location of sites sampled in Indiana and Ohio, map numbers 116 through 147.

provide a backup sample in case reanalysis was needed. Ten percent of the duplicates were analyzed by GC/MS for quality-assurance purposes. These analyses are identified in table 7 at the end of this report by a sample-type code of "L." In addition, blind duplicates were collected and analyzed by both GC/MS and immunoassay for about 5 percent of the samples. These samples were disguised in such a way that laboratories doing the analyses were not aware they were duplicates. Analyses of these samples are identified in table 7 by a sample-type code of "B." About 2 percent of the samples analyzed by GC/MS were spikes prepared in organic-free distilled water by the U.S. Geological Survey's National Water Quality Laboratory (NWQL) in Arvada, Colorado. Analyses of these spikes are given in table 3. Also, included in the table are the herbicide concentrations added to the samples and an analysis of spikes by the NWQL. Analyses in table 3 identified by a sample-type code of "S" were known to be spikes by the analyzing laboratory. Some spikes were disguised as water samples. Results for these samples are identified in table 3 by a sample-type code of "BS."

## DISCUSSION

Analytical results are presented in tables 6 and 7 at the end of this report. These results

indicate that large concentrations of herbicides were flushed from cropland and transported through the surface-water system in response to late spring and early summer rainfall (Thurman and others, 1992). Of the 11 herbicides and 2 atrazine metabolites analyzed, atrazine, alachlor, cyanazine, and metolachlor, were present in the largest percentage of samples, as shown in table 4. During the 1989 post-application sampling, atrazine was detected (greater than 0.05 µg/L) in 98 percent of the samples. Alachlor, metolachlor, and deethylatrazine, an atrazine metabolite, were detected (greater than 0.05 µg/L) in water in more than 80 percent of the samples, and cyanazine was detected (greater than 0.20 µg/L) in more than 60 percent of the samples. Similar seasonal distributions were observed for the other major herbicides. It should be noted that the two atrazine metabolites (deethylatrazine and deisopropylatrazine) also can be produced by simazine and propazine. However, because these two herbicides are generally present in low concentrations relative to atrazine, their contribution to the formation of these two metabolites should be minimal.

During the late spring and summer (May or June, 1989-90), concentrations of one or more herbicides exceeded U.S. Environmental Protection Agency (USEPA) Maximum Contaminant Levels (MCL) or lifetime Health Advisories

**Table 3. Concentrations of selected herbicides and two atrazine metabolites in quality-assurance samples, 1989-90**

[DEA, deethylatrazine; DIA, desopropylatrazine; ametryn, prometryn, terbutryn were not detected; S, spike sample; BS, blind spiked sample; spike, spiked concentration; NWQL, analysis by U.S. Geological Survey National Water Quality Laboratory, Arvada, Colo.; --, no data; <, less than]

Laboratory identification number	Quality assurance identification	Round <sup>1</sup>	Sample type	Concentrations, in micrograms per liter						
				Alachlor	Atrazine	Cyana-zine	DEA	DIA	Metala-chlor	Metri-buzin
Organic-free water										
MT-101A	0DG0	1	BS	<.05	<.20	<.05	<.05	<.05	<.05	<.05
MT-102	0DG0	1	BS	<.05	<.20	<.05	<.05	<.05	<.05	<.05
MT-1217A	0DG0	3	BS	<.05	<.20	<.05	<.05	<.05	<.05	<.05
MT-2077	0DG0	4	S	<.05	<.20	<.05	<.05	<.05	<.05	<.05
Mean				<.05	<.20	<.05	<.05	<.05	<.05	<.05
Theoretical	IDG1	--	--	<.05	<.05	<.05	<.05	<.05	<.05	<.05
MT-35A	IDG1	1	BS	.14	.07	<.20	<.05	<.05	.40	<.05
MT-67	IDG1	1	S	.25	.13	.24	<.05	<.05	.14	.43
MT-7	IDG1	1	S	.22	.14	<.20	<.05	<.05	.10	.61
MT-8	IDG1	1	S	.20	.13	<.20	<.05	<.05	.11	.47
Mean				.20	.12	<.20	<.05	<.05	.09	.48
Theoretical	IDG1	--	--	.26	.11	.19	<.05	<.05	.14	.30
NWQL	IDG1	--	--	.26	.11	.19	--	--	<.10	.30
MT-3	IDG2	1	S	.54	.38	<.20	<.05	<.05	.22	.83
MT-4	IDG2	1	S	.59	.46	<.20	<.05	<.05	.25	.90
MT-69	IDG2	1	S	.56	.33	.46	<.05	<.05	.23	.71
Mean				.56	.39	.22	<.05	<.05	.23	.81
Theoretical	IDG2	--	--	.66	.32	.57	<.05	<.05	.27	.75
NWQL	IDG2	--	--	--	--	--	--	--	--	--
MT-5	IDG3	1	S	1.1	1.4	<.20	<.05	<.05	.58	1.4
MT-6	IDG3	1	S	1.1	1.3	<.20	<.05	<.05	.58	1.3
MT-65	IDG3	1	S	1.3	1.4	<.20	<.05	<.05	.64	1.5
MT-97A	IDG3	1	BS	.93	1.4	<.20	<.05	<.05	.60	1.7
MT-97B	IDG3	1	BS	.97	1.3	<.20	<.05	<.05	.54	1.4
Mean				1.1	1.4	<.20	<.05	<.05	.59	1.5
Theoretical	IDG3	--	--	1.3	1.1	.95	<.05	<.05	.68	1.5
NWQL	IDG3	--	--	1.2	1.0	1.1	--	--	.70	1.2

**Table 3.** Concentrations of selected herbicides and two atrazine metabolites in quality-assurance samples, 1989-90--Continued

Laboratory identification number	Quality assurance identification	Round <sup>1</sup>	Sample type	Concentrations, in micrograms per liter									
				Aalachlor	Atrazine	Cyana-zine	DEA	DIA	Metola-chlor	Metri-buzin	Pro-meton	Pro-pazine	Simazine
<u>Organic-free water--Continued</u>													
MT-2	1DG4	1	S	3.0	<.05	2.1	<.05	.05	2.3	2.6	<.05	2.3	0.68
MT-6B	1DG4	1	S	3.9	<.05	1.8	<.05	.05	2.9	3.2	<.05	2.7	.75
Mean				3.5	<.05	2.0	<.05	.05	2.6	2.9	<.05	2.5	.72
Theoretical	1DG4	--	--	3.9	3.8	1.9	<.05	.05	2.7	3.0	<.05	1.7	.71
NWQL	1DG4	--	--	--	--	--	--	--	--	--	--	--	
MT-10	1DG5	1	S	4.8	5.1	5.7	<.05	.05	4.4	4.8	<.05	4.6	1.4
MT-66	1DG5	1	S	6.0	5.7	5.5	<.05	.05	5.2	6.5	<.05	5.3	1.6
MT-9	1DG5	1	S	4.7	4.9	5.5	<.05	.05	4.3	4.8	<.05	4.5	1.3
Mean				5.2	5.2	5.6	<.05	.05	4.6	5.4	<.05	4.8	1.4
Theoretical	1DG5	--	--	6.5	4.3	5.7	<.05	.05	5.4	6.0	<.05	3.4	1.4
NWQL	1DG5	--	--	--	--	--	--	--	--	--	--	--	
MT-275A	2DG1	2	S	.44	.40	<.20	<.05	.05	5.6	1.5	<.05	1.5	2.3
Mean				.44	.40	<.20	<.05	.05	5.6	1.5	<.05	1.5	2.3
Theoretical	2DG1	--	--	.40	.27	<.20	<.05	.05	5.5	1.5	<.05	1.4	2.1
NWQL	2DG1	--	--	.30	.22	--	<.05	.05	6.1	1.4	--	1.3	2.0
MT-276A	2DG2	2	S	1.3	1.1	.80	<.05	.05	1.7	<.05	<.05	5.3	5.9
MT-612A	2DG2	2	BS	1.1	1.1	.58	<.05	.05	1.5	<.05	<.05	5.0	5.4
MT-612B	2DG2	2	BS	1.1	1.1	.70	<.05	.05	1.5	<.05	<.05	5.0	5.4
Mean				1.2	1.1	.69	<.05	.05	1.6	<.05	<.05	5.1	5.6
Theoretical	2DG2	--	--	1.3	.90	1.0	<.05	.05	2.0	--	--	5.1	5.7
NWQL	2DG2	--	--	1.0	.90	1.5	--	--	1.5	--	--	5.2	6.2
MT-277A	2DG3	2	S	5.4	4.9	6.4	<.05	.05	.26	.37	<.05	.54	.51
MT-506A	2DG3	2	BS	1.7	3.5	<.20	<.05	.05	.07	<.05	<.05	.38	.26
Mean				3.6	4.2	3.3	<.05	.05	.17	.20	<.05	.46	.39
Theoretical	3DG3	--	--	5.2	3.8	5.7	<.05	.05	.30	.30	<.05	.42	.40
NWQL	3DG3	--	--	4.6	3.8	4.8	--	--	.30	.30	--	.50	.40

**Table 3.** Concentrations of selected herbicides and two atrazine metabolites in quality-assurance samples, 1989-90--Continued

Quality assurance identification number	Round <sup>1</sup>	Sample type	Alachlor	Atrazine	Cyanazine	DEA	DIA	Metolachlor	Metrybuzin	Prometon	Propazine	Simazine	Concentrations, in micrograms per liter	
													Organic-free water--Continued	
MT-1203A	3DG1	3	S	0.33	0.53	1.6	0.13	<.05	3.7	3.8	<.05	0.61	4.4	
MT-1219B	3DG1	3	BS	.34	.56	1.6	.11	<.05	4.7	3.7	<.05	.66	5.3	
MT-1221A	3DG1	3	BS	.33	.51	1.9	.10	<.05	3.3	4.0	<.05	.62	4.2	
MT-1221B	3DG1	3	BS	.30	.73	2.1	.11	<.05	3.6	7.3	<.05	.66	4.9	
MT-1225A	3DG1	3	BS	.37	.56	1.7	.11	<.05	3.3	3.9	<.05	.63	4.4	
MT-1294	3DG1	3	S	.30	.69	3.0	<.05	<.05	4.5	4.0	<.05	.64	4.6	
Mean				.33	.60	2.0	.11	<.05	3.9	4.5	<.05	.64	4.6	
Theoretical NWQL	3DG1	--	--	.30	.50	4.0	.10	<.05	3.5	3.0	<.05	.50	4.5	
	3DG1	--	--	.27	.44	2.4	.19	--	2.3	2.7	--	.45	3.3	
MT-1204A	3DG2	3	S	3.3	3.8	.70	.38	<.05	1.3	.57	<.05	1.8	1.5	
MT-1218B	3DG2	3	BS	3.1	5.4	1.1	.39	<.05	1.2	.70	<.05	1.9	1.5	
MT-1220B	3DG2	3	BS	3.2	5.1	.71	.32	<.05	1.1	.56	<.05	1.8	1.3	
MT-1224A	3DG2	3	BS	3.1	3.8	1.0	.37	<.05	1.2	.73	<.05	1.8	1.7	
MT-1293	3DG2	3	S	4.2	5.1	1.8	.36	<.05	1.5	.65	<.05	1.9	1.5	
Mean				3.4	4.6	1.1	.36	<.05	1.3	.64	<.05	1.8	1.5	
Theoretical NWQL	4DG2	--	--	3.0	4.0	1.5	.40	<.05	1.1	.50	<.05	1.5	1.5	
	4DG2	--	--	2.7	3.8	.94	.49	--	1.0	.41	--	1.4	1.1	
MT-1964	4DG1	4	S	.34	.25	.38	<.05	<.05	.21	<.05	<.05	<.05	<.05	
MT-2155	4DG1	4	S	.33	.27	<.20	<.05	<.05	.19	<.05	<.05	<.05	<.05	
MT-3138	4DG1	4	S	.35	.26	<.20	<.05	<.05	.19	<.05	<.05	<.05	<.05	
Mean				.34	.26	<.20	<.05	<.05	.20	<.05	<.05	<.05	<.05	
Theoretical NWQL	4DG1	--	--	.30	.20	.05	<.05	<.05	.20	<.05	<.05	<.05	<.05	
	4DG1	--	--	.30	.20	--	--	--	.20	--	--	--	--	

**Table 3.** Concentrations of selected herbicides and two atrazine metabolites in quality-assurance samples, 1989-90--Continued

Laboratory identification number	Quality assurance identification	Round <sup>1</sup>	Sample type	Alachlor	Atrazine	Cyanazine	DEA	DIA	Metolachlor	Metribuzin	Prometon	Propazaine	Simazine	Concentrations, in micrograms per liter	
														Organic-free water--Continued	
MT-1963	4DG2	4	S	0.78	1.4	<0.20	<0.05	<0.05	0.48	0.23	<0.05	<0.05	0.55		
MT-2156	4DG2	4	S	.86	1.5	.20	<.05	<.05	.50	.17	<.05	<.05	.50		
MT-3139	4DG2	4	S	.99	1.5	.50	<.05	<.05	.51	.28	<.05	<.05	.59		
Mean				.88	1.5	.27	<.05	<.05	.50	.23	<.05	<.05	.55		
Theoretical	4DG2	--	--	.80	1.2	.50	<.05	<.05	.50	.20	<.05	<.05	.50		
NWQL	4DG2	--	--	.60	1.0	.50	--	--	.40	.20	--	--	.40		

<sup>1</sup>Round-1, 1989 pre-application (March or April);

2, 1989 post-application (May or June);

3, 1989 fall low flow (October or November); and

4, 1990 pre-application (March or April).

**Table 4.** *Herbicide compounds analyzed and percentage of detections in pre-application, post-application, and fall low-flow samples during 1989-90*

[ $\mu\text{g/L}$ , micrograms per liter; N, number of samples analyzed; <, less than. 1990 samples are from a stratified random subset of sites sampled in 1989 and are biased toward a slightly higher detection rate; ametryn, prometryn, terbutryn were not detected]

Herbicide or metabolite	Reporting limit ( $\mu\text{g/L}$ )	Percentage of detections				
		1989 pre-application (N=55)	1989 post-application (N=132)	1989 fall low flow (N=145)	1990 pre-application (N=52)	1990 post-application (N=52)
Alachlor	0.05	18	86	12	50	92
Atrazine	.05	91	98	76	96	100
Cyanazine	.20	5	63	0	8	81
Deethylatrazine	.05	54	86	47	86	98
Deisopropylatrazine	.05	9	54	0	44	90
Metolachlor	.05	34	83	44	67	96
Metribuzin	.05	2	53	0	0	58
Prometon	.05	0	23	6	10	17
Propazine	.05	0	40	<1	0	65
Simazine	.05	7	55	3	15	67

Levels (HAL) for drinking water (U.S. Environmental Protection Agency, 1988; Federal Register, 1989, 1990) in more than one-half of the samples (table 5). The majority of exceedences occurred during the post-application samplings (table 5). However, in order for these concentrations to be in violation of the Safe Drinking Water Act, the average annual concentrations must exceed the MCLs.

Analytical results indicate that atrazine was the most frequently detected and the most persistent of the major herbicides followed by deethylatrazine and metolachlor (Goolsby and others, 1990). Alachlor and cyanazine were detected primarily during the late spring and summer (table 7 at the end of this report). Atrazine was detected in three-fourths of the streams sampled during the fall. Resampling of about 50 streams in 1990 before and after herbicide application produced results similar to those obtained in 1989.

Concentrations of alachlor, atrazine, cyanazine, and metolachlor for the five sampling periods (data from table 7) are shown in figures 6-9. These figures also show that atrazine had the largest concentration, followed by alachlor, metolachlor, and cyanazine.

ELISA proved to be a rapid, reliable, and inexpensive method for analysis of herbicides in surface water. The presence or absence of triazine herbicides at concentrations of about 0.5  $\mu\text{g/L}$  can be determined by visually comparing water samples with standard solutions containing atrazine. When a spectrophotometer is used to quantify immunoassay results, triazine concentrations as small as 0.2  $\mu\text{g/L}$  can be detected. No false-positive identifications of atrazine were made by immunoassay, and no false-negative detections resulted from samples containing 0.5  $\mu\text{g/L}$  or more of atrazine concentrations. The immunoassay results were useful in determining the occurrence and seasonal distribution of atrazine in the corn and soybean belt of the midwestern United States (Goolsby and others, 1991b).

**Table 5.** Comparison of herbicide samples collected during 1989-90 with Maximum Contaminant Levels and Health Advisory Levels for drinking water in effect at the time of this study

[MCL, U.S. Environmental Protection Agency Maximum Contaminant Level; HAL, Health Advisory Levels; µg/L, micrograms per liter; --, no data. 1990 samples are from a stratified random subset of sites sampled in 1989 and are biased toward slightly higher concentrations. HAL for cyanazine was 10 µg/L in 1989-90; current (1993) HAL is 1 µg/L; MCL for simazine was 1 µg/L in 1989-90; current (1993) MCL is 4 mg/L]

Herbicide	Percentage of samples above MCL or HAL							
	1990 MCL (µg/L)	1990 Lifetime HAL (µg/L)	1989 pre- application (March or April)	1989 post- application (May or June)	1989 fall low flow (October or November)	1990 pre- application (March or April)	1990 post- application (May or June)	
Alachlor	2.0	--	0	36	0	0	40	
Atrazine	3.0	3.0	0	52	.60	1.6	74	
Cyanazine	--	10	0	48	0	0	63	
Simazine	1.0	4.0	5.6	8.8	0	3.1	12	

## REFERENCES CITED

- Bushway, R.J., Perkins, Bill, Savage, S.A., Lekousi, S.L., and Ferguson, B.S., 1988, Determination of atrazine residues in water and soil by enzyme immunoassay: Bulletin of Environmental Contamination Toxicology, v. 40, p. 647-654.
- Federal Register, 1989, National Primary and Secondary Drinking Water Regulations: Washington, D.C., v. 54, no. 97, May 22, 1989, p. 22062-22160.
- , 1990, National Primary and Secondary Drinking Water Regulations: Washington, D.C., v. 55, no. 143, July 25, 1990, p. 30370-30447.
- Fishman, M.J., and Friedman, L.C., 1989, Methods for the determination of inorganic substances in water and fluvial sediments: U.S. Geological Survey Techniques of Water-Resources Investigations, book 5, chap. A1, 545 p.
- Gianessi, L.P., and Puffer, C.M., 1990, Herbicide use in the United States--National Summary Report, revised April 1991: Washington, D.C., Resources for the Future, 128 p.
- Goolsby, D.A., Coupe, R.C., and Markovchick, D.J., 1991a, Distribution of selected herbicides and nitrates in the Mississippi River and its major tributaries--April-June 1991: U.S. Geological Survey Water-Resources Investigations Report 91-4163, 35 p.
- Goolsby, D. A., Thurman, E. M., Clark, M.L., and Pomes, M.L., 1991b, Immunoassay as a screening tool for atrazine herbicides in streams--Comparison with gas chromatographic/mass spectrometric methods, in Vanderlaan, Martin, Stanker, L.H., Watkins, B.E., and Roberts, D.W., eds., Immunoassays for trace chemical analysis: American Chemical Society, ACS Symposium Series 451, chap. 8, p. 86-99.
- Goolsby, D. A., Thurman, E. M., and Kolpin, D. W., 1990, Geographic and temporal distribution of herbicides in surface waters of the upper midwestern United States, 1989-90: U.S. Geological Survey Water-Resources Investigations Report 91-4034, p. 86-99.
- Meyer, M.T., Thurman, E.M., and Pedersen, M.J., 1993, Automated solid-phase extraction of herbicides from water for gas chromatographic-mass spectrometric analysis: Journal of Chromatography, v. 629, p. 55-59.




Figure 6. Concentrations of alachlor during 1989-90 reconnaissance study.




Figure 7. Concentrations of atrazine during 1989-90 reconnaissance study.





Figure 8. Concentrations of cyanazine during 1989-90 reconnaissance study.



**Figure 9.** Concentrations of metolachlor during 1989-90 reconnaissance study.

Thurman, E.M., Goolsby, D.A., Meyer, M.T., Mills, M.S., Pomes, M.L., and Kolpin, D.W., 1992, A reconnaissance study of herbicides and their metabolites in surface water of the midwestern United States using immunoassay and gas chromatography/mass spectrometry: Environmental Science and Technology, v. 26, p. 2440-2447.

Thurman, E.M., Meyer, Michael, Pomes, Michael, Perry, C.A., and Schwab, A.P.,

1990, Enzyme-linked immunosorbent assay compared with gas chromatography/mass spectrometry for the determination of triazine herbicides in water: Analytical Chemistry, v. 62, p. 2043-2048.

U.S. Environmental Protection Agency, 1988, Drinking water health advisory--Pesticides: Chelsea, Mich., Lewis Publishers, 819 p.

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90**

[ft<sup>3</sup>/s, cubic feet per second; µS/cm, microsiemens per centimeter at 25 degrees Celsius; mg/L, milligrams per liter; µg/L, micrograms per liter; --, no data; <, less than; >, greater than; ELISA, enzyme-linked immunosorbent assay; GC/MS, gas chromatography/mass spectrometry]

Map no (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream- flow (ft <sup>3</sup> /s)	Specific conduct- ance (µS/cm)	pH (standard units)	Nitrite plus nitrate as nitrogen (mg/L)	Atrazine ELISA-	Atrazine Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
							Iowa <sup>1</sup> (µg/L)		
<b>South Dakota</b>									
1	Big Sioux River near Watertown	04/04/89	71	360	8.5	0.44	<.20	--	--
		06/29/89	1.5	570	9.5	<.10	<.20	--	0.08
		11/03/89	.74	790	8.0	.14	<.20	--	--
2	Big Sioux River near Castlewood	04/04/89	32	730	8.4	.45	.20	--	--
		06/29/89	4.0	980	9.7	<.10	1.5	--	1.5
		11/03/89	6.2	1,130	8.0	3.8	.40	--	--
		04/10/90	8.4	1,190	9.6	.30	--	.07	.07
		08/27/90	12	990	8.0	.50	--	--	.14
3	Big Sioux River near Brookings	04/04/89	312	600	8.4	.83	<.20	--	--
		06/29/89	60	885	9.2	.17	.90	--	.82
		11/03/89	24	1,110	8.5	2.9	<.20	--	.06
4	Big Sioux River at N. Cliff Avenue	04/04/89	550	560	8.4	1.2	.30	--	--
		06/27/89	33	1,130	8.4	1.0	2.5	--	1.6
		11/02/89	19	1,770	8.0	17	.20	--	--
5	James River near Scotland	04/03/89	1,460	520	7.9	.88	.70	0.20	.55
		06/27/89	139	1,230	8.3	<.10	<.20	--	.19
		11/02/89	43	2,000	8.0	<.10	<.20	--	<.05

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90.-Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream flow (ft <sup>3</sup> /s)	Specific conductance (µS/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA- Iowa <sup>1</sup> (µg/L)	Atrazine ELISA- Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
<u>South Dakota-Continued</u>									
6	Vermillion River near Vermillion	04/03/89 06/27/89 11/02/89	111 19 5.0	980 1,340 1,660	8.7 8.2 7.7	<0.10 <.10 <.10	<0.20 <.20 <.20	0.20 -- --	0.17 <.05 <.05
7	Brule Creek near Elk Point	04/03/89 06/27/89 11/02/89	18 7.8 2.8	1,000 950 1,000	8.9 8.4 8.1	.43 .12 <.10	<.20 .30 <.20	-- -- --	-- .36 --
8	Big Sioux River at Akron	04/03/89 06/27/89 11/02/89	2,140 223 107	500 740 1,290	8.5 9.1 9.0	1.8 <.10 2.6	<.20 .90 <.20	.30 -- --	.46 .59 <.05
<u>Nebraska</u>									
9	Bow Creek near St. James	04/05/89 06/26/89 10/18/89	48 449 30.0	696 580 666	8.4 8.4 8.3	2.5 1.3 1.8	<.20 <.20 <.20	-- -- --	-- .20 <.05
10	Elkhorn River near Atkinson	04/05/89 06/26/89 10/23/89	43 22.3 12.2	231 211 254	8.4 8.9 8.4	1.5 1.0 2.0	<.20 .20 1.9	-- -- --	-- -- 1.2
11	Clearwater Creek near Clearwater	04/05/89 06/26/89 10/18/89	30.0 13.4 22.0	316 307 285	8.3 8.2 8.4	.16 .12 <.10	<.20 <.20 <.20	-- -- --	-- .20 <.05

**Table 6.** Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued

Map no (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream flow (ft <sup>3</sup> /s)	Specific conductance (µS/cm)	pH (standard units)	Nitrate plus nitrogen (mg/L)	Atrazine ELISA- Iowa <sup>1</sup> (µg/L)	Atrazine ELISA- Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
Nebraska--Continued									
12	Mira Creek near Northloup	04/11/89 06/25/89 10/20/89	0.70 .71 .23	708 544 585	8.4 7.9 7.7	<0.10 <.10 <.10	<0.20 <.20 .70	<0.30 -- --	0.27 .15 .47
13	Shell Creek near Columbus	04/05/89 06/26/89 10/19/89	18.0 137 14.5	605 300 696	8.4 7.3 8.1	.57 4.5 .25	<.20 >5.0 .50	-- -- --	-- 14 .28
14	Maple Creek near Nickerson	04/05/89 06/26/89 10/19/89	15.0 333 13.0	588 269 556	8.5 7.6 8.1	.65 2.8 .38	<.20 >5.0 <.20	<.20 -- --	.07 8.7 .08
15	Mud Creek at Sweetwater	04/11/89 06/25/89 10/23/89	21.0 27.8 15.0	576 434 584	8.6 8.1 8.2	.18 1.1 <.10	<.20 1.9 <.20	-- -- --	-- 1.7 <.05
16	Big Blue River at Surprise	04/05/89 06/26/89 10/31/89	3.5 71.0 3.0	570 375 475	8.4 7.5 7.5	1.1 1.2 .23	.20 >5.0 1.3	-- -- --	-- 17 .70
17	Wahoo Creek at Itica	04/04/89 06/08/89 10/19/89 03/20/90 06/08/90	40.0 161 26.0 32.0 126	821 248 812 870 548	8.4 7.5 8.1 8.5 7.8	1.1 2.3 1.2 1.5 1.9	<.20 >5.0 <.20 -- --	<.20 -- -- -- --	<.05 52 <.05 .07 9.6

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream- flow (ft <sup>3</sup> /s)	Specific conduct- ance (µS/cm)	pH (standard units)	Nitrite plus nitrate as nitrogen (mg/L)	Atrazine ELISA- Iowa <sup>1</sup> (µg/L)	Atrazine ELISA- Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
<u>Nebraska--Continued</u>									
18	W. Fork Big Blue River near Dorchester	04/04/89 06/27/89 10/31/89	70.0 1,830 56.0	632 130 572	8.1 8.8 7.9	1.7 4.0 .84	<.20 <.20 .50	<.20 -- --	0.16 23 .34
19	Salt Creek at Roca	04/06/89 06/26/89 10/31/89 03/27/90 06/07/90	10.5 53.0 10.5 19.0 59.0	735 497 990 1,232 581	8.8 7.7 7.7 8.8 8.2	<.10 1.8 .60 .20 .20	<.20 >5.0 2.5 -- --	-- -- -- -- --	-- 13 1.7 .12 2.1
20	Muddy Creek near Arapahoe	03/31/89 06/25/89 11/07/89	6.74 4,160 5.58	552 101 595	8.4 7.8 8.0	1.5 .80 2.0	<.20 4.2 <.20	-- -- --	-- 6.6 .11
21	Little Blue River near Fairbury	04/05/89 06/26/89 10/30/89 03/21/90 05/17/90	116 6,590 106 131 258	495 162 438 513 381	8.5 7.5 7.9 8.4 8.0	.94 .60 1.1 1.4 1.7	<.20 >5.0 .40 -- --	-- -- -- -- --	-- 22 .31 .21 15
22	Little Nemaha River at Auburn	04/06/89 06/26/89 10/24/89	57.0 696 53.0	586 217 612	8.4 7.6 8.2	.36 1.2 .12	<.20 >5.0 .20	-- -- --	-- 10 .24

**Table 6.** Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued

Map no (figs. 2-5)	Site name	Date of collection (month/day/year)	Stream flow (ft <sup>3</sup> /s)	Specific conductance ( $\mu$ S/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA-Iowa <sup>1</sup> ( $\mu$ g/L)	Atrazine ELISA-Kans. <sup>2</sup> ( $\mu$ g/L)	Atrazine GC/MS <sup>3</sup> ( $\mu$ g/L)
<b>Nebraska--Continued</b>									
23	Big Blue River at Barneston	04/06/89 06/27/89	226 3,380	696 310	8.9 7.9	2.0 2.1	2.1 >5.0	1.7 .90	0.91 16
		10/23/89	579	684	8.9	.55	--	--	.51
		03/21/90	330	708	8.7	2.1	--	--	.51
		05/17/90	801	378	7.9	2.0	--	--	14
24	Big Nemaha River at Fall City	04/06/89 06/26/89 10/24/89 03/27/90 05/17/90	56.0 656 56.0 123 1,447	668 291 746 726 356	8.4 7.9 8.2 8.5 7.9	<.10 1.4 <.10 .30 1.8	<.20 >5.0 .50 -- --	-- -- -- -- --	-- 20 .54 .81 5.6
<b>Kansas</b>									
25	Turkey Creek near Seneca	03/21/89 05/22/89 10/12/89	2.64 13.0 6.40	712 630 724	8.6 7.9 8.3	<.10 <.10 <.10	<.20 .30 <.20	<.20 .40 --	<.05 .22 .23
26	Black Vermillion River near Frankfort	03/16/89 06/27/89 10/04/89 03/22/90 05/16/90	12.2 588 7.30 24.6 2,560	540 205 572 606 268	8.4 7.8 8.4 8.2 7.6	<.10 1.4 <.10 .48 1.6	<.20 >5.0 .30 -- --	-- -- -- -- --	-- 16 .35 .47 6.1

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/day/year)	Stream flow (ft <sup>3</sup> /s)	Specific conductance (µS/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA-Iowa <sup>1</sup> (µg/L)	Atrazine ELISA-Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
<b>Kansas--Continued</b>									
27	Delaware River near Muscotah	03/15/89	9.87	569	8.8	<0.10	<0.20	--	--
		05/22/89	54.3	664	8.0	.19	>5.0	6.0	10
		10/11/89	3.00	629	8.2	<.10	.90	--	.82
28	Salt Creek near Ada	03/21/89	4.58	2,960	8.2	<.10	<.20	<.20	<.05
		05/22/89	550	307	8.1	.97	.40	.40	.27
		10/12/89	2.10	3,280	8.0	<.10	.40	--	.49
29	Mill Creek near Paxico	03/14/89	4.29	789	8.3	<.10	<.20	--	--
		09/05/89	330	230	7.8	.50	.30	--	--
		10/06/89	22.5	661	8.2	<.10	<.20	--	<.05
30	Kansas River at Topeka	03/14/89	842	1,050	8.8	<.10	.80	--	--
		08/30/89	3,690	440	8.1	.87	3.8	--	--
		10/06/89	2,360	546	8.4	.76	2.3	--	1.8
<b>Minnesota</b>									
31	Knife River near Mora	04/21/89	114	116	8.1	.17	<.20	--	--
		07/19/89	9.20	242	7.8	.17	<.20	--	.09
		10/31/89	1.50	265	8.1	.37	<.20	--	<.05
		05/30/90	61.0	158	8.2	.10	--	--	<.05

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream flow (ft <sup>3</sup> /s)	Specific conductance (µS/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA- Iowa <sup>1</sup> (µg/L)	Atrazine ELISA- Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
<b>Minnesota--Continued</b>									
32	Sauk River near St. Cloud	03/23/89 05/25/89 11/08/89	300 700 --	720 581 --	7.9 8.2 --	2.0 .51 .96	0.30 .60 .40	-- -- --	-- -- 0.17
33	Pomme de Terre River at Appleton	05/04/89 07/18/89 10/25/89	197 66.0 22.0	950 925 680	8.3 8.4 8.2	<.10 .21 <.10	<.20 <.20 <.20	0.40 -- --	.06 .32 <.05
34	Rum River near St. Francis	04/19/89 07/19/89 10/31/89	790 165 150	221 357 346	8.6 8.4 8.1	.34 <.10 .27	<.20 <.20 <.20	-- -- --	-- <.05 <.05
<b>Wisconsin</b>									
35	St. Croix River at St. Croix Falls	04/24/90 05/31/89 10/11/89 04/23/90 07/30/90	7,590 9,910 1,710 5,450 5,600	121 117 208 168 174	7.2 -- 8.2 7.8 7.8	.14 .13 <.10 <.10 .30	<.20 <.20 <.20 -- --	-- -- -- -- --	-- .14 <.05 <.05 .16
<b>Minnesota</b>									
36	Crow River at Rockford	04/19/89 05/25/89 10/26/89	465 390 47.0	544 672 970	9.4 8.3 8.1	<.10 .74 <.10	.20 3.3 <.20	.30 3.4 --	.33 2.8 <.05

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/day/year)	Stream flow (ft <sup>3</sup> /s)	Specific conductance ( $\mu$ S/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA-Iowa <sup>1</sup> ( $\mu$ g/L)	Atrazine ELISA-Kans. <sup>2</sup> ( $\mu$ g/L)	Atrazine GC/MS <sup>3</sup> ( $\mu$ g/L)
Minnesota -- Continued									
37	Minnesota River near Jordan	03/22/89	1,892	590	8.2	2.8	0.40	0.20	0.17
		07/19/89	508	868	8.1	<.10	<.20	--	.29
		10/29/89	230	995	8.2	<.10	5.0	--	3.1
38	Redwood River near Redwood Falls	05/04/89	62.0	1,670	8.3	<.10	<.20	.20	.08
		06/27/89	93.0	1,200	8.1	3.7	.30	--	.42
		10/24/89	8.70	1,440	8.3	<.10	.30	--	.11
39	Cottonwood River near New Ulm	03/21/89	126	450	7.9	3.7	<.20	--	--
		07/18/89	58.0	642	8.4	<.10	<.20	--	.15
		11/08/89	17.0	1,000	8.2	.11	.20	--	<.05
		05/01/90	71.0	962	8.6	<.10	--	--	.33
		06/05/90	316	1,223	8.4	8.5	--	--	.26
40	Le Sueur River near Rapidan	04/20/89	101	657	8.7	2.6	<.20	--	--
		07/18/89	36.0	631	8.4	<.10	<.20	--	.27
		11/07/89	14.0	775	8.3	.18	<.20	--	<.05
41	Straight River near Faribault	04/05/89	204	660	8.4	6.0	.20	.30	.29
		07/19/89	30.0	678	8.8	<.10	<.20	--	.32
		11/08/89	49.0	721	8.4	1.1	--	--	<.05
42	N. Fork Whitewater River near Elba	04/05/89	38.0	546	8.2	1.6	<.20	--	--
		07/19/89	342	531	8.4	1.9	--	--	--
		11/08/89	22.0	521	8.1	2.6	--	--	.05

**Table 6.** Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued

Map no (figs. 2-5)	Site name	Date of collection (month/day/year)	Stream flow (ft <sup>3</sup> /s)	Specific conductance ( $\mu$ S/cm)	pH (standard units)	Nitrite plus nitrate as nitrogen (mg/L)	Atrazine ELISA Iowa <sup>1</sup> ( $\mu$ g/L)	Atrazine ELISA Kans. <sup>2</sup> ( $\mu$ g/L)	Atrazine GC/MS <sup>3</sup> ( $\mu$ g/L)
Minnesota -- Continued									
43	Rock River at Luverne	05/03/89 06/27/89	87.0 200	790 510	8.1 7.5	1.8 4.0	<0.20 3.3	-- --	-- 2.9
		10/23/89	12.0	440	8.7	1.1	<.20	--	.08
		05/08/90	36.3	730	8.4	.60	--	--	.08
		06/19/90	212	390	8.2	2.1	--	--	3.5
44	Des Moines River at Jackson	05/03/89 06/27/89	143 40.0	1,010 1,110	8.9 8.7	<.10 <.10	<.20 1.4	-- --	-- 1.0
		10/23/89	1.50	835	8.3	<.10	.70	--	.61
		05/08/90	46.0	1,290	8.8	<.10	--	--	.11
		06/19/90	424	925	8.5	4.9	--	--	1.4
45	Rush Creek near Rushford	04/05/89 05/25/89 11/07/89	48.0 55.0 39.0	490 445 479	8.2 7.9 8.2	2.1 1.8 1.8	<.20 1.7 --	-- -- --	2.1 .06
Iowa									
46	Ocheyedan River near Spencer	03/28/89 05/24/89 10/12/89	89.2 -- 4.55	460 -- 640	7.6 -- 8.3	2.1 4.0 <.10	<.20 >5.0 <.20	0.20 8.0 --	.13 12 <.05
47	Des Moines River at Estherville	03/28/89 10/12/89	818 1.30	440 2,950	7.4 8.0	2.0 .40	<.20 .20	.20 --	.14 .16

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/day/year)	Stream-flow (ft <sup>3</sup> /s)	Specific conductance ( $\mu\text{S}/\text{cm}$ )	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA-Iowa <sup>1</sup> ( $\mu\text{g}/\text{L}$ )	Atrazine ELISA-Kans. <sup>2</sup> ( $\mu\text{g}/\text{L}$ )	Atrazine GC/MS <sup>3</sup> ( $\mu\text{g}/\text{L}$ )
<u>Iowa -- Continued</u>									
48	Winnebago River at Mason City	03/20/89 10/11/89	163 10.7	492 740	7.6 7.6	6.3 <.10	0.20 <.20	0.40 --	0.51 <.05
49	Little Cedar River near Ionia	03/21/89 10/16/89	66.5 9.70	350 430	7.7 7.7	2.4 <.10	.70 .20	-- --	-- .14
50	Turkey River at Spillville	03/31/89 10/04/89 03/22/90 06/14/90	50.0 10.4 99.8 212	494 550 565 430	8.3 7.8 8.4 7.9	3.1 1.7 8.3 <.10	<.20 <.20 -- --	-- -- -- --	<.05 .43 8.2
51	Upper Iowa River near Dorchester	03/31/89 05/31/89 10/05/89	348 300 133	453 430 430	8.4 7.6 8.3	2.3 <.10 <.10	<.20 -- <.20	-- -- --	.33 10 .15
52	Little Sioux River at Correctionville	03/23/89 05/25/89 10/02/89 03/16/90 06/14/90	509 1,620 42.0 264 788	540 308 970 575 622	7.9 6.9 8.7 8.5 7.1	3.1 .10 <.10 1.2 6.5	<.20 >5.0 <.20 -- --	-- -- -- -- --	-- 9.9 .13 .19 7.4
53	North Raccoon River near Sac City	03/24/89 05/24/89 10/12/89	163 -- 12.6	630 -- 1,120	7.7 -- 8.8	4.6 7.0 <.10	4.2 .80 <.20	-- -- --	-- 1.1 <.05

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream flow (ft <sup>3</sup> /s)	Specific conductance (µS/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA- Iowa <sup>1</sup> (µg/L)	Atrazine ELISA- Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
Iowa -- Continued									
54	Des Moines River at Fort Dodge	03/27/89 05/24/89	1,170 --	430 --	7.6 --	2.4 7.8	<0.20 .20	--	--
		10/13/89	57.1	690	7.6	<.10	<.20	2.8	2.3
		03/16/90	746	625	7.8	8.8	--	--	<.05
		06/15/90	3,880	505	7.7	11	--	--	.12
									1.8
55	Iowa River near Rowan	03/23/89 10/11/89	122 8.22	530 930	7.3 7.2	3.2 <.10	<.20 <.20	--	--
									<.05
56	Cedar River at Cedar Falls	03/20/89 11/09/89	2,870 402	348 550	8.2 8.9	3.4 1.4	.90 <.20	--	--
									.15
57	Black Hawk Creek at Hudson	03/22/89 10/18/89	102 2.93	460 720	7.4 7.3	3.9 .10	.30 .40	--	--
									<.05
58	Wapsipinicon River at Independence	03/20/89 11/09/89 03/22/90 05/09/90 06/22/90	854 174 988 1,270 2,210	240 555 488 358 578	8.1 8.5 8.3 8.5 8.0	3.0 9.7 13 8.1 <.10	2.7 <.20 -- -- --	1.5 -- -- -- --	1.7 .39 .63 22 3.9
59	Maple River at Mapleton	03/23/89 05/24/89	128 852	700 303	8.0 6.6	6.40 .10	-- >5.0	--	--
									34

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream- flow (ft <sup>3</sup> /s)	Specific conduct- ance (µS/cm)	pH (standard units)	Nitrite plus nitrate as nitrogen (mg/L)	Atrazine ELISA- Iowa <sup>1</sup> (µg/L)	Atrazine ELISA- Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
Iowa -- Continued									
59	Maple River at Mapleton--Continued	10/02/89	44.4	680	8.5	3.0	0.80	--	0.59
		03/16/90	150	680	8.4	6.8	--	--	.40
		06/13/90	4,960	242	6.7	3.6	--	--	13
60	Boyer River at Logan	03/23/89	158	750	7.8	5.9	<.20	<0.20	.14
		05/24/89	2,070	323	6.5	.10	>5.0	10	45
		10/02/89	50.3	630	8.5	.12	<.20	--	.23
		03/16/90	461	540	8.3	4.1	--	--	.55
		06/13/90	15,700	220	6.6	2.1	--	--	8.0
61	Raccoon River at Van Meter	03/21/89	516	512	8.1	3.3	<.20	--	--
		05/25/89	2,700	452	8.6	2.0	.70	--	1.1
		11/08/89	201	605	8.6	3.2	<.20	--	.32
		05/03/90	534	584	8.7	2.2	--	--	.12
		05/10/90	3,420	500	8.3	8.2	--	--	2.3
62	Indian Creek near Mingo	03/20/89	11.0	765	8.2	1.0	.40	--	--
		05/24/89	1,010	205	8.1	2.8	3.8	--	5.0
		10/10/89	1.30	580	8.1	<.10	<.20	--	<.05
63	Iowa River near Marengo	03/24/89	693	432	7.7	2.5	.40	.40	.42
		05/25/89	516	405	7.4	<.10	>5.0	9.4	15
		11/07/89	134	540	9.5	<.10	<.20	--	.15

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream- flow (ft <sup>3</sup> /s)	Specific conduct- ance (µS/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA- Iowa <sup>1</sup> (µg/L)	Atrazine ELISA- Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
Iowa -- Continued									
64	Old Mans Creek near Iowa City	03/24/89 05/25/89 11/07/89	3.40 557 3.80	476 250 565	7.4 7.2 9.4	1.2 5.3 <.10	>5.0 .30	-- --	-- 72 .69
65	N. Fork Maquoketa River at Fulton	03/27/89 10/05/89	174 107	580 570	7.6 8.3	3.4 3.1	.20 <.20	-- --	-- <.05
66	S. Skunk River near Oskaloosa	03/22/89 05/24/89 10/16/89	160 2,090 55.6	508 258 680	8.1 7.2 8.2	1.6 6.3 .44	0.30 >5.0 <.20	-- -- --	-- 48 .19
67	N. Skunk River near Sigourney	03/22/89 05/24/89 10/02/89 03/15/90 05/21/90	65.0 638 55.1 3,100 776	400 362 475 295 490	7.9 7.5 8.2 7.3 8.2	1.5 3.4 2.0 8.1 13	1.0 >5.0 .40 -- --	-- -- -- -- --	-- 42 .26 .89 1.1
68	Nishnabotna River above Hamburg	03/24/89 06/05/89 10/10/89	345 738 673	560 304 570	7.7 7.2 7.7	2.5 3.9 3.2	.30 >5.0 1.0	-- -- --	-- 37 1.0
69	Chariton River near Chariton	03/20/89 05/30/89 10/02/89	2.70 228 .10	330 151 262	8.1 8.0 7.9	.64 1.9 <.10	<.20 >5.0 1.7	0.30 7.2 --	.21 13 1.3

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream flow (ft <sup>3</sup> /s)	Specific conductance (µS/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA- Iowa <sup>1</sup> (µg/L)	Atrazine ELISA- Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
<u>Iowa -- Continued</u>									
70	S. Fork Chariton River near Promise City	03/20/89	1.90	315	8.8	0.20	0.50	--	--
		05/30/89	26.0	238	7.0	1.2	>5.0	--	16
		10/03/89	.40	438	7.9	<.10	1.0	--	.96
71	Skunk River at Augusta	03/22/89	373	360	7.9	2.0	.90	0.40	.69
		05/26/89	3,200	240	7.6	7.9	>5.0	10	30
		11/13/89	165	630	8.8	<.10	.40	--	.36
		03/08/90	470	615	8.4	3.9	--	--	.36
		06/21/90	26,800	205	7.5	4.3	--	--	6.5
<u>Missouri</u>									
72	Nodaway River near Graham	03/15/89	331	265	7.8	.22	1.2	1.0	1.0
		06/23/89	4,490	125	7.2	1.6	>5.0	--	7.8
		10/03/89	414	395	7.9	.86	.20	--	.21
		04/10/90	592	430	8.2	3.7	--	--	.14
		05/22/90	612	612	8.4	4.7	--	--	1.5
73	South Fabius River near Taylor	04/04/89	95.8	479	8.1	<.10	<.20	--	--
		10/05/89	3.70	370	8.1	<.10	.50	--	.57
74	Grand River near Sumner	03/08/89	325	430	7.4	2.4	.20	--	--
		05/19/89	2,420	210	7.4	1.5	>5.0	--	--
		10/04/89	241	390	7.7	<.10	.50	--	.35

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90-Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream flow (ft <sup>3</sup> /s)	Specific conductance (µS/cm)	pH (standard units)	Nitrite plus nitrate as nitrogen (mg/L)	Atrazine ELISA- Iowa <sup>1</sup> (µg/L)	Atrazine ELISA- Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
<b>Missouri-Continued</b>									
75	Middle Fork Sal River at Paris	04/03/89	17.5	842	8.0	<.10	0.40	<.20	0.53
		05/30/89	3,670	113	7.1	.66	2.3	1.0	3.0
		10/06/89	3.10	310	7.9	.36	.80	--	.65
76	Blackwater River at Blue Lick	03/16/89	28.0	685	7.9	1.3	.30	--	--
		05/20/89	3,040	165	7.4	.93	>5.0	--	11
		10/04/89	13.7	580	7.6	<.10	.90	--	.50
77	Missouri River at Hermann	03/12/89	52,300	490	7.9	<.10	<.20	<.20	.18
		06/07/89	62,200	--	--	<.10	--	--	1.5
		10/11/89	45,300	665	8.2	<.10	<.20	--	.26
78	Mississippi River at Grafton	03/10/89	30,000	436	9.0	<.10	.30	.20	.24
		06/05/89	81,900	--	--	<.10	--	--	2.0
		11/13/89	57,000	430	8.5	<.10	<.20	--	.21
<b>Wisconsin</b>									
79	Prairie River near Merrill	04/21/89	213	104	8.1	.21	<.20	--	--
		05/05/89	178	119	8.2	.11	<.20	--	<.05
		10/23/89	72.0	204	8.6	<.10	<.20	--	<.05
80	Trempealeau River at Dodge	03/14/89	2,320	250	7.4	.99	--	--	--
		05/30/89	810	242	--	1.9	>5.0	--	26
		10/17/89	346	300	8.0	1.3	<.20	--	.15

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90-Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/day/year)	Stream-flow (ft <sup>3</sup> /s)	Specific conductance (µS/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA-Iowa <sup>1</sup> (µg/L)	Atrazine ELISA-Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
Wisconsin - Continued									
81	Black River near Galesville	03/15/89 05/30/89 10/18/89	2,680 4,100 539	166 92 155	7.2 — 7.6	1.0 .39 .47	<.20 2.4 <.20	— — —	— 3.2 <.05
82	Wisconsin River at Muscada	03/31/89 06/02/89 10/05/89 03/20/90 06/22/90	18,800 19,100 2,420 20,700 15,400	230 180 290 280 195	8.2 7.6 8.5 7.7 7.9	.82 .57 .32 .82 .60	— .30 <.20 — —	— — — — —	— .65 .21 .31 .77
83	Grant River near Burton	03/23/89 05/31/89 11/02/89	285 85.0 72.0	558 586 630	8.2 8.3 8.4	3.4 1.9 2.5	<.20 1.1 <.20	0.30 — —	.32 1.7 .15
84	Pecatonic River at Martintown	04/11/89 07/19/89 10/31/89	382 298 249	600 565 610	8.3 8.3 8.3	3.1 1.8 2.1	.20 <.20 <.20	— — —	— .11 .14
85	Rock River at Afton	04/12/89 07/19/89 10/31/89 04/12/90 07/06/90	3,420 747 852 3,350 2,540	500 665 690 515 600	9.0 8.6 8.8 9.1 8.6	1.5 1.6 1.2 1.0 .50	<.20 .40 .30 — —	.20 — — — —	.34 .51 .15 .39 .72
86	Fox River at Wakesha	04/20/89 07/10/89 11/03/89	160 77.0 43.0	1,070 780 1,070	8.2 8.1 8.1	.92 1.6 3.4	<.20 .50 <.20	— — —	— — .12

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90-Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream flow (ft <sup>3</sup> /s)	Specific conductance (µS/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Nitrite plus nitrate as nitrogen (mg/L)	Atrazine ELISA- Iowa <sup>1</sup> (µg/L)	Atrazine ELISA- Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
<b>Wisconsin--Continued</b>										
87	Root River at Racine	03/22/89 07/10/89 11/03/89 03/27/90 07/19/90	116 10.0 41.0 188 31.0	985 990 1,020 808 760	7.8 8.4 8.1 8.1 8.6	3.1 <.10 2.2 3.4 2.2	<.20 1.2 <.20 -- --	<.20 -- -- -- --	0.08 .88 .11 .17 .44	
<b>Illinois</b>										
88	Des Plains River at Russell	04/05/89 05/25/89 11/01/89	100 27.4 5.50	990 452 1,260	7.7 7.7 7.7	7.4 .24 1.4	<.2 .40 .20	-- -- --	-- .23 .15	
89	Nippersink Creek at Spring Cove	04/05/89 05/25/89 11/01/89	95.0 95.5 43.0	735 659 780	8.6 8.0 8.0	3.1 .96 1.7	.30 1.3 <.20	<.20 -- --	.37 .90 .38	
90	S. Br. Kishwaukee at Fairdale	03/22/89 05/25/89 10/24/89 04/12/90 05/14/90	145 76.7 76.0 294 1,280	770 696 857 755 --	7.8 7.6 5.6 8.6 7.8	.30 >5.0 .40 11 19	-- -- -- -- --	-- -- -- -- --	-- .11 .24 .25 2.1	
91	Kiswaukee River near Perryville	03/22/89 05/25/89 10/24/89	445 225 269	720 722 796	7.6 7.8 --	5.3 1.2 3.2	<.20 4.0 <.20	<.20 6.0 --	.12 4.9 .15	

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90-Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/day/year)	Stream-flow (ft <sup>3</sup> /s)	Specific conductance ( $\mu$ S/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA-Iowa <sub>1</sub> ( $\mu$ g/L)	Atrazine ELISA-Kans. <sub>2</sub> ( $\mu$ g/L)	Atrazine GC/MS <sub>3</sub> ( $\mu$ g/L)
Illinois -- Continued									
92	Elkhorn Creek near Penrose	03/22/89 05/25/89 10/24/89	48.0 40.4 30.0	671 643 704	7.3 7.7 --	7.9 4.9 6.6	<0.20 2.0 .30	-- -- --	-- 1.9 .15
93	Illinois River near Marseilles	03/21/89 05/25/89 10/31/89	9,940 6,390 3,200	1,025 850 758	7.9 7.7 8.9	5.7 8.9 4.0	.30 2.1 .40	-- -- --	-- -- 2.5
94	Fox River near Dayton	03/21/89 05/25/89 10/31/89	2,100 638 698	768 803 781	7.8 8.4 8.6	3.8 .51 <.10	.30 .50 .24	<0.20 1.2 --	.20 .80 .26
95	Dupage River near Shorwood	04/06/89 05/25/89 11/02/89 04/04/90 06/22/90	248 269 110 334 --	1,555 1,290 1,770 1,360 --	9.0 7.5 8.7 8.6 --	7.8 7.2 10 6.7 6.3	<.20 .60 <.20 -- --	<.20 1.0 -- -- --	<.05 .77 <.05 .07 .37
96	Iroquois River near Chebanse	04/03/89 05/25/89 11/03/89 04/04/90 05/17/90	1,600 1,530 477 2,410 5,560	668 721 770 693 685	8.4 7.7 8.2 8.2 8.2	11 18 5.0 9.0 15	.30 3.5 .20 -- --	.40 5.0 -- -- --	.47 2.8 .16 .19 4.2
97	Edwards River near New Boston	03/22/89 10/23/89	57.0 55.0	567 680	7.3 --	2.3 4.3	<.20 .20	<.20 --	.15 .34

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream- flow (ft <sup>3</sup> /s)	Specific conduct- ance (µS/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA- Iowa <sup>1</sup> (µg/L)	Atrazine ELISA- Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
Illinois--Continued									
98	Spoon River at London Mills	04/21/89 06/02/89 10/17/89 04/10/90 05/14/90	58.0 1,620 81.0 490 3,500	820 361 700 750 666	8.6 7.5 8.3 8.1 8.1	<.10 5.1 4.2 12 14	.20 >5.0 .30 -- --	.20 -- -- -- --	0.23 108 .24 .18 .20
99	La Moine River at Colmar	04/20/89 06/03/89 10/17/89	8.90 12.1 2.00	700 648 520	8.3 7.4 8.2	<.10 .10 .10	.30 >5.0 .40	-- -- --	-- 11 .29
100	Sangamon River at Oaksford	04/20/89 05/23/89 10/19/89	2,340 2,620 346	669 664 870	8.2 8.4 8.2	7.8 9.2 1.9	<.20 2.5 .60	-- -- --	-- 2.7 .25
101	Lake Fork near Cornland	03/21/89 05/22/89 10/16/89	72.0 120 32.0	622 727 742	7.6 8.1 8.3	6.0 13 2.7	<.20 .40 <.20	-- -- --	-- .64 <.05
102	Sangamon River at Riverton	03/21/89 05/22/89 10/12/89 03/26/90 05/14/90	806 1,450 77.0 1,010 5,730	755 675 1,716 682 575	7.7 8.1 8.4 7.9 8.0	5.9 12 1.8 .60 7.6	.80 -- -- -- --	-- -- -- -- --	2.6 -- .33 .55 3.8
103	Illinois River at Hardin	03/10/89 06/04/89 10/16/89	14,500 27,500 4,760	-- -- 675	-- -- 8.2	<.10 <.10 2.9	<.20 -- .70	<.20 -- --	.24 3.1 .25

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream flow (ft <sup>3</sup> /s)	Specific conductance (µS/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA-Iowa (µg/L)	Atrazine ELISA-Kans. (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
Illinois -- Continued									
104	Macoupin Creek near Kane	03/21/89	2,890	299	7.5	2.4	0.80	0.70	0.89
		05/23/89	128	537	7.4	2.5	>5.0	10	20
		10/16/89	2,70	519	7.4	.10	1.1	--	.62
		03/28/90	262	758	8.5	.82	--	--	.79
		05/14/90	4,020	329	8.2	3.1	--	--	16
105	Kaskaskia River near Cowden	03/20/89	1,460	447	7.8	1.7	.20	--	--
		05/22/89	1,300	527	6.9	9.2	2.0	--	2.7
		10/26/89	7.10	555	7.0	2.3	2.2	--	1.3
		03/27/90	214	553	8.3	2.7	--	--	1.3
		05/14/90	2,420	244	8.5	4.2	--	--	33
106	Kaskaskia River at Vandalia	03/20/89	4,780	253	7.5	1.2	.50	--	--
		05/23/89	1,600	532	7.4	7.4	3.5	--	4.9
		10/11/89	42.1	612	7.5	.10	.80	--	.76
107	Embarra River at Ste. Marie	04/13/89	2,320	529	7.4	12	.60	--	--
		05/22/89	1,900	540	7.5	11	>5.0	--	8.6
		10/18/89	231	663	8.2	2.9	.50	--	.33
108	Shoal Creek near Breeze	03/20/89	2,040	208	7.2	1.1	.40	.30	.61
		05/23/89	152	429	7.0	1.4	>5.0	8.4	15
		10/11/89	13.8	436	7.0	.10	2.7	--	2.1
		03/28/90	196	700	8.2	.53	--	--	1.0
		05/04/90	607	458	7.5	.70	--	--	26

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream- flow (ft <sup>3</sup> /s)	Specific conduct- ance (µS/cm)	pH (standard units)	Nitrite plus nitrate as nitrogen (mg/L)	Atrazine ELISA- Iowa <sup>1</sup> (µg/L)	Atrazine ELISA- Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
Illinois -- Continued									
109	Silver Creek near Freeburg	03/20/89 05/23/89 10/11/89	490 131 2,00	361 550 595	7.1 7.8 6.7	1.6 1.8 .10	0.30 >5.0 .50	-- -- --	19 .52
110	Bonpas Creek at Browns	04/12/89 05/22/89 11/27/89 04/17/90 07/10/90	104 77.0 12.0 112 2,30	472 493 824 631 26	6.9 6.8 7.8 7.4 10.1	2.6 5.0 2.2 1.7 1.6	.50 >5.0 .50 -- --	-- -- -- -- --	.41 1.8 9.2
Indiana									
111	Wabash River near New Harmony	03/29/89 06/22/89 10/31/89	49,300 29,700 12,700	450 527 612	7.6 7.8 8.2	3.8 4.3 2.6	<.20 4.2 1.1	-- -- --	6.7 .68
Illinois									
112	Little Wabash River at Carmi	04/12/89 05/22/89 11/27/89 04/17/90 07/10/90	18,300 1,110 125 6,580 440	160 218 725 381 540	6.6 7.0 7.7 7.2 7.5	.49 1.5 .88 1.0 1.8	.60 >5.0 2.0 -- --	-- -- -- -- --	34 .69 3.8 9.0
113	Big Muddy River near Murphysboro	04/11/89 05/24/89 10/16/89	12,800 1,210 76.2	310 995 1,160	6.7 7.0 7.2	.22 1.0 .63	.60 3.9 .70	-- -- --	.51 .57

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream flow (ft <sup>3</sup> /s)	Specific conductance (µS/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA- Iowa <sup>1</sup> (µg/L)	Atrazine ELISA- Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
<b>Missouri</b>									
114	Mississippi River at Thebes	03/15/89	173,000	523	7.9	<.10	0.20	0.20	0.27
		06/10/89	184,000	--	--	<.10	--	--	1.8
		11/08/89	89,800	520	8.0	.73	.50	--	.37
<b>Illinois</b>									
115	Ohio River near Grand Chain	03/16/89	721,000	232	7.3	<.10	<.20	<.20	<.05
		06/11/89	309,000	--	--	<.10	--	--	3.1
		10/11/89	275,000	209	7.6	.10	<.20	--	<.05
<b>Indiana</b>									
116	Kankakee River at Shelby	03/22/89	2,020	606	8.2	3.5	<.20	<.20	.05
		05/31/89	2,420	608	7.7	2.0	2.7	4.4	2.8
		10/16/89	871	654	7.9	.59	<.20	--	<.05
117	Tippecanoe River near Ora	03/22/89	807	631	8.2	4.8	<.20	--	--
		05/31/89	650	612	7.9	3.3	1.0	--	1.2
		10/16/89	292	638	8.3	.43	.30	--	.23
118	Iroquois River near Foresman	03/22/89	531	746	8.3	13	.20	--	--
		05/31/89	1,640	542	7.3	15	>5.0	--	5.9
		10/16/89	162	762	7.9	2.0	<.20	--	.11
119	Eel River near Logansport	03/22/89	1,190	558	8.3	7.7	<.20	.20	.26
		05/22/89	604	690	8.2	2.7	>5.0	--	8.0
		10/16/89	170	716	8.4	1.5	<.20	--	.12
		03/26/90	660	658	7.8	4.2	--	--	.19
		05/14/90	1,550	619	7.5	7.6	--	--	9.6

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/day/year)	Stream-flow (ft <sup>3</sup> /s)	Specific conductance ( $\mu$ S/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA-Iowa <sup>1</sup> ( $\mu$ g/L)	Atrazine ELISA-Kans. <sup>2</sup> ( $\mu$ g/L)	Atrazine GC/MS <sup>3</sup> ( $\mu$ g/L)
Indiana -- Continued									
120	Wabash River at Linn Grove	03/23/89 05/26/89 10/17/89	350 2,500 20.0	682 229 1,140	8.1 7.3 7.9	13 3.3 .35	0.30 >5.0 .50	--	--
121	Wildcat Creek near Lafayette	03/23/89 05/27/89 10/16/89 03/26/90 05/14/90	648 11,500 234 54.0 1,220	739 288 767 721 564	8.4 7.6 8.3 7.8 7.7	8.1 5.5 1.9 5.6 11	<.20 >5.0 .40 -- --	0.20 10 -- -- --	.22 27 .26 .22 14
122	Wildcat Creek near Jerome	03/23/89 05/22/89 10/17/89 03/26/90 05/14/90	95.0 91.0 33.0 90.0 560	695 685 718 641 588	8.3 8.1 7.7 8.0 7.4	12 13 2.7 3.1 15	0.40 1.0 .40 -- --	--	--
123	Wabash River at Covington	03/24/89 05/22/89 10/16/89	5,930 8,080 2,700	663 621 678	8.5 8.0 8.3	6.5 8.9 2.6	>5.0 -- 1.5	.20 5.2 --	.18 8.8 .68
124	White River near Nora	03/27/89 05/26/89 11/01/89 03/26/90 05/14/90	829 7,600 544 959 6,430	727 362 793 693 444	8.4 7.6 8.1 7.9 7.7	5.1 4.3 2.4 3.6 6.6	<.20 >5.0 .30 -- --	.30 8.0 -- -- --	.18 9.8 .29 .32 17

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream- flow (ft <sup>3</sup> /s)	Specific conduct- ance (µS/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA- Iowa <sup>1</sup> (µg/L)	Atrazine ELISA- Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
Indiana-Continued									
125	Big Blue River at Carthage	03/27/89 05/23/89 10/17/89	136 1,680 93.0	685 335 727	8.3 7.6 8.0	5.2 4.4 3.7	<0.20 >5.0 <.20	-- -- --	-- 13 .12
126	Whitewater River near Alpine	03/23/89 05/23/89 10/17/89 03/27/90 05/15/90	613 4,440 340 570 1,880	625 377 682 646 555	8.2 7.2 8.0 8.0 8.1	6.7 3.8 3.2 4.3 6.2	<.20 >5.0 .20 -- --	-- -- -- -- --	-- 20 .25 .14 .82
127	Big Walnut Creek near Reelsville	03/24/89 05/23/89 10/31/89	388 420 240	563 488 580	8.3 8.2 8.5	5.5 3.0 1.9	.20 4.0 .50	0.20 5.6 --	.25 6.6 .41
128	White River near Centerton	03/28/89 05/23/89 10/30/89	1,870 8,900 1,490	807 435 821	7.7 7.6 7.8	5.2 3.0 3.5	<.20 4.4 .60	-- -- --	-- 8.7 .36
129	Sugar Creek near Edinburgh	03/24/89 05/25/89 10/30/89 03/27/90 05/15/90	774 1,620 126 295 2,191	617 500 711 645 463	8.2 7.8 8.2 8.3 7.9	8.2 8.7 1.1 6.0 7.1	<.20 >5.0 <.20 -- --	-- -- -- -- --	-- 15 <.05 .14 .20
130	Flatrock River at Columbus	03/24/89 05/25/89 10/30/89	1,070 1,820 162	578 471 635	8.1 7.9 8.0	9.7 9.1 2.9	<.20 >5.0 .20	-- -- --	-- 15 .17

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream-flow (ft <sup>3</sup> /s)	Specific conductance (µS/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA- Iowa <sup>1</sup> (µg/L)	Atrazine ELISA- Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
Indiana--Continued									
131	Busseron Creek near Carlisle	03/29/89 06/22/89 10/31/89	179 22.0 74.0	718 1,320 1,100	7.5 7.6 7.7	.88 .59 .30	0.20 2.5 .30	-- -- --	3.5 .22
132	E. Fork White River near Bedford	03/28/89 05/25/89 10/30/89 03/27/90 05/15/90	10,800 9,820 1,140 4,550 11,590	419 264 605 512 373	7.7 7.5 8.1 8.0 7.4	3.6 2.2 1.8 3.0 3.5	<.20 >5.0 .50 -- --	-- -- -- -- --	20 .28 .22 11
133	Muscatatuck River near Deputy	03/28/89 05/25/89 10/30/89	175 303 43.0	377 238 406	8.0 7.7 7.8	.84 .28 .50	<.20 2.1 4.6	<0.20 -- --	.06 2.7 .37
134	Blue River at Fredricksburg	03/28/89 05/25/89 10/30/89 03/27/90 05/15/90	313 185 51 231 575	374 370 438 348 323	8.1 7.9 8.0 8.2 7.6	3.5 2.4 1.9 2.0 2.3	<.20 3.6 .50 -- --	-- -- -- -- --	5.1 .30 .16 5.1
Ohio									
135	Tiffin River at Stryker	03/14/89 06/02/89 10/25/89 03/21/90 05/14/90	217 4,570 100 780 966	600 300 745 565 555	8.1 7.5 7.3 8.4 7.2	2.5 6.3 1.3 4.0 13	<.20 >5.0 .50 -- --	.30 10 -- -- --	.38 28 .66 .45 35

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/day/year)	Stream flow ( $\text{ft}^3/\text{s}$ )	Specific conductance ( $\mu\text{S}/\text{cm}$ )	pH (standard units)	Nitrate plus nitrate as standard nitrogen (mg/L)	Atrazine ELISA Iowa 1 ( $\mu\text{g}/\text{L}$ )	Atrazine ELISA Kans. 2 ( $\mu\text{g}/\text{L}$ )	Atrazine GC/MS <sup>3</sup> ( $\mu\text{g}/\text{L}$ )
Ohio--Continued									
136	Maumee River at Waterville	03/29/89 05/27/89 10/25/89	5,060 34,300 1,430	600 370 560	8.2 8.0 9.2	7.9 6.0 .49	0.30 >5.0 1.0	-- -- --	7.0 1.1
137	Sandusky River near Fremont	03/24/89 05/24/89 10/25/89	909 4,640 215	640 620 780	8.2 7.8 8.3	11 7.9 3.8	.50 3.5 1.9	-- -- --	5.9 1.7
138	Auglaize River near Fort Jennings	03/14/89 05/27/89 10/25/89 03/21/90 05/14/90	164 4,550 65.0 200 2,100	770 300 770 700 440	8.2 7.4 7.2 8.7 7.9	6.2 7.5 2.5 6.4 13	.30 >5.0 .90 -- --	-- -- -- -- --	15 .53 .30 16 --
139	Olentangy River at Claridon	03/15/89 06/14/89 10/24/89 03/21/90 05/14/90	117 94.0 11.0 60.0 1,380	642 620 760 630 385	8.2 7.9 7.8 8.4 7.4	6.3 4.3 .70 2.7 11	.20 3.3 .60 -- --	0.40 -- -- -- --	.49 2.9 .43 .24 32
140	Scioto River near Prospect	03/21/89 06/14/89 10/19/89 03/23/90 05/15/90	487 344 20.0 248 2,540	780 810 1,050 890 485	8.1 8.0 7.8 8.0 7.5	7.6 5.9 5.5 5.2 14	>5.0 1.5 .30 -- --	-- -- -- -- --	-- 1.9 .21 .26 27

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/ day/year)	Stream- flow (ft <sup>3</sup> /s)	Specific conduct- ance (µS/cm)	pH (standard units)	Nitrite plus nitrate as nitrogen (mg/L)	Atrazine ELISA- Iowa <sup>1</sup> (µg/L)	Atrazine ELISA- Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
Ohio--Continued									
141	Kokosing River at Mount Vernon	03/13/89 06/14/89 10/12/89	293 1,200 47	420 350 525	8.4 7.7 7.8	2.9 1.4 1.2	<0.20 3.4 1.5	-- -- --	4.7 .90
142	Mad River at Eagle City	03/23/89 06/14/89 10/12/89 03/22/90 05/14/90	330 835 168 323 1,100	730 620 770 770 650	8.5 8.0 8.6 8.3 8.1	4.9 4.7 4.8 3.9 7.7	<.20 >5.0 <.20 -- --	-- -- -- -- 11	-- 10 <.05 .05 11
143	Little Miami River near Oldtown	03/23/89 06/14/89 10/12/89 03/22/90 05/14/90	183 1,160 35 125 608	700 410 760 750 570	8.5 8.0 8.5 8.6 8.2	7.7 4.0 <10 4.6 14	<.20 >5.0 <.20 -- --	-- -- -- -- 26	-- 26 <.05 .20 26
144	Big Darby Creek at Darbyville	03/14/89 06/14/89 10/26/89 03/22/90 05/15/90	791 481 77 295 3,090	625 735 864 702 452	8.2 8.3 8.2 8.6 8.2	8.0 6.4 1.1 3.8 10	.30 2.0 .20 -- --	-- -- -- -- 18	-- 1.4 .20 .18 18
145	Clear Creek near Rockbridge	03/14/89 06/14/89 10/16/89 03/22/90 05/26/90	165 245 43 78 1,250	398 380 467 383 210	8.2 8.0 7.9 7.4 <.10	4.2 5.6 1.3 1.4 --	<.20 >5.0 <.20 -- --	<.20 -- -- -- --	.13 19 <.05 .11 9.0

**Table 6. Results of analyses of water-quality samples collected from Midwest streams and comparison of laboratory analyses for atrazine by ELISA and GC/MS, 1989-90--Continued**

Map no (figs. 2-5)	Site name	Date of collection (month/day/year)	Stream flow (ft <sup>3</sup> /s)	Specific conductance ( $\mu$ S/cm)	pH (standard units)	Nitrate plus nitrate as nitrogen (mg/L)	Atrazine ELISA- <sup>1</sup> Iowa <sup>1</sup> (µg/L)	Atrazine ELISA- <sup>2</sup> Kans. <sup>2</sup> (µg/L)	Atrazine GC/MS <sup>3</sup> (µg/L)
<u>Ohio--Continued</u>									
146	Scioto River at Higby	03/23/89	11,600	574	7.9	5.7	0.30	--	--
		06/19/89	9,910	438	8.2	3.7	2.8	--	4.1
		10/03/89	948	830	8.2	3.3	1.1	--	.95
		03/26/90	3,440	690	8.1	3.3	--	--	.24
		05/15/90	15,400	465	7.9	6.8	--	--	13
147	Little Miami River at Milford	03/23/89	2,930	560	8.3	5.0	1.0	0.40	.48
		06/14/89	2,230	620	8.2	3.4	2.0	--	2.3
		10/17/89	347	590	8.4	2.1	.40	--	.30

<sup>1</sup>Atrazine concentration analyzed by ELISA at U.S. Geological Survey laboratory in Iowa City, Iowa.

<sup>2</sup>Atrazine concentration analyzed by ELISA at U.S. Geological Survey laboratory in Lawrence, Kansas.

<sup>3</sup>Atrazine concentration analyzed by GC/MS at U.S. Geological Survey laboratory in Lawrence, Kansas.

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90**

[ $\mu\text{g/L}$ , micrograms per liter; DEA, deethylatrazine; DIA, desopropylatrazine; --, no data; <, less than; >, greater than; R, regular sample; L, laboratory duplicate; B, blind duplicate; X, extra sample]

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala- chlor ( $\mu\text{g/L}$ )	Atra- zine <sup>2</sup> ( $\mu\text{g/L}$ )	Cyana- zine ( $\mu\text{g/L}$ )	DEA ( $\mu\text{g/L}$ )	DIA ( $\mu\text{g/L}$ )	Meto- lachlor ( $\mu\text{g/L}$ )	Metri- butin ( $\mu\text{g/L}$ )	Pro- meton- azine ( $\mu\text{g/L}$ )	Pro- simazine ( $\mu\text{g/L}$ )
South Dakota												
1	Big Sioux River near Watertown	06/29/89	R	<.05	0.08	<.20	<.05	<.05	<.05	<.05	<.05	<.05
2	Big Sioux River near Castlewood	06/29/89	R	<.05	1.5	<.20	.17	<.05	<.05	.93	<.05	<.05
		04/10/90	L	<.05	.07	<.20	<.05	<.05	<.05	.33	<.05	<.05
		04/10/90	R	<.05	.07	<.20	<.05	<.05	<.05	.26	<.05	<.05
		08/27/90	R	<.05	.14	<.20	<.05	<.05	<.05	.50	<.05	.20
3	Big Sioux River near Brookings	06/29/89	L	.11	.77	<.20	.11	<.05	<.05	.29	.19	<.05
		06/29/89	R	.12	.82	<.20	.12	<.05	<.05	.18	<.05	.05
		11/03/89	R	<.05	.06	<.20	.06	<.05	<.05	<.05	<.05	<.05
4	Big Sioux River at N. Cliff Ave.	06/27/89	R	<.05	1.6	<.20	<.05	<.05	<.05	<.05	<.05	<.05
5	James River near Scotland	04/03/89	R	<.05	.55	<.20	<.05	<.05	<.05	<.05	<.05	<.05
		06/27/89	R	<.05	.19	<.20	<.05	<.05	<.05	<.05	<.05	<.05
		11/02/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05
6	Vermillion River near Vermillion	04/03/89	R	<.05	.17	<.20	<.05	<.05	<.05	<.05	<.05	<.05
		06/27/89	L	<.05	.11	<.20	<.05	<.05	<.05	<.05	<.05	<.05
		06/27/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05
		11/02/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05
7	Brule Creek near Elk Point	06/27/89	R	.09	.36	<.20	.06	<.05	<.05	<.05	<.05	<.05
8	Big Sioux River at Akron	04/03/89	B	<.05	.43	<.20	.08	<.05	<.05	<.05	<.05	<.05
		04/03/89	B	<.05	.46	<.20	.07	<.05	<.05	<.05	<.05	<.05
		04/03/89	L	<.05	.49	<.20	.12	<.05	.06	<.05	<.05	<.05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams,  
1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala- chlor (µg/L)	Atra- zine <sup>2</sup> zinc (µg/L)	Cyana- zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto- lachlor (µg/L)	Metri- buzin (µg/L)	Pro- meton (µg/L)	Pro- pazine (µg/L)	Sima- zine (µg/L)
South Dakota--Continued													
8	Big Sioux River at Akron--Continued	04/03/89	R	<.05	.46	<.20	.10	<.05	<.05	<.05	<.05	<.05	<.05
		06/27/89	B	<.05	.56	<.20	<.05	<.05	.12	<.05	<.05	<.05	<.05
		06/27/89	B	<.05	.62	<.20	<.05	<.05	.14	<.05	<.05	<.05	<.05
		06/27/89	R	<.05	.59	<.20	<.05	<.05	.13	<.05	<.05	<.05	<.05
		11/02/89	B	<.05	.06	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		11/02/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
Nebraska													
9	Bow Creek near St. James	06/26/89	R	.08	.20	.27	<.05	<.05	.12	<.05	<.05	<.05	<.05
		10/18/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
10	Elkhorn River near Atkinson	10/23/89	R	<.05	1.2	<.20	<.05	<.05	2.2	<.05	<.05	<.05	<.05
11	Clearwater Creek near Clearwater	06/26/89	R	.06	.20	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		10/18/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
12	Mira Creek near Northloup	04/11/89	R	<.05	.27	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		06/25/89	R	<.05	.15	<.20	.07	<.05	<.05	<.05	<.05	<.05	<.05
		10/20/89	R	<.05	.47	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
13	Shell Creek near Columbus	06/26/89	R	4.7	14	6.2	2.5	1.8	.74	.30	.20	.18	.15
		10/19/89	R	<.05	.28	<.20	.10	<.05	<.05	<.05	<.05	<.05	<.05
14	Maple Creek near Nickerson	04/05/89	R	<.05	.07	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		06/26/89	R	4.3	8.7	1.3	.97	1.4	.50	.35	<.05	.16	.12
		10/19/89	R	<.05	.08	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/day/year)	Sample type <sup>1</sup>	Alachlor <sup>1</sup> ( $\mu\text{g/L}$ )	Atrazine <sup>2</sup> ( $\mu\text{g/L}$ )	Cyanazine <sup>2</sup> ( $\mu\text{g/L}$ )	DEA ( $\mu\text{g/L}$ )	DIA ( $\mu\text{g/L}$ )	Metolachlor ( $\mu\text{g/L}$ )	Metribuzin ( $\mu\text{g/L}$ )	Propazine ( $\mu\text{g/L}$ )	Simazine ( $\mu\text{g/L}$ )
Nebraska--Continued												
15	Mud Creek at Sweetwater	06/25/89 10/23/89	R R	0.29 <.05	1.7 <.20	0.22 <.05	<.05 <.05	0.20 <.05	<.05 <.05	0.41 <.05	<.05 <.05	<.05 <.05
16	Big Blue River at Surprise	06/26/89 10/31/89	R R	3.1 .11	17 .70	5.9 <.20	1.7 .44	1.4 <.05	3.8 <.05	.41 <.05	.64 <.05	.25 <.05
17	Wahoo Creek at Itica	04/04/89 06/08/89 10/19/89 03/20/90 03/20/90	R R R L R	<.05 2.1 <.05 .05 <.05	<.20 52 <.20 .07 .07	<.05 4.8 <.05 <.20 <.20	<.05 1.9 <.05 <.05 <.05	<.05 3.0 <.05 <.05 <.05	<.05 7.3 <.05 <.05 <.05	<.05 5.9 <.05 <.05 <.05	<.05 .58 <.05 <.05 <.05	<.05 .31 <.05 <.05 <.05
18	W. Fork Big Blue River near Dorchester	05/14/90 05/14/90 06/08/90	X X R	.28 .31 .48	1.7 1.8 9.6	1.8 2.2 6.0	.12 .13 .83	.10 .13 .72	.34 .34 1.3	<.05 <.05 .38	<.05 <.05 .12	<.05 <.05 .05
19	Salt Creek at Rocca	06/26/89 10/31/89 03/27/90 05/14/90 06/07/90	R R R X R	1.0 <.05 <.05 .07 .08	13 .16 .12 .86 2.1	<.20 <.20 <.20 <.20 <.20	<.05 <.05 <.05 <.05 <.05	<.05 3.0 2.0 1.9 .83	<.05 1.1 1.1 .24 .24	<.05 .30 .12 .12 .12	<.05 .34 .27 .27 .27	<.05 .31 .05 .05 .05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/day/year)	Sample type <sup>1</sup>	Ala-chlor <sup>1</sup> ( $\mu\text{g/L}$ )	Atra-zine <sup>2</sup> ( $\mu\text{g/L}$ )	Cyana-zine <sup>2</sup> ( $\mu\text{g/L}$ )	DEA ( $\mu\text{g/L}$ )	DIA ( $\mu\text{g/L}$ )	Meto-lachlor ( $\mu\text{g/L}$ )	Metri-buzin ( $\mu\text{g/L}$ )	Pro-metonazin ( $\mu\text{g/L}$ )	Pro-pazine ( $\mu\text{g/L}$ )	Sima-zine ( $\mu\text{g/L}$ )
Nebraska--Continued													
20	Muddy Creek near Arapahoe	06/25/89 11/07/89	R R	1.4 <.05	6.6 .11	1.2 <.20	.09	<.05	.10	<.05	.07	0.08	0.06
21	Little Blue River near Fairbury	06/26/89 10/30/89 03/21/90 05/17/90 05/17/90	R R R L R	2.3 <.05 <.05 3.4 3.5	22 .31 .21 15 15	1.9 <.20 <.20 .66 .73	3.7 .12 .10 1.1 1.1	2.2 <.05 <.05 .58 .55	1.6 .30 .06 3.1 3.3	.30 <.05 <.05 .06 .06	.06 <.05 <.05 .19 .20	.32 <.05 <.05 .19 .20	1.0 <.05 <.05 .84 .81
22	Little Nemah River at Auburn	06/26/89 10/24/89	R R	.62 <.05	10 .24	.67 <.20	.67 <.05	.88 <.05	1.1 <.05	.30 <.05	<.05 <.05	<.05 <.05	<.05
23	Big Blue River at Barneston	04/06/89 04/06/89 04/06/89 04/06/89 06/27/89	B B L R B	<.05 <.05 <.05 <.05 1.2	.87 .88 .83 .91 16	<.20 <.20 <.20 <.20 .87	<.05 <.05 <.05 <.05 2.1	.58 .48 .41 .59 1.6	<.05 <.05 <.05 <.05 1.3	<.05 <.05 <.05 <.05 .37	.23 <.05 <.05 <.05 .12	.23 <.05 <.05 <.05 .15	
		06/27/89 06/27/89 10/23/89 10/23/89 03/21/90	L R B R B	1.2 1.2 <.05 <.05 <.05	18 16 .47 .51 .65	.87 .83 <.20 <.20 <.20	2.1 2.1 .16 .19 .10	1.5 1.5 <.05 <.05 <.05	.41 .36 <.05 <.05 <.05	.15 .12 <.05 <.05 <.05	.22 .22 <.05 <.05 <.05	.09 .13 .18 .21 .09	
		03/21/90 05/17/90 05/17/90	R R B	<.05 5.7 5.3	.51 14 13	<.20 1.8 1.7	.08 1.1 1.0	<.05 .53 .55	<.05 3.2 2.8	<.05 .29 .26	<.05 <.05 <.05	.05 .20 .16	.05 2.9 2.6

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams,  
1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala- chlor (µg/L)	Atra- zine <sup>2</sup> (µg/L)	Cyana- zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto- lachlor (µg/L)	Metri- buzin (µg/L)	Pro- meton- bazine (µg/L)	Simazine (µg/L)
Nebraska--Continued												
24	Big Nemaha River at Fall City	06/26/89	R	2.1	20	0.33	2.2	1.5	2.9	0.30	0.05	0.23
		10/24/89	R	.07	.54	<.20	.12	<.05	.10	<.05	<.05	<.05
		03/27/90	R	<.05	.81	<.20	.09	<.05	.17	<.05	<.05	<.05
		05/17/90	R	1.1	5.6	.80	.61	.25	1.5	.21	<.05	.06
<u>Kansas</u>												
25	Turkey Creek near Seneca	03/21/89	B	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05
		03/21/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05
		05/22/89	B	<.05	.26	<.20	.06	<.05	.05	<.05	<.05	<.05
		05/22/89	R	<.05	.22	<.20	<.05	<.05	<.05	<.05	<.05	<.05
		10/12/89	B	<.05	.23	<.20	.07	<.05	.06	<.05	<.05	<.05
		10/12/89	R	<.05	.23	<.20	<.05	<.05	<.05	<.05	<.05	<.05
26	Black Vermillion River near Frankfort	06/27/89	L	1.3	17	<.20	2.3	1.6	2.1	.65	.13	.17
		06/27/89	R	1.6	16	<.20	1.8	.77	3.5	.60	<.05	.12
		10/04/89	L	<.05	.39	<.20	.11	<.05	.07	<.05	<.05	<.05
		10/04/89	R	<.05	.35	<.20	<.05	<.05	.08	<.05	<.05	<.05
		03/22/90	R	.06	.47	<.20	.14	.06	.05	<.05	<.05	<.05
		05/16/90	R	1.6	6.1	.41	.58	.24	1.3	.20	<.05	.05
27	Delaware River near Muscotah	05/22/89	L	.30	10	.22	.54	.28	4.3	<.05	.06	.09
		05/22/89	L	.34	10	<.20	.28	<.05	3.6	<.05	<.05	.10
		05/22/89	R	.33	10	<.20	.59	.35	4.2	<.05	.07	.09
		10/11/89	R	<.05	.82	<.20	.18	<.05	.16	<.05	<.05	<.05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams,  
1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup> (µg/L)	Ala- chlor (µg/L)	Atra- zine <sup>2</sup> (µg/L)	Cyana- zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto- lachlor (µg/L)	Metri- buin (µg/L)	Pro- pazine (µg/L)	Pro- simazine (µg/L)
Kansas--Continued												
28	Salt Creek near Ada	03/21/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05
		05/22/89	R	<.05	.27	<.20	.11	<.05	<.05	<.05	.14	<.05
		10/12/89	R	<.05	.49	<.20	.14	<.05	<.05	<.05	<.05	<.05
29	Mill Creek near Paxico	10/06/89	L	<.05	.09	<.20	.06	<.05	<.05	<.05	<.05	<.05
		10/06/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05
30	Kansas River at Topeka	10/06/89	R	.10	1.8	<.20	.39	<.05	.20	<.05	.08	<.05
Minnesota												
31	Knife River near Mora	07/19/89	R	<.05	.09	<.20	<.05	<.05	<.05	<.05	<.05	<.05
		10/31/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05
		05/30/90	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05
32	Sauk River near St Cloud	05/25/89	R	.21	.56	<.20	.11	<.05	.22	<.05	<.05	<.05
		11/08/89	R	<.05	.17	<.20	.10	<.05	<.05	<.05	<.05	.10
33	Pomme de Terre River at Appleton	05/04/89	R	<.05	.06	<.20	<.05	<.05	<.05	<.05	<.05	<.05
		07/18/89	L	<.05	.28	<.20	<.05	<.05	.06	<.05	.19	<.05
		07/18/89	R	<.05	.32	<.20	<.05	<.05	.07	<.05	.21	<.05
		10/25/89	R	<.05	<.05	<.20	.06	<.05	<.05	<.05	<.05	<.05
		10/25/89	X	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05
34	Rum River near St. Francis	07/19/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05
		10/31/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala- chlor (µg/L)	Atra- zine <sup>2</sup> (µg/L)	Cyana- zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto- lachlor (µg/L)	Metri- buzin (µg/L)	Pro- meton azine (µg/L)	Pro- pazine (µg/L)	Sima- zine (µg/L)
<b>Wisconsin</b>													
35	St. Croix River at St. Croix Falls	05/31/89	R	<0.05	0.14	<0.20	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
		10/11/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		04/23/90	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		07/30/90	R	<.05	.16	<.20	.06	<.05	<.05	<.05	<.05	<.05	<.05
<b>Minnesota</b>													
36	Crow River at Rockford	04/19/89	B	<.05	.28	<.20	.09	<.05	<.05	<.05	<.05	<.05	<.05
		04/19/89	R	<.05	.33	.21	.10	<.05	<.05	<.05	<.05	<.05	<.05
		05/25/89	B	1.1	2.9	2.1	.15	.08	.22	<.05	.08	<.05	<.05
		05/25/89	L	1.1	2.9	2.3	.15	.09	.22	<.05	<.05	<.05	<.05
		05/25/89	R	1.1	2.8	2.1	.14	.09	.23	<.05	.07	<.05	<.05
		10/26/89	B	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		10/26/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
37	Minnesota River near Jordan	03/22/89	R	.11	.17	<.20	<.05	<.05	.19	<.05	<.05	<.05	<.05
		07/19/89	R	.08	.29	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		10/29/89	R	<.05	3.1	<.20	.07	<.05	3.4	<.05	<.05	<.05	<.05
38	Redwood River near Redwood Falls	05/04/89	L	<.05	.07	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		05/04/89	R	<.05	.08	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		06/27/89	R	.26	.42	<.20	<.05	<.05	.18	<.05	<.05	<.05	<.05
		10/24/89	R	<.05	.11	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
39	Cottonwood River near New Ulm	07/18/89	R	<.05	.15	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		11/08/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		05/01/90	R	.64	.33	<.20	<.05	<.05	1.0	<.05	<.05	<.05	<.05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/day/year)	Sample type <sup>1</sup>	Ala-chlor <sup>1</sup> ( $\mu\text{g/L}$ )	Atra-zine <sup>2</sup> ( $\mu\text{g/L}$ )	Cyana-zine ( $\mu\text{g/L}$ )	DEA ( $\mu\text{g/L}$ )	DIA ( $\mu\text{g/L}$ )	Meto-lachlor ( $\mu\text{g/L}$ )	Metri-lachlor ( $\mu\text{g/L}$ )	Pro-meton ( $\mu\text{g/L}$ )	Pro-pazine ( $\mu\text{g/L}$ )	Sima-zine ( $\mu\text{g/L}$ )
Minnesota--Continued													
39	Cottonwood River near New Ulm--Continued	06/05/90 06/05/90	R L	.26 .25	2.0 1.8	0.07 .07	.22 .20	0.25 .19	<.05 <.05	0.26 .25	<.05 <.05	<.05 <.05	<.05 <.05
40	Le Sueur River near Rapidan	07/18/89 11/07/89	R R	<.05 <.05	.27 <.20	<.20 <.05	<.05 <.05	.05 .06	<.05 <.05	<.05 <.05	<.05 <.05	<.05 <.05	<.05 <.05
41	Straight River near Faribault	04/05/89 07/19/89 11/08/89	R R R	<.05 .09 <.05	.29 .32 <.20	<.20 .20 <.05	.08 .05 <.05	<.05 <.05 <.05	.05 .05 <.05	<.05 <.05 <.05	<.05 <.05 <.05	<.05 <.05 <.05	<.05 <.05 <.05
42	N. Fork Whitewater River near Elba	11/08/89	R	<.05	.05	<.20	.09	<.05	<.05	<.05	<.05	<.05	<.05
43	Rock River at Luverne	06/27/89 10/23/89 05/08/90 06/19/90	R R R R	.55 <.05 <.05 .63	2.9 .08 .08 3.5	2.2 <.20 <.20 4.0	.53 <.05 .06 .38	.47 <.05 <.05 .30	1.3 .09 .10 2.1	.24 <.05 <.05 <.05	<.05 <.05 <.05 <.05	<.05 <.05 <.05 <.05	<.05 <.05 <.05 <.05
44	Des Moines River at Jackson	06/27/89 06/27/89 10/23/89 10/23/89 05/08/90	L R L R R	.11 .08 <.05 <.05 .16	1.2 1.0 .61 .61 .11	5.0 3.9 <.20 <.20 <.20	.24 .21 <.05 <.05 .06	<.05 <.05 <.05 <.05 <.05	.47 .37 .09 .09 .16	<.05 <.05 <.05 <.05 <.05	<.05 <.05 <.05 <.05 <.05	<.05 <.05 <.05 <.05 <.05	<.05 <.05 <.05 <.05 <.05
		05/08/90 06/19/90 06/19/90	L L R	.15 .05 <.05	.10 .14 .14	<.20 .34 .32	.05 .17 .17	.16 .17 .16	<.05 .17 .17	<.05 .12 .12	<.05 .12 .12	<.05 .05 .05	<.05 <.05 <.05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala- chlor (µg/L)	Atra- zine <sup>2</sup> (µg/L)	Cyana- zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto- lachlor (µg/L)	Metri- buin (µg/L)	Pro- meton (µg/L)	Pro- pazine (µg/L)	Sima- zine (µg/L)
Minnesota--Continued													
45	Rush Creek near Rushford	05/25/89 11/07/89	R R	1.1 <.05	2.1 .06	1.6 <.20	0.15 .09	<.05 <.05	0.89 <.05	<.05 <.05	<.05 <.05	<.05 <.05	<.05 <.05
46	Ocheyedan River near Spencer	03/28/89 05/24/89 10/12/89 10/12/89	R R L R	<.05 12 <.05 <.05	.13 12 <.20 <.20	<.20 .31 <.05 <.05	<.05 .36 <.05 <.05	.19 11 <.05 <.05	<.05 2.2 <.05 <.05	<.05 <.05 <.05 <.05	<.05 <.05 <.05 <.05	<.05 <.05 <.05 <.05	<.05 .06
47	Des Moines River at Estherville	03/28/89 10/12/89	R R	<.05 <.05	.14 .16	<.20 <.20	<.05 <.05	<.05 <.05	.13 .05	<.05 .11	<.05 <.05	<.05 <.05	<.05 <.05
48	Winnebago River at Mason City	03/20/89 03/20/89 10/11/89	L R R	<.05 .05 <.05	.52 .51 <.20	<.20 .13 <.05	<.05 <.05 <.05	.36 .33 .05	<.05 <.05 <.05	<.05 <.05 <.05	<.05 <.05 <.05	<.05 <.05 <.05	<.05 <.05
49	Little Cedar River near Ionia	10/16/89	R	<.05	.14	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
50	Turkey River at Spillville	08/23/89 10/04/89 03/22/90 05/19/90 06/14/90	X R R X R	<.05 <.05 .10 1.2 6.7	<.20 <.20 .43 2.0 8.2	.51 <.05 <.20 1.3 3.4	<.05 <.05 -.36 .35 1.4	<.05 <.05 .10 .09 .84	<.05 <.05 .09 .88 2.5	<.05 <.05 .05 .05 .83	<.05 <.05 .05 .05 .13	<.05 <.05 <.05 <.05 .05	<.05 <.05 <.05 <.05 .05
51	Upper Iowa River near Dorchester	03/31/89 03/31/89 05/31/89 10/05/89	L R R R	<.05 <.05 6.4 <.05	.36 .33 10 .15	<.20 <.20 2.8 <.20	<.05 <.05 .52 <.05	<.05 <.05 .23 <.05	<.05 <.05 1.5 <.05	<.05 <.05 .05 <.05	<.05 <.05 .05 <.05	<.05 <.05 .10 <.05	<.05 <.05 .10 <.05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala- chlor (µg/L)	Atra- zine <sup>2</sup> (µg/L)	Cyana- zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto- lachlor (µg/L)	Metri- buzin (µg/L)	Pro- meton (µg/L)	Pro- pazine (µg/L)	Simazine (µg/L)
Iowa--Continued													
52	Little Sioux River at Correctionville	05/25/89	R	9.1	9.9	1.3	.61	.66	8.2	5.7	<.05	<.05	<.05
		10/02/89	R	<.05	.13	<.20	<.05	<.05	.07	<.05	<.05	<.05	<.05
		03/16/90	R	.09	.19	<.20	<.05	<.05	.22	<.05	<.05	<.05	<.05
		05/19/90	X	2.9	5.8	.25	.46	.37	6.8	2.0	<.05	<.05	<.05
		06/14/90	L	.18	7.4	3.1	.46	.37	1.3	.14	.11	.11	.08
		06/14/90	R	.20	7.4	3.4	.49	.43	1.3	.16	.11	.12	.09
53	North Raccoon River near Sac City	05/24/89	L	5.5	.80	<.20	.07	<.05	.72	<.05	<.05	<.05	<.05
		05/24/89	R	5.5	1.1	2.4	.09	.12	.69	.29	<.05	<.05	<.05
		10/12/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
54	Des Moines River at Fort Dodge	05/24/89	L	12	2.5	2.8	.17	<.05	7.6	1.0	<.05	<.05	<.05
		05/24/89	R	12	2.3	5.4	.14	.14	7.7	.97	.09	<.05	<.05
		10/13/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		03/16/90	R	.12	.12	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		05/14/90	X	.47	.19	<.20	.10	.14	1.3	<.05	.10	<.05	<.05
		05/19/90	X	1.6	.34	.77	.14	.16	2.3	.22	<.05	<.05	<.05
		06/15/90	L	.18	2.0	.26	.19	.17	1.5	<.05	.06	<.05	<.05
		06/15/90	R	.18	1.8	.25	.20	.17	1.4	<.05	.07	<.05	<.05
55	Iowa River near Rowan	10/11/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
56	Cedar River at Cedar Falls	08/23/89	X	<.05	.14	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		11/09/89	R	<.05	.15	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
57	Black Hawk Creek at Hudson	10/18/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala- chlor (µg/L)	Atra- zine <sup>2</sup> (µg/L)	Cyana- zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto- lachlor (µg/L)	Metri- buzin (µg/L)	Pro- meton (µg/L)	Pro- pazine (µg/L)	Sima- zine (µg/L)
<b>Iowa -- Continued</b>													
58	Wapsipinicon River at Independence	03/20/89	R	0.19	1.7	0.26	0.39	0.18	0.53	<0.05	<0.05	<0.05	<0.05
		11/09/89	L	.16	.38	<.20	.26	<.05	.21	<.05	<.05	<.05	<.05
		11/09/89	R	<.05	.39	<.20	.26	<.05	.22	<.05	<.05	<.05	<.05
		03/22/90	R	.17	.63	<.20	.35	.09	.11	<.05	<.05	<.05	<.05
		05/09/90	L	15	23	4.2	1.0	.69	6.6	<.05	<.05	.28	<.05
		05/09/90	R	19	22	2.5	1.0	.77	6.8	<.05	<.05	<.05	<.05
		06/22/90	L	.77	2.4	.60	.73	.51	.72	.34	.27	.35	.11
		06/22/90	R	.89	3.9	.51	.83	.40	.77	.09	<.05	.05	<.05
59	Maple River at Mapleton	05/24/89	R	5.3	34	30	.98	.77	26	7.6	<.05	.50	.26
		10/02/89	R	.49	.59	<.20	<.05	<.05	.19	<.05	<.05	<.05	<.05
		03/16/90	R	.18	.40	<.20	.17	<.05	.31	<.05	<.05	<.05	<.05
		05/19/90	X	1.8	6.8	15	.63	.52	9.8	1.1	<.05	.06	<.05
		06/13/90	R	.90	13	5.3	1.0	.76	4.3	.50	<.05	.18	.09
60	Boyer River at Logan	03/23/89	R	<.05	.14	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		05/24/89	R	13	45	27	1.3	1.1	32	5.6	<.05	.32	.26
		10/02/89	R	<.05	.23	<.20	<.05	<.05	.10	<.05	<.05	<.05	<.05
		03/16/90	R	.12	.55	<.20	.21	<.05	.25	<.05	.06	<.05	<.05
		05/10/90	X	.08	.09	<.20	<.05	<.05	.19	<.05	<.05	<.05	<.05
		05/20/90	X	4.1	4.6	5.8	.51	.35	7.0	.80	<.05	.06	<.05
		06/13/90	R	.75	8.0	2.8	.48	.65	3.0	.07	<.05	.37	.07
61	Raccoon River at Van Meter	05/25/89	R	.95	1.1	1.3	.09	.07	1.0	.22	<.05	<.05	<.05
		11/08/89	R	<.05	.32	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		05/03/90	R	.08	.12	.23	.08	<.05	.18	<.05	<.05	<.05	<.05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams,  
1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala- chlor (µg/L)	Atra- zine <sup>2</sup> (µg/L)	Cyana- zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto- lachlor (µg/L)	Metri- buzin (µg/L)	Pro- meton (µg/L)	Pro- pazine (µg/L)	Sima- zine (µg/L)
Iowa--Continued													
61	Raccoon River at Van Meter--Continued	05/03/90	B	0.08	0.13	<0.20	0.09	<0.05	0.19	<0.05	<0.05	<0.05	<0.05
		05/10/90	L	3.0	2.4	<.20	.24	.25	3.2	.20	<.05	<.05	.06
		05/10/90	R	3.1	2.3	<.20	.23	.24	3.3	.20	<.05	<.05	.06
		05/10/90	B	3.1	2.2	<.20	.21	.20	3.2	<.05	<.05	<.05	<.05
		06/14/90	X	.42	2.3	1.3	.25	.25	1.1	.21	<.05	<.05	<.05
62	Indian Creek near Mingo	05/24/89	R	3.4	5.0	7.9	.40	<.05	14	.62	<.05	.05	<.05
		10/10/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
63	Iowa River near Marengo	03/24/89	R	.06	.42	<.20	.07	<.05	.30	<.05	<.05	<.05	<.05
		05/25/89	R	13	15	10	.77	.63	13	.33	<.05	.19	.14
		11/07/89	R	<.05	.15	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
64	Old Mans Creek near Iowa City	05/25/89	R	51	72	36	3.7	3.2	33	4.2	<.05	.91	.43
		11/07/89	R	<.05	.69	<.20	.16	<.05	<.05	<.05	<.05	<.05	<.05
		03/15/90	X	.10	.78	<.20	.40	.16	.08	<.05	<.05	<.05	<.05
65	N. Fork Maquoketa River at Fulton	08/04/89	X	<.05	.21	<.20	.16	<.05	1.2	<.05	<.05	<.05	<.05
		10/05/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
66	S. Skunk River near Oskaloosa	05/24/89	R	41	48	45	3.1	3.1	23	4.9	.12	.64	.42
		10/16/89	R	<.05	.19	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
67	N. Skunk River near Sigourney	05/24/89	R	40	42	61	2.4	2.0	4.9	2.7	<.05	.51	.41
		10/02/89	R	<.05	.26	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		03/15/90	R	.12	.89	<.20	.49	.26	.23	<.05	<.05	<.05	<.05
		05/21/90	R	.66	1.1	1.6	.22	.10	1.6	<.05	<.05	<.05	<.05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams,  
1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala-chlor (µg/L)	Atra-zine <sup>2</sup> (µg/L)	Cyana-zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto-lachlor (µg/L)	Metri-buzin (µg/L)	Pro-meton (µg/L)	Pro-pazine (µg/L)	Sima-zine (µg/L)	
Iowa--Continued														
68	Nishnabotna River above Hamburg	06/05/89 10/10/89	R R	.20	1.0	37	34	2.2	2.0	19	1.5	0.05	0.51	0.39
69	Chariton River near Chariton	03/20/89 05/30/89 10/02/89	R R R	<.05 2.9 <.05	.21 13 1.3	<.20 7.8 <.20	<.05 1.2 <.05	<.05 1.5 <.05	<.05 3.9 .25	<.05 1.9 <.05	<.05 .06 <.05	<.05 .18 <.05	<.05 .16 <.05	
70	S. Fork Chariton River near Promise City	05/30/89 05/30/89 10/03/89	L R R	2.2 2.4 <.05	15 16 .96	8.0 8.7 <.20	1.3 1.5 .30	.88 1.4 <.05	7.7 7.9 .30	1.0 1.2 <.05	<.05 <.05 <.05	.17 .21 <.05	.12 .16 <.05	
71	Skunk River at Augusta	03/22/89 05/26/89 11/13/89 03/08/90 05/24/90 06/21/90	R R R R X R	<.05 8.9 .14 .10 2.0 1.5	.69 30 .36 .36 1.6 6.5	<.20 31 <.20 <.20 3.0 <.20	.12 1.5 .10 .15 .29 1.4	<.05 1.8 <.05 .17 .31 1.6	.50 5.0 .14 .11 2.8 2.1	<.05 1.6 <.05 .11 .12 .26	<.05 .06 <.05 <.05 <.05 <.05	<.05 .06 <.05 <.05 <.05 <.05	<.05 .36 <.05 <.05 <.05 <.05	
Missouri														
72	Nodaway River near Graham	03/15/89 03/15/89 06/23/89 10/03/89 10/03/89	B R R B R	<.05 <.05 .30 <.05 <.05	.78 1.0 7.8 .20 .21	<.20 <.20 5.6 <.20 <.20	.17 .22 .82 <.05 <.05	.08 .09 .58 <.05 <.05	.20 .27 1.7 .07 .06	<.05 <.05 .42 <.05 <.05	<.05 <.05 .07 <.05 <.05	<.05 <.05 .07 <.05 <.05	<.05 <.05 .06 <.05 <.05	

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala- chlor (µg/L)	Atra- zine <sub>2</sub> (µg/L)	Cyana- zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto- lachlor (µg/L)	Metri- buzin (µg/L)	Pro- meton (µg/L)	Pro- pazine (µg/L)	Sima- zine (µg/L)
Missouri--Continued													
72	Nodaway River near Graham--Continued	04/10/90 05/22/90	R R	<0.05 .12	0.14 1.5	<0.20 1.5	.10 .19	<0.05 .14	.50 .50	<0.05 <.05	<0.05 <.05	<0.05 <.05	<0.05 <.05
73	South Fabius River near Taylor	08/30/89 10/05/89	X R	<.05 <.05	.82 .57	<.20 <.20	<.05 .18	<.05 <.05	.17 .09	<.05 <.05	<.05 <.05	<.05 <.05	<.05 <.05
74	Grand River near Summer	10/04/89	R	<.05	.35	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
75	Middle Fork Salr River at Paris	04/03/89 05/30/89 05/30/89 10/06/89 10/06/89	R L R L R	<.05 .93 .87 .05 .05	.53 3.1 3.0 .42 .65	<.20 1.6 1.4 <.20 <.20	.08 .30 .28 .11 .17	<.05 .39 .34 .08 <.05	.17 .62 .60 .08 <.05	<.05 <.05 <.05 <.05 <.05	<.05 <.05 <.05 <.05 <.05	<.05 <.05 <.05 <.05 <.05	
76	Blackwater River at Blue Lick	05/20/89 10/04/89	R R	2.0 <.05	11 .50	<.20 <.20	.30 .21	<.05 <.05	2.4 .09	.54 <.05	<.05 <.05	<.05 <.05	<.05 <.05
77	Missouri River at Hermann	03/12/89 06/07/89 10/11/89	R R R	<.05 .47 <.05	.18 1.5 .26	<.20 .94 <.20	<.05 .08 <.05	<.05 .05 .12	.05 1.3 .20	<.05 .12 <.05	<.05 .20 <.05	<.05 .05 <.05	<.05 <.05 <.05
78	Mississippi River at Grafton	03/10/89 06/05/89 11/13/89	R R R	.22 .97 .18	.24 2.0 .21	<.20 .14 <.20	.07 .13 .09	<.05 .09 <.05	.05 .87 .16	<.05 .16 <.05	<.05 .05 <.05	<.05 .05 <.05	<.05 <.05 <.05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala- chlor (µg/L)	Atra- zine <sup>2</sup> (µg/L)	Cyana- zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto- lachlor (µg/L)	Metri- buzin (µg/L)	Pro- meton pazine (µg/L)	Pro- simazine (µg/L)
<b>Wisconsin</b>												
79	Prairie River near Merrill	05/05/89 10/23/89	R R	<.05 <.05	<.05 <.05	<.20 <.20	<.05 <.05	<.05 <.05	<.05 <.05	<.05 <.05	<.05 <.05	<.05 <.05
80	Trempealeau River at Dodge	05/30/89 10/17/89	R R	4.5 <.05	26 .15	21 <.20	.84 <.05	.69 <.05	9.5 <.05	<.05 <.05	.37 <.05	.40 <.05
81	Black River near Galesville	05/30/89 10/18/89	R R	2.0 <.05	3.2 <.05	.94 <.20	.11 <.05	.05 <.05	.57 <.05	<.05 <.05	<.05 <.05	.05 <.05
82	Wisconsin River at Muscada	06/02/89 10/05/89 10/05/89 03/20/90 06/22/90	R L R R R	.38 <.05 <.05 .06 .20	.65 .22 .21 .31 .77	<.20 <.20 <.20 <.20 .73	.06 .08 .05 .15 .13	.05 .05 .05 .05 .15	<.05 <.05 <.05 <.05 .19	<.05 <.05 <.05 <.05 .05	<.05 <.05 <.05 <.05 <.05	
83	Grant River near Burton	03/23/89 03/23/89 05/31/89 05/31/89 11/02/89	B R L R B	<.05 <.05 .17 .15 <.05	.32 .32 .14 .17 .22	<.20 <.20 .68 .61 <.20	.12 .11 .16 .07 <.05	<.05 <.05 .09 .09 <.05	<.05 <.05 .53 .56 <.05	<.05 <.05 <.05 <.05 <.05	<.05 <.05 <.05 <.05 <.05	
84	Pecatonic River at Martintown	07/19/89 10/31/89	R R	<.05 <.05	.11 .14	<.20 <.20	.14 .17	<.05 <.05	<.05 <.05	<.05 <.05	<.05 <.05	<.05 <.05
85	Rock River at Afton	04/12/89 04/12/89	L R	<.05 <.05	.29 .34	<.20 <.20	.10 .11	<.05 <.05	<.05 <.05	<.05 <.05	<.05 <.05	<.05 <.05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/day/year)	Sample type <sup>1</sup>	Alachlor (µg/L)	Atrazine <sup>2</sup> (µg/L)	Cyana-zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto-lachlor (µg/L)	Metri-buzin (µg/L)	Pro-meton (µg/L)	Pro-pazine (µg/L)	Sima-zine (µg/L)
<b>Wisconsin--Continued</b>													
85	Rock River at Afton--Continued	07/19/89	L	<.05	.59	<.20	.27	<.05	.09	<.05	<.05	<.05	<.05
		07/19/89	R	<.05	.51	<.20	.19	<.05	<.05	<.05	<.05	<.05	<.05
		10/31/89	R	<.05	.15	<.20	.10	<.05	<.05	<.05	<.05	<.05	<.05
		04/12/90	R	<.05	.39	<.20	.27	.06	<.05	<.05	<.05	<.05	<.05
		07/06/90	R	.08	.72	.29	.25	<.05	.05	<.05	<.05	<.05	<.05
86	Fox River at Wakesha	11/03/89	R	<.05	.12	<.20	.11	<.05	<.05	<.05	<.05	<.05	<.05
87	Root River at Racine	03/22/89	R	<.05	.08	<.20	<.05	<.05	.16	<.05	<.05	<.05	<.05
		07/10/89	L	.10	.82	.32	.15	<.05	.66	<.05	.34	<.05	<.05
		07/10/89	R	.09	.88	.61	.16	<.05	.69	.14	.38	<.05	<.05
		11/03/89	R	<.05	.11	<.20	.09	<.05	.06	<.05	<.05	.09	<.05
		03/27/90	R	.07	.17	<.20	.11	<.05	.09	<.05	<.05	<.05	<.05
		07/19/90	R	.06	.44	.40	.17	.21	.43	<.05	.07	<.05	<.05
<b>Illinois</b>													
88	Des Plains River at Russell	05/25/89	R	.18	.23	<.20	.10	<.05	<.05	<.05	<.05	<.05	<.05
		11/01/89	R	<.05	.15	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
89	Nippersink Creek at Spring Cove	04/05/89	R	<.05	.37	<.20	.16	.11	<.05	<.05	<.05	<.05	<.05
		05/25/89	L	.52	1.7	<.20	.19	<.05	.46	<.05	<.05	<.05	.51
		05/25/89	R	.32	.90	<.20	.12	.10	.16	<.05	<.05	<.05	.33
		11/01/89	R	<.05	.38	<.20	.30	<.05	<.05	<.05	<.05	<.05	<.05
90	S. Br. Kishwaukee River at Fairdale	05/25/89	R	.44	11	14	.75	<.05	.62	<.05	<.05	<.05	<.05
		10/24/89	L	<.05	.23	<.20	.09	<.05	.14	<.05	.11	<.05	<.05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/day/year)	Sample type <sup>1</sup>	Alachlor <sup>1</sup> ( $\mu\text{g/L}$ )	Atrazine <sup>2</sup> ( $\mu\text{g/L}$ )	Cyanazine <sup>2</sup> ( $\mu\text{g/L}$ )	DEA ( $\mu\text{g/L}$ )	DIA ( $\mu\text{g/L}$ )	Metolachlor <sup>1</sup> ( $\mu\text{g/L}$ )	Metricuzin <sup>1</sup> ( $\mu\text{g/L}$ )	Prometonazin <sup>1</sup> ( $\mu\text{g/L}$ )	Simazine <sup>2</sup> ( $\mu\text{g/L}$ )
Illinois--Continued												
90	S. Br. Kishwaukee River at Fairdale--Continued	10/24/89 04/12/90 05/14/90	R R R	<.05 .07 1.6	.24 .25 2.1	<.20 <.20 <.20	<.05 .29 .70	<.05 .10 <.05	.16 .08 1.0	<.05 <.05 <.05	.11 .05 <.05	<.05 <.05 .06
91	Kishwaukee River near Perryville	03/22/89 05/25/89 10/24/89	R R R	<.05 .57 <.05	.12 4.9 .15	<.20 <.20 <.20	.09 .16 .12	<.05 .08 <.05	.05 2.9 .31	<.05 <.05 <.05	.05 .05 <.05	<.05 <.05 <.05
92	Elkhorn Creek near Penrose	05/25/89 10/24/89	R R	1.2 <.05	1.9 .15	1.7 <.20	.54 .19	<.05 <.05	.76 .08	<.05 <.05	<.05 <.05	<.05 <.05
93	Illinois River near Marseilles	05/25/89 10/31/89	R R	.91 <.05	2.5 .15	1.1 <.20	.12 <.05	<.05 <.05	.68 .09	<.05 <.05	<.05 .18	.15 <.05
94	Fox River near Dayton	03/21/89 05/25/89 10/31/89	R R R	<.05 .23 <.05	.20 .80 .26	<.20 1.5 <.20	.08 .12 .17	<.05 .09 <.05	<.05 1.9 <.05	<.05 .11 <.05	<.05 .07 <.05	<.05 .17 .11
95	Dupage River near Shorwood	04/06/89 05/25/89 05/25/89 11/02/89 11/02/89	R B R B R	<.05 .71 .30 <.05 <.05	<.05 .91 .77 <.05 <.05	<.20 1.7 .42 <.20 <.20	<.05 .12 .07 <.05 <.05	<.05 .13 .05 <.05 <.05	.57 .21 .28 <.05 <.05	<.05 .21 .05 <.05 <.05	<.05 .08 .05 <.05 <.05	.06 .26 .13 <.05 <.05
		04/04/90 06/22/90 06/22/90	R L R	<.05 <.05 <.05	.07 .30 .37	<.20 <.20 <.20	.07 .10 .12	.06 .12 .08	<.05 .12 .15	<.05 .05 .05	<.05 .05 .05	<.05 .14 .17

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/day/year)	Sample type <sup>1</sup>	Ala-chlor (µg/L)	Atra-zine <sup>2</sup> (µg/L)	Cyana-zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto-lachlor (µg/L)	Metri-buzin (µg/L)	Pro-meton (µg/L)	Pro-pazine (µg/L)	Simazine (µg/L)
Illinois--Continued													
96	Iroquois River near Chebanse	04/03/89	R	0.13	0.47	0.52	<0.05	<0.05	0.34	<0.05	<0.05	<0.05	<0.05
		05/19/89	X	.43	1.2	.62	<.05	<.05	.43	<.05	<.05	<.05	<.05
		05/25/89	L	.34	1.1	.62	.06	<.05	.42	<.05	<.05	<.05	<.05
		05/25/89	R	1.0	2.8	2.4	.13	.09	.77	.16	<.05	<.05	<.05
		11/03/89	R	<.05	.16	<.20	.13	<.05	.14	<.05	<.05	<.05	<.05
		04/04/90	R	.07	.19	<.20	.12	.07	.14	<.05	<.05	<.05	<.05
		05/17/90	R	4.7	4.2	8.2	.43	.41	1.9	<.05	<.05	.13	.09
97	Edwards River near New Boston	03/22/89	R	<.05	.15	<.20	<.05	<.05	<.05	.16	<.05	<.05	<.05
		10/23/89	R	<.05	.34	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
98	Spoon River at London Mills	04/21/89	R	<.05	.23	<.20	<.05	<.05	.16	<.05	<.05	<.05	<.05
		06/02/89	L	47	109	11	4.2	2.5	40	1.4	<.05	1.4	1.7
		06/02/89	R	47	108	11	4.4	2.2	40	1.5	<.05	1.4	1.9
		10/17/89	R	<.05	.24	<.20	.12	<.05	.24	<.05	<.05	<.05	<.05
		04/10/90	R	.06	.18	<.20	.15	<.05	.19	<.05	<.05	<.05	<.05
		05/14/90	R	1.6	2.0	<.20	.45	.27	4.3	<.05	<.05	<.05	.08
99	La Moine River at Colmar	06/03/89	R	.87	11	11	.80	.45	1.0	<.05	<.05	.08	.15
		10/17/89	R	<.05	.29	<.20	.09	<.05	.08	<.05	<.05	<.05	<.05
100	Sangamon River at Oaksford	05/23/89	R	.74	2.7	.28	.20	<.05	1.3	<.05	<.05	<.05	.12
		10/19/89	R	<.05	.25	<.20	.19	<.05	<.05	<.05	<.05	<.05	<.05
101	Lake Fork near Cornland	05/22/89	R	.18	.64	<.20	<.05	<.05	.26	.15	<.05	<.05	<.05
		10/16/89	L	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		10/16/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala- chlor (µg/L)	Atra- zine <sup>2</sup> (µg/L)	Cyana- zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto- lachlor (µg/L)	Metri- buzin (µg/L)	Pro- meton (µg/L)	Pro- pazine (µg/L)	Sima- zine (µg/L)
<b>Illinois--Continued</b>													
102	Sangamon River at Riverton	05/22/89	R	0.91	2.6	0.94	0.27	<.05	1.9	<.05	0.06	<.05	0.09
		10/12/89	R	<.05	.33	<.20	.11	<.05	.44	<.05	<.05	<.05	<.05
		03/26/90	R	.09	.55	<.20	.27	.13	.20	<.05	<.05	<.05	<.05
		05/14/90	R	1.2	3.8	2.4	.37	.19	1.3	.20	.27	.05	.11
103	Illinois River at Hardin	03/10/89	R	<.05	.24	<.20	.05	<.05	.16	<.05	<.05	<.05	<.05
		06/04/89	R	.48	3.1	.74	.25	.17	.20	<.05	<.05	.07	.07
		10/16/89	R	<.05	.25	<.20	.10	<.05	.13	<.05	.13	<.05	<.05
104	Macoupin Creek near Kane	03/21/89	R	<.05	.89	<.20	.18	.18	.20	<.05	<.05	<.05	<.05
		05/23/89	R	.94	20	2.0	1.5	.61	5.2	.34	.07	.94	<.05
		10/16/89	R	<.05	.62	<.20	.15	<.05	.15	<.05	<.05	<.05	<.05
		03/28/90	R	.26	.79	<.20	.16	.06	.23	<.05	.08	<.05	<.05
		05/14/90	B	2.2	19	2.8	1.8	1.4	5.1	.80	<.05	.20	<.05
		05/14/90	B	1.6	17	3.7	1.4	.79	4.8	.68	<.05	.18	.11
		05/14/90	R	1.3	16	4.6	1.2	.52	3.8	.33	<.05	.17	.09
105	Kaskaskia River near Cowden	05/22/89	R	.23	2.7	<.20	.16	<.05	1.0	.23	<.05	<.05	.55
		10/26/89	R	<.05	1.3	<.20	.34	<.05	.28	<.05	<.05	<.05	<.05
		03/27/90	B	<.05	1.4	<.20	.57	.27	.29	<.05	1.0	<.05	.08
		03/27/90	R	<.05	1.3	<.20	.45	.20	.25	<.05	.07	<.05	<.05
		05/14/90	R	1.4	33	<.20	2.4	1.6	12	2.2	<.05	.41	4.4
106	Kaskaskia River at Vandalia	05/23/89	R	.43	4.9	<.20	.37	.16	2.0	.23	<.05	.05	.55
		10/11/89	L	<.05	.81	<.20	.20	<.05	.20	<.05	<.05	<.05	<.05
		10/11/89	R	<.05	.76	<.20	<.05	<.05	.17	<.05	<.05	<.05	<.05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala-chlor (µg/L)	Atra-zine <sub>2</sub> (µg/L)	Cyana-zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto-lachlor (µg/L)	Metri-buzin (µg/L)	Pro-meton (µg/L)	Pro-pazine (µg/L)	Sima-zine (µg/L)
<b>Illinois -- Continued</b>													
107	Embarras River at Ste. Marie	05/22/89	L	4.4	11	<.20	.21	<.05	4.1	<.05	<.05	<.05	<.05
		05/22/89	R	4.9	8.6	<.20	.22	<.05	3.9	<.05	<.05	<.05	<.05
		10/18/89	R	<.05	.33	<.20	<.05	<.05	.08	<.05	<.05	<.05	<.05
108	Shoal Creek near Breese	03/20/89	R	.11	.61	<.20	.10	<.05	.14	<.05	<.05	<.05	<.05
		05/23/89	R	5.5	15	4.4	.79	.58	6.0	.69	<.05	<.05	.41
		10/11/89	R	<.05	2.1	<.20	.42	<.05	.21	<.05	<.05	<.05	<.05
		03/28/90	R	<.05	1.0	<.20	.27	.12	.09	<.05	<.05	<.05	<.05
		05/04/90	R	1.4	26	.40	.65	1.6	1.3	<.05	<.05	<.05	<.05
109	Silver Creek near Freeburg	05/23/89	R	4.8	19	6.4	1.5	.72	8.2	1.4	<.05	.23	.92
		10/11/89	R	<.05	.52	<.20	.18	<.05	.29	<.05	<.05	<.05	<.05
110	Bonpas Creek at Browns	05/22/89	R	16	52	4.5	3.8	2.1	14	.81	<.05	.75	7.0
		11/27/89	R	<.05	.41	<.20	.22	<.05	.09	<.05	<.05	<.05	<.05
		04/17/90	R	.09	1.8	<.20	.20	.12	.68	<.05	<.05	<.05	.13
		07/10/90	R	1.1	9.2	1.3	2.8	1.6	2.7	.13	.05	.14	.35
	<b>Indiana</b>												
111	Wabash River near New Harmony	06/22/89	R	1.1	6.7	.39	.46	<.05	3.0	<.05	<.05	<.05	.44
		10/31/89	R	.12	.68	<.20	.32	<.05	.37	<.05	<.05	<.05	<.05
	<b>Illinois</b>												
112	Little Wabash River at Carmi	05/22/89	R	15	34	3.1	2.5	1.7	8.1	1.4	<.05	.46	4.9
		11/27/89	R	.21	.69	<.20	.19	<.05	.21	<.05	<.05	<.05	<.05
		04/17/90	R	.41	3.8	.50	.40	.34	1.1	<.05	<.05	<.05	1.4
		07/10/90	R	3.9	9.0	2.4	3.2	2.0	3.0	.79	<.05	.16	.17

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala-chlor (µg/L)	Atra-zine <sup>2</sup> (µg/L)	Cyana-zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto-lachlor (µg/L)	Metri-buzin (µg/L)	Pro-meton (µg/L)	Pro-pazine (µg/L)	Sima-zine (µg/L)
Illinois--Continued													
113	Big Muddy River near Murphyboro	05/24/89	R	1.4	5.1	2.1	0.25	0.13	0.29	<0.05	<0.05	<0.05	0.46
		10/16/89	R	<.05	.57	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
114	Mississippi River at Thebes	03/15/89	R	.44	.27	<.20	.08	<.05	<.05	<.05	<.05	<.05	<.05
		06/10/89	R	.65	1.8	.45	.15	.05	.80	.15	<.05	<.05	.06
		11/08/89	R	.13	.37	<.20	.11	<.05	.15	<.05	<.05	<.05	<.05
115	Ohio River near Grand Chain	03/16/89	R	<.05	<.05	<.20	.05	<.05	<.05	<.05	<.05	<.05	<.05
		06/11/89	R	.57	3.1	.47	.29	.16	1.1	.18	<.05	<.05	.21
		10/11/09	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
Indiana													
116	Kankakee River at Shelby	03/22/89	R	<.05	.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		05/31/89	R	1.0	2.8	<.20	.17	<.05	1.4	<.05	<.05	<.05	.05
		10/16/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
117	Tippecanoe River near Ora	05/31/89	R	.42	1.2	<.20	.13	<.05	.16	.14	<.05	<.05	.43
		10/16/89	R	<.05	.23	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
118	Iroquois River near Foresman	05/31/89	R	2.1	5.9	4.5	.39	.14	1.8	.30	<.05	.07	.16
		10/16/89	R	.09	.11	<.20	.08	<.05	<.05	<.05	<.05	<.05	<.05
119	Eel River near Logansport	03/22/89	R	.08	.26	<.20	.06	<.05	<.05	<.05	<.05	<.05	<.05
		05/22/89	L	1.3	7.4	2.3	.16	.09	4.7	<.05	.06	.05	.34
		05/22/89	R	1.7	8.0	2.5	.17	<.05	5.3	.24	<.05	.06	.37
		10/16/89	R	<.05	.12	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		03/26/90	R	.05	.19	<.20	.09	<.05	.15	<.05	<.05	<.05	<.05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala-chlor ( $\mu\text{g/L}$ )	Atra-zine <sup>2</sup> ( $\mu\text{g/L}$ )	Cyana-zine ( $\mu\text{g/L}$ )	DEA ( $\mu\text{g/L}$ )	DIA ( $\mu\text{g/L}$ )	Meto-lachlor ( $\mu\text{g/L}$ )	Metri-buzin ( $\mu\text{g/L}$ )	Pro-pazine ( $\mu\text{g/L}$ )	Sima-zine ( $\mu\text{g/L}$ )
Indiana--Continued												
119	Eel River near Logansport--Continued	05/14/90	R	4.3	9.6	10	0.91	0.46	3.3	<0.05	0.12	0.13
120	Wabash River at Linn Grove	05/26/89 10/17/89	R R	7.7 <.05	8.6 .27	1.4 <.20	.57 .08	.23 <.05	2.8 .25	.74 <.05	.08 <.05	.09 <.05
121	Wildcat Creek near Lafayette	03/23/89 05/27/89 05/27/89 10/16/89 03/26/90	R L R R R	.07 11 10 <.05 <.05	.22 29 27 .26 .22	<.20 6.9 6.1 <.20 <.20	.05 1.5 1.4 <.05 .15	<.05 .86 .78 <.05 .06	.35 17 16 .14 .12	<.05 1.5 1.3 <.05 <.05	<.05 .19 .16 <.05 <.05	<.05 .14 .25 <.05 <.05
		05/14/90	R	5.4	14	3.6	1.2	.53	8.6	.53	<.05	.20
122	Wildcat Creek near Jerome	05/22/89 10/17/89 03/26/90 05/14/90 05/14/90	R R R L R	.53 <.05 <.05 7.9 7.9	1.5 .35 .27 9.8 10	<.20 <.20 <.20 4.8 5.2	.13 .14 .21 .98 1.0	<.05 <.05 .09 .43 .45	1.3 .32 .16 8.1 8.4	<.05 <.05 <.05 1.2 1.2	<.05 <.05 <.05 1.2 <.05	<.05 <.05 <.05 .14 .14
123	Wabash River at Covington	03/24/89 05/22/89 10/16/89 10/16/89	R R L R	<.05 3.1 <.05 .13	.18 8.8 .79 .68	<.20 2.7 <.20 <.20	<.05 .55 .26 .20	<.05 .23 <.05 <.05	.14 4.9 .33 .29	<.05 .34 <.05 <.05	<.05 .07 <.05 <.05	<.05 .07 <.05 <.05
124	White River near Nora	03/27/89 03/27/89 05/26/89	B R B	<.05 <.05 3.3	.22 .18 9.8	<.20 <.20 4.2	.08 .07 .61	<.05 <.05 .35	.05 4.6 .19	<.05 <.05 .19	<.05 <.05 .06	<.05 .07 .55

**Table 7.** Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams,  
1989-90--Continued

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala-chlor (µg/L)	Atra-zine <sup>2</sup> (µg/L)	Cyana-zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto-lachlor (µg/L)	Metri-luzin (µg/L)	Pro-meton buzin (µg/L)	Pro-pazine (µg/L)	Sima-zine (µg/L)
Indiana--Continued													
124	White River near Nora--Continued	05/26/89	R	3.1	9.8	4.0	0.57	0.33	4.5	0.19	<0.05	0.08	0.51
		11/01/89	B	<.05	.28	<.20	.17	<.05	.14	<.05	<.05	<.05	<.05
		11/01/89	R	.07	.29	<.20	.19	<.05	.14	<.05	<.05	<.05	<.05
		03/26/90	B	<.05	.21	<.20	.13	.07	.09	<.05	<.05	<.05	<.05
		03/26/90	R	.06	.32	<.20	.20	.10	.14	<.05	.05	<.05	<.05
		05/14/90	B	10	19	5.2	1.6	1.0	6.0	1.3	.09	.29	.70
		05/14/90	R	9.8	17	13	1.4	.86	5.6	1.2	<.05	.27	.62
125	Big Blue River at Carthage	05/23/89	L	13	13	3.9	.67	<.05	.18	.28	.15	.13	.87
		05/23/89	R	13	13	4.5	.78	<.05	.20	.30	.17	.13	.92
		10/17/89	R	<.05	.12	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
126	Whitewater River near Alpine	05/23/89	R	9.9	20	14	1.0	.53	5.7	.38	.09	.23	.83
		10/17/89	R	<.05	.25	<.20	.16	<.05	.14	<.05	<.05	<.05	<.05
		03/27/90	L	<.05	.13	<.20	.12	<.05	<.05	<.05	<.05	<.05	<.05
		03/27/90	R	<.05	.14	<.20	.13	<.05	<.05	<.05	<.05	<.05	<.05
		05/15/90	R	3.2	8.2	<.20	1.1	.57	2.6	.31	<.05	.11	.78
		05/15/90	L	2.2	5.9	2.6	.84	.38	2.1	.18	<.05	<.05	.48
127	Big Walnut Creek near Reelsville	03/24/89	R	<.05	.25	<.20	.11	<.05	<.05	<.05	<.05	<.05	<.05
		05/23/89	R	.73	6.6	<.20	.34	.13	1.8	<.05	<.05	.06	1.2
		10/31/89	R	<.05	.41	<.20	.22	<.05	<.05	<.05	<.05	<.05	<.05
128	White River near Centerton	05/23/89	R	1.7	8.7	<.20	0.23	<.05	1.6	<.05	<.05	.10	1.7
		10/30/89	R	<.05	.36	<.20	.21	<.05	.17	<.05	<.05	<.05	<.05

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup> (µg/L)	Ala-chlor (µg/L)	Atra-zine <sup>2</sup> (µg/L)	Cyana-zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto-lachlor (µg/L)	Metri-buzin (µg/L)	Pro-meton (µg/L)	Pro-pazine (µg/L)	Sima-zine (µg/L)
Indiana--Continued													
129	Sugar Creek near Edinburgh	05/25/89	L	4.4	14	9.7	.89	.86	2.2	<.05	<.05	0.37	0.23
		05/25/89	R	4.3	15	7.7	.91	.32	2.1	.27	<.05	.18	.24
		10/30/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		03/27/90	R	<.05	.14	<.20	.11	<.05	<.05	<.05	<.05	<.05	<.05
		05/15/90	R	7.0	20	7.5	2.0	1.0	4.2	.34	<.05	.27	.26
130	Flatrock River at Columbus	05/25/89	L	8.1	14	6.6	.65	<.05	1.9	<.05	<.05	.17	.92
		05/25/89	R	8.5	15	6.9	.96	<.05	2.2	.41	<.05	.15	.78
		10/30/89	R	<.05	.17	<.20	.12	<.05	<.05	<.05	<.05	<.05	<.05
131	Busseron Creek near Carlisle	06/22/89	R	.17	3.5	<.20	.24	<.05	.43	<.05	<.05	<.05	<.05
		10/31/89	R	<.05	.22	<.20	.13	<.05	.05	<.05	<.05	<.05	<.05
132	E. Fork White River near Bedford	05/25/89	R	4.0	20	9.4	1.1	.59	6.8	.29	.09	.23	1.5
		10/30/89	R	<.05	.28	<.20	.12	<.05	<.05	<.05	<.05	<.05	<.05
		03/27/90	R	<.05	.22	<.20	.15	<.05	.05	<.05	<.05	<.05	<.05
		05/15/90	R	2.2	11	4.7	1.2	.69	3.1	<.05	<.05	.14	.64
133	Muscatatuck River near Deputy	03/28/89	R	<.05	.06	<.20	<.05	<.05	<.05	<.05	<.05	<.05	<.05
		05/25/89	R	.23	2.7	2.8	.37	.24	1.0	.14	<.05	<.05	.46
		10/30/89	R	<.05	.37	<.20	.17	<.05	.11	<.05	<.05	<.05	<.05
134	Blue River at Fredricksburg	05/25/89	R	.14	5.1	2.8	.42	.19	4.0	<.05	<.05	.06	.36
		10/30/89	R	.23	.30	<.20	.38	<.05	.07	<.05	<.05	<.05	<.05
		03/27/90	R	<.05	.16	<.20	.15	<.05	<.05	<.05	<.05	<.05	<.05
		05/15/90	R	.17	5.1	1.4	.49	.23	.34	<.05	<.05	.09	.11

**Table 7.** Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams,  
1989-90.-Continued

Map no. (figs. 2-5)	Site name	Date of collection (month/day/year)	Sample type <sup>1</sup>	Ala-chlor (µg/L)	Atra-zine <sup>2</sup> (µg/L)	Cyana-zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto-lachlor (µg/L)	Metri-buzin (µg/L)	Pro-meton (µg/L)	Pro-pazine (µg/L)	Sima-zine (µg/L)
<u>Ohio</u>													
135	Tiffin River at Stryker	03/14/89	R	<.05	.38	<.20	.16	<.05	<.05	<.05	<.05	<.05	<.05
		06/02/89	R	11	28	8.7	2.5	1.9	11	4.7	<.05	.27	.60
		10/25/89	R	.30	.66	<.20	.20	<.05	.54	<.05	<.05	<.05	<.05
		03/21/90	R	.08	.45	<.20	.28	.14	.08	<.05	<.05	.07	.07
		05/14/90	L	22	35	44	3.7	3.4	17	4.5	<.05	.56	.57
		05/14/90	R	10	.14	22	1.4	.85	7.0	1.5	<.05	.14	.12
136	Maumee River at Waterville	05/27/89	R	4.5	7.0	9.0	.30	.12	2.0	1.2	<.05	.05	.39
		10/25/89	R	.23	1.1	<.20	.45	<.05	.63	<.05	.23	<.05	<.05
137	Sandusky River near Fremont	05/24/89	R	.96	5.9	.38	.56	<.05	3.0	.21	<.05	<.05	.21
		10/25/89	R	.50	1.7	<.20	.37	<.05	1.1	<.05	<.05	<.05	<.05
138	Auglaize River near Fort Jennings	05/27/89	L	3.4	12	4.6	.87	.54	7.0	1.2	.12	.14	.26
		05/27/89	R	4.2	15	3.4	.74	.66	7.8	<.05	<.05	.20	.34
		10/25/89	R	<.05	.53	<.20	<.05	<.05	1.5	<.05	.17	<.05	<.05
		03/21/90	R	.07	.30	<.20	.22	.12	1.1	<.05	<.05	.12	.12
		05/14/90	R	4.5	16	11	1.6	.64	12	1.1	<.05	.22	.16
139	Olentangy River at Claridon	03/15/89	R	.19	.49	<.20	.12	<.05	.13	<.05	<.05	<.05	.08
		06/14/89	R	.75	2.9	.35	.33	.18	1.4	.21	<.05	<.05	.10
		10/24/89	L	.12	.48	<.20	.18	<.05	.24	<.05	.15	<.05	<.05
		10/24/89	R	<.05	.43	<.20	.15	<.05	.21	<.05	.07	<.05	<.05
		03/21/90	L	<.05	<.05	<.20	.10	<.05	.06	<.05	<.05	<.05	.30
		03/21/90	R	<.05	.24	<.20	.14	<.05	.10	<.05	.05	<.05	<.05
		05/14/90	R	15	32	12	2.3	<.05	23	<.05	.57	.57	.85

**Table 7. Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams, 1989-90--Continued**

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala-chlor (µg/L)	Atra-zine <sup>2</sup> (µg/L)	Cyana-zine (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto-lachlor (µg/L)	Metronuzin (µg/L)	Pro-pazine (µg/L)	Sima-zine (µg/L)
<u>Ohio--Continued</u>												
140	Scioto River near Prospect	06/14/89	R	0.49	1.9	<.20	0.14	<.05	0.95	<.05	<.05	0.10
		10/19/89	R	<.05	.21	<.20	<.05	<.05	.23	<.05	<.05	0.28
		03/23/90	R	.10	.26	<.20	.14	.13	.23	<.05	<.05	<.05
		05/15/90	R	7.9	27	19	2.1	1.1	14	1.9	<.05	.31
141	Kokosing River at Mount Vernon	06/14/89	R	.63	4.7	1.0	.37	.16	2.2	.23	<.05	.92
		10/12/89	R	<.05	.90	<.20	<.05	<.05	.34	<.05	<.05	<.05
142	Mad River at Eagle City	06/14/89	R	2.1	10	2.4	.97	.46	4.0	.90	<.05	.12
		10/12/89	R	<.05	<.05	<.20	<.05	<.05	.09	<.05	<.05	.17
		03/22/90	R	<.05	.05	<.20	.06	<.05	<.05	<.05	<.05	<.05
		05/14/90	R	2.9	11	5.2	.96	.41	4.1	.28	<.05	.10
143	Little Miami River near Oldtown	06/14/89	R	17	26	9.3	1.0	<.05	1.7	1.1	<.05	.12
		10/12/89	R	<.05	<.05	<.20	<.05	<.05	<.05	<.05	<.05	<.05
		03/22/90	R	<.05	.20	<.20	.19	<.05	.13	<.05	<.05	<.05
		05/14/90	R	20	26	22	2.4	1.5	9.8	2.3	<.05	.42
144	Big Darby Creek at Darbyville	06/14/89	R	.14	1.4	1.0	.14	<.05	.51	<.05	<.05	.33
		10/26/89	L	<.05	.20	<.20	<.05	<.05	.10	<.05	<.05	<.05
		10/26/89	R	<.05	.20	<.20	<.05	<.05	.10	<.05	<.05	<.05
		03/22/90	R	.05	.18	<.20	.15	.10	.05	<.05	<.05	<.05
		05/15/90	R	2.7	18	18	1.8	.98	7.8	.58	<.05	.23
145	Clear Creek near Rockbridge	03/14/89	B	<.05	.12	<.20	.07	<.05	<.05	<.05	<.05	.05
		03/14/89	R	<.05	.13	<.20	.08	<.05	<.05	<.05	<.05	<.05
		06/14/89	B	3.6	14	8.3	1.6	1.5	8.5	.91	<.05	.15
		06/14/89	R	3.6	19	8.3	1.5	1.5	8.8	.96	<.05	.08

**Table 7.** Results of analyses of water-quality samples for selected herbicides and two atrazine metabolites in Midwest streams,  
1989-90--Continued

Map no. (figs. 2-5)	Site name	Date of collection (month/ day/year)	Sample type <sup>1</sup>	Ala- chlor (µg/L)	Atra- zine <sup>2</sup> (µg/L)	Cyana- zine <sup>2</sup> (µg/L)	DEA (µg/L)	DIA (µg/L)	Meto- lachlor (µg/L)	Metri- buzin (µg/L)	Pro- meton azine (µg/L)	Pro- pazine (µg/L)	Sima- zine (µg/L)
Ohio--Continued													
145	Clear Creek near Rockbridge--Continued	10/16/89	B	<.05	.17	<.20	<.05	<.05	.09	<.05	<.05	<.05	<.05
		10/16/89	L	<.05	.24	<.20	<.05	<.05	.19	<.05	<.05	<.05	<.05
		10/16/89	R	<.05	<.05	<.20	<.05	<.05	.09	<.05	<.05	<.05	<.05
		03/21/90	B	<.05	.09	<.20	.08	<.05	<.05	<.05	<.05	<.05	.05
		03/22/90	R	<.05	.11	<.20	.10	<.05	.05	<.05	<.05	<.05	.07
		05/14/90	B	.26	.47	.47	.31	.16	<.05	<.05	.06	.06	.73
		05/15/90	X	.23	4.8	4.4	.46	.28	1.7	<.05	<.05	.05	.74
		05/15/90	X	.29	4.8	4.8	.48	.32	1.7	<.05	<.05	.06	.72
		05/26/90	L	1.8	13	3.7	2.2	2.1	7.2	.98	<.05	.22	2.2
		05/26/90	R	1.7	9.0	5.2	1.5	1.5	4.2	.12	<.05	.12	2.2
146	Scioto River at Higby	06/19/89	R	.73	4.1	<.20	.18	<.05	1.6	.05	.05	.05	<.05
		10/03/89	R	<.05	.95	<.20	.19	<.05	.40	<.05	<.05	<.05	<.05
		03/26/90	L	<.05	.49	<.20	.14	.10	.64	<.05	<.05	<.05	2.1
		03/26/90	R	<.05	.24	<.20	.08	.07	.31	<.05	<.05	<.05	<.05
		05/15/90	R	2.6	13	12	.98	.63	5.1	.34	.07	.07	1.0
147	Little Miami River at Milford	03/23/89	R	<.05	.48	<.20	.25	<.05	.21	<.05	<.05	<.05	.12
		06/14/89	R	.37	2.3	1.8	.22	.13	.56	<.05	<.05	<.05	.33
		10/17/89	R	<.05	.30	<.20	<.05	<.05	.12	<.05	<.05	<.05	<.05

<sup>1</sup>Sample type: B, blind sample;  
L, duplicate sample;  
R, regular sample; and  
X, extra sample.

<sup>2</sup>Analyses by gas chromatography/mass spectrometry at U.S. Geological Survey in Lawrence, Kansas.