

Chapter 10. References

- Bakun, W. H., and Lindh, A. G., 1977, Local magnitudes, seismic moments and coda durations for earthquakes near Oroville, California, Bulletin of the Seismological Society of America, 67, p. 615-629.
- Draper, N. R., and Smith, H., 1966, "Applied regression analysis." 407 pp., John Wiley and Sons, New York.
- Eaton, J. P., 1969, HYPOLAYR, a computer program for determining hypocenters of local earthquakes in an earth consisting of uniform flat layers over a half space, U.S. Geological Survey Open-File Report, 155 p.
- Eaton, J. P., 1970, Harmonic magnification of the complete telemetered seismic system, from seismometer to film viewer screen, U.S. Geological Survey Open File Report, 23 pp.
- Geiger, L., 1912, Probability method for the determination of earthquake epicenters from the arrival time only, (translated from Geiger's 1910 German article) Bulletin of St. Louis University, 8 (1), p. 56-71.
- Jeffreys, H., 1961, Theory of Probability, (3rd ed.) Oxford at the Clarendon Press, London, p. 214-216.
- Klein, F. W., 1985, User's guide to HYPOINVERSE, a program for VAX and Professional 350 computers to solve for earthquake locations, U.S. Geological Survey Open-File Report 85-515, 53 p.
- Lahr, J. C., Page, R. A., and Thomas, J. A., 1975, Catalog of earthquakes in south central Alaska, April-June 1972, U.S. Geological Survey, Open-File Report, 30 p.
- Lahr, J. C., 1980, HYPOELLIPSE/MULTICS: A computer program for determining local earthquake hypocentral parameters, magnitude, and first motion pattern, U.S. Geological Survey Open-File Report 80-59, 59 p.
- Lahr, J. C., 1980, SQUASH/MULTICS: A computer program to be used in conjunction with HYPOELLIPSE to generate an augmented phase data archive, U.S. Geological Survey Open-File Report 80-375, 11 p.
- Lahr, J. C., 1984, HYPOELLIPSE/VAX: A computer program for determining local earthquake hypocentral parameters, magnitude, and first-motion pattern, U.S. Geological Survey Open-File Report 84-519, 76 p.
- Lahr, J. C., 1989, HYPOELLIPSE/Version 2.0: A computer program for determining local earthquakes hypocentral parameters, magnitude, and first-motion pattern, U.S. Geological Survey Open-File Report 89-116, 92 p.

Lee, W. H. K., Bennett, R. E., and Meagher, K. L., 1972, A method of estimating magnitude of local earthquakes from signal duration: U.S. Geological Survey Open-File Report, 28 p.

Lee, W. H. K., and Lahr, J. A., 1972, HYPO71: a computer program for determining hypocenter, magnitude and first motion pattern of local earthquakes, U.S. Geological Survey Open-File Report, 100 p.

Lienert, B. R., Berg, E., and Frazer, L. N., 1986, HYPOCENTER: An earthquake location method using centered, scaled, and adaptively damped least squares, Bulletin of the Seismological Society of America, v. 76, p. 771-783.

Madansky, A., 1959, The fitting of straight lines when both variables are subject to error, Journal of the American Statistical Association, V. 54, p. 173-205.

Pavlis, G. L. and Booker, J. R., 1983, Progressive multiple event location (PMEL): Bulletin of the Seismological Society of America, V. 73, p. 1753-1777.

Pavlis, G. L., and Hokanson, N. B., 1985, Separated earthquake location, Journal of Geophysical Research, V. 90, p. 12777-12789.

Richter, C. F., 1958, "Elementry Seismology." Freeman, San Francisco, California.

Rogers, J. A., Maslak, Sam, and Lahr, J. C., 1980, A seismic electronic system with automatic calibration and crystal reference, U. S. Geological Survey Open-File Report 80-324, 130p.

Uhrhammer, R. A., and Collins, E. R., 1990, Synthesis of Wood-Anderson seismograms from broadband digital records, Bulletin of the Seismological Society of America, v. 80, p. 702-716.