

**Makah Formation--A Deep-Marginal-Basin Sedimentary  
Sequence of Late Eocene and Oligocene Age in the  
Northwestern Olympic Peninsula, Washington**

---

**GEOLOGICAL SURVEY PROFESSIONAL PAPER 1162 - B**



Makah Formation--A Deep-Marginal-Basin Sequence of  
Late Eocene and Oligocene Age in the  
Northwestern Olympic Peninsula, Washington

By P.D. SNAVELY, JR., A.R. NIEM, N.S. MACLEOD, J.E. PEARL, and W.W. RAU

SHORTER CONTRIBUTIONS TO STRATIGRAPHY

---

GEOLOGICAL SURVEY PROFESSIONAL PAPER 1162 - B

*A study of stratigraphy, petrology, paleontology, and paleogeology  
of a marine sedimentary sequence of the Olympic Peninsula*



**UNITED STATES DEPARTMENT OF THE INTERIOR**

**CECIL D. ANDRUS**, *Secretary*

**GEOLOGICAL SURVEY**

**H. William Menard**, *Director*

## CONTENTS

	Page		Page
Abstract.....	1	Physical characteristics -- Continued	
Introduction.....	1	Falls Creek unit.....	15
Acknowledgments.....	2	Unnamed sandstone unit.....	15
Geologic setting and contact relations.....	3	Petrography.....	15
Physical characteristics.....	7	Reservoir potential of turbidite	
Baada Point Member.....	9	sandstone members.....	16
Dtokoah Point Member.....	12	Age and regional correlations.....	17
Carpenters Creek Tuff Member.....	12	Depositional environment.....	21
Klachopis Point Member.....	13	Paleogeology.....	22
Third Beach Member.....	13	Paleodispersal pattern.....	24
Jansen Creek Member.....	14	Source areas.....	25
Other units.....	15	Summary.....	26
		References cited.....	27

## ILLUSTRATIONS

	Page
Figure 1. Sketch map of northern Olympic Peninsula showing area of report and distribution of upper Eocene and Oligocene strata.....	2
2. Generalized bedrock geologic map of the northwest Olympic Peninsula showing the relation of the Makah Formation to other Tertiary units.....	4
3. Generalized geologic map showing the distribution of the Makah Formation at its composite reference section along the Hoko River.....	6
4. Composite type section of the Makah Formation from Waadah Island and Baada Point to Kydaka Point.....	7
5. Composite reference section of the Makah Formation along the Hoko River.....	8
6-8. Photographs showing:	
6. Northeast-trending sandstone dikes cutting a sandstone channel and thin-bedded siltstone and sandstone in the upper part of the Makah Formation.....	9
7. Aerial view toward southeast of prominent headlands formed by the four turbidite members in the lower part of the Makah Formation.....	9
8. Medium- to very thick-bedded turbidite sandstone beds of the Baada Point Member of the Makah Formation.....	10
9. Generalized stratigraphic sections of the four turbidite members of the Makah Formation at their type sections.....	11
10-11. Photograph showing:	
10. Flute casts at base of thick amalgamated sandstone of the Dtokoah Point Member at its type section.....	12
11. Thick beds of turbidite sandstone in the Klachopis Point Member.....	13
12. Ternary diagram showing composition of sandstones in the Makah Formation.....	15
13-14. Photograph showing:	
13. Calcite-cemented biotite-bearing arkosic sandstone in the Third Beach Member.....	16
14. Lithic arkosic wacke from Baada Point Member.....	16
15. Paleogeographic map showing the inferred margins of the Tofino-Fuca basin relative to other deep-water marginal basin facies in the Pacific Northwest.....	23
16. Rose diagrams showing orientations of flute casts and groove casts in the Makah Formation and three of its members.....	24

## TABLES

Table 1. Checklist of foraminifers from the Waadah Island section of the Makah Formation showing frequency of occurrence.....	18
2. Checklist of foraminifers from the Hoko River section of the Makah Formation showing frequency of occurrence.....	20



# Makah Formation--A Deep-Marginal-Basin Sedimentary Sequence of Late Eocene and Oligocene Age in the Northwestern Olympic Peninsula, Washington

By P.D. Snavelly, Jr., A.R. Niem, N.S. MacLeod, J.E. Pearl, and W.W. Rau

## ABSTRACT

The Makah Formation of the Twin River Group crops out in a northwest-trending linear belt in the northwesternmost part of the Olympic Peninsula, Wash. This marine sequence consists of 2800 meters of predominantly thin-bedded siltstone and sandstone that encloses six distinctive newly named members--four thick-bedded amalgamated turbidite sandstone members, an olistostromal shallow-water marine sandstone and conglomerate member, and a thin-bedded water-laid tuff member. A local unconformity of submarine origin occurs within the lower part of the Makah Formation except in the central part of the study area, where it forms the contact between the older Hoko River Formation and the Makah. Foraminiferal faunas indicate that the Makah Formation ranges in age from late Eocene (late Narizian) to late Oligocene (Zemorrian) and was deposited in a predominantly lower to middle bathyal environment.

The Makah Formation is part of a deep-marginal-basin facies that crops out in the western part of the Olympic Peninsula, in southwesternmost Washington and coastal embayments in northwestern Oregon, and along the central part of the coast of western Vancouver Island. On the basis of limited subsurface data from exploratory wells, correlative deep-marginal-basin deposits underlie the inner continental shelf of Oregon and the continental shelf (Tofino basin) along the southwestern side of Vancouver Island.

Directional structures in the Makah Formation indicate that the predominantly lithic arkosic sandstone that forms the turbidite packets was derived from the northwest. A possible source of the clastic material is the dioritic, granitic, and volcanic terranes in the vicinity of the Hesquiat Peninsula and Barkley Sound on the west coast of Vancouver Island. Vertical and lateral variations of turbidite facies suggest that the four packets of sandstone were formed as depositional lobes on an outer submarine fan. The thin-bedded strata between the turbidite packets have characteristics of basin-plain and outer-fan fringe deposits.

## INTRODUCTION


The Makah Formation is part of a thick deep-water marine sequence of upper Eocene to lower Miocene sandstone, siltstone, turbidite sandstone, and conglomerate that crops out in the northwestern part of the Olympic Peninsula (fig. 1) in a northwest-trending linear belt more than 100 km long. The Makah Formation was named and briefly described by Snavelly, Niem, and Pearl (1978) from detailed geologic mapping in the western 35 kilometers of this belt, which includes the Makah Indian Reservation in the northwestern part of the study area. The Makah consists chiefly of well-bedded siltstone and thin-bedded turbidite sandstone and contains six lithologically distinctive and mappable members, described and formally named in this report, and two informal sandstone units. The named members are four thick-bedded amalgamated turbidite sandstone members, a thin but distinctive tuff member, and a penecontemporaneously deformed allochthonous sandstone and conglomerate member that initially was deposited in shallow water.

The type locality of the Makah Formation, designated by Snavelly, Niem, and Pearl (1978) as the shore cliffs and wave-cut platform exposures along the Strait of Juan de Fuca from Waadah Island and Baada Point to Kydaka Point (fig. 2), is herein redesignated the type section. Rocks exposed in the lower reaches of the Sekiu and Hoko Rivers were selected as reference sections (Snavelly and others, 1978).

The Makah Formation is equivalent to the middle member of the Twin River Formation of Brown and Gower (1958). Snavelly, Niem, and Pearl (1978) raised the Twin River Formation to group rank and divided the Twin River Group into three new formations, from oldest to youngest, the Hoko River, the Makah, and the Pysht Formations.

Excellent exposures in the mapped area make it possible to collect detailed sedimentological data on dispersal patterns and lateral and vertical facies changes in the turbidite sandstone units. These data are used in this report to interpret the depositional environment, provenance, and paleogeology of the Makah Formation during late Eocene and Oligocene time.

## MAKAH FORMATION--A DEEP-MARGINAL-BASIN SEQUENCE



## EXPLANATION

- |  |   |
|--|---|
|  | Marine sandstone, siltstone, and conglomerate; minor nonmarine beds in upper part (early Miocene)   |
|  | Thin-bedded marine sandstone, siltstone, turbidite sandstone, and conglomerate (early Miocene to late Eocene)   |
|  | Massive to thin-bedded siltstone, sandstone, and conglomerate; contains lens of mudflow breccia (late and middle Eocene)  |
|  | Melange and broken formation consisting of thick-bedded sandstone, conglomerate, and thin-bedded siltstone and sandstone; contains olistostromal blocks of pre-Tertiary rocks at Point of the Arches and Eocene(?) pillow basalt at Portage Head (late and middle Eocene) |
|  | Pillow lava and breccia with interbedded basaltic sandstone and siltstone (middle and early Eocene)   |

Figure 1.--Sketch map of northern Olympic Peninsula showing area of report and distribution of upper Eocene and Oligocene strata (dot pattern) in relation to other stratigraphic units.

## ACKNOWLEDGMENTS

The valuable data on the age and depositional environment of mollusks in the Makah Formation was provided by Warren O. Addicott. His contribution to this report is gratefully acknowledged. Holly C. Wagner and Howard D. Gower improved the quality of the report

by their helpful suggestions. Mobil Oil Canada, Ltd. generously lent the authors slides of foraminifers and thin sections of correlative stratigraphic units from Vancouver Island, Canada. The authors express their appreciation to Diane Lander and Wendy Niem, who assisted in the preparation of illustrations and editing.

## GEOLOGIC SETTING AND CONTACT RELATIONS

The Makah Formation occurs in the western part of a broad belt of uppermost Eocene and Oligocene strata that crops out along the northern Olympic Peninsula (fig. 1). The Makah is a steeply northward-dipping homoclinal sequence of strata in the upper part of a middle Eocene to lower Miocene marine sedimentary sequence that is more than 6000 m thick. This marine sequence overlies lower and middle Eocene submarine pillow basalt and breccia of the Crescent Formation (fig. 2) that are interpreted as oceanic ridge basalts and associated seamounts by Snavely and his coworkers (Snavely and others, 1968; MacLeod and Snavely, 1973; Snavely and MacLeod, 1977) and Glassley (1974).

The Makah Formation overlies the upper Eocene Hoko River Formation. The contact between the Makah Formation and the Hoko River Formation (Snavely and others, 1978) is mostly covered. This covered interval is less than 50 m wide along Carpenters Creek, Ozette Road, and Little Hoko River. The massive, hackly fractured iron-stained siltstone and very thin bedded sandstone of the Hoko River (fig. 2) are readily distinguished from the well-bedded siltstone and turbidite sandstone of the Makah. Subordinate thick lenses of dark-gray lithic (phyllite- and basalt-rich) sandstone are interstratified with thick-bedded channelized lithic conglomerate and minor pebbly mudstone that occur locally in the Hoko River Formation. These lithologies are uncommon in the Makah Formation.

East of the mapped area, in Pysht, Lake Crescent, Joyce, and Port Angeles quadrangles, Brown and Gower (1958) and Gower (1960) indicate a conformable contact between the lower and middle members (Hoko River Formation and Makah Formation equivalents) of their Twin River Formation. Similarly, in the Little Hoko River and in roadcuts along nearby logging roads in the center of sec. 34, T. 32 N., R. 13 W. (D on fig. 3), the exposed lower contact of the Makah Formation appears to be conformable and gradational over a 50-m interval. At these two localities, two to three units (10 m or thicker) of thin-bedded strata of the Makah containing clastic dikes alternate with beds of massive siltstone (10 m or thicker) which are more typical of the underlying Hoko River Formation. Although beds of massive siltstone like that of the Hoko River Formation occur higher in the Makah, well-bedded turbidite strata dominate the Makah.

A local unconformity occurs within the lower part of the Makah Formation from a point about 2 km west of Jansen Creek eastward to near Charley Creek (fig. 2). From Jansen Creek to the Hoko River, it forms the contact between the Hoko River and Makah Formations. In the area of the Hoko and the Little Hoko Rivers, the unconformity is intraformational, truncating a small syncline and faults that involve the strata in the lower part of the Makah Formation (fig. 3). The unconformity appears to extend eastward almost as far as Charley Creek but was not observed along the Clallam River in the easternmost part of the study area. West of Jansen Creek, the local unconformity probably does not extend as far west as Rasmussen Creek, for it was not observed in the almost continuous sequence of

the Makah and Hoko River Formations exposed in the stream bed.

The local unconformity is readily apparent in the central part of the mapped area (fig. 2), where it forms the contact between the Hoko River and Makah Formations. There the Makah laps onto a broad north-east-trending anticlinal high in the underlying sedimentary and volcanic rocks. Several members of the Makah that are exposed elsewhere in the lower part of the Makah are missing, whereas members above the unconformity extend uninterrupted across the structure. The absence of basal conglomerate, the local nature of the unconformity, and the occurrence of deep-water upper Eocene strata directly above and below the unconformity suggest that it is of submarine origin.

The small northeast-trending faulted syncline that lies between the Hoko and Little Hoko Rivers (figs. 2 and 3) is interpreted as a small flexure that developed on the east flank of the broad northeast-trending anticlinal fold. The structural trends of strata in both the Hoko River and Makah Formations define this small asymmetric syncline that formed prior to the local unconformity. A thrust fault in the upper part of the Hoko River Formation along the western flank of the syncline may account for the asymmetry of the fold (fig. 3). The axial part of the syncline is complicated by several small thrust faults. These and other faults not recognized in the poorly exposed axial part of the syncline probably account for the thicker section of strata apparent between the lowest member of the Makah and the local unconformity (figs. 3 and 5).

The Makah Formation underlies the upper Oligocene and lower Miocene Pysht Formation--a sequence of thick-bedded sandstone, boulder-and-pebble conglomerate, and massive siltstone and mudstone (figs. 2 and 3). In most places in the study area, the contact between the Makah and Pysht Formations is masked by glacial drift. The contact between them is conformable in a cut on a side road 0.8 km southwest of Eagle Point (NW1/4 sec. 14, T. 32 N., R. 13 W.), where very thick bedded sandstone and channelized boulder-to-pebble conglomerate of the Pysht Formation are intercalated with thin-bedded siltstone and sandstone typical of the Makah. Where exposed in quarry and roadcuts immediately north of the airport 0.6 km northwest of Sekiu, a submarine channel conglomerate and sandstone of the Pysht Formation rests unconformably upon an irregular surface cut into thin-bedded siltstone and sandstone of the Makah with as much as a meter of relief; these relations indicate that submarine erosion has taken place locally.

A thickness of more than 2800 m for the Makah Formation is estimated at the type section from the base of the unit near the town of Neah Bay to the upper contact at Kydaka Point (figs. 2 and 4). The minimum thickness of the composite section along the Hoko and Little Hoko Rivers is approximately 2500 m (figs. 3 and 5). The Makah thins over the broad anticlinal high in the central part of the mapped area, where about 550 meters of strata in the lower part of the formation is missing.

MAKAH FORMATION--A DEEP-MARGINAL-BASIN SEQUENCE




Figure 2.--Generalized bedrock geology of northwest Olympic Peninsula showing the relation of the Makah Formation to other Tertiary units. Based on geology by P. D. Snavelly, Jr., N. S. MacLeod, J. E. Pearl, and A. R. Niern, 1970 to 1978.


EXPLANATION


DESCRIPTION OF MAP UNITS

- CLALLAM FORMATION
- PYSHT FORMATION
- MAKAH FORMATION
  - Falls Creek unit
  - Jansen Creek Member
  - Third Beach Member
  - Klachopis Point Member
  - Carpenters Creek Tuff Member
  - Dtokoah Point Member
  - Baada Point Member
- HOKO RIVER FORMATION
- LYRE FORMATION
- ALDWELL FORMATION
- MELANGE AND BROKEN FORMATION
- SANDSTONE AND SILTSTONE
- CRESCENT FORMATION


- Contact
- Local unconformity
- Fault—Dotted where concealed; queried where doubtful;  
U, upthrown side; D, downthrown side
- Left-lateral strike-slip fault
- Thrust fault—Dotted where concealed; queried where doubtful;  
sawteeth on upper plate
- Anticline
- Syncline
- Synform
- Strike and dip of beds
  - Inclined
  - Overturned

CORRELATION OF MAP UNITS





**EXPLANATION**  
CORRELATION OF MAP UNITS



**DESCRIPTION OF MAP UNITS**

**TWIN RIVER GROUP:**

- Lower part of Pysht Formation—Siltstone, sandstone, and conglomerate
- Makah Formation
- Falls Creek unit
- Klachopis Point Member
- Carpenters Creek Tuff Member
- Unnamed tuff
- Baada Point Member
- Unnamed sandstone unit
- Hoko River Formation—Siltstone with sandstone and conglomerate interbeds

- Contact
- Local unconformity
- Fault—D, downthrown side; U, upthrown side
- Thrust fault—Sawteeth on upper plate
- Strike and dip of beds

0 2 KILOMETERS

Figure 3.--Generalized geologic map showing the distribution of the Makah Formation at its composite reference section along the Hoko River from point A (1.6 km east of its mouth) to point B (near the center of section 28) and near the Little Hoko River (between points C and D).


PHYSICAL CHARACTERISTICS

The most common lithologic type in the Makah Formation is siltstone in beds 1 to 10 cm thick rhythmically alternating with thin beds of turbidite sandstone (figs. 4 and 5). The sandstone is fine to very fine grained, medium gray, and quartzo-feldspathic. It is generally parallel laminated to micro-cross laminated, has sharp bottom and top contacts, and forms resistant ledges or ribs in outcrop. In wave-cut exposures, the thin sandstone beds typically display Bouma (1962) turbidite sequences that lack the basal "a" division (T<sub>b-e</sub> through T<sub>d-e</sub>), although some beds are entirely ripple micro-trough cross laminated throughout the bed thickness. The siltstone is medium light gray and hackly fractured and commonly contains very fine grained sandstone stringers and carbonaceous laminae. Siltstone interbeds do not stand out as boldly as the numerous sandstone beds that form ribs and ledges. Thin-bedded strata of this typical lithology occur in the Makah between several thick-bedded amalgamated sandstone members and are well exposed in the roadcuts along Ozette Road adjacent to the Hoko River and on the wave-cut platform along the Strait of Juan de Fuca (figs. 2 and 3). In the stream beds of Rasmussen Creek and the Clallam River, thick units of laminated siltstone occur in the lower part of the Makah and beds of sandstone are rare.

The thick- to very thick bedded amalgamated sandstone units commonly contain Bouma sequences that start with the basal "a" division. Typically, these sandstone beds consist of a thick structureless "a" division overlain by a much thinner carbonaceous parallel-laminated "b" division and less commonly an overlying convolute or rarely micro-cross-laminated "c" division. Some very thick beds form a series of amalgamated Bouma "a" divisions.

Calcareous concretions are found throughout the Makah Formation; they range in size from small spheroids to 60 to 150 mm in diameter in the thin-bedded units to resistant zones of ellipsoidal to interconnected elongate concretions 0.5 to 4 m in length in the thick amalgamated sandstone units. Some spheroidal concretions in the siltstone contain concentrations of worm burrows; others have formed around pseudomorphic crystals of calcite similar to those described by Boggs (1972). Many small calcareous concretions lack obvious nuclei, but a few contain fragments of crustaceans or carbonized plant debris. Scattered throughout the thick-bedded amalgamated sandstone at the type section are tiny spherical pyrite concretions with oxidized rims.

Clastic dikes occur throughout the Makah Formation (fig. 6). The sandstone in these dikes is fine to medium grained, moderately well sorted, and quartzo-feldspathic in composition. Most dikes are nearly vertical and pinch and swell along strike. Some have been injected along small faults or fractures that had minor displacements, whereas others have been emplaced in an en echelon pattern along fractures with no apparent displacement. The general trend of the clastic dikes perpendicular to the strike of the Makah suggests that the dikes were intruded along small tear faults formed when segments of the upper part of the



EXPLANATION

- Siltstone
- Sandstone
- Concretionary sandstone
- Conglomerate
- Clastic dike
- Allochthonous penecontemporaneous deformed strata

Figure 4.--Composite type section of the Makah Formation from Waadah Island and Baada Point to Kydaka Point. Numbers in parenthesis are foraminifer sample locality numbers (table 1), Waadah Island.

MAKAH FORMATION--A DEEP-MARGINAL-BASIN SEQUENCE




Figure 5.--Composite reference section of the Makah Formation along the Hoko River from point A in section 14 (1.6 km southeast of its mouth) to point B (near the center of section 28) and between points C and D near the Little Hoko River. Locations are shown on figure 3. Numbers in parenthesis are foraminifer sample locality numbers (table 2).




Figure 6.--Northeast-trending sandstone dikes cutting a sandstone channel (0.7 m thick at axis) and thin-bedded siltstone and sandstone in the upper part of the Makah Formation on wave-cut platform 0.4 km west of the mouth of the Sekiu River. Man for scale in left center.

thick sedimentary prism crept basinward or that the sand was injected along fractures perpendicular to the direction of minimum compressional stress in a manner similar to that of basalt dikes (Nakamura, 1977).

The sandstone dikes range in width from 20 mm to 1 m and commonly form carbonate-cemented resistant ridges that stand above the adjacent strata. On the wave-cut platform on the southern shore of Waadah Island, a distinctive set of resistant dikes, 0.9 m and 0.75 m wide, can be traced for more than 300 m southwestward across the lowest thick-bedded turbidite member and the underlying thin-bedded strata before they extend so far below sea level that they can no longer be seen. At Klachopis Point, a 1-m-wide quartzo-feldspathic sandstone dike can be traced for 37 m before it disappears beneath the Strait of Juan de Fuca.

Zones of penecontemporaneously deformed strata to several meters thick occur locally in the thin-bedded sandstone and siltstone of the Makah sequence. The disrupted turbidite sandstone beds in these zones appear to have been hydroplastically deformed. Broad slump folds, pull-apart structures, and small recumbent folded sandstone blocks in structureless concretionary siltstone are the most common features.

Several diastems with angular discordance of as much as 10° are present within the thin-bedded sequences of the Makah exposed in roadcuts adjacent to the Hoko River. These unconformities involve several tens of meters of thin-bedded turbidite strata. The angular discordances probably resulted from minor



Figure 7.--Aerial view toward southeast of prominent headlands formed by the four turbidite members in lower part of the Makah Formation. Headland in foreground is formed by turbidite of the Baada Point Member. The second headland is formed by the Dtokoah Point Member; the third, the Klachopis Point Member; the fourth, the Third Beach Member.

slumping and rotation of large cohesive blocks of sediments that were later covered by hemipelagic silt and turbidite sand. Some angular discordance between juxtaposed sequences of thin-bedded strata may be caused by broad channeling and deposition of turbidite sand beds within the channels with initial dips different than those of the underlying truncated strata.

The six newly named members of the Makah Formation differentiated on the map (fig. 2) are described below from oldest to youngest.

#### BAADA POINT MEMBER

The stratigraphically lowest packet of thick-bedded amalgamated turbidite sandstone in the Makah Formation is here named the Baada Point Member for exposures at its type section at Baada Point. The type section is designated as the wave-cut platform and headland exposures at Baada Point (SE1/4NW1/4 sec.12, T. 33 N., R. 15 W.; fig. 7). The member is also well exposed on the wave-cut platform of Waadah Island northwest of Baada Point (fig. 8). At its type section, the Baada Point member is 120 m thick and occurs approximately 450 m above the base of the Makah Formation. Here the member displays gradational contacts with the overlying and underlying thin- to medium-bedded strata of the Makah; the contacts are arbitrarily chosen at the lowest and uppermost amalgamated sandstone beds that exceed 1.2 m in thickness.

The Baada Point Member can be mapped southeastward for about 12.5 km to where it abruptly terminates against the west side of an anticlinal high and is un-



Figure 8.--Medium- to very thick-bedded turbidite sandstone beds of the Baada Point Member of the Makah Formation exposed on the northwest wave-cut platform of Waadah Island along the Strait of Juan de Fuca north of Baada Point. Note dark-colored zones of calcareous concretions. Man for scale in left center foreground.

conformably overlapped by younger strata of the Makah in the central part of the mapped area (fig. 2). East of the Hoko River, the Baada Point Member crops out again in a northeast-trending faulted syncline and extends relatively undeformed southeastward for a distance of about 13 km to the eastern limits of the mapped area and beyond. The member forms a mappable couplet with a thinner unnamed amalgamated sandstone unit that occurs 75 m below it in the area between the Hoko River and Charley Creek (figs. 2, 3, and 5).

The member thins from 120 m at the type section to 43 m in the Clallam River in the easternmost part of the mapped area. Between these two localities, the sandstone-to-siltstone ratio decreases from 2.5:1 to 1:1. On Waadah Island and at Baada Point, 3- to 5-m-thick amalgamated sandstone beds are common in the

member, whereas in the Clallam River, the thickest sandstone bed is 1.2 m. More than 100 individual medium to thick beds of sandstone occur at the type section. Significant variations in total thickness and number of sandstone beds in the member occur along strike between the type locality and the Clallam River, perhaps related to channeling and lensing of some of the sandstone beds.

The sandstone in the Baada Point Member is typically light olive gray on fresh surfaces and weathers to a "dirty appearing" yellow brown. It is porous, quartzo-feldspathic in composition, predominately fine to medium grained, and moderately well sorted.

The member consists of several ridge-forming units of thick-bedded amalgamated turbidite sandstone and intervening sequences of less resistant thin-bedded




Figure 9.--Generalized stratigraphic sections of the four turbidite members of the Makah Formation at their type sections. Beds represented by R and S referred to in text. See figure 4 for position of the members in type section of the Makah.

sandstone and siltstone (fig. 9). In shoreline exposures, the sandstones form prominent ribs and thin-bedded sandstone and siltstone form inlets that are partly covered during high tides (figs. 7 and 8). Layers of resistant light-brown calcareous concretions are common throughout the thick amalgamated sandstone (fig. 8). Concretions are lenticular to spheroidal in shape, are light medium gray when freshly broken, and display a case-hardened honeycomb weathering pattern on the wave-cut platform.

Bedding in the thick amalgamated sandstone is defined by surface indentations produced by differential erosion of less resistant thin units of laminated to convolute carbonaceous sandstone and rare siltstone (fig. 8). Normal grading in the amalgamated beds is characterized by an upward change from massive fine-

grained sandstone to laminated or convolute-bedded very fine grained carbonaceous sandstone that forms the upper several centimeters of each 0.5- to 3-m-thick bed. In the northwestern part of the outcrop area, scattered dark coarse-grained lithic fragments (coarse tail grading) and concentrations of small light-gray siltstone rip-ups serve to differentiate thinner turbidite beds within seemingly structureless thick amalgamated sandstones. Bouma *Ta*, *Tab*, and *Tabc* sequences are common in thick beds, whereas in the thin sandstone beds in the intervening sandstone-siltstone units, Bouma *Tbed* and *Tb-e* sequences are most common. The sharp bases of many thick sandstone beds contain load casts and, less commonly, burrow, flute, and groove casts.

### DTOKOAH POINT MEMBER

The packet of turbidite sandstone that occurs approximately 165 m above the Baada Point Member is here named the Dtokoah Point Member for exposures at Dtokoah Point (figs. 2 and 7). The type section is designated as the exposures on the point and wave-cut platform in the NW1/4SE1/4 sec. 12, T. 33 N., R. 15 W. The Dtokoah Point Member is separated from the underlying Baada Point Member by 165 meters of very thin bedded siltstone and very fine grained sandstone of the Makah (fig. 4), exposed during low tides. At its type section the Dtokoah Point Member is 65 m thick and displays gradational lower and upper contacts with the thinner bedded strata enclosing it. The bottom and top of the member are arbitrarily defined as the lowest and uppermost sandstone beds greater than 0.3 m thick. The member has been mapped for 9 km southeast to where it apparently pinches out east of Bullman Creek (fig. 2).

The sandstone in the Dtokoah Point Member is fine to very fine grained, moderately well sorted, and weathers to olive gray. The sandstone appears to be compositionally and texturally similar to sandstone in the Baada Point Member; it is rich in matrix, contains lithic fragments, and is quartzo-feldspathic. Sandstone in the Dtokoah Point Member is commonly much thinner bedded than that in the Baada Point Member, the beds generally ranging in thickness from 20 mm to 0.4 m, and contains only a few thick amalgamated beds (fig. 9).

At the type section of the Dtokoah Point Member, it forms a low, commonly tide-covered wave-cut platform composed of numerous thin- to medium-bedded sandstone ribs with intervening swales of less resistant laminated medium-gray siltstone beds, 20- to 70-mm-thick. The more than 170 sandstone beds in the member commonly display sharp bottom and top contacts. Common features are parallel laminations, convolute bedding, micro-trough or micro-ripple cross-laminations, and graded bedding. Bouma *Tb-e*, *Tbc*, and *Tde* sequences predominate.

A 2-m-thick amalgamated sandstone bed near the top of the unit forms a broad lens that pinches and swells laterally (R on fig. 9). Graded beds within this sandstone are defined by concentrations of siltstone rip-ups, shell fragments, convolute and parallel laminations, and phyllite- and basalt-rich grits. Overlying this sandstone lens is 2 to 4 meters of thin turbidite beds that are channelized, pinch out laterally, and display cross-cutting stratigraphic relations (fig. 9).

A distinctive 3-m-thick amalgamated sandstone marker bed occurs near the base of the member (S on fig. 9). This concretionary sandstone forms a prominent ridge and contains well-developed flute marks that indicate a dispersal pattern to the southeast (fig. 10). Spheroidal calcareous concretions to 0.2 m in diameter are scattered throughout the member.




Figure 10.--Flute casts (above hammer) at base of thick amalgamated sandstone of the Dtokoah Point Member at its type section showing southeastward (arrow) dispersal pattern.

### CARPENTERS CREEK TUFF MEMBER

The Carpenters Creek Tuff Member, here named for exposures in Carpenters Creek, is generally 1 m thick and consists of seven thin water-laid tuff beds 40 to 150 mm thick intercalated with 30- to 150-mm-thick siltstone beds. Its type section is in the NW1/4NW1/4 sec. 17, T. 32 N., R. 13 W. It was recognized only in the eastern part of the study area, where it has been mapped for a distance of about 20 km (fig. 2). Near the Little Hoko River, the tuff member occurs approximately 325 m stratigraphically above the Baada Point Member (fig. 5).

The tuff is calcified and silicified, even bedded, forms thin resistant ledges, weathers to light-yellowish-gray blocks and chips, and displays sharp bottom and top contacts with the intervening, less resistant, hackly fractured siltstone. Individual tuff beds are generally structureless to faintly laminated. More rarely, they are cross laminated. This member may be significant in correlating other units in the Makah along the north side of the Olympic Peninsula because it defines a time horizon.

The Carpenters Creek Tuff Member is best exposed near the top of a 20-m-high roadcut in thin-bedded strata of the Makah along Ozette Road east of the Hoko River in the NE1/4 sec. 28, T. 32 N., R. 13 W. Other good exposures of the member occur in the Little Hoko River and along an abandoned railroad grade adjacent to the west bank of the Hoko River (figs. 2 and 3).

Another tuff unit consisting of two to three 20- to 50-mm-thick light-yellowish-gray beds with thin intervening siltstone occurs 100 m below the Carpenters Creek Tuff Member (fig. 5); it can be traced for approximately 3 km east of Ozette Road (fig. 3). In the easternmost part of the study area along the Clallam River, several additional tuff beds occur below the Carpenters Creek Tuff Member in the lower part of the Makah Formation. Since these tuff beds have not been recognized west of the Clallam River, the source of the ash probably was from the east.

### KLACHOPIS POINT MEMBER

The packet of thick-bedded turbidite sandstone that occurs 155 m stratigraphically above the Dtokoah Point Member (fig. 4) is here named the Klachopis Point Member for exposures at Klachopis Point. This 73-m-thick member forms the third wave-resistant platform and headland east of Neah Bay at Klachopis Point, the type section for the member (NE1/4SE1/4 sec. 12, T. 33 N., R. 15 W.; figs. 7, and 9). The member displays a gradational contact with the underlying thin-bedded siltstone and sandstone of the Makah Formation. At the type section, the upper contact is covered with beach sand. The contacts are arbitrarily drawn at the lowest and uppermost sandstone beds that exceed 0.7 m in thickness.

The Klachopis Point Member consists of 40 or more thick to very thick beds of amalgamated sandstone that form a strike ridge entirely across the mapped area, a distance of 32 km (fig. 2). It gradually thins eastward. At Charley Creek, in the eastern part of the area (fig. 2), the member is 49 m thick. It is well exposed on Bullman and Rasmussen Creeks, along the Hoko River (fig. 11) and Sekiu River, and along State Highway 112. Sandstone of the Klachopis Point Member is characteristically more micaceous and feldspathic and better sorted than that of the Baada Point and Dtokoah Point Members. Coarse-grained flakes of muscovite and biotite are ubiquitous. The sandstone typically is iron stained, fine to medium grained, moderately well sorted, and porous. At the type section on the coast, this unit, like the Baada Point Member, contains resistant layers of interconnected dark olive-black-stained calcareous concretions.

The lower one-third to one-half of the member typically consists of well-laminated micaceous, carbonaceous sandstone with siltstone interbeds (fig. 9). Varying concentrations of carbonized plant debris and muscovite and biotite flakes along laminae impart a distinctive platy character to the sandstone. Each sandstone bed has sharp upper and lower contacts with the intercalated medium-dark-gray siltstone and very fine grained carbonaceous sandstone.

The upper one-half to two-thirds of the member forms a thick resistant ridge composed of several even-bedded, predominantly structureless amalgamated sandstone beds (fig. 9). These beds are 1 to 5 m thick and are separated by very thin intervals of less resistant laminated siltstone and thin- to medium-bedded sandstone. Sandstone-to-siltstone ratios vary from 10:1 to 60:1.

Each amalgamated sandstone bed is composed of two or more predominantly structureless parts that range in thickness from 0.15 m to 1 m. Stratification is defined by minor concentrations of siltstone rip-ups, scattered coarse sand-size lithic fragments, and thin beds of less resistant very fine grained carbonaceous sandstone. The carbonaceous sandstone that commonly forms the thin upper part of many structureless beds is parallel laminated to convolute laminated, rarely cross laminated. Bouma *T<sub>1</sub>*, *T<sub>2</sub>*, and *T<sub>3</sub>* sequences are common.

Other sedimentary features in the sandstone beds of the Klachopis Point Member are sharp bottom contacts and gradational upper contacts, bifurcating *Scalari-tuba* worm burrows, rare dish structures, and flute, frondescant load, and groove casts.

### THIRD BEACH MEMBER

The 43-m-thick packet of turbidite sandstone that occurs 290 m stratigraphically above the Klachopis Point Member is here named the Third Beach Member for exposures in the headland east of Third Beach (fig. 7). The type section of the Third Beach Member is designated as the resistant wave-cut terrace and headland immediately east of the beach (SE1/4SW1/4 sec. 7, T. 33 N., R. 14 W.; fig. 9). The Klachopis Point and Third Beach Members are separated by an interval of rhythmically alternating very thin- to thin-bedded siltstone and very fine grained sandstone that is seasonally covered by beach sand.



Figure 11.--Thick beds of turbidite sandstone in the Klachopis Point Member exposed at boundary between secs. 21 and 28, T. 32 N., R. 13 W., along the Ozette Road, Hoko River area.

The Third Beach Member forms a spectacular linear dip-slope seacliff from the type section 2 km south-eastward to Sail River. The member terminates abruptly at Bullman Creek (fig. 2), possibly the result of post-depositional uplift and slumping, as biotite-rich arkosic sandstone blocks similar in composition to sandstone in the Third Beach Member occur in a younger penecontemporaneously deformed siltstone unit 5 km farther east near Jansen Creek (fig. 2). The position of the siltstone unit approximately 100 m stratigraphically above the Third Beach Member precludes their origin by submarine landsliding of sandstone of the Third Beach Member alone; rather, the sandstone was buried and indurated before it was uplifted and slumped in the basin.

Because the lower and upper contacts on the Third Beach Member are generally covered by beach sand at the type section, the contacts are arbitrarily placed at the uppermost and lowest amalgamated sandstone beds more than 3 m thick.

The sandstone in the Third Beach Member is similar to that in the upper half of the underlying Klachopis Point Member, being predominantly thick- to very thick-bedded, concretionary, clean, micaceous, and feldspathic (fig. 9). It differs by containing more abundant and larger (to 1.5 mm), ubiquitous black biotite flakes that impart a distinctive salt-and-pepper appearance to fresh hand specimens. More than 45 sandstone beds occur in the member. The sandstone weathers to an iron-stained grayish orange to light gray and is fine to medium grained.

At its type section, the Third Beach Member consists of three very thick bedded amalgamated sandstone ridges separated by a few beds of less resistant thin- to medium-bedded turbidite sandstone and siltstone (fig. 9). Each sandstone ridge consists of three or more 0.5- to 6-m-thick predominantly structureless concretionary beds. Less resistant laminated and convolute-bedded very fine grained carbonaceous sandstone in the upper part of each bed delineates stratification within the amalgamated sandstones. Trough cross-laminations, large penecontemporaneous slump folds, siltstone rip-ups, and rare dish structures are present. Bouma  $T_1$ ,  $T_{1b}$ , and less commonly,  $T_{2c}$  sequences, occur.

Between the amalgamated sandstone ridges, several thin rib-forming sandstone beds are intercalated with laminated siltstone and very fine grained carbonaceous sandstone. The fine- to medium-grained sandstone beds have sharp bottom and gradational upper contacts, even bedding, load casts, rare siltstone rip-ups, and  $T_{1b}$ ,  $T_{2c}$ , and  $T_{3-e}$  Bouma sequences.

### JANSEN CREEK MEMBER

The Jansen Creek Member, here named for exposures along the shoreline adjacent to the mouth of Jansen Creek (SE1/4SE1/4 sec. 26, T. 33 N., R. 14 W.), is composed of large tabular and penecontemporaneously deformed strata of shallow-water marine conglomerate and fossiliferous sandstone enclosed in deep-water marine siltstone and sandstone. The type section of this olistostromal unit is designated as the discontinuous exposures in the sea cliffs, wave-cut plat-

forms, and sea stacks from the headland 0.5 km east of Bullman Creek to a point 0.7 km southeast of Brush Point (NE1/4NW1/4 sec. 6, T. 32 N., R. 13 W.; fig. 2). The Jansen Creek Member is approximately 200 m thick and is separated from the underlying Third Beach Member by about 250 meters of thin-bedded siltstone and sandstone. In the type section of the Jansen Creek Member, large tabular blocks, 5 to 100 meters in length, of interbedded fossiliferous shallow-marine basaltic sandstone and pebble conglomerate are common. These blocks are infolded or are aligned sub-parallel to the enclosing deep-water marine thin-bedded turbidite sandstone and siltstone and massive hackly fractured concretionary siltstone. The type section contains very thick bedded fossiliferous shallow-water marine sandstone that forms massive resistant overhanging cliffs that can be traced for several kilometers along strike. In places the sandstone beds are folded into broad antiforms and synforms. The yellowish-gray basaltic to feldspathic sandstone is mottled and bioturbated and contains irregular olive-gray calcareous concretionary masses. Scattered fossils in the allochthonous fine- to medium-grained sandstone include the mollusks *Conchocele*, *Lucinoma*, and *Ostrea*, and a few bryozoans and *Teredo*-bored carbonized wood fragments. Several thin coquina-like beds consist of pelecypods, gastropods, and encrusting stromatolitic calcareous algae.

The sandstone beds are interbedded with, and overlain by, well-stratified basalt-pebble conglomerate as much as 10 m thick. The calcite-cemented olive-black conglomerate beds are best exposed at low tide on the wave-cut terraces, where they form resistant ribs or large loose blocks. Individual conglomerate layers are moderately to poorly sorted and range in thickness from 80 mm to 0.2 m. The well-rounded to subrounded pebbles and cobbles consist predominantly of finely crystalline to aphanitic basalt, discoid to spherical in shape and locally imbricated.

On the wave-cut platform from Seal Rock to Shipwreck Point, basaltic pebble conglomerate and shallow-marine sandstone form a series of disharmonic folds with axes dipping as much as 50° to the north or south. The amplitude of these infolds is as much as 100 m. Locally, undeformed clastic dikes cut across the penecontemporaneously deformed strata.

The basal contact of the allochthonous shallow-marine sandstone and conglomerate blocks with the underlying undeformed deep-water marine strata of the Makah Formation is planar to irregular with little or no shearing. In the wave-cut platform at the mouth of Rasmussen Creek, the contact between the Jansen Creek Member and older strata is sharp and appears conformable. In several roadcuts along State Highway 112 between Bullman Creek and Rasmussen Creek, the basal contact is discordant. Thin siltstone beds within overturned blocks of the massive shallow-marine sandstone display 60° to 90° disparity in attitude with the underlying strata. The Jansen Creek Member is overlain by undeformed thin-bedded siltstone and sandstone of the Makah that contain a few broad channels of sandstone. This upper contact is sharp in the few places where it is exposed during very low tides.

Near Brush Point the member is overlain by interbedded basaltic sandstone and pebble conglomerate of the Makah Formation that contain displaced shallow-water mollusks. These basaltic strata are exposed at Brush Point and eastward on the wave-cut platform as far east as the mouth of the Sekiu River. In these coastal outcrops, they are gently flexed and are interbedded with typical thin-bedded siltstone and feldspathic sandstone of the Makah. The similarity in composition of the basaltic sandstone and conglomerate to that of some detached blocks and infolds in the Jansen Creek Member suggests that the basaltic sediments were derived from a source area similar to that of the basaltic beds of the olistostromal Jansen Creek Member but were deposited by normal sedimentary processes.

OTHER UNITS

Falls Creek unit

A poorly exposed sequence of amalgamated turbidite sandstone beds, informally called the Falls Creek unit, crops out in the northeastern part of the mapped area (fig. 2). This thick-bedded lithic arkosic sandstone forms a dip slope along the southwestern margin of Clallam Bay and is well exposed in Falls Creek 1.1 km southeast of Sekiu Point. It is overlain by numerous beds of thin- to medium-bedded siltstone and sandstone, exposed along the beach at low tide. The Falls Creek unit is more than 30 m thick and is separated from the underlying Jansen Creek Member by an estimated 450 meters of poorly exposed thin-bedded sandstone and siltstone of the Makah Formation (fig. 4). Although the unit is poorly exposed, it probably extends along strike for about 5 km from Falls Creek westward to the junction of State Highway 112 and the Ozette Road. Here a poorly exposed sandstone bed more than 5 m thick is exposed in the roadcut.

Unnamed sandstone unit

A 15-m-thick unnamed unit, composed of approximately 11 beds 0.15- to 1.5-m-thick of fine-grained sandstone, crops out between the Hoko and Little Hoko Rivers (fig. 3) approximately 75 m stratigraphically below the Baada Point Member (fig. 5). The sandstone beds display sharp to amalgamated contacts and convolute bedding. The unit may correlate with a similar thick 11-bed unit that crops out in the Clallam River about 8 km east and perhaps with thicker sandstone units exposed below the Baada Point Member in Charley Creek about 5.5 km east.

PETROGRAPHY

Sandstone of the Makah Formation and its members ranges in composition from basaltic to arkosic wacke and arenite but is predominantly lithic arkose (fig. 12). It is generally fine to medium grained and is compositionally and texturally immature to sub-mature. Most sandstone in the Makah is moderately sorted (range: very poorly sorted to moderately well

sorted) and is composed of angular to subangular grains.

Matrix abundance of both detrital and diagenetic varieties ranges from a trace to 22 percent. The more abundant diagenetic matrix was formed from *in situ* chemical alteration and crushing of volcanic-rock grains (pseudomatrix of Dickinson, 1970). The matrix consists of celadonite, chlorite, and scattered angular silt-size quartz, feldspar, and mica flakes.

Common cementing materials are calcite in the concretionary sandstone (10 to 50 percent; fig. 13) and diagenetic clays (including chlorite and celadonite) in the more abundant wacke. Other cements include hematite, limonite, and sparse laumontite. Sparry calcite cement apparently formed early in the burial history of the concretionary sandstone prior to formation of diagenetic clay matrix. Framework clasts are supported and partially replaced and embayed by the surrounding pore-filling calcite cement. The number of grain-to-grain contacts and percentage of clay matrix are abnormally low. As a result, the calcareous sandstone is compositionally arenite. Minor chlorite and/or celadonite cements are present as coatings on lithic grains and as pore fillings in some sandstone. Plagioclase grains in some samples are partially laumontized, and laumontite occurs locally as cement that preserves chlorite or celadonite coats.

Using a classification modified from Williams, Turner, and Gilbert (1954), sandstone in the Baada Point and Dtokoah Point members and in the intervening thin-bedded strata of the Makah is chiefly lithic




Figure 12.--Ternary diagram showing composition of sandstone in the Makah Formation and its members. Modified from Williams, Turner, and Gilbert (1954).




Figure 13.--Calcite-cemented biotite-bearing arkosic sandstone in the Third Beach Member. Horizontal field of view, 3.5 mm (crossed nicols).

arkosic arenite and wacke (figs. 12 and 14). Major framework constituents are quartz, feldspar, volcanic rock fragments, and micas together with minor amounts of heavy minerals and metamorphic rock fragments.

Subangular to subrounded quartz is the most abundant grain type, making up 10 to 34 percent of the sandstone. Quartz varieties include strained and unstrained monocrystalline quartz, polycrystalline quartz, and chert. Tiny fluid inclusions containing oscillating bubbles (2-phase fluid inclusions) are present in some monocrystalline quartz grains.

Feldspar constitutes 13 to 31 percent of the sandstone; it includes both albite-twinning plagioclase ( $An_{3-64}$ ) and minor potassium feldspar. The plagioclase-to-potassium feldspar ratio is approximately 4:1. Orthoclase and microcline are the most abundant potassium feldspars. Lithic fragments are mainly basalt and andesite (3 to 10 percent), celadonite-replaced amygdaloidal basalt, metabasalt (greenschist facies), and silicic volcanic fragments. Other lithic fragments include minor elongate carbonaceous phyllite clasts, quartz mica schist, and metaquartzite fragments. Biotite flakes form 1 to 4 percent of the sandstone, chlorite flakes 1 to 7 percent. Traces of muscovite are present. Heavy minerals are dominated by clinozoisite, actinolite, epidote, zircon, pink and colorless garnet, hypersthene, apatite, magnetite, and pyrite.

Sandstone in the Klachopis Point and Third Beach Members ranges from micaceous arkosic to lithic arkosic arenite and wacke (fig. 12). The sandstone contains less matrix, has a somewhat higher abundance of quartz, micas, and feldspar, and contains fewer lithic fragments than sandstone in the Baada Point and Dtokoah Point Members (compare figs. 13 and 14). In general, the composition and relative abundance of the different lithic and mineral types in all four of these members are nearly the same except that sandstone in the Third Beach Member consistently contains more coarse-grained biotite flakes than sandstone in the other members.

The shallow marine sandstone in the Jansen Creek Member differs significantly from the turbidite sand-




Figure 14.--Lithic arkosic wacke from Baada Point Member. Horizontal field of view, 1.25 mm (crossed nicols).

stone members in composition and texture; it is volcanic arenite (fig. 12) and is generally coarser grained. Metabasalt (greenschist facies) and basalt fragments make up as much as 24 percent of this sandstone. Other characteristic grain types are monocrystalline quartz, plagioclase, silicic volcanic rock fragments, actinolite (to 5 percent), clinozoisite, chlorite, and molluscan shell fragments. Minor constituents include biotite, augite, pyrite, magnetite, epidote, and algal stromatolites. Calcite cement is ubiquitous and constitutes as much as 53 percent of the rock.

Granule-and-pebble basalt clasts in conglomerate of the Jansen Creek Member are rounded to subrounded. Basalt, gabbro, and metabasalt are the predominant clast types. Other pebble constituents include minor silicic volcanic rock, metatuff, and molluscan shell fragments.

The Carpenter Creek Tuff Member is composed of abundant sickle and rarer bubble wall-glass shards set in a devitrified siliceous clay matrix. Shards are commonly altered to a cherty groundmass. Other constituents include scattered grains of plagioclase, monocrystalline quartz, biotite, muscovite, and chlorite flakes, rare cherty siliceous volcanic fragments, hypersthene, ilmenite grains altering to limonite, and clinozoisite. Patches of secondary sparry-calcite cement are locally abundant. Deposition of this rhyodacitic ash in a marine environment is indicated by the presence of a few foraminifers.

## RESERVOIR POTENTIAL OF TURBIDITE SANDSTONE MEMBERS

Three surface samples from thick amalgamated turbidite sandstone of the Baada Point and Klachopis Point Members have relatively low permeabilities and moderately low effective porosities on the basis of quantitative laboratory analyses. The low porosities and permeabilities probably result from the abundance of detrital and diagenetically formed clay matrix and the calcite and less common limonite, hematite, and laumontite cements that have almost completely filled

the pore spaces between framework grains. This make-up is particularly evident in the two matrix-rich samples of the Baada Point Member, each with moderately low porosity (20.4 to 20.7 percent) and low permeability 2.0 to 7.5 millidarcies), and probably in the calcareous concretionary zones of all the members.

The samples from the Klachopis Point Member has the highest effective porosity and a moderately high permeability (24.6 percent and 657 millidarcies) and also tends to be more friable and cleaner in thin section than the sandstone from the Baada Point and Dtokoah Point Members. It could act as a permeable reservoir rock in the subsurface. At one locality, a 1-m-thick sandstone bed in the Klachopis Point Member emitted a petroliferous odor when freshly broken. The arkosic sandstone of the Third Beach Member, like the sandstone of the Klachopis Point Member, generally contains less pore-filling clay matrix than the more lithic sandstone of the Baada Point, Dtokoah Point, and Jansen Creek Members and may represent an important permeable unit.

All the amalgamated sandstone members are enclosed by thick (hundreds of meters) sequences of thin- to very thin-bedded siltstone and turbidite sandstone that could act as both cap rocks and contiguous source rocks in the subsurface. Hydrocarbon analyses of surface samples by George Claypool (reported in Snively, Pearl, and Lander, 1977, table 3), however, indicate that upper Eocene and Oligocene siltstone in this area is generally geothermally immature, containing only minor quantities of light hydrocarbons.

The Klachopis Point Member, being among the thickest, most widespread, and highest in permeability and porosity, and to a much lesser extent the Baada Point Member, probably have the highest reservoir potential. The Third Beach and Dtokoah Point Members have much more limited distribution and therefore lower reservoir capacity (particularly the thinner bedded sandstone of the Dtokoah Point Member). The olistostromal shallow-marine sandstone and conglomerate of the Jansen Creek Member are too discontinuous, chaotically arranged, and tightly cemented by calcite to represent an important reservoir unit in the subsurface.

Onshore, where the homoclinal sequence is breached by erosion, the petroleum potential of the four northward-dipping turbidite members is limited. Stratigraphic traps may exist in the lower part of the Makah Formation, where it laps onto the broad anticlinal high in the central part of the study area. As these units dip and strike beneath the Strait of Juan de Fuca, structural and/or stratigraphic traps may exist in this area.

## AGE AND REGIONAL CORRELATIONS

Foraminiferal assemblages, sparse and varied throughout the Makah Formation, indicate that the sequence ranges in age from late Eocene to late Oligocene. Foraminifers from the lower part of the Makah at Waadah Island (table 1) indicate that the lower 325 meters of section exposed below the Baada Point Member (fig. 4) ranges from the upper Narizian Stage of

Mallory (1959) to the Refugian Stage of Schenck and Kleinpell (1936).

On the basis of the known range of species within the Tertiary of the Pacific Northwest (Rau, 1958, 1964, 1966; Fulmer, 1975; Armentrout and Berta, 1977), the 100 meters of the Makah stratigraphically below the Baada Point Member on Waadah Island is best referred to the Refugian (fig. 4). This age assignment is based on the presence of *Melonis halkyardi*, *Elphidium californicum*, *Sigmomorphina* cf. *S. schencki*, and in particular *Ceratobulimina washburni* (table 1). Foraminifers from the interval 100 to 325 m below the Baada Point Member on the east side of Waadah Island have greater affinities for the Narizian Stage than for the Refugian Stage. The high occurrence of *Vulvulina curta*, *Anomalina garzaensis*, *Pleurostomella nutalli*, and *Quinqueloculina goodspeedi* all signify a Narizian age.

The upper part of the Makah Formation, strata between the Third Beach Member and the base of the Pysht Formation (fig. 4), are Oligocene (Zemorrian) in age. The checklist of foraminifers (table 2) shows the occurrence of species in the middle and upper parts of the Makah in the Hoko River reference section (A to B on figs. 3 and 5). *Dentalina quadrulata*, *Uvigerina* cf. *U. gesteri*, and *U. gallowayi* are among diagnostic Zemorrian species occurring in the upper part of the section at Hoko River between localities 7190 and 7609 (table 2, fig. 5).

The precise boundary between the Zemorrian and Refugian Stages is difficult to define in the Hoko River section, as several key species found elsewhere in the Pacific Northwest (Rau, 1958, 1964, and 1966) that define the Zemorrian Stage and the Refugian Stage occur together in a 1200-m-thick stratigraphic interval (table 2, fig. 5, localities 7387 to 7190). If the total assemblage is considered, the Refugian-Zemorrian boundary is best placed between localities 5789 and 5788 (table 2, fig. 5).

Several siltstone samples collected on Waadah Island and near the Little Hoko River contain foraminifers indicative of Ulatisian and early Narizian ages. These anomalous foraminifers are probably reworked from older Eocene strata. Submarine erosion across growing broad anticlinal highs in the pre-Makah strata, as at Cape Flattery and in the central part of the study area (fig. 2), undoubtedly contributed reworked older sediments and their Ulatisian to early Narizian microfossils to the younger strata that were being deposited in synclinal low areas.

Foraminiferal assemblages from the Makah Formation (see tables 1 and 2) clearly indicate deep-water open-sea conditions. Almost all of those foraminifers consistently occurring throughout the formation suggest bathyal conditions. Moreover, most of these taxa support no less than middle bathyal depths. Many, such as *Gyroidina soldanii*, *Stilostomella*, hispid uvigerinids, *Bulimina alsatica*, and *Pullenia bulloides*, probably thrived at lower bathyal depths. The minor but consistent occurrence throughout the formation of planktonic taxa (*Globigerina*) supports open-sea conditions.

MAKAH FORMATION--A DEEP-MARGINAL-BASIN SEQUENCE

Table 1.--Checklist of foraminifers from the Waadah Island section of the Makah Formation

[Symbols of frequency of occurrence: C, common; F, few; R, rare; ?, questionable identification]

Species	Pacific Northwest												
	Reference	Collection No.											
	8278	8277	8322	7135	8330	7250	7134B	8321	7247	7133	7132	8276	
Quinqueloculina spp. . . . .	R	R	R	R	R	R	-	R	R	-	R	R	-
Dentalina cf. D. pauperata (d'Orbigny) . . . . .	F	R	-	R	R	?	-	-	-	R	-	-	-
Stilostomella sp. (large final chamber) . . . . .	R	-	-	-	-	-	-	-	-	-	-	-	-
Nodosaria longiscata d'Orbigny . . . . .	R	R	F	R	R	R	C	F	F	R	F	F	F
Pseudoglandulina inflata (Bornemann) . . . . .	F	?	-	-	-	?	?	-	?	-	R	?	-
Gyroidina soldanii d'Orbigny . . . . .	C	-	-	-	R	-	R	R	F	-	-	R	-
Melonis pompilioides (Fichtel and Moll) . . . . .	?	-	-	-	-	-	-	?	-	-	-	-	-
Fursenkoina (Virgulina) sp. . . . .	R	-	-	-	-	-	-	-	-	-	-	-	-
*Melonis halkyardi (Cushman) . . . . .	F	-	-	-	-	-	-	-	-	-	-	-	-
Cibicides cf. C. lobatulus (Walker and Jacob) . . . . .	R	-	-	?	?	-	-	?	-	-	-	-	-
*Elphidium californicum Cook MS. . . . .	R	-	-	-	-	-	-	-	-	-	-	-	-
Globigerina spp. . . . .	R	F	R	F	R	R	F	-	F	R	R	R	R
Guttulina irregularis d'Orbigny . . . . .	R	-	-	-	-	-	-	-	-	-	-	-	-
Eggerella bradyi (Cushman) . . . . .	?	-	R	?	R	-	-	R	?	R	-	R	-
Karrerriella chilostoma (Reuss) . . . . .	R	R	-	-	-	-	-	-	-	-	-	-	?
*Cassidulina galvinensis Cushman and Frizzell . . . . .	?	-	-	-	?	-	-	-	-	-	-	-	-
Globocassidulina globosa (Hantken) . . . . .	?	-	-	-	-	-	-	R	?	-	-	R	-
Cibicides cf. C. elmaensis Rau . . . . .	R	F	-	R	-	-	-	F	R	-	R	-	-
*Ceratobulimina washburni Cushman and Schenck . . . . .	R	-	-	-	-	-	-	-	-	-	-	-	-
Cibicides spp. . . . .	R	R	R	R	R	R	R	-	-	R	-	-	-
Spiroloculina cf. S. texana Cushman and Ellisor . . . . .	-	R	-	-	-	-	-	-	R	R	-	-	-
*Sigmomorphina cf. S. schencki Cushman and Ozawa . . . . .	-	R	-	-	-	-	-	-	-	-	-	-	-
Vaginulinopsis saundersi (Hanna and Hanna) . . . . .	-	R	-	-	-	-	-	-	-	-	-	-	-
Bulimina cf. B. instabilis Cushman and Parker . . . . .	-	R	-	-	F	-	?	-	-	?	-	-	-
?Alabamina sp. . . . .	-	R	-	-	-	-	-	-	-	-	-	-	-
Anomalina cf. A. californiensis Cushman and Hobson . . . . .	-	R	-	-	R	-	-	-	R	F	R	R	-
Eponides umbonatus (Reuss) . . . . .	-	?	-	-	?	-	R	R	R	R	-	F	-
Pullenia bulloides d'Orbigny . . . . .	-	?	-	-	R	-	?	R	F	R	-	F	-
Globobulimina cf. G. pacifica Cushman . . . . .	-	R	?	R	-	?	-	-	-	-	-	-	-
Martinottiella cf. M. nodulosa Cushman . . . . .	-	-	R	-	-	-	-	-	-	-	R	-	-
Sigmoilina sp. . . . .	-	-	R	-	-	-	-	-	-	-	-	-	-
Cornuspira sp. . . . .	-	-	R	-	-	-	-	-	-	-	-	-	-
Stilostomella cf. S. sanctaecrucis (Kleinpell) . . . . .	-	-	F	R	F	?	F	F	R	R	R	R	R
Chilostomelloides cf. C. cyclostoma (Rzehak) . . . . .	-	-	R	-	R	R	-	-	F	F	F	R	-
Cancris joaquinensis Smith . . . . .	-	-	R	?	-	-	?	R	-	-	-	-	-
Gyroidina soldanii d'Orbigny var. (rounded edges) . . . . .	-	-	-	R	-	-	-	R	-	R	F	-	-
*Vulvulina curta Cushman and Siegfus . . . . .	-	-	-	-	-	?	-	F	F	F	-	-	-
Nodosaria clavaeformis Neugeboren . . . . .	-	-	-	-	R	-	-	-	-	-	-	-	-
Nodosaria pyrula d'Orbigny . . . . .	-	-	-	-	R	-	-	-	-	-	-	-	-
Praeglobobulimina cf. P. pyrula (d'Orbigny) . . . . .	-	-	-	-	R	-	-	R	-	-	-	-	-
*Anomalina garzaensis Cushman and Siegfus . . . . .	-	-	-	-	R	-	-	?	-	-	-	-	?

Mollusks are sparse in the deep-water sedimentary rocks of the Makah Formation and occur chiefly in olistostromal blocks of shallow-water basaltic sandstone and conglomerate in the Jansen Creek Member. Addicott (written commun., 1973-1977) identified the following species collected from the Jansen Creek Member on the coast near the southwest corner of sec. 22, T. 33 N., R. 14 W.:

Gastropods:

- Acrilla n. sp. ? aff. A. olympicensis Durham
- Naticid
- ?Perse sp.
- Turritella cf. T. porterensis Weaver
- Undet. fragments - 3 spp.

Table 1.--Checklist of foraminifers from the Waadah Island section of the Makah Formation -- Continued

Species	Pacific Northwest Reference Collection No.										
	8278	8277	8322	7135	8330	7250	7134 <sup>B</sup>	8321	7247	7133	8276
<i>Valvulineria jacksonensis welcomensis</i> Mallory . . . . .	-	-	-	-	-	R	-	-	-	-	-
<i>Plectofrondicularia</i> sp. . . . .	-	-	-	-	-	R	R	-	-	-	-
* <i>Pleurostomella nuttalli</i> Cushman and Siegfus . . . . .	-	-	-	-	-	R	-	-	-	R	?
<i>Gyroidina condoni</i> (Cushman and Schenck) . . . . .	-	-	-	-	-	-	R	-	-	?	-
<i>Lenticulina</i> spp. . . . .	-	-	-	-	-	-	-	R	R	R	-
<i>Plectofrondicularia</i> cf. <i>P. gracilis</i> H. P. Smith . . . . .	-	-	-	-	-	-	-	R	-	?	-
<i>Cibicides spiropunctatus</i> Galloway and Morrey . . . . .	-	-	-	-	-	-	-	?	-	-	-
<i>Cibicides</i> sp. (large, variable, incised sutures, coarse perforate) . . . . .	-	-	-	-	-	-	-	R	F	-	-
<i>Cassidulina</i> sp. (discoid, rounded perf.) . . . . .	-	-	-	-	-	-	-	R	-	-	-
? <i>Fursenkoina</i> ( <i>Virgulina</i> ) sp. (narrow) . . . . .	-	-	-	-	-	-	-	R	-	-	-
<i>Pyrgo</i> sp. . . . .	-	-	-	-	-	-	-	-	R	-	-
<i>Quinqueloculina imperialis</i> Hanna and Hanna . . . . .	-	-	-	-	-	-	-	-	R	-	-
* <i>Quinqueloculina goodspeedi</i> Hanna and Hanna . . . . .	-	-	-	-	-	-	-	-	?	?	R
<i>Pullenia salisburyi</i> R. E. and K. C. Stewart . . . . .	-	-	-	-	-	-	-	-	-	?	-
<i>Praeglobbulimina</i> cf. <i>P. ovata</i> (d'Orbigny) . . . . .	-	-	-	-	-	R	-	-	F	-	?
*Key species											
	Refugian					Narizian					
	Late Eocene										

Pelecypods:

- Conchocele* cf. *C. disjuncta* (Gabb)
- Xrenella* sp.
- Cyclocardia hannibali* (Clark)
- Lucinoma* cf. *L. columbiana* (Clark and Arnold)
- Nemocardium* cf. *N. weaveri* (Anderson and Margin)
- Ostrea* sp.

Scaphopods:

- Dentalium porterenensis* Weaver

Addicott assigned a late Eocene age (Lincoln Stage of Weaver, 1937) to this collection and suggested a middle or outer sublittoral environment of deposition for the basaltic sandstone. Foraminiferal assemblages in the Jansen Creek Member indicate a Refugian (late Eocene) age, a Lincoln equivalent, and a shallow-water depositional environment. The penecontemporaneously deformed member is both underlain and overlain by younger (Zemorrian) strata of the Makah that were deposited at lower to middle bathyal depths. This relation supports the interpretation that the shallow-marine sandstone and conglomerate of the Jansen Creek Member are allochthonous deposits derived from

uplifted older (Refugian) strata along an ancient narrow shelf that bordered the deep marginal basin of the Makah to the north. These shallow-water Refugian deposits were transported into the deep marginal basin by a large submarine slide during Zemorrian time.

The Makah Formation represents part of an upper Eocene to Oligocene deep marginal marine facies that crops out in the northwestern part of the Olympic Peninsula, on western Vancouver Island, in southwesternmost Washington, and in northwestern Oregon but is also widespread on the inner continental shelf of Oregon (Snively and others, 1977). The Makah Formation is in part correlative with upper Eocene and Oligocene strata of the Hoh rock assemblage in the Hoh-Quillayette area of the western coastal area of the Olympic Peninsula (Rau, 1973, 1975, 1979).

Strata similar to the Makah Formation in lithology, age, and depositional environment are exposed in the Hesquiat Peninsula and Nootka Island area along the central part of the west coast of Vancouver Island, British Columbia. This sequence, described in detail by Jeletzky (1954, 1973, 1975) and Cameron (1971, 1972, 1973, 1975, 1979), was named the Hesquiat Formation by Jeletzky (1975).

Shallow-water sandstone, siltstone, and conglomerate that correlate with the Makah Formation fringe the south side of Vancouver Island from Sombrio Point westward to near the entrance to Nitinat Lake. These

Table 2.--Checklist of foraminifers from the Hoko River section of the Makah Formation

[Symbols of frequency of occurrence: C, common; F, few; R, rare; ?, questionable identification.]

Species	Pacific Northwest Reference Collection No.																									
	7609	5190	5423	5783	5784	5785	5786	5182	5187	7190	7191	5787	7192	7193	7194	5788	5789	5790	7195	5791	5726	5186	5792	8242	7387	
<i>Lenticulina</i> cf. <i>L. inornatus</i> (d'Orbigny)	C	R	-	-	-	R	-	-	?	-	R	-	-	-	R	R	-	-	-	-	-	-	R	R	-	-
<i>Marginulina glabra</i> d'Orbigny	?	R	-	-	-	-	-	-	-	-	?	?	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Stilostomella</i> cf. <i>S. sanctaerucis</i> (Kleinpell)	R	-	-	R	?	?	-	-	-	-	R	-	-	R	R	R	R	-	?	-	-	-	?	-	-	-
<i>Dentalina pauperata</i> (d'Orbigny)	?	F	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Dentalina consobrina</i> d'Orbigny	?	F	-	?	-	-	-	-	-	-	-	-	?	-	-	-	-	-	-	?	-	-	?	?	-	-
<i>Nodosaria</i> cf. <i>N. soluta</i> (Reuss)	R	-	-	-	-	-	-	-	-	-	-	-	R	-	-	-	-	-	-	-	-	R	-	-	-	-
* <i>Dentalina quadrulata</i> Cushman and Laiming	F	F	-	R	-	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Sigmoilina</i> sp.	R	-	-	-	R	R	-	-	-	-	-	-	-	-	R	R	-	-	-	-	-	-	-	-	R	-
<i>Stilostomella frizzelli</i> (Rau)	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Stilostomella advena</i> (Cushman and Laiming)	R	R	-	-	-	-	-	-	-	-	-	-	-	?	?	-	-	-	-	-	-	-	-	-	?	-
<i>Eponides umbonatus</i> (Reuss)	?	?	-	-	R	?	-	-	-	-	R	-	-	-	F	R	-	-	-	-	-	-	-	-	-	F
* <i>Anomalina californiensis</i> Cushman and Laiming	F	F	-	R	?	-	-	C	F	F	-	F	?	-	-	-	-	-	?	-	?	-	?	-	-	-
<i>Guttulina</i> cf. <i>G. problema</i> d'Orbigny	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Gyroidina soldanii</i> d'Orbigny	R	F	-	F	-	?	?	?	F	-	F	F	R	R	R	R	-	R	R	R	R	F	R	R	R	-
<i>Praeglobobulimina</i> cf. <i>P. ovata</i> (d'Orbigny)	R	-	-	R	?	-	-	-	-	-	?	-	?	R	-	?	C	F	-	-	F	R	-	-	-	-
* <i>Uvigerina</i> cf. <i>U. gesteri</i> Barbat and von Estorff	R	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Uvigerina garzaensis</i> Cushman and Siegfus	C	R	-	-	F	-	-	-	-	R	-	-	?	R	R	R	R	F	-	-	R	-	C	-	-	-
* <i>Sphaeroidina variabilis</i> Reuss	C	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	R	R	-	-	-	-	-	-	-	-
<i>Globigerina</i> sp.	R	-	R	R	-	R	-	F	R	R	-	R	-	-	-	-	-	-	-	-	R	F	R	R	R	-
<i>Globobulimina</i> cf. <i>G. pacifica</i> Cushman	R	R	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	F	-	-	-	-	-	-	-
<i>Nodosaria longiscata</i> d'Orbigny	-	R	-	R	R	-	R	R	-	R	R	-	R	-	F	R	R	R	F	R	R	R	R	R	R	-
<i>Pseudoglandulina inflata</i> (Bornemann)	-	F	R	R	R	R	R	R	-	-	R	R	-	R	R	F	-	-	-	-	-	-	-	-	-	R
<i>Melonis pompilioides</i> (Fichtel and Moll)	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Fissurina</i> sp.	-	R	-	R	-	-	-	R	R	-	-	-	-	-	-	-	R	-	-	-	R	-	R	-	-	-
* <i>Bulimina alsatica</i> Cushman and Parker	-	F	-	-	R	R	-	-	R	R	R	-	R	R	-	R	R	-	R	?	F	-	F	-	-	-
* <i>Florulis</i> cf. <i>F. incisum</i> (Cushman)	-	F	-	?	?	F	-	-	-	-	?	R	-	-	-	-	-	-	-	-	-	-	R	-	-	-
<i>Quinqueloculina weaveri</i> Rau	-	?	-	-	-	-	-	-	-	-	-	-	F	-	-	-	-	-	-	-	R	-	-	-	-	-
<i>Cassidulina crassipunctata</i> Cushman and Hobson	-	C	-	F	-	?	-	-	?	C	-	F	?	-	-	-	R	-	-	-	-	-	-	-	?	-
<i>Fursenkoina</i> ( <i>Virgulina</i> ) sp.	-	?	-	-	-	-	-	-	-	-	-	-	-	-	-	-	R	-	-	-	-	-	-	-	-	-
<i>Martinottiella</i> cf. <i>M. nodulosa</i> (Cushman)	-	-	R	R	R	-	-	R	-	-	-	-	-	-	-	R	-	-	-	-	R	-	-	-	-	-
<i>Pyrgo</i> spp.	-	-	R	R	-	-	-	-	-	-	-	-	-	-	F	-	R	R	-	F	-	R	-	-	-	-
<i>Epistomina eocenica</i> (Cushman and Hanna)	-	-	?	?	F	-	-	R	-	-	-	-	-	-	-	-	-	-	-	R	-	-	-	-	-	-
<i>Cibicides</i> spp.	-	-	-	R	-	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	R
* <i>Uvigerina gallowayi</i> Cushman	-	-	-	F	C	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
* <i>Cassidulinoides</i> sp.	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	R	-
* <i>Buccella mansfieldi oregonensis</i> (Cushman, R. E. and Stewart, K. C.)	-	-	-	-	?	-	-	?	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

strata were referred to the Carmanah Formation by Clapp (1912) and are mapped and described by Muller (1971, 1977).

The Makah Formation correlates in part with the Lincoln Creek Formation (Weaver, 1937; Beikman and others, 1967) of southwestern Washington, the deep-water marine Blakeley Formation of Weaver (1912) (Fulmer, 1975; McLean, 1977) of the Puget Sound area. In the southern and central Oregon Coast Range, the time-stratigraphic equivalents are the shallow-marine Eugene Formation on the southeastern flank of

the range and the deep-water marine Alsea Formation along the central Oregon coast (Snively and others, 1975). In the northern Oregon Coast Range, the Makah Formation is in part coeval with the shallow-marine Pittsburg Bluff Formation (Schenck, 1927; Moore, 1976), the upper member of the Keasey Formation (Schenck, 1927; Moore and Vokes, 1953) in the north-eastern part of the range, and the lower part of the deep-water marine mulstone of Oswald West on the northwestern flank of the range (Niem and Van Atta, 1973).

Table 2.--Checklist of foraminifers from the Hoko River section of the Makah Formation -- Continued

Species	Pacific Northwest																									
	Reference Collection No.																									
	7609	5190	5423	5783	5784	5785	5786	5182	5187	7190	7191	5787	7192	7193	7194	5788	5789	5790	7195	5791	5726	5186	5792	8242	7387	
<i>Quinqueloculina</i> cf. <i>Q. imperialis</i> Hanna and Hanna . . . . .	-	-	-	-	-	R	-	-	-	-	-	?	-	-	-	R	-	-	?	?	-	-	-	-	-	-
<i>Spiroloculina</i> cf. <i>S. texana</i> Cushman and Ellisor . . . . .	-	-	-	-	-	R	R	-	-	-	-	-	-	-	F	R	-	-	-	-	-	-	-	-	-	-
<i>Chilostomelloides</i> sp. . . . .	-	-	-	-	-	R	-	-	-	-	-	R	-	-	-	R	F	-	-	-	-	-	-	-	-	-
<i>Plectofrondicularia</i> cf. <i>P. gracilis</i> Smith . . . . .	-	-	-	-	-	F	R	-	-	-	-	-	-	-	R	R	R	R	-	R	R	R	-	-	-	-
<i>Plectofrondicularia vaughani</i> Cushman . . . . .	-	-	-	-	-	R	R	-	-	-	R	R	R	-	R	-	R	R	-	?	R	-	R	-	-	-
<i>Bolivina</i> sp. . . . .	-	-	-	-	-	R	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Nodosaria pyrula</i> d'Orbigny . . . . .	-	-	-	-	-	-	R	R	-	-	-	-	-	-	-	-	-	-	-	-	-	R	?	-	-	R
<i>Eggerella bradyi</i> (Cushman) . . . . .	-	-	-	-	-	-	?	-	-	-	-	R	-	-	R	-	-	-	-	-	-	-	-	-	-	-
<i>Pullenia bulloides</i> (d'Orbigny) . . . . .	-	-	-	-	-	-	R	R	-	?	-	-	-	-	R	-	?	R	R	-	R	-	-	R	-	R
<i>Cassidulina</i> cf. <i>C. globosa</i> Hantken . . . . .	-	-	-	-	-	-	R	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Siphonodosaria</i> (?) <i>verneuili</i> (d'Orbigny) . . . . .	-	-	-	-	-	-	-	-	-	F	F	-	-	-	-	-	?	-	F	-	-	-	-	-	-	-
<i>Cassidulina</i> cf. <i>C. kernensis</i> Smith . . . . .	-	-	-	-	-	-	-	-	-	-	R	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Lenticulina</i> cf. <i>L. texana</i> (Cushman and Applin) . . . . .	-	-	-	-	-	-	-	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-
* <i>Lenticulina</i> cf. <i>L. calcar</i> (Linne) . . . . .	-	-	-	-	-	-	-	-	-	-	-	R	-	-	-	-	-	-	?	-	-	-	-	-	-	-
<i>Cibicides</i> cf. <i>C. pseudougerianus</i> <i>evolutus</i> Cushman and Hobson . . . . .	-	-	-	-	-	-	-	-	-	-	-	R	F	?	-	R	F	?	?	?	?	-	-	-	-	-
<i>Gaudryina alazanensis</i> Cushman . . . . .	-	-	-	-	-	-	-	-	-	-	-	-	F	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Guttulina</i> cf. <i>G. franki</i> Cushman and Ozawa . . . . .	-	-	-	-	-	-	-	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Pseudoglandulina</i> aff. <i>P. inflata</i> (Bornemann) . . . . .	-	-	-	-	-	-	-	-	-	-	-	-	F	-	-	-	-	-	-	-	R	-	-	-	-	-
<i>Plectofrondicularia</i> sp. (thick) . . . . .	-	-	-	-	-	-	-	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Quinqueloculina</i> spp . . . . .	-	-	-	-	-	-	-	-	-	-	-	-	-	R	-	-	R	R	R	R	R	R	-	-	R	-
<i>Uvigerina</i> sp. (small, fine costae) . . . . .	-	-	-	-	-	-	-	-	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Karrerella washingtonensis</i> Rau . . . . .	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	R	R	-	?	-	-	-	-	-	-
<i>Lagena</i> cf. <i>L. becki</i> Sullivan . . . . .	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	R	R	-	-	-	-	-	-	-	-
<i>Cyclogyra</i> ( <i>Cornuspira</i> ) cf. <i>C. byramensis</i> (Cushman) . . . . .	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
* <i>Karrerella chilostoma</i> (Reuss) . . . . .	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	R	-	-	-	-
? <i>Elphidium</i> sp . . . . .	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Guttulina problema</i> d'Orbigny . . . . .	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	R	-	-	-	-	-	-
<i>Eponides?</i> sp. . . . .	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	R	-	-	-	-
*? <i>Cibicides hodgei</i> Cushman and Schenck . . . . .	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	R	-	-	-
<i>Glomospira charoides corona</i> Cushman and Jarvis . . . . .	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	R	-
? <i>Anomalina</i> sp . . . . .	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	R
*Key species																										
	Zemorrian											?	Refugian													
	Oligocene											?	Late Eocene													

DEPOSITIONAL ENVIRONMENT

The Makah Formation consists of a 2800-m-thick sequence of thin-bedded turbidite sandstone and siltstone punctuated by four distinct, areally widespread packets of thick-bedded amalgamated turbidite sand-

stone, one olistostromal unit of penecontemporaneously deformed shallow-marine strata, and a thin but distinctive tuff unit. The microfossil assemblages from siltstone beds throughout the formation indicate deposition in an open-marine lower to middle bathyal environment. The paleoecology, total thickness, turbidite

nature, and geometry of this upper Eocene and Oligocene sedimentary sequence indicate rapid deposition in a submarine-fan setting.

Applying the criteria developed by Mutti and Ricci Lucchi (1972, 1975) for recognizing turbidite facies in ancient submarine-fan sequences, the vertical and lateral turbidite facies variations of the Makah Formation best fit the depositional lobe setting of an outer submarine fan. The amalgamated sandstone members appear to have been deposited as lobes; the thick sequences of thin-bedded strata between the sandstone packets have the sedimentary characteristics of basin-plain and outer-fan fringe deposits.

In the turbidite facies scheme of Walker and Mutti (1973), the sandstone strata of the Baada Point, Klachopis Point, and Third Beach Members appear to be a sequence of alternating proximal C and B turbidite facies with minor intervening D facies. The high sandstone-to-siltstone ratio, several thickening-upward cycles (as in the Baada Point Member), abundant  $Tu$ ,  $Tub$ , and  $Tubc$  Bouma sequences, thick even bedding, coarse tail grading, and overall sheetlike geometry are characteristic of turbidite sands deposited as a depositional lobe of an outer submarine fan (Mutti and others, 1978). The minor sequences of thin-bedded turbidite within each thick amalgamated sandstone member (facies D of Walker and Mutti, 1973) may reflect deposition between shifting lobes. Variations in individual sandstone thickness along strike within the members may reflect the influence of submarine topography and other depositional factors (Mutti and others, 1978).

The Dtokoah Point Member, being thinner bedded and less amalgamated (fig. 9) and having more  $Tbcd$  turbidite sequences and a lower sandstone-to-siltstone ratio than the other turbidite sandstone members, may represent a transition from sedimentation on a depositional lobe to sedimentation on an outer-fan fringe. The local thin-bedded channelized strata in this member appear to be a channel margin of interchannel facies of a depositional lobe as described by Mutti (1977). The very thick bedded amalgamated facies B turbidite sandstone units of the Klachopis Point and Third Beach Members at their type sections (fig. 9) with thick intervals of Bouma "a" divisions and some dish structures may reflect a transition from depositional-lobe to a middle-fan channelized facies.

In the thick sequences of thin-bedded sandstone and siltstone strata between the amalgamated sandstone members, the sandstone-to-siltstone ratio is low. Thin to very thin, sharp, even turbidite sandstone beds alternate with siltstone beds, and internal sedimentary structures occur in Bouma  $Tb-e$  through  $Td-e$  sequences. No systematic upward thickening or thinning cycles are recognized. These features are characteristic of sediments deposited in the basin-plain or outer-fan fringe environment (turbidite facies D and G of Walker and Mutti, 1973; Mutti and Ricci Lucchi, 1972; Ricci Lucchi, 1975). Deposition on a slope and subsequent overloading and slumping of the sedimentary pile are indicated by minor intraformational unconformities and by clastic dikes and prolated bedding observed throughout the formation.

Higher in the section, conglomerate in the olistostromal blocks of the Jansen Creek Member that contains a displaced Refugian fauna consists of well-rounded basalt clasts that are well bedded, moderately sorted, and imbricated and contain scattered broken molluscan valves. These features suggest that the conglomerate was originally deposited in a littoral zone in Refugian time. Massive bioturbated basaltic sandstone that is interstratified with the conglomerate contains coquina of disarticulated fossil shells and scattered articulated gastropods, pelecypods, particularly *Ostrea*, and stromatolitic algae. These characteristics and faunas (Addicott, written communis., 1972-1977) indicate that sand deposition occurred under fluctuating high-energy conditions at neritic water depths.

At the present time, no lithologically similar shelf facies of Refugian age is exposed on the southwestern part of Vancouver Island, but the Oligocene section there is incompletely preserved. The nearest outcrops of compositionally similar thick-bedded basaltic sandstone and conglomerate containing molluscan coquinas and metabasalt clasts are in the younger Sooke Formation of southern Vancouver Island (Clapp, 1912, 1913; Clapp and Cooke, 1917), an upper Oligocene and lower Miocene strandline formation, also derived from erosion of the underlying Eocene Metchosin Volcanics (correlative of the Crescent Formation). The Sooke Formation depositional setting may therein be analogous to the environment that produced the compositionally and lithologically similar, but older, Refugian Jansen Creek Member that slid into the Makah marginal basin in Zemorrian time.

The olistostromal blocks and penecontemporaneous folds of the Jansen Creek Member form a thin stratigraphic unit that is traceable for 9 km. The linear trend of this olistostromal belt parallel to the presumed shelf margin suggests that the lithified blocks slid off an ancient fault scarp or elongate high along the northern margin of the basin of deposition of the Makah Formation.

## PALEOGEOLOGY

During late Eocene and Oligocene time, deep-water sediments were deposited in the Pacific Northwest in several marginal basins that may have been interconnected but had different sediment-source areas (Snively and Wagner, 1963). The Makah Formation is part of this deep marginal basin facies that now crops out only in the northwestern Olympic Peninsula, in southwestern Washington and northwestern Oregon, and along the coast of central western Vancouver Island. On the basis of limited subsurface data from exploration wells, correlative deep-marginal-basin deposits of late Eocene and Oligocene age also underlie the inner continental shelf of Oregon (Snively and others, 1977) and the Tofino basin along the western side of Vancouver Island (Shouldice, 1971). In all but the northernmost basin, Tofino-Fuca basin (fig. 15), the upper Eocene and Oligocene strata consist chiefly of deep-water siltstone with minor sandstone interbeds. The Tofino-Fuca basin in which the Makah occurs was unique because of the abundance of turbidite sands deposited in it.




Figure 15.--Paleogeographic map showing the inferred margins of the Tofino-Fuca basin relative to other deep-water marginal basin facies in the Pacific Northwest during the late Eocene and Oligocene. Modified from Snively and others (1975). B, Brooks Peninsula; N, Nootka Island; H, Hesquiat Peninsula; BS, Barkley Sound; PSJ, Port San Juan; LR, Lyre River; O, Ozette Island; MCB, Minter Creek Basin; and F, Forks. Numbers represent Shell Canada Ltd. off-shore exploration wells: 1, Apollo; 2, Zeus I-65; 3, Zeus D-14; 4, Pluto; and 5, Prometheus.

The Tofino-Fuca basin (fig. 15), an elongate deep, narrow depositional basin or trough, is inferred to have extended from the Kyuquot uplift near the Brooks Peninsula in northern Vancouver Island (Tiffin and others, 1972) southeastward to, and perhaps east of, the Lyre River on the central part of the northern flank of the Olympic Peninsula, a distance of more than 350 km (fig. 15). The southwestern margin of the basin, though now difficult to delineate because of post-early Miocene deformation, erosion, and cover by younger strata, may once have been along a welt of lower and middle Eocene volcanic rocks. This welt may have been defined by the Prometheus magnetic high off southwestern Vancouver Island (Shouldice, 1971; MacLeod and others, 1977) and a ridge formed by volcanic rocks of the Crescent Formation along the northwestern flank of the Olympic Mountains (figs. 1 and 2). Upper Eocene and Oligocene deep-water marine turbidite strata on Nootka Island and the Hesquiat Peninsula (Cameron, 1973, 1975) suggest that the northeastern margin of the late Eocene and Oligocene basin probably was several kilometers shoreward of the present coastline of Vancouver Island between the Brooks Peninsula and Barkley Sound (Tiffin and others, 1972). Southeast of the sound, this margin must have been southwest of the present shoreline, for neritic strata of late Eocene and Oligocene age (Carmanah Formation of Clapp, 1912; see also Muller, 1971) crop out locally along the Vancouver coast between Barkley Sound and Sombrio Point. This deep marginal basin would therefore encompass the Tofino basin on the Vancouver shelf (Shouldice, 1971) and the Fuca basin along the northwestern flank of the Olympic Mountains (fig. 15).

Paleobathymetry based on foraminiferal assemblages suggests that the Fuca part of this marginal basin was deep from late Eocene through late Oligocene time during deposition of the Makah Formation (fig. 2) and the overlying Pysht Formation (fig. 2). Shoaling and filling in early Miocene time were marked by deposition of shallow-water and nonmarine strata of the overlying Clallam Formation (Gower, 1960; Addicott, 1976).

#### PALEODISPERSAL PATTERN

Dispersal of clastic material into the Olympic part of the Tofino-Fuca basin during Makah time was from two distinct directions. The turbidite sandstone was introduced from the northwest by longitudinal filling from sources that may have existed between Barkley Sound and the Hesquiat Peninsula (fig. 15). The other more northerly dispersal direction is represented by submarine landslide blocks in the Jansen Creek Member that were derived from slumping of shallow-water marine strata off the southern Vancouver Island shelf margin. The abundant basalt and metabasalt clasts in sandstone and conglomerate of the Jansen Creek Member were most likely derived from the locally metamorphosed basalt in the Metchosin Volcanics that crops out along the southern margin of Vancouver Island.

A northwestern source area for the four thick amalgamated turbidite members of the Makah Formation and the intervening thin-bedded strata is indicated by




Figure 16.--Rose diagrams showing orientation of flute casts and groove casts in the Makah Formation and three of its members. Bearings are plotted to the nearest 30° after correcting for tectonic tilt of beds.

a general coarsening and thickening of turbidite sandstone beds and increase of total thickness of each member in that direction and because paleocurrent directions of flute marks in these units display a south to southeastward dispersal pattern parallel to that of groove cast orientation (fig. 16). It is possible that post-Makah tectonic rotation may have had minor effect on these observed paleodispersal trends.

Projecting the dispersal pattern and thickening trend of the Makah turbidites northwestward, thick turbidite sandstone beds are expected to occur in the Tofino basin. The Hesquiat Formation, which crops out on the Hesquiat Peninsula and Nootka Island area of western Vancouver Island (the northeasternmost part of the late Eocene and Oligocene Tofino-Fuca basin; fig. 15), lies along this northwest trend. This correlative unit consists predominantly of a sequence of sandstone and siltstone more than 1500 m thick and minor channel conglomerate (Jeletzky, 1954, 1973, 1975; Cameron, 1971, 1972, 1973, 1975).

Foraminifers from stratigraphic sections from the Hesquiat Formation along the Vancouver Island coast kindly loaned to the U.S. Geological Survey by Mobil Oil Canada, Ltd., are late Eocene (Refugian) and Oligocene (Zemorrian) in age. These microfossils indicate cold deep-water conditions that ranged from upper bathyal to middle bathyal.

The sandstones of the Hesquiat and Makah Formations have similar amounts of quartz, feldspar, muscovite, and biotite. The proportions of lithic components are nearly identical; basalt, andesite, and felsic volcanic-rock fragments predominate over phyllite and schist clasts.

The marked similarities in age, depositional environment, and lithology of the two units suggest that these strata are coeval and were deposited in the same deep marginal basin. Accordingly, from paleocurrent patterns in the Makah Formation and from age and lithologic data, the turbidite sandstone of the Makah Formation was transported along the axis of the Tofino-Fuca deep marginal basin, possibly from sources between the Hesquiat Peninsula and Barkley Sound area (fig. 15). The Fuca part of the basin in which the Makah Formation occurs was apparently deeper, as it acted as a sink for turbidite deposits and as a result is 1000 m thicker than the Hesquiat sequence that Cameron (1979) considers to represent proximal turbidites.

A difficulty with this paleogeographic reconstruction is that thick sandstone units are absent in two offshore wells drilled in the west-central part of the Tofino basin between the outcrops of upper Eocene and Oligocene sandstone on the Hesquiat and Olympic Peninsulas. These two wells, Shell Canada Ltd. Zeus I-65 and Pluto (wells 2 and 4 on fig. 15), penetrated a predominantly deep marine siltstone section of late Eocene and Oligocene age (Shouldice, 1971). The absence of upper Eocene and Oligocene turbidite sandstone in these two test wells can be interpreted in three ways: (1) a structural high existed along the southwestern margin of the Tofino basin during the late Eocene and Oligocene, and turbidite sands were not deposited across it; (2) most sands from the Hesquiat Peninsula source area bypassed the Tofino basin and were deposited as a turbidite facies in the deeper part of the marginal basin (Fuca basin) to the southeast. Because the upper Eocene and Oligocene sequence in the study area is almost twice as thick as that reported by Shouldice (1971) for the correlative sequence in the Tofino basin, this part of the deep marginal basin must have subsided at a faster rate than the Tofino basin during the late Eocene and Oligocene, forming a natural depositional sink for turbidite sand transported through the Tofino basin; (3) post middle Miocene left-lateral strike-slip movement along the offshore extension of the Calawah fault (fig. 2), which lies along the seaward margin of the Tofino-Fuca basin (MacLeod and others, 1977), offset correlative turbidite sandstone of the Makah south of the fault to the southeast and brought a predominantly siltstone sequence into the position of Shell Canada's Zeus I-65 and Pluto wells (fig. 15).

The third hypothesis is supported by the presence of infolded strata and blocks (broken formation) of Oligocene and lower Miocene turbidite sandstone that occur in melange along the northwestern Olympic coast as on Ozette Island and in Minter Creek basin, 14 km south of Forks (fig. 15). Some of these infolded sandstone masses and blocks are compositionally similar to the micaceous quartzo-feldspathic sandstone of the Makah Formation. We suggest that these Oligocene

turbidite sandstone units were originally deposited along the southwestern margin of the Tofino basin and were offset to their present locations by left-lateral movement along the Calawah fault (fig. 2) in post-middle Miocene time. If this interpretation is correct, then the Oligocene section penetrated in Shell Canada's Pluto and Zeus I-65 exploratory wells (Shouldice, 1971) would lie south of the offshore extension of the Calawah fault (MacLeod and others, 1977) and was deposited farther northwest in the basin, away from the major source of the coarse clastic detritus derived in the Hesquiat Peninsula source area.

A 24-channel seismic reflection profile collected by the senior author aboard the U.S. Geological Survey research vessel *S. P. Lee* across the offshore extension of the Calawah fault about 40 km northwest of Cape Flattery further supports the strike-slip hypothesis. This profile shows more than 2500 meters of northeastward-dipping strata on the northeast side of the fault to be unconformably overlain by a gently deformed unit of Miocene(?) and Pliocene(?) age. The 2500-m section lies along the strike of the thick sedimentary sequences of the Hoko and Makah Formations on land. On the southwest side of the fault, this sequence of probable late Eocene and Oligocene age is missing, and the Miocene and Pliocene strata rest unconformably on a high-velocity acoustic basement, probably basalt of the Crescent Formation that forms the Prometheus magnetic anomaly (MacLeod and others, 1977). Shell Canada's Prometheus test well (fig. 15) drilled 30 km north of the seismic profile (near the axis of the Prometheus magnetic anomaly) penetrated 1785 meters of Miocene and younger strata unconformably overlying basalt similar to that of the Crescent Formation (Shouldice, 1971). Southwest of the Calawah fault, the upper Eocene and Oligocene turbidite strata either were never deposited, were removed by erosion following post-Oligocene uplift (Shouldice, 1971; Tiffin and others, 1972), or were displaced southeastward to the west coast of the Olympic Peninsula by left-lateral movement along the Calawah fault prior to the unconformable deposition of the upper Miocene and younger strata on the Crescent volcanic basement.

#### SOURCE AREAS

The abundant strained and unstrained monocrystalline quartz, coarse-grained flakes of biotite, especially in the Klachopis Point and Third Beach Members, and grains of sodic and calcic plagioclase, orthoclase, and microcline in the turbidite sandstone indicate a volumetrically important dioritic to granitic source for the Makah Formation. Extensive Eocene plutons and the middle Jurassic Island Intrusions (Muller, 1971, 1977; Carson, 1972) crop out on Vancouver Island east of the Barkley Sound-Hesquiat Peninsula area. These rocks may have been the granodiorite and gneissic sources for the quartzo-feldspathic and micaceous turbidite sandstone of the Makah. Pearl (1977), in a scanning electron microscope study, noted the close similarity of Fe/Mg ratios of biotite in the turbidite sandstone beds of the Makah and biotite from a granodiorite intrusive body midway between Barkley Sound and Port San Juan.

The subordinate amount of basalt, andesite, meta-basalt, and silicic volcanic fragments and associated heavy minerals in the turbidite sandstone units of the Makah also indicates a contribution of lithic detritus from a volcanic terrane. On Vancouver Island, possible sources for the Makah include the Paleozoic Sicker and the Mesozoic Bonanza and Karmutsen Groups that surround the granodiorite intrusive bodies north of the San Juan fault (Muller, 1971, 1977). A southwestward dispersal pattern noted in the coeval Hesquiat Formation (Cameron, oral commun., 1978) suggests that west-central Vancouver Island probably was a major source area for the lithologically similar turbidite sandstone of the Makah (fig. 15).

The only other nearby source area likely to have been exposed during the late Eocene and Oligocene lay due north or northeast across the Strait of Juan de Fuca on southern Vancouver Island. If the paleodispersal of the turbidite sandstone of the Makah had been from the northeast before Olympic tectonism, graphitic schist and phyllite detritus from the Paleozoic Leech River Formation and basalt from the Eocene Metchosin Volcanics that form much of southern Vancouver Island (Muller, 1971, 1977) should be more abundant in these sandstone beds. This terrane does appear to be a source of detritus in the underlying Hoko River Formation, which contains abundant phyllite and basalt clasts in sandstone and conglomerate channels. Although these source rocks probably were exposed during Makah time, as in places they are major constituents in the coeval nearshore sedimentary rocks on southern Vancouver Island and the basaltic sandstone of the Jansen Creek Member, the mineralogy of the sandstone of the Makah suggests little contribution from these sources. A source area directly to the north is unlikely for the turbidite of the Makah because the coeval neritic facies rocks that crop out along the coast between Port San Juan and Barkley Sound (fig. 15) are predominantly of neritic massive tuffaceous siltstone and well-bedded to massive concretionary sandstone that contain abundant mollusks.

A volumetric problem exists when one attempts to explain the high abundance of framework minerals derived from dioritic and granitic rocks relative to volcanic and metamorphic clasts in the sandstone of the Makah Formation because the crystalline basement of Vancouver Island consists chiefly of Paleozoic and Mesozoic volcanic and metamorphic rocks (Muller, 1971, 1977). Possibly the fine grain size of the sandstone of the Makah precludes the presence of many lithic fragments from other Vancouver source rocks, since metamorphic, volcanic, and intrusive clasts commonly are chemically altered to clays before they are reduced to a fine sand size (Pettijohn, 1975). The large Mesozoic Coast Range batholith on the British Columbia mainland may be an alternative source for the thick micaceous quartzo-feldspathic sandstone of the Makah. Detritus shed from this upland area in late Eocene and Oligocene time may have been transported via a major westward flowing river(s) across a low-lying ancestral Vancouver terrane to the late Eocene and Oligocene shoreline somewhere east of the present Hesquiat Peninsula-Barkley Sound area, and hence by turbidity flows into the Tofino-Fuca basin.

## SUMMARY

In late Eocene and Oligocene time, a submarine fan that now forms the Makah Formation prograded over the hemipelagic siltstone and rarer interbedded phyllitic sandstone and conglomerate channels of the underlying upper Eocene (Narizian) sequence (Hoko River Formation, fig. 2), forming a gradational contact. The geometry of the sandstone packets, together with the paleocurrent orientations, suggests that growth of the fan was by sand and silt turbidity flows that swept southeastward down the axis of the Tofino-Fuca basin (fig. 15).

A well-developed active submarine channel system on the upper and middle fan (presumably located northwest of the study area) temporarily funneled many sheetlike high-density flows into the outer-fan depositional environment in rapid succession to form thick widespread depositional lobes. Several changes in the channel system led to shifting of lobe sedimentation over outer-fan deposits to form the interstratification of thin-bedded outer-fan and basin-plain strata with four thick amalgamated sandstone members. The farthest extent of the prograding depositional lobe was the Klachopis Point Member, which spread across and beyond the study area.

A local submarine unconformity in the lower part of the Makah Formation that formed prior to deposition of the Klachopis Point Member was produced by minor compressional folding and faulting of underlying Eocene sedimentary and volcanic rocks as well as the lower part of the Makah Formation locally.

In middle(?) Zemorrian time, deposition of the Makah submarine-fan--basin-plain strata was interrupted by a large landslide that carried olistostromal blocks of Refugian basaltic conglomerate and sandstone into the basin from the continental shelf to the north, thus producing the deformed zone of sedimentary rocks, the Jansen Creek Member. Basin-plain--outer-fan sedimentation of the Makah resumed after the Jansen Creek Member episode and was temporarily interrupted by progradation of a minor depositional lobe that formed the unnamed amalgamated turbidite unit at Falls Creek.

Outer-fan and basin-plain deposition of the upper part of the Makah Formation was followed by the accumulation of the conformably overlying upper Oligocene conglomerate, sandstone, and siltstone of the Pysht Formation (fig. 2). Clasts in these deep-water channel conglomerates and thick-bedded sandstones were derived from metamorphic and volcanic sources and from a neritic fossiliferous sandstone that lay northward across the Strait of Juan de Fuca on southern Vancouver Island.

## REFERENCES CITED

- Addicott, W. O., 1976, Molluscan paleontology of the lower Miocene Clallam Formation, northwestern Washington: U.S. Geological Survey Professional Paper 976, 44 p.
- Armentrout, J. M., and Berta, Annalisa, 1977, Eocene-Oligocene foraminiferal sequence from the northeast Olympic Peninsula, Washington: *Journal of Foraminiferal Research*, v. 7, no. 3, p. 216-233.
- Arnold, Ralph, and Hannibal, Harold, 1913, The marine Tertiary stratigraphy of the North Pacific Coast of America: *American Philosophical Society Proceedings*, v. 52, no. 212, p. 559-605.
- Beikman, H. M., Rau, W. W., and Wagner, H. C., 1967, The Lincoln Creek Formation, Grays Harbor basin, southwestern Washington: U.S. Geological Survey Bulletin 1244-I, 14 p.
- Boggs, Sam, Jr., 1972, Petrography and geochemistry of rhombic, calcite pseudomorphs from mid-Tertiary mudstones of the Pacific Northwest, U.S.A.; *Sedimentology*, v. 19, p. 219-235.
- Bouma, A. H., 1962, *Sedimentology of some flysch deposits*: Amsterdam, Elsevier, 168 p.
- Brown, R. D., Jr., and Gower, H. D., 1958, Twin River Formation (redefinition), northern Olympic Peninsula, Washington: *American Association of Petroleum Geologists Bulletin*, v. 42, no. 10, p. 2492-2512.
- Cameron, B. E. B., 1971, Tertiary stratigraphy and microfaunas from the Hesquiat-Nootka area, west coast, Vancouver Island (92E): *Geological Survey of Canada Paper 71-1, Pt. B*, p. 91-94.
- \_\_\_\_\_, 1972, Tertiary foraminiferal succession of the western Cordillera and Pacific margin: *Geological Survey of Canada Paper 72-1, Pt. A*, p. 198-201.
- \_\_\_\_\_, 1973, Tertiary stratigraphy and microfaunas from the Pacific margin, west coast Vancouver Island: *Geological Survey of Canada Paper 73-1, Pt. A*, p. 19-20.
- \_\_\_\_\_, 1975, Geology of the Tertiary rocks north of latitude 49°, west coast of Vancouver Island: *Geological Survey of Canada Paper 75-1, Pt. A*, p. 17-19.
- \_\_\_\_\_, 1979, Early Cenozoic paleogeography of Vancouver Island, British Columbia, in Armentrout, J. M., Cole, M. R., TerBest, H., Jr., eds., *Pacific Coast Paleogeography Symposium*, v. 3, Cenozoic paleogeography of the Western United States: *Pacific Section Society of Economic Paleontologists and Mineralogists*, Los Angeles, California, p. 326.
- Carson, D. J. T., 1972, The plutonic rocks of Vancouver Island, British Columbia - their petrography, chemistry, age, and emplacement: *Geological Survey of Canada Paper 72-44*, 70 p.
- Clapp, C. H., 1912, *Southern Vancouver Island: Geological Survey of Canada Memoir 13*, 208 p.
- \_\_\_\_\_, 1913, *Geology of the Victoria and Saanich map areas, Vancouver Island, British Columbia: Geological Survey of Canada Memoir 36*, 143 p.
- Clapp, C. H., and Cooke, H. C., 1917, *Sooke and Duncan map areas, Vancouver Island, British Columbia: Geological Survey of Canada Memoir 96*, 445 p.
- Dickinson, W. R., 1970, Interpreting detrital modes of graywacke and arkose: *Journal of Sedimentary Petrology*, v. 40, p. 695-707.
- Fulmer, C. V., 1975, Stratigraphy and paleontology of the type Blakeley and Blakeley Harbor Formations, in Weaver, D. W., Hornaday, G. R., and Tipton, Ann, eds., *Conference on future energy horizons of the Pacific Coast, Paleogene symposium and selected technical papers: American Association of Petroleum Geologists, Society of Economic Paleontologists and Mineralogists, and Society of Economic Geophysicists, Annual Meeting, Pacific Sections, Proceedings*, p. 210-271.
- Glassley, William, 1974, *Geochemistry and tectonics of the Crescent volcanic rocks, Olympic Peninsula, Washington: Geological Society of America Bulletin* 85, p. 785-794.
- Gower, H. D., 1960, *Geologic map of the Pysht quadrangle, Washington: U.S. Geological Survey Quadrangle Map GQ-129, scale 1:62,500*.
- Jeletzky, J. A., 1954, Tertiary rocks of the Hesquiat-Nootka area, west coast of Vancouver Island, British Columbia: *Geological Survey of Canada Paper 53-17*, 65 p.
- \_\_\_\_\_, 1973, Age and depositional environments of Tertiary rocks of Nootka Island, British Columbia, (92-E): Mollusks versus foraminifers: *Canadian Journal of Earth Sciences*, v. 10, no. 3, p. 331-365.
- \_\_\_\_\_, 1975, Hesquiat Formation (new): a neritic channel and interchannel deposit of Oligocene age, western Vancouver Island, British Columbia: *Geological Survey of Canada Paper 75-32*, 54 p.
- Kleinpell, R. M., 1938, *Miocene stratigraphy of California: Tulsa, American Association of Petroleum Geologists*, 450 p.
- Mallory, V. S., 1959, *Lower Tertiary biostratigraphy of the California Coast Ranges: Tulsa, American Association of Petroleum Geologists*, 416 p.
- MacLeod, N. S., and Snaveley, P. D., Jr., 1973, Volcanic and intrusive rocks of the central part of the Oregon Coast Range: *Oregon Department of Geology and Mineral Industries Bulletin 77*, p. 47-74.
- MacLeod, N. S., Tiffin, D. L., Snaveley, P. D., Jr., and Currie, R. G., 1977, Geologic interpretation of magnetic and gravity anomalies in the Strait of Juan de Fuca, U.S. - Canada: *Canadian Journal of Earth Sciences*, v. 14, no. 2, p. 223-238.
- McLean, Hugh, 1977, Lithofacies of the Blakeley Formation, Kitsap County, Washington - a submarine fan complex?: *Journal of Sedimentary Petrology*, v. 47, no. 1, p. 78-88.
- Moore, E. J., 1976, Oligocene marine mollusks from the Pittsburg Bluff Formation in Oregon: *U.S. Geological Survey Professional Paper 922*, 66 p.
- Moore, R. C., and Vokes, H. E., 1953, Lower Tertiary crinoids from northwestern Oregon: *U.S. Geological Survey Professional Paper 233-E*, p. 113-148.
- Muller, J. E., 1971, *Revised geological reconnaissance map of Vancouver Island and Gulf Islands, B.C.: Geological Survey of Canada Open-File 61*.
- \_\_\_\_\_, 1977, *Evolution of the Pacific Margin, Vancouver Island, and adjacent regions: Canadian Journal of Earth Sciences*, v. 14, no. 9, p. 2062-2085.

- Mutti, Emiliano, 1977, Distinctive thin-bedded turbidite facies and related depositional environments in the Eocene Hecho Group (South-central Pyrenees, Spain): *Sedimentology*, v. 24, p. 107-131.
- Mutti, Emiliano, Nilsen, T. H., and Ricci Lucchi, Franco, 1978, Outer fan depositional lobes of the Laga Formation (upper Miocene and lower Pliocene), east-central Italy, *in* Stanley, D. J., and Kelling, G., eds., *Sedimentation in submarine canyons, fans, and trenches*: Stroudsburg, Pa., Dowden, Hutchinson and Ross, p. 210-223.
- Mutti, Emiliano, and Ricci Lucchi, Franco, 1972, Le turbiditi dell'Appennino settentrionale: introduzione all'analisi di facies: *Societa Geological Italiana Memorie*, v. 11, p. 161-199.
- \_\_\_\_\_, 1975, Turbidite facies and facies associations, *in* Examples of turbidite facies and facies associations from selected formations of the Northern Appennines: *International Sedimentological Congress, Nice 1975, Field trip guidebook A-11 9*, p. 21-36.
- Nakamura, K., 1977, Volcanoes as possible indicators of tectonic stress orientation; principle and proposal: *Journal Volcanology and Geothermal Research*, v. 2, no. 1, p. 1-16.
- Niem, A. R., and Van Atta, R. O., 1973, Cenozoic stratigraphy of northwestern Oregon and adjacent southwestern Washington, *in* Beaulieu, J. D., ed., *Geologic field trips in northern Oregon and southern Washington*: Oregon Department of Geology and Mineral Industries Bulletin 77, p. 75-132.
- Pearl, J. E., 1977, Petrology of Tertiary sedimentary rocks in the northernmost part of the Olympic Peninsula, Washington: San Jose State University, San Jose, California, M.S. thesis, 91 p.
- Pettijohn, F. J., 1975, *Sedimentary rocks*, 3d ed.: New York, Harper and Row, 628 p.
- Rau, W. W., 1958, Stratigraphy and foraminiferal zonation in some of the Tertiary rocks of southwestern Washington: U.S. Geological Survey Oil and Gas Investigations Chart OC-57, 2 sheets.
- \_\_\_\_\_, 1964, Foraminifera from the northern Olympic Peninsula, Washington: U.S. Geological Survey Professional Paper 374-G, p. 1-33, 7 pls.
- \_\_\_\_\_, 1966, Stratigraphy and Foraminifera of the Satsop River area, southern Olympic Peninsula, Washington: Washington Division of Mines and Geology Bulletin 53, 66 p.
- \_\_\_\_\_, 1973, Geology of the Washington coast between Point Grenville and the Hoh River: Washington Division of Natural Resources Bulletin 66, p. 1-58.
- \_\_\_\_\_, 1975, Geological map of the Destruction Island and Taholah quadrangles, Washington: Washington Division of Geology and Earth Resources Geologic Map GM-13, scale 1:62,500.
- \_\_\_\_\_, in press, Geologic map in the vicinity of the lower Bogachiel and Hoh River Valleys, and the Washington Coast: Washington Division of Geology and Earth Resources Geology Map GM-24.
- Ricci Lucchi, Franco, 1975, Depositional cycles in two turbidite formations of Northern Appennines (Italy): *Journal of Sedimentary Petrology*, v. 45, no. 1, p. 3-43.
- Schenck, H. G., 1927, Marine Oligocene of Oregon: California University Publications, Department of Geology Bulletin, v. 16, no. 12, p. 449-460.
- Schenck, H. G., and Kleinpell, R. M., 1936, Refugian stage of the Pacific coast Tertiary: *American Association of Petroleum Geologists Bulletin*, v. 20, no. 2, p. 215-225.
- Shouldice, D. H., 1971, Geology of the western Canadian continental shelf: *Canadian Petroleum Geology Bulletin*, v. 19, no. 2, p. 405-436.
- Snavely, P. D., Jr., and MacLeod, N. S., 1977, Evolution of Eocene continental margin of western Oregon and Washington: *Geological Society of America Abstracts with Programs*, v. 9, no. 7, p. 1183.
- Snavely, P. D., Jr., MacLeod, N. S., Rau, W. W., Addicott, W. O., and Pearl, J. E., 1975, Alsea Formation - an Oligocene marine sedimentary sequence in the Oregon Coast Range: *U.S. Geological Survey Bulletin 1395-F*, 21 p.
- Snavely, P. D., Jr., MacLeod, N. S., and Wagner, H. C., 1968, Tholeiitic and alkalic basalts of the Eocene Siletz River Volcanics, Oregon Coast Range: *American Journal of Science*, v. 266, no. 6, p. 454-481.
- Snavely, P. D., Jr., Niem, A. R., and Pearl, J. E., 1978, Twin River Group (upper Eocene to lower Miocene)--defined to include the Hoko River, Makah, and Pysht Formations, Clallam County, Washington: U.S. Geological Survey Bulletin 1457-A, p. A111-A120.
- Snavely, P. D., Jr., Pearl, J. E., and Lander, D. L., 1977, Interim report on petroleum resources potential and geologic hazards in the outer continental shelf - Oregon and Washington Tertiary Province: U.S. Geological Survey Open-File Report 77-282, 64 p.
- Snavely, P. D., Jr., and Wagner, H. C., 1963, Tertiary geologic history of western Oregon and Washington: Washington Division of Mines and Geology Report of Investigations 22, 25 p.
- Tiffin, D. L., Cameron, B. E. B., and Murray, J. W., 1972, Tectonics and depositional history of the continental margin off Vancouver Island, British Columbia: *Canadian Journal of Earth Sciences*, v. 9, no. 3, p. 280-296.
- Walker, R. G., and Mutti, Emiliano, 1973, Turbidite facies and facies associations, *in* Middleton, G. V., and Bouma, A. H., eds., *Turbidites and deep-water sedimentation*: Society of Economic Paleontologists and Mineralogists, Pacific Section Short Course, p. 119-158.
- Weaver, C. E., 1912, A preliminary report on the Tertiary paleontology of western Washington: Washington Geological Survey Bulletin 15, 80 p.
- \_\_\_\_\_, 1937, Tertiary stratigraphy of western Washington and northwestern Oregon: Washington University Publications in Geology, v. 4, 266 p.
- Williams, Howell, Turner, F. J., and Gilbert, C. M., 1954, *Petrography*: San Francisco, W. H. Freeman, 406 p.