

MF 31 / GQ 71

RESOURCE COMPILATION SECTION

(200)
767
C. C. Cater

G E O L O G Y O F T H E
D A V I S M E S A Q U A D R A N G L E , C O L O R A D O

By Fred W. Cater, Jr.,
with a section on "The Mines" by Leonid Bryner

GEOLOGICAL SURVEY
RECEIVED
MAR 3 1954
Denver

*Not to be resource
carded
Rosemary Robbins
5-4-54*

Trace Elements Memorandum Report 694

UNITED STATES DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY





(200)
T67mm

IN REPLY REFER TO:

an

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY
WASHINGTON 25, D. C.

FEB 26 1954

AEC - 490/4

Dr. Phillip L. Merritt, Assistant Director
Division of Raw Materials
U. S. Atomic Energy Commission
P. O. Box 30, Ansonia Station
New York 23, New York


Dear Phil:

Transmitted herewith is one copy of TEM-694, "Geology of the Davis Mesa quadrangle, Colorado," by Fred W. Cater, Jr., with a section on "The Mines" by Leonid Bryner, November 1953.

On December 9, 1953, Mr. Hosted approved our plan to publish this report in the Quadrangle Map Series.

Sincerely yours,

Dwight M. Lammont
for W. H. Bradley
Chief Geologist



FEB - 1 2001

Geology and Mineralogy

This document consists of 29 pages,
plus 1 figure.
Series A

UNITED STATES DEPARTMENT OF THE INTERIOR

GEOLOGICAL SURVEY

GEOLOGY OF THE DAVIS MESA QUADRANGLE, COLORADO*

By

Fred W. Cater, Jr.,

with a section on "The Mines" by

Leonid Bryner

November 1953

Trace Elements Memorandum Report 694

This preliminary report is distributed
without editorial and technical review
for conformity with official standards
and nomenclature. It is not for public
inspection or quotation.

*This report concerns work done partly on behalf of the Division
of Raw Materials of the U. S. Atomic Energy Commission.

USGS - TEM-694

GEOLOGY AND MINERALOGY

<u>Distribution (Series A)</u>	<u>No. of copies</u>
Division of Raw Materials, New York	1
Division of Raw Materials, Washington	1
Exploration Division, Grand Junction Operations Office	1
Grand Junction Operations Office	1
Technical Information Service, Oak Ridge	1
U. S. Geological Survey:	
Geochemistry and Petrology Branch, Washington	1
Geophysics Branch, Washington	1
Mineral Deposits Branch, Washington	1
R. P. Fischer, Grand Junction	1
W. A. Fischer, Washington	1
TEPCO, Denver	1
TEPCO, RPS, Washington	2
(Including master)	<u>13</u>

CONTENTS

	Page
Abstract	4
Introduction	5
Regional geology	6
Stratigraphy	6
Hermosa formation	7
Paradox member	7
Limestone member	7
Cutler formation	8
Moenkopi formation	8
Chinle formation	9
Glen Canyon group	10
Wingate sandstone	10
Kayenta formation	11
Navajo sandstone	11
San Rafael group	12
Carmel formation and Entrada sandstone	12
Summerville formation	13
Morrison formation	14
Salt Wash sandstone member	14
Brushy Basin shale member	15
Burro Canyon formation	16
Dakota sandstone	17
Quaternary deposits	17
Structure	18
Regional setting	18
Structure in Davis Mesa quadrangle	19
Structural history	19
Mineral deposits	22
Mineralogy	22
Ore bodies	23
Origin of ore	23
Suggestions for prospecting	25
The mines	26
Club mines	26
Tramp No. 2 mine	27
Mines in the Long Park area	27
Other deposits	28
Literature cited	28

ILLUSTRATION

Preliminary geologic map and section of the Davis Mesa quadrangle, Colorado	In envelope
--	-------------

GEOLOGY OF THE DAVIS MESA QUADRANGLE, COLORADO

by Fred W. Cater, Jr., with a section on "The Mines"

by Leonid Bryner

ABSTRACT

The Davis Mesa quadrangle is one of eighteen 7 1/2-minute quadrangles covering the principal carnotite-producing area of southwestern Colorado. The geology of these quadrangles was mapped by the U. S. Geological Survey for the Atomic Energy Commission as part of a comprehensive study of carnotite deposits. The rocks exposed in the eighteen quadrangles consist of crystalline rocks of pre-Cambrian age and sedimentary rocks that range in age from late Paleozoic to Quaternary. Over much of the area the sedimentary rocks are flat lying, but in places the rocks are disrupted by high-angle faults, and northwest-trending folds. Conspicuous among the folds are large anticlines having cores of intrusive salt and gypsum.

Most of the carnotite deposits are confined to the Salt Wash sandstone member of the Jurassic Morrison formation. Within this sandstone, most of the deposits are spottily distributed through an arcuate zone known as the "Uravan Mineral Belt". Individual deposits range in size from irregular masses containing only a few tons of ore to large, tabular masses containing many thousands of tons. The ore consists largely of sandstone selectively impregnated and in part replaced by uranium and vanadium minerals. Most of the deposits appear to be related to certain sedimentary structures in sandstones of favorable composition.

INTRODUCTION

The U. S. Geological Survey mapped the geology of the Davis Mesa quadrangle, Colorado, as part of a comprehensive study of carnotite deposits. The study, covering the principal carnotite-producing area in southwestern Colorado, included detailed examination of mines and geologic mapping of eighteen 7 1/2-minute quadrangles, of which the Davis Mesa quadrangle is one. Parts of the texts accompanying these maps have been standardized; these parts comprise some descriptions of geologic formations and general descriptions of regional structural setting, geologic history, and ore deposits. A comprehensive report presenting in greater detail the geologic features of the entire area and interpretations of these features is in preparation. Work was started in the area in 1939 as a cooperative project with the State of Colorado Geological Survey Board and the Colorado Metal Mining Fund and was continued through 1945 as a wartime strategic minerals project. Since 1947 the Geological Survey has been continuing this geologic study on behalf of the Division of Raw Materials of the Atomic Energy Commission. A part of the Davis Mesa quadrangle was mapped in 1941; the rest was mapped in 1948.

The Davis Mesa quadrangle covers about 59 square miles in Montrose County, Colo., and lies in the Canyon Lands division of the Colorado Plateau physiographic province. The quadrangle, except for the floor of Paradox Valley, is a rugged area of mesas and canyons. Total relief within the quadrangle is slightly more than 2,270 feet; altitudes range from about 4,850 feet in the canyon of the Dolores River to over 7,120 feet on Skein Mesa. The Dolores and San Miguel Rivers and their tributaries drain all the area in the quadrangle.

No accurate information on rainfall is available, but the annual precipitation is probably between 10 and 15 inches; the area is semiarid and supports a moderate growth of juniper and piñon on rocky terrain and abundant sagebrush where soils are thick. Cacti and sparse grass are widely distributed. Most of the quadrangle is accessible over a system of dry-weather roads.

REGIONAL GEOLOGY

Rocks exposed in the 18 quadrangles mapped consist of crystalline pre-Cambrian rocks and sedimentary rocks that range in age from late Paleozoic to Quaternary. Crystalline rocks crop out only in the northeastern part of the area along the flanks of the Uncompahgre Plateau; the rest of the area is underlain by sedimentary rocks. The latest Paleozoic and earliest Mesozoic beds wedge out northeastward against the crystalline pre-Cambrian rocks, but later Mesozoic units were deposited on top of the pre-Cambrian rocks. Over most of the region the sedimentary beds are flat-lying, but in places they are disrupted by high-angle faults, or folded into northwest-trending monoclines, shallow synclines, and strongly developed anticlines. The largest of the folds is the Uncompahgre Plateau uplift, a fold nearly 100 miles long that traverses the northeastern part of the area. Well developed anticlines having intrusive cores of salt and gypsum underlie Sinbad Valley, Paradox Valley, and Gypsum Valley in the central part of the area; the Dolores anticline in the southwestern part of the area probably has a salt-gypsum core, although it is not exposed.

The Davis Mesa quadrangle lies in the central part of the area astride Paradox Valley.

STRATIGRAPHY

The oldest rocks exposed in the Davis Mesa quadrangle are of early Pennsylvanian age and are exposed in the central part of Paradox Valley. Rocks of late Pennsylvanian and Permian age crop out on the floor and along the sides of the valley. Triassic rocks are exposed on the floor and the sides of Paradox Valley and in the canyon of the Dolores River. Jurassic rocks crop out along the sides of Paradox Valley and in the canyon walls and on the benches and slopes below the mesas. Cretaceous rocks cap the higher mesas. Recent deposits of wind-blown material and sheet wash are widely distributed on top of the mesas, along the benches, and on the valley floors.

The stratigraphic sequence is similar to that studied by Baker (1933) and Dane (1935) in nearby areas in Utah; most of the formations can be traced continuously from the Davis Mesa quadrangle into Utah.

Hermosa formation

The Pennsylvanian Hermosa formation comprises two members in this area; these are the lower or Paradox member consisting largely of intrusive salt and gypsum, and the upper or limestone member.

Paradox member

The Paradox member, where exposed, consists largely of cellular and earthy gypsum and minor amounts of limestone, black shale, and sandstone. At depth, more than half the formation is rock salt. All known surface occurrences of the Paradox are intrusive, and the beds are complexly folded and contorted. The true, undisturbed thickness of the member is not known, but a well, drilled in the center of Paradox Valley a short distance southeast of the Davis Mesa quadrangle, penetrated over 10,800 feet of intrusive beds without reaching pre-Paradox strata. Baker (1933, p. 17-18) and Dane (1935, p. 27-29) assigned the Paradox to the lower Pennsylvanian.

Limestone member

Presumably the upper or limestone member of the Hermosa formation conformably overlies the Paradox member. Only a few hundred feet of steeply dipping beds in the upper part of the member are exposed in the Davis Mesa quadrangle; these beds consist almost entirely of thick-bedded gray marine limestone. A well, drilled in the southwest flank of the Paradox Valley anticline a few miles west of the Davis Mesa quadrangle, penetrated about 2,300 feet of beds belonging to the limestone member of the Hermosa formation before reaching anhydrite thought to be uppermost Paradox. The limestone member is probably of late Pennsylvanian age.

Cutler formation

The Permian (?) Cutler formation consists of maroon, purple, red, and mottled light-red arkosic sandstone and conglomerate and small quantities of reddish-brown sandy mudstone. The arkosic beds are derived largely from pre-Cambrian crystalline rocks that formed the ancestral Uncompahgre highland, a site now occupied in part by the Uncompahgre Plateau. The poorly sorted, rudely bedded layers and lenses consist of grains of quartz, fresh feldspar, and dark minerals, and pebbles and cobbles of granite, gneiss, schist, and quartzite. Interlayered with the arkosic material are thin irregular beds of sandy mudstone. At the base of the formation where it is in undisturbed contact with the Hermosa formation, beds of marine limestone and arkosic material alternate. These beds are lithologically similar to beds of the Rico formation in the San Juan Mountains and occupy the same relative stratigraphic position. Because the sequence is thin and the age uncertain, it has been mapped with the Cutler formation. The thickness of the Cutler formation in this area is about 3,500 feet, but it thickens to the northeast before wedging out on the southwest flank of the Uncompahgre Plateau.

Moenkopi formation

The Lower Triassic Moenkopi formation unconformably overlies the Cutler formation. In this area the formation consists of three members, a lower member of poorly sorted sandy mudstone, a middle member characterized by beds of arkosic conglomerate and conglomeratic sandstone, and an upper member of thinly bedded shale and lesser amounts of fine-grained sandstone.

The lower member of the Moenkopi formation consists almost entirely of brick-red poorly sorted sandy mudstone and silty sandstone in which grains range in size from clay particles to granules as much as 3 millimeters across. The base of the member contains seams of gypsum and, in a few places, thin beds of gypsum.

The middle member of the Moenkopi formation consists of purple and reddish-brown ledge-forming beds of arkosic conglomerate and conglomeratic sandstone separated by beds of reddish-brown and chocolate-colored sandstone and shale. The conglomeratic beds are cross-laminated, fill channels cut in underlying beds, and show current lineations and ripple marks. Shale beds show fossil mud cracks.

The upper member of the Moenkopi formation consists predominantly of reddish-brown and chocolate-brown shale and sandstone. The shale is micaceous; some bedding planes are sheeted with minute flakes of mica. Individual laminae in the shale are paper-thin, and ripple-bedding is pervasive. Thin beds of coarse-grained sandstone are scattered throughout the member.

The lithologic characteristics of the formation indicate the lower member of the formation was deposited by dumping of material in a body of standing water, the middle member by deposition under fluvial conditions, and finally, in the upper member, a return to deposition in a shallow body of water. In the Davis Mesa quadrangle the Moenkopi formation attains a maximum thickness of about 500 feet divided about equally among the three members. Toward the center of the valley near the salt core of the Paradox Valley anticline, the formation thins greatly and only a small part of the middle member is present.

Chinle formation

The Upper Triassic Chinle formation consists of red to orange-red siltstone, with interbedded red, fine-grained sandstone, shale, and limestone pebble and clay pellet-conglomerate. These lithologic units are lenticular and discontinuous. The lower part of the formation contains numerous lenses of a very distinctive limestone pebble and clay pellet-conglomerate; in places the lowermost lenses contain quartz pebbles or consist of a relatively clean quartz grit. These quartz-bearing lenses are probably the stratigraphic equivalent of the Shinarump conglomerate, which is widely distributed in eastern Utah and northern Arizona. Much of the Chinle formation consists of indistinctly bedded red siltstone that breaks into angular fragments. Evenly bedded shale is rare. The sandstone layers differ in bedding characteristics; some layers are massive, whereas others are cross-bedded, and still others are conspicuously

ripple-bedded. Almost everywhere the formation crops out as a steep slope broken in places by more resistant ledges of sandstone and conglomerate.

The Chinle formation in the Davis Mesa quadrangle ranges from 425 feet to 450 feet in thickness, except where it thins abruptly on the flanks of the Paradox Valley anticline.

Glen Canyon group

The Glen Canyon group of Jurassic (?) age comprise, in ascending order, the Wingate sandstone, the Kayenta formation, and the Navajo sandstone.

Wingate sandstone

The Wingate sandstone conformably overlies the Chinle formation. The sandstone is a massive, fine-grained rock composed of clean, well-sorted quartz sand. It typically crops out as an impressive red or dark brown wall, stained and streaked in places with a surficial red and black desert varnish. Vertical joints cut the sandstone from top to bottom; the spalling of vertically jointed slabs largely causes the recession of the cliff. The sandstone is divided into horizontal layers by extensive bedding planes spaced 2 to 50 feet apart. Within each horizontal layer the sandstone is cross-bedded on a magnificent scale; great sweeping tangential cross-beds of eolian type, in places extending across the entire thickness of the horizontal layer are disposed in all directions. The sandstone is rather poorly cemented and crumbles easily; this quality probably accounts for the readiness with which the rock disintegrates in faulted areas.

In the Davis Mesa quadrangle the Wingate sandstone ranges in thickness from 250 to 325 feet, except along the walls of Paradox Valley where it thins markedly.

Kayenta formation

The Kayenta formation conformably overlies the Wingate sandstone; the contact between the two formations is gradational in most places. The formation is notable for its variety of rock types. Sandstone, red, buff, gray, and lavender in color, is the most abundant type; but the formation also contains considerable quantities of red siltstone, thin-bedded shale, and conglomerate. The conglomerate contains pebbles of sandstone, shale, and limestone. The sandstone is composed of rounded to sub-rounded quartz grains, and minor quantities of mica, feldspar, and dark minerals. Most of the sandstone is thin-bedded, cross-bedded in part, and flaggy; some is massive. Individual sandstone beds are lenticular and discontinuous and interfinger with shale and, in places, with conglomerate. The Kayenta typically crops out in a series of benches and ledges. The ledges in many places overhang recesses where softer beds have eroded back. The lower part of the formation is more firmly cemented and forms resistant, thick ledges that protect the underlying Wingate sandstone from erosion.

The Kayenta formation in the Davis Mesa quadrangle ranges in thickness from 180 to 240 feet, except on the walls of Paradox Valley where it thins greatly. Abrupt local changes in thickness of 10 to 20 feet are common. The irregular bedding, channel filling, and range of thickness all indicate a fluvial origin.

Navajo sandstone

The Navajo sandstone conformably overlies the Kayenta formation. The Navajo is a gray to buff massive fine-grained clean quartz sandstone. Tangential cross-beds of tremendous size leave little doubt of the eolian origin of the sandstone. The sandstone weathers by disintegration and tends to develop rounded topographic forms where exposed on slopes or benches, and vertical cliffs where protected by overlying rocks.

The Navajo sandstone ranges in thickness from a maximum of about 190 feet in Wild Steer Canyon to a knife-edge under Davis Mesa.

San Rafael group

In this area the San Rafael group, of Middle and Late Jurassic age, comprises, in ascending order, the Carmel formation (Middle and Upper Jurassic), the Entrada sandstone, (Upper Jurassic), and the Summerville formation (Upper Jurassic). The group crops out in a narrow band along canyon walls and on the sides of Paradox Valley. The Carmel formation and the Entrada sandstone were mapped together because in most places they form a narrow outcrop.

Carmel formation and Entrada sandstone

The Carmel formation consists largely of red to buff, soft, horizontally bedded siltstone, mudstone, and sandstone. In some localities the basal beds consist of reworked Navajo sandstone. Pebbles and angular fragments of white and gray chert, as much as an inch across, are scattered rather abundantly through the lower part of the formation and less abundantly through the upper part. These chert pebbles and angular fragments are sufficiently abundant locally to form layers of conglomerate. Included in these layers are scattered greenish-gray, red, and yellow quartzite pebbles and boulders as large as 5 by 8 inches in diameter. In many places the upper part of the formation contains scattered barite nodules as much as an inch across.

The Carmel formation ranges from 10 feet or less to 75 feet in thickness. This large range appears to be due chiefly to deposition on irregular, eroded surfaces of Navajo sandstone or beds of the Kayenta formation. No definite evidence indicates that the Carmel formation of this area is of marine origin as is the Carmel of central Utah, but the probabilities are that the Carmel of southwestern Colorado was deposited in shallow water marginal to a sea.

The Carmel formation grades upward, in most places without prominent breaks, into the Entrada sandstone. The Entrada sandstone, known locally as the "slick rim" because of its appearance, is perhaps the most strikingly picturesque of all the formations in the plateau region of Colorado. The smoothly rounded, in places bulging, orange, buff, and white cliffs formed by this sandstone are a distinctive and scenic feature of the region. Horizontal rows of pits resulting from differential weathering and ranging from a few inches to a foot or more across are characteristic of these cliffs.

The Entrada consists of alternating horizontally bedded units and sweeping, eolian-type cross-bedded units. The horizontally bedded units are most common in the basal part and in the uppermost, lighter-colored part of the Entrada, whereas the cross-bedded units are dominant in the middle part. The Entrada sandstone differs from the somewhat similar Wingate sandstone and Navajo sandstone by the sorting of sand into two distinct grain sizes. Subrounded to subangular quartz grains mostly less than 0.15 mm in diameter make up the bulk of the sandstone. The sandstone also contains larger grains, which are well-rounded, have frosted surfaces, and range from 0.4 to 0.8 mm in diameter; most of these grains are of quartz, but grains of chert are scattered among them. Most of the larger grains are distributed in thin layers along bedding planes.

The Entrada sandstone is 120 to 180 feet thick, except along the flanks of the Paradox Valley anticline, where it thins to less than 100 feet.

Summerville formation

The Summerville formation generally crops out as a steep, debris-covered slope, with very few good exposures. Where exposed the Summerville exhibits a remarkably even, thin, horizontal bedding. Beds are predominantly red of various shades, although some beds are green, brown, light yellow, or nearly white. Sandy and silty shale are the most abundant kinds of rock but all gradations from claystone to clean, fine-grained sandstone are interbedded with them. Well-rounded, amber-colored quartz grains with frosted or matte surfaces are disseminated throughout most of the formation, including beds consisting almost entirely of claystone. Thin beds of authigenic red and green chert are widespread. A thin, discontinuous bed of dark-gray dense fresh-water limestone occurs in the upper part of the formation. Sandstone beds are thicker and sandstone is more abundant in the lower part of the formation than in the upper part. Commonly the sandstone beds are ripple-marked, and in places they show small-scale low-angle cross-bedding.

The Summerville formation rests conformably on the Entrada sandstone, and, although a sharp lithologic change marks the contact, no cessation of deposition separated the two formations. Regionally the upper part of the Entrada and the lower part of the Summerville intertongue, and the contact does not occur everywhere at the same stratigraphic horizon. The upper contact of the Summerville is uneven and channeled, and the channels are filled by the overlying basal sandstones of the Morrison formation. Locally, however, the contact is difficult to determine, because the overlying shales and mudstones of the Morrison formation are similar to beds of the Summerville.

The Summerville formation has a moderately uniform thickness of about 105 feet except where it thins on the flanks of the Paradox Valley anticline.

Morrison formation

The Upper Jurassic Morrison formation is of special interest economically because of the uranium- and vanadium-bearing deposits it contains. The formation comprises two members in this area; the lower is the Salt Wash sandstone member, and the upper is the Brushy Basin shale member. In the Davis Mesa quadrangle the Morrison formation ranges in thickness from 650 to 720 feet. The Salt Wash sandstone member and the Brushy Basin shale member in general are of approximately equal thickness. In some areas, however, their thicknesses vary independently whereas in other areas a thinning in one member is accompanied by a thickening in the other.

Salt Wash sandstone member

The Salt Wash sandstone member ordinarily crops out above the slope-forming Summerville formation as a series of thick, resistant ledges and benches. Sandstone predominates and ranges from nearly white to gray, light-buff, and rusty red. Interbedded with the sandstone are red shale and mudstone and locally a few thin lenses of dense gray limestone. Sandstone commonly occurs as strata traceable as ledges for considerable distances along the outcrop, but within each stratum individual beds are lenticular and discontinuous; beds wedge out laterally, and other beds occupying essentially the same stratigraphic position wedge in. Thus, any relatively continuous sandstone stratum ordinarily consists of numerous

interfingering lenses, with superposed lenses in many places filling channels carved in underlying beds. Lenses are separated in places by mudstone and contain mudstone seams. Most of the sandstone is fine- to medium-fine-grained, cross-bedded, and massive; single beds or lenses may attain a maximum thickness of 120 feet. Ripple marks, current lineations, rill marks, and cut-and-fill structures indicate the Salt Wash was deposited under fluvial conditions.

The sandstone consists largely of subangular to subrounded quartz grains, but orthoclase, microcline, and albite grains occur in combined amounts of 10 to 15 percent. Chert and heavy-mineral grains are accessory. Considerable quantities of interstitial clay and numerous clay pellets occur in places, especially near the base of some of the sandstone lenses. Fossil wood, carbonaceous matter, and saurian bones occur locally.

The Salt Wash sandstone member ranges from 280 to 350 feet in thickness and, unlike the underlying formations, does not thin along the flanks of the Paradox Valley anticline. Local changes in thickness of as much as 30 feet are common.

Brushy Basin shale member

The Brushy Basin shale member contrasts strongly in overall appearance with the underlying Salt Wash sandstone member. Although the lithologic differences are marked, the contact between the two members is gradational. The mapped contact, taken as the base of the lowermost layer of conglomerate lenses, is arbitrary in many respects, and probably does not mark an identical stratigraphic horizon in all localities.

The Brushy Basin shale member consists predominantly of varicolored bentonitic shale and mudstone, with intercalated beds and lenses of conglomerate and sandstone, and a few thin layers of limestone. Because of its high proportion of soft, easily eroded bentonitic shale and mudstone, the Brushy Basin member forms smooth slopes covered with blocks and boulders weathered from the more resistant layers of the member and form the overlying formations. The shales and mudstones are thin-bedded and range in color from pure white to pastel tints of red, blue, and green. Exposed surfaces of the rock are covered with a loose, fluffy layer of material several inches thick, caused by

the swelling of the bentonitic material during periods of wet weather. Scattered through the shale and mudstone are thin beds of fine-grained hard silicified rock that breaks with a conchoidal fracture. The silica impregnating these beds may have been released during the devitrification of volcanic debris in adjacent beds. Beds of chert pebble-conglomerate a few inches to 25 feet thick occur at intervals throughout the member. These conglomerate beds are commonly dark rusty red and form conspicuous resistant ledges. Silicified saurian bones and wood are much more abundant in the Brushy Basin shale member than in the Salt Wash sandstone member, especially in some of the conglomerate beds.

The Brushy Basin shale member, like the Salt Wash sandstone member, undoubtedly was deposited under fluvial conditions. The conglomerate and sandstone lenses mark stream channels that crossed flood plains on which were deposited the fine-grained sediments now represented by the mudstone and shale.

The Brushy Basin shale member ranges from 290 to 410 feet in thickness; erratically distributed local variations in thickness of 20 to 30 feet are prevalent throughout the quadrangle.

Burro Canyon formation

The name Burro Canyon formation was proposed by Stokes and Phoenix (1948) for the heterogeneous sequence of Lower Cretaceous conglomerate, sandstone, shale, and thin lenses of limestone that overlies the Morrison formation. The Burro Canyon characteristically crops out as a cliff or a series of thick, resistant ledges. The bulk of the formation consists of white, gray, and red sandstone and conglomerate that form beds up to 100 feet thick. These beds are massive, irregular, and lenticular. Cross-bedding and festoon-bedding are prevalent throughout the formation. The sandstone is poorly sorted and consists of quartz and lesser amounts of chert. The conglomerate consists largely of chert pebbles, but intermixed are pebbles of quartz, silicified limestone, quartzite, sandstone, and shale. In places beds are highly silicified. A considerable part of the formation consists of bright-green mudstone and shale, and locally these predominate over the sandstone and conglomerate. Thin, discontinuous beds of dense, gray limestone crop out in a few scattered localities. The formation was undoubtedly deposited under fluvial conditions. The lower contact is indistinct in many places and appears to interfinger with the

upper part of the Brushy Basin shale member; elsewhere local erosion surfaces intervene and the contact is sharp. The upper contact is an erosion surface of regional extent.

In the Davis Mesa quadrangle the Burro Canyon formation ranges in thickness from 110 to 170 feet.

Dakota sandstone

The Dakota sandstone of Early and Late Cretaceous age crops out as the capping beds on Skein Mesa. The Dakota consists principally of gray, yellow, and buff flaggy sandstone; less abundant are conglomerate, carbonaceous shale, and impure coal. Some of the sandstone is fine-grained and thin-bedded, but much of it is coarse-grained and cross-bedded. Scattered through the sandstone are irregular, discontinuous beds and lenses of conglomerate containing chert and quartz pebbles as much as 2 inches across. Interfingering with the sandstone beds are thin-bedded gray and black carbonaceous shales and thin coal seams and beds. Plant impressions abound in both the sandstone and the shale. The entire thickness of the Dakota sandstone is not exposed in the quadrangle; the upper beds have been stripped off by erosion; but in Dry Creek Basin in the eastern part of the quadrangle the beds that remain attain a maximum thickness of 180 feet.

Quaternary deposits

The deposits of Quaternary age consist of lithified fanglomerate, wind-deposited material, alluvium, talus, and landslides. A hard, thoroughly cemented fanglomerate crops out over a small area in the center of Paradox Valley and at several localities on the northeast side of the floor of the valley; the outcrops in these latter localities were too small to map. The fanglomerate consists of a poorly sorted, rudely bedded accumulation of sand and angular fragments and boulders derived from the older formations in the walls of the valley. The age of this fanglomerate is uncertain but it may be as old as Pliocene. Extensive deposits of light-red sandy and silty material mantle the benches and mesa tops. This material appears to be mostly wind-deposited, although much of it has been reworked by water and intermixed with sheet wash. These deposits have not been mapped where they are spotty, discontinuous, or less than a foot thick; the greatest observed thickness is about 10 feet. The floors of Paradox Valley

are covered with soils and alluvium that generally differ markedly from the wind-deposited material on the mesas. These valley soils are derived not only from wind-blown material but also from the disintegration of the rocks exposed on the valley walls and floors. Considerable alluvium has accumulated along the Dolores River because of ponding in Paradox Valley during periods of high water. Talus covers many of the steeper slopes. Except for the conglomerate these deposits are in places difficult to differentiate and they have not, therefore, been separated on the geologic map. Landslides consisting largely of Brushy Basin debris are prominent along the sides of Paradox Valley.

STRUCTURE

Regional setting

Many geologic structures on the Colorado Plateau are so large that a 7 1/2-minute quadrangle covers only a small part of any complete structural unit. The larger structural units consist of salt anticlines, 45 to 80 miles long; uplifted blocks, 50 to 125 miles long, bounded by monoclinial folds; and domical uplifts, 8 to 20 miles across, around stock-like and laccolithic intrusions.

The salt anticlines trend northwest and lie in a group between eastward-dipping monoclines on the west side of the Plateau and westward-dipping monoclines on the east side of the Plateau. The cores of these anticlines consist of relatively plastic salt and gypsum, derived from the Paradox member of the Hermosa formation and intruded into overlying late Paleozoic and early Mesozoic rocks. All the anticlines are structurally similar in many respects, but each exhibits structural peculiarities not common to the rest; furthermore, all are more complex than their seemingly simple forms would suggest. Faults, grabens, and collapse and slump structures alter the forms of the anticlines. Erosion has removed much of the axial parts of these anticlines, leaving exposed large intrusive masses of the Paradox member and forming valleys such as Sinbad Valley, Paradox Valley, and Gypsum Valley in Colorado and similar valleys in Utah. Alternating with these anticlines are broad, shallow, simple synclines.

Structure in Davis Mesa quadrangle

The Davis Mesa quadrangle lies across the central part of the Paradox Valley anticline. In the northeast and southwest corners of the quadrangle the rock strata are nearly horizontal, but near the edges of Paradox Valley the rocks are sharply upturned along the flanks of the anticline that underlies the valley. The pre-Morrison formations thin against the salt-gypsum core of the anticline, and the older pre-Morrison formations dip more steeply than the younger.

Particularly striking are the complex systems of faults that cut the sides of Paradox Valley. In general the blocks and slivers formed by these faults are downthrown toward the valleys, but some blocks form small horsts. A small graben, about 1,000 feet wide and several miles long, lies along the southwest edge of Davis Mesa. A similar graben has formed on the northeast flank of the anticline near the head of Hieroglyphic Canyon.

One of the most remarkable structural features in the region is the nearly circular downdropped block of rocks north of Saucer Basin. This block, which is nearly three-quarters of a mile across, is encircled by faults, and the central part of the block has dropped about 900 feet below its normal position. This structural feature probably overlies a nearly circular salt plug from which much of the salt has been removed either by solution or salt flowage or both.

Structural history

In order to understand the structural history of the Davis Mesa quadrangle, it is necessary to understand the structural history of the adjoining part of southwestern Colorado. Parts of this history are still in doubt, because no clear record remains of some events; the record of other events, although legible, is subject to different interpretations. All the events described in the following discussion affected the Davis Mesa quadrangle either directly or indirectly, although the evidence for some of them is not visible within the boundaries of the quadrangle.

Mild compressive forces, which probably began in early Pennsylvanian times, gently warped the region. This warping gave rise to the ancestral Uncompahgre highland, an element of the ancestral Rocky Mountains, and to the basin in which the Paradox member of the Pennsylvanian Hermosa formation was deposited. These major structural features controlled the pattern and the prevailing northwest-trending grain of the smaller structures later superimposed on them. The boundary between the highland and the basin, which is closely followed by the southwest margin of the present-day Uncompahgre Plateau, was a steep northwest-trending front, possibly a fault scarp, along which were deposited arkosic fanglomerates during late Pennsylvanian and Permian time. The older fanglomerates inter-finger with Pennsylvanian marine sedimentary rocks of the Hermosa formation. The bulk of the fanglomerates probably is of Permian age and belongs to the Cutler formation. Intrusion of salt from the Paradox member, probably initiated by gentle regional deformation, began sometime during deposition of the Permian Cutler formation. Isostatic rise of salt ruptured the overlying Hermosa and Cutler formations, and after the Cutler was deposited salt broke through to the surface. From then until flowage ceased, late in the Jurassic, the elongate salt intrusions such as those in Paradox Valley and Gypsum Valley stood as actual topographic highs at one place or another along their lengths. The rate of upwelling of additional salt, perhaps accelerated by the increase of the static load of sediments accumulating in the surrounding areas, balanced or slightly exceeded the rate of removal of salt by solution and erosion at the surface. Consequently, all the Mesozoic formations to the base of the Morrison formation wedge out against the flanks of the salt intrusions. Salt flowage was not everywhere continuous and at a uniform rate; rather, in many places it progressed spasmodically. Local surges of comparatively rapid intrusion gave rise to cupolas at different times and in different places along the salt masses. At the beginning of deposition of the Morrison sediments finally covered the salt intrusions, perhaps because the supply of salt underlying the areas between the intrusions was exhausted. Relative quiescence prevailed throughout the remainder of the Mesozoic and probably through the early part of the Tertiary.

The second major period of deformation occurred in the Tertiary--probably during the Eocene (Hunt, written communication), but the date cannot be determined accurately. The region of the salt intrusions was compressed into a series of broad folds, guided and localized by the pre-existing salt intrusions. Although salt flowage was renewed, it seems unlikely that any considerable amount of new salt was forced into the intrusions; flowage probably consisted largely of redistribution of the salt already present. By the end of this period of deformation these folds had attained approximately their present structural form, except for modifications imposed by later collapse of the anticlines overlying the salt intrusions. Owing to the mobility of the rocks in the cores of the anticlines, normal faulting took place along the crests of the anticlines, probably during relaxation of compressive stresses after folding ceased. At this time the crests of the anticlines in places were dropped, as grabens, several hundred to a few thousand feet. A period of crustal quiescence followed, during which the highlands overlying the anticlines and domes were reduced by erosion and topographic relief became low throughout the area.

Then, during the middle Tertiary, the entire Colorado Plateau was uplifted. This uplift rejuvenated the streams and increased ground water circulation. The crests of the anticlines were breached, and the underlying salt was exposed to rapid solution and removal. With the abstraction of salt, renewed collapse of the anticlines began. Although much of the collapse was due directly to removal of salt by solution, it seems unlikely that all the collapse can be attributed to this process, as was believed by earlier workers in the area. Rather, much of the collapse apparently was caused by flowage of salt from the parts of the anticlines still overlain by thick layers of sediments to the parts from which the overlying sediments had been removed. Once the crests of the anticlines had been breached, the relatively plastic salt offered little support for the beds overlying the Paradox member of the Hermosa formation in the flanks of the anticlines; consequently these essentially unsupported beds slumped, probably along fractures and joints formed during earlier flexures. Small faults and folds in Quaternary deposits may indicate that collapse and local readjustments are still continuing.

MINERAL DEPOSITS

The only commercially important mineral deposits in the Davis Mesa quadrangle are those that contain uranium, vanadium, and radium. Although deposits containing these metals were discovered in 1899 near Roc Creek, about 6 miles north of the Davis Mesa quadrangle, intensive mining of these ores did not begin in the Plateau region until 1911. Thereafter the ores were mined primarily for their radium content until 1923, when the Belgian Congo pitchblende deposits began to supply radium. The mines were mostly idle from 1923 until 1937, but since 1937 they again have been exploited intensively, first for vanadium and in more recent years for both vanadium and uranium.

Most of the deposits are restricted to the upper layer of sandstone lenses in the Salt Wash sandstone member, but within this layer the deposits have a spotty distribution. Ore bodies range from small irregular masses containing only a few tons of ore to large tabular masses containing many thousands of tons. The ore consists mainly of sandstone impregnated with uranium- and vanadium-bearing minerals.

Mineralogy

The most common ore minerals are carnotite and a fine-grained, vanadium-bearing micaceous mineral. Carnotite ($K_2(UO_2)_2(VO_4)_2 \cdot 3H_2O$), is a yellow, fine-grained, earthy or powdery material. Tyuyamunite ($Ca(UO_2)_2(VO_4)_2 \cdot nH_2O$), the calcium analogue of carnotite, is also present and is nearly indistinguishable from carnotite. The micaceous vanadium mineral, which formerly was thought to be roscoelite, is now considered to be related to the nontronite or montmorillonite group of clay minerals. It forms aggregates of minute flakes coating or partly replacing sand grains and filling pore spaces in the sandstone. It colors the rock gray. Other vanadium ore minerals present are montroseite ($nFeO \cdot nV_2O_4 \cdot nV_2O_3 \cdot nH_2O$), corvusite ($V_2O_4 \cdot 6V_2O_5 \cdot nH_2O$), and hewettite ($CaO \cdot 3V_2O_5 \cdot 9H_2O$). Corvusite and montroseite occur together, forming compact masses of bluish-black ore, whereas hewettite commonly forms stringers and veinlets along joints and fractures. Recent deeper drilling and mining in the Plateau have indicated that below the zone of oxidation black oxides of uranium and vanadium, accompanied by pyrite and perhaps other sulfides, are more abundant, and uranyl vanadates are scarce or absent.

Ore bodies

The ore consists mostly of sandstone selectively impregnated and in part replaced by uranium and vanadium minerals; but rich concentrations of carnotite and the micaceous vanadium clay mineral are also associated with thin mudstone partings, beds of mudstone pebbles, and carbonized fossil plant material. Many fossil logs replaced by nearly pure carnotite have been found. In general the ore minerals were deposited in irregular layers that roughly followed the sandstone beds. In most deposits the highest-grade concentrations of ore minerals occur in sharply bounded, elongate concretionary structures, called "rolls" by the miners. These rolls are encompassed by rich, veinlike concentrations of the micaceous vanadium-bearing clay mineral that curve across bedding planes. Within these rolls this mineral generally is distributed as diffusion layers, the richer layers commonly lying nearer the margins of the rolls; the distribution of carnotite in the rolls is less systematic. Margins of ore bodies may be vaguely or sharply defined. Vaguely defined margins may have mineralized sandstone extending well beyond the limits of commercial ore; on the other hand, sharply defined margins, such as occur along the surfaces of rolls ordinarily mark the limits of both the mineralized sandstone and the commercial ore.

Although many rolls are small and irregular, the larger ones are elongate and may extend with little change of direction for more than 100 feet. The elongate rolls in an ore body or group of ore bodies in a given area generally have a common orientation. This orientation is roughly parallel to the elongation of the ore bodies.

Origin of ore

The origin of the uranium-vanadium ores in the Morrison formation is uncertain and controversial. In some respects the deposits are unique, and much of the evidence concerning the genesis of the ore is either not conclusive or appears to be contradictory. In this brief account only a small amount of evidence can be presented and the hypotheses can only be summarized.

Most of the deposits are closely associated with certain sedimentary features. Layers of ore lie essentially parallel to the bedding; most of the deposits occur in the thicker parts and commonly near the base of these sandstone lenses; the trend of the long direction of the deposits and the trend of the ore rolls in the sandstone are roughly parallel to the trend of the fossil logs in the sandstone and to the average or resultant dip of the cross-bedding in the sandstone. These relations strongly suggest the primary structures in the sediments were instrumental in localizing most of the ore deposits.

Recent investigations have revealed new data bearing on the origin of the ores (Waters and Granger, 1953). Below the zone of oxidation some of the ore consists chiefly of oxides, such as pitchblende and low-valent oxides of vanadium, and small quantities of sulfides such as pyrite, bornite, galena, and chalcopyrite; fully oxidized and fully hydrated minerals are either rare or non-existent. A hard variety of uraninite, previously reported only from hydrothermal deposits, has been found in the Gray Daun mine in San Juan County, Utah (Razor, 1952), and in the Happy Jack mine in White Canyon, Utah. Studies of lead-uranium ratios in ores from the Colorado Plateau indicate that, regardless of where or in what formation found, all the ores are of roughly the same age, and this age is no older than latest Cretaceous (Stieff and Stern, 1952). Some geologists believe field relations in pre-Morrison formations at White Canyon (Benson, and others, 1952) and Temple Mountain in Utah, indicate that the deposits may be genetically related to faults and fractures. At the Rajah mine near Roc Creek in Colorado, ore occurs along a fault and horsetails out into the wall rock.

Two main hypotheses have arisen to explain the origin of the ores. The oldest and probably the most widely held is the hypothesis that the ores are penesynthetic and were formed soon after the enclosing rocks were deposited (Coffin, 1921; Hess, 1933; Fischer, 1937, 1942, 1950, and Fischer and Hilpert, 1952). Later movements of ground water may have dissolved and reprecipitated the ore constituents, but the essential materials were already present in the host rocks or in the waters permeating them. Although this hypothesis offers a reasonable explanation for the relation of ores to sedimentary features, it faces some difficulty in explaining: (1) the discrepancy between the age of the uranium and the age of the enclosing rock; (2) the broad stratigraphic distribution of uranium occurrences and association of ores with fractures in a few localities; and (3) the hydrothermal aspect

of the mineral suites in some ores. The second hypothesis, and the one the author favors, is essentially a telethermal hypothesis and assumes the ore to have originated from a hypogene source. Proponents of this hypothesis believe that ore-bearing solutions, originated at depth from an igneous source and ascended along fractures. After these solutions mingled with circulating ground waters the minerals were precipitated in favorable beds as much as several miles from fractures. This hypothesis explains more readily the difficulties inherent in the penesynthetic hypothesis but poses two other difficulties: (1) the hypothetical location of igneous source rocks and (2) the difficulty of proving the connection between fractures and faults and the ore deposits. A third hypothesis, advanced by some geologists, suggests that the source of the ore metals was the volcanic material in the beds overlying the ore-bearing sandstones and that these metals were subsequently leached and redeposited in the beds that now contain the ore. This hypothesis encounters not only most of the difficulties in the penesynthetic hypothesis, but it presents some additional ones of its own.

Suggestions for prospecting

Regardless of the actual origin of the deposits, certain habits of the deposits--habits that have been recognized through geologic mapping and exploration experience--are useful as guides for finding ore (Weir, 1952). In southwestern Colorado most of the deposits are in the uppermost sandstone stratum in the Salt Wash member of the Morrison formation. Generally the central or thicker parts of the sandstone lenses are more favorable--many deposits are in sandstone that is 40 feet or more thick, few deposits are in sandstone less than 20 feet thick. Cross-bedded, relatively coarse-grained sandstone is more favorable than thinly or evenly bedded, fine-grained sandstone. Light-yellow-brown sandstone speckled with limonite stain is more favorable than red or reddish-brown sandstone. Sandstone that contains or is underlain by a considerable amount of gray, altered mudstone is more favorable than sandstone containing and underlain by red, unaltered mudstone--this guide is perhaps the most useful in diamond-drill exploration. If the deposits have a hypogene origin, then localities where favorable host rocks are near or coextensive with areas of more intense deformation may be especially favorable for finding ore.

Because of the presence of favorable host rocks the northeastern part of the Davis Mesa quadrangle, (the part lying east of the Dolores River and north of Paradox Valley), is probably the most likely area for finding ore.

The largest mines are in the Long Park and Club Mesa areas. Neither of these areas is far removed from zones of intense deformation. Elsewhere in the quadrangle the rocks are generally unfavorable hosts for ore.

The mines

by Leonid Bryner

The mines in the Davis Mesa quadrangle are among the most productive on the Colorado Plateau. Many of the ore bodies are large and contain tens of thousands of tons of ore. All the larger deposits are in the upper layer of sandstone lenses in the Salt Wash sandstone member of the Morrison formation.

Recently, drilling programs have discovered or extended deposits within the quadrangle, especially around Club Mesa and Long Park. Much of the exploration near old workings and in the shallower drilling areas has been conducted by private firms, but where deeper drilling and higher costs have discouraged private enterprise, the exploration has been undertaken by the Geological Survey on behalf of the Atomic Energy Commission.

Club mines

The Club mines, the most productive in the Davis Mesa quadrangle, are in a zone of ore deposits that extends from the Club mines on the northeast to the Tramp No. 2 mine on the southwest. The ore bodies are large and reach a maximum length of 2,000 feet and a width of 600 feet. They are aligned in a N. 70° E. direction parallel to the trend of the zone of deposits in this area, a trend that is also followed by the rolls and to a lesser extent by the fossil logs in the deposits. Deposits consist of closely grouped rolls interconnected by one or more thin mineralized layers. In general the largest rolls are in the largest deposits. The chief ore minerals are carnotite and the micaceous vanadium clay mineral, but minor quantities of corvusite and montroseite are also present. Some of the fossil

logs and plant material are richly replaced with carnotite, and in many places mudstone pebbles are rich in vanadium minerals.

Tramp No. 2 mine

The Tramp No. 2 mine, another highly productive mine, lies at the southwest end of the same zone of deposits on which the Club mines are located. The trend of the zone of deposits and the rolls in this locality is east. Some of the ore bodies are several hundred feet long and as much as 10 feet thick; most of the ore is confined to rolls; the mineralized rock between the rolls being thin. The ore minerals are chiefly carnotite and the micaceous vanadium clay mineral, but minor quantities of corvusite and montroseite also are found. Although a number of fossil logs have been found during mining operations, few of them were mineralized.

Mines in the Long Park area

A large number of mines are concentrated in the Long Park area. Many of these have been very productive and the total amount of ore taken from the area has been large. Ore deposits occur in several different lenses in the upper sandstone stratum of the Salt Wash. In some mines ore is found at more than one level, either in separate, overlapping sandstone lenses, or in both the upper and lower parts of a single lens where the lens is divided by a mudstone layer or split. Ore bodies are of several types; some consist largely of rolls, others are large tabular masses of disseminated ore, still others are irregular mineralized accumulations of fossil plant material and "trash pockets", and others consist of various combinations of the preceding types. Many of the ore bodies are large and are several hundred feet long. The trend of ore bodies, rolls, and logs in general is N. 70°-80° W., although some rolls in the Sharkey mine trend northeastward. The ore minerals are carnotite, the micaceous vanadium clay mineral, corvusite, montroseite, and hewettite. These minerals are found in varying quantities in different mines, but commonly carnotite and vanadium clay are most important. Commonly the richest concentrations of carnotite are associated with mudstone seams, carbonaceous material, and "trash pockets"; whereas vanadium clay minerals are disseminated through the sandstone or occur

abundantly in the rolls. Corvusite and montroseite occur as local concentrations and in the Coloradium mine they are important ore minerals. Several deposits have been cut by faults; these faults offset the ore bodies, and accordingly they are probably post-ore; in any event, the most recent movement along them has been post-ore.

Other deposits

As a result of recent deep exploratory drilling conducted by the Geological Survey on behalf of the Atomic Energy Commission, a number of deposits rivaling in size those in the Club mines have been discovered in Long Park and on Club Mesa between the Club mine and the Tramp No. 2 mine. Elsewhere in the general area of Club Mesa and Long Park a large number of small deposits have been worked. These are similar in most respects to the larger deposits but are generally of lower grade.

LITERATURE CITED

- Baker, A. A., 1933, Geology and oil possibilities of the Moab district, Grand and San Juan Counties, Utah: U. S. Geol. Survey Bull. 841.
- Benson, W. E., Trites, A. F., Jr., Beroni, E. P., and Feeger, J. A., 1952, Preliminary report on the White Canyon area, San Juan County, Utah: U. S. Geol. Survey Circ. 217.
- Coffin, R. C., 1921, Radium, uranium, and vanadium deposits of southwestern Colorado: Colorado Geol. Survey Bull. 16.
- Dane, C. H., 1935, Geology of the Salt Valley anticline and adjacent areas, Grand County, Utah: U. S. Geol. Survey Bull. 863.
- Fischer, R. P., 1937, Sedimentary deposits of copper, vanadium-uranium, and silver in southwestern United States: Econ. Geology, v. 32, no. 7, p. 906-951.
- _____, 1942, Vanadium deposits of Colorado and Utah, a preliminary report: U. S. Geol. Survey Bull. 936-P, p. 363-394.
- _____, 1950, Uranium-bearing sandstone deposits of the Colorado Plateau: Econ. Geology, v. 45, no. 1, p. 1-11.
- Fischer, R. P., and Hilpert, L. S., 1952, Geology of the Uravan mineral belt: U. S. Geol. Survey Bull. 988-A, p. 1-13.
- Hess, F. L., 1933, Uranium, vanadium, radium, gold, silver and molybdenum sedimentary deposits: in Ore deposits of the Western States (Lindgren volume), p. 450-481, Am. Inst. Min. Met. Eng.

LITERATURE CITED--Continued

- Rasor, C. A., 1952, Uraninite from the Grey Dawn mine, San Juan County, Utah: *Science*, v. 116, no. 3004, p. 89-90.
- Stieff, L. R., and Stern, T. W., 1952, Lead-uranium ages of some uraninites from Triassic and Jurassic sedimentary rocks of the Colorado Plateau (abstract): *Geol. Soc. America Bull.*, v. 63, no. 12, part 2, p. 1299-1300.
- Stokes, W. L., and Phoenix, D. A., 1948, Geology of the Egnar-Gypsum Valley area, San Miguel and Montrose Counties, Colorado: U. S. Geol. Survey Prelim. Oil and Gas Inv., Map 93.
- Waters, A. C., and Granger, H. C., 1953, Volcanic debris in uraniferous sandstones, and its possible bearing on the origin and precipitation of uranium: U. S. Geol. Survey Circ. 224.
- Weir, D. B., 1952, Geologic guides to prospecting for carnotite deposits on Colorado Plateau: U. S. Geol. Survey Bull. 988-B, p. 14-27.