



Approximate position of underground workings and map at right from H. L. Bauer, Jr., 1952, Fluorspar deposits, north end of Spors Mountain, Thomas Range, Juab County, Utah; University of Utah unpublished thesis, figure 4



EXPLANATION

<p>Quaternary</p> <ul style="list-style-type: none"> <li> Alluvium</li> <li> Intrusive rhyolitic breccia</li> </ul> <p>Tertiary</p> <ul style="list-style-type: none"> <li> Thursdays dolomite</li> <li> Lost Sheep dolomite</li> <li> Harrisite dolomite</li> </ul> <p>Silurian</p> <ul style="list-style-type: none"> <li> High-grade fluorspar</li> </ul>	<ul style="list-style-type: none"> <li> Low-grade fluorspar</li> <li> Fault breccia</li> <li> Contact <i>Dashed where approximately located</i></li> <li> Fault, showing dip <i>Dashed where approximately located</i></li> <li> Vertical fault</li> <li> Concealed fault</li> <li> Calcite vein, showing dip</li> <li> Vertical calcite vein</li> <li> Strike and dip of beds</li> </ul>	<ul style="list-style-type: none"> <li> Strike and dip of joints</li> <li> Strike of vertical joints</li> <li> Underground workings</li> <li> Portal and opencut</li> <li> Dump</li> <li> Foot of raise</li> <li> Position of channel sample SB-77-50; 0.033 Sample number; percent of uranium</li> </ul>
--	---	---



Geology by H. L. Bauer, Jr. and V. R. Wilmarth, August 1950 and July 1951

GEOLOGIC MAP AND SECTIONS OF THE BLOWOUT PROPERTY, JUAB COUNTY, UTAH