


EXPLANATION

<p>QUATERNARY</p> <p>CRETACEOUS OR YOUNGER</p>	<p>Alluvium</p> <p>Coastal Plain sedimentary rocks</p> <p>Diabase dike</p> <p>Adamellite</p> <p>Gabbro</p>
<p>TRIASSIC</p>	<p>Richtex Formation a, phyllite and slate; gneiss and schist east of Winnsboro Mills b, amphibolite and greenstone; amphibolite east of Winnsboro Mills</p>
<p>PRECAMBRIAN OR PALEOZOIC</p>	<p>Persimmon Fork Formation a, tuffaceous phyllite b, tuffaceous phyllite with blue quartz fragments</p> <p>Persimmon Fork and Wildhorse Branch Formations, undifferentiated Include mafic and felsic tuffite, bedded argillite, laminated phyllite and other undifferentiated slate-belt rocks east of Ridgeway and Boney</p> <p>Wildhorse Branch Formation</p>
<p>Contact Dashed where approximately located</p> <p>Fault</p> <p>Small-stream drainage basin and sample location</p> <p>RC 291</p> <p>Sampled domestic well</p> <p>Stream-gaging station</p>	

Base from U.S. Geological Survey
1:250,000, Spartanburg 1953



Interior—Geological Survey, Reston, Virginia—1975—G75153
Modified from Secor and Wagener (1968)

GEOLOGIC MAP OF THE CEDAR CREEK-BLYTHEWOOD AREA, SOUTH CAROLINA