


U.S. DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY

CORRELATION OF MAP UNITS AND MINERALIZATION




DESCRIPTION OF MAP UNITS


(All units may not appear on all maps)
Gravel, sand and conglomerate (Holocene to Miocene)—Alluvium filling intermontane basins, on pediments in alluvial aprons and stream terraces, and along water courses.
Basalts (Pliocene to Miocene)—Lava flows and cinder deposits.
Sedimentary and volcanic rocks, undivided (Pliocene to Eocene)—Elastic to andesitic lava and tuff, and some interbedded conglomerate, sandstone, and shale.
Rhyolitic rocks (Miocene and Oligocene)—Includes lava flows, tuffs, and tuffaceous sandstone.
Rhyolitic tuffs (Miocene and Oligocene)—Aerial tuff, ash-flow tuff, tuff breccia, welded tuff, and some sedimentary rocks.
Rhyolitic lava flows (Miocene and Oligocene)—May include some intrusive bodies.
Intrusive rocks (Miocene and Oligocene)—Dikes and plugs.
Granite (Oligocene)—Stocks.
Andesitic rocks (Oligocene)—Lava flows, breccia deposits, and interbedded sedimentary rocks.
Intrusive rocks (Eocene to Late Cretaceous)—Mainly Eocene to Late Cretaceous granite, monzonite, granodiorite, and diorite; some Oligocene to Late Cretaceous peridotite and gabbro (mostly granite).
Volcanic and sedimentary rocks (Eocene to Upper Cretaceous)—Andesitic lava flows and breccia sheets, rhyolitic tuff and welded tuff, and volcaniclastic sedimentary rocks.
Andesite (Eocene to Late Cretaceous)—Plugs, dikes, and stocks.
Rhyolite (Eocene to Late Cretaceous)—Plugs and dikes.
Sedimentary and volcanic rocks, undivided (Upper Cretaceous)—Volcaniclastic conglomerate, sandstone, lacustrine shale, and some andesitic and rhyolitic tuff.
Rhyolite (Upper Cretaceous)—Lava flows, tuffs, and interbedded conglomerate and sandstone.
Rhyolitic tuff (Upper Cretaceous)—Includes aerial and ash-flow tuffs, tuff breccia, welded tuff, and sedimentary rocks.
Rhyolitic lava flows (Upper Cretaceous)—Includes some full and sedimentary rocks.
Andesite (Upper Cretaceous)—Lava flows, breccia sheets, and interbedded conglomerate and sandstone.
Babbler Group (Lower Cretaceous)—Mainly gray shale and siltstone, and some sandstone, conglomerate, and limestone.
Basaltic andesite and andesite (Lower Cretaceous)—Lava flows, cinder deposits, and some dikes, sills, and plugs.
Babbler and Temporal Formations, undivided (Lower Cretaceous)—Andesitic to rhyolitic rocks, conglomerate, and sandstone.
Intrusive rocks (Jurassic)
Granite stocks
Rhyolite plugs
Volcanic and sedimentary rocks (Jurassic to Upper Triassic)—Rhyolite, welded tuff, lava flows, andesitic lava flows, andesitic sandstone, and redbeds. Includes Walnut Gap Formation, Catalina Hills Volcanics, and Gardner Canyon and Mount Wilson Formations.
Metamorphic rocks (Paleozoic or Middle Proterozoic)—Metasandstone, quartzite, and calc-silicate carbonate rocks.
Neo Group (Lower Permian and Pennsylvanian)—Mainly meta-siltstone and dolomite; some siltstone, sandstone, and marble.

Mt Sedimentary rocks (Mississippian)—Generally only Escabrosa Limestone; to the east unit also includes Paradise Formation; mostly shale.
Pl Lower Paleozoic formations, undivided (Upper Devonian to Middle Cambrian)—Mainly limestone and dolomite; some sandstone, shale, and conglomerate. Includes Pinal Schist, Pinal, Sonolite, Martin, El Paso, and Abajo Formations, Coronado Sandstone, and Bobsa Quartzite.
Dikes (Middle Proterozoic)—Includes some metadiorite, in sills, dikes, and plugs; line shows more acidic rocks.
Apache Group (Middle Proterozoic)—Sandstone, shale, argillite, some conglomerate, and possibly some limestone.
Yg Intrusive rocks (Middle Proterozoic)—Granite, granodiorite, and some diorite, sills, and leucogabbro.
Gneissic rocks (Middle and Early Proterozoic)—Metamorphosed granite and older schist or gneiss.
Xp Pinal Schist (Early Proterozoic)—Schist, phyllite, metaquartzite, metagabbro, and meta-igneous rocks.
Contact—Dotted where concealed, queried where uncertain.
Fault—Showing dip; dotted where concealed or intruded, queried where uncertain. Where solid line becomes dotted line within a map unit, that unit is a composite of several formations, of which a younger one conceals faulting in an older one.
Normal fault—But and bar on downthrown side; dotted where concealed, queried where uncertain.
Thrust fault—Swallow on upper plate.
Glide fault—Open swallow on glide plane.
Complex fault—Earlier thrust fault on which later glide faulting has taken place.
Strike-slip fault—Arrow couple shows relative movement, queried where uncertain.
Oblique-slip fault—Composite of strike-slip and normal movement locally, but either type of movement may have occurred without the other.
Fold axis—Dotted where concealed, arrow shows direction of plunge.
Anticline to foliation.
Overturned anticline—Side of closure of arrow ends is side of fold crest relative to fold axis.
Syncline to foliation.
Overturned syncline—Side of closure of arrow ends is side of fold trough relative to fold axis.
Strike and dip of beds.
Horizontal.
Inclined.
Vertical.
Overturned.
Strike and dip of foliation.
Inclined.
Vertical.
Cinder cone—Queried where uncertain.


Weights of base relative importance: the higher the number, the more important, as based on a balance between total production and frequency of occurrence.



Santa Rita and Atascosa-Cobre-Coches-Pajarito-San Luis-Tumacacori Forest units



INDEX MAP SHOWING LOCATION OF CORONADO NATIONAL FOREST (GRAY AREAS).
Forest units shown on this plate are dark gray.



Whetstone and Patagonia-Huachuca-Canelo Forest units

EXPLANATION OF GEOCHEMICAL DATA
Vector diagram showing range of abundance for eight elements—Shortest vector indicates an abundance equivalent to background; medium vector indicates an abundance at or near threshold; longest vector indicates an anomalous abundance. Range given in table below, see chapter C, table 2, this volume, for threshold values. N, not detected at abundance shown; L, detected below abundance shown; G, greater than abundance shown.

Geochemically anomalous areas discussed in text—Arrows labeled a-e (boundaries in text) are subareas of Forest unit discussed in text as C1, D2e, and so on.

(D1) Forest unit discussed in text

Element	CONCENTRATION (PARTS PER MILLION)		
	Background values	High background values at or near threshold	Anomalous value
Au	NID 0.023-1.0 (0.02)	0.002	0.003-4.6
Ag	NID 0.045-0.11	0.12-0.16	0.17-24
As	NID 0.5-1.9	0.0-1.9	0.0-400
Sb	NID 0.5-1.1	1.2-1.6	1.7-250
Cu	1.1-3.9	40-59	60-51,000
Pb	4.3-5.9	60-79	80-10,000
Zn	4.3-5.9	60-79	80-10,000
Mo	NID 0.99-1.6	1.7-2.6	2.7-130

INTRODUCTION
These maps show the distribution of sample localities and abundances determined by inductively coupled plasma spectrometry, except Au, determined by fluorescence atomic absorption spectrophotometry of Au, Ag, As, Cu, Mo, Pb, Sb, and Zn in RASS and NURE stream-sediment and soil samples from the Santa Rita and Atascosa-Cobre-Coches-Pajarito-San Luis-Tumacacori Forest units (area D) and the Whetstone and Patagonia-Huachuca-Canelo Forest Units (area D) of Coronado National Forest and adjacent areas, southeastern Arizona and southwestern New Mexico.
Areas outlined in black on the map indicate regions in which elements having anomalous (or, in some cases, high abundance) form patterns (geochemical signatures). Anomalous regions outlined on the map are based largely on data for the elements and sample methods described above but may have been modified in light of emission-spectrographic data for stream-sediment and paired-concentrate samples (see plate 5); the elements that characterize the geochemical signature in each region are indicated. Some elements, such as chromium, nickel, and strontium, whose abundances are lithologically controlled, are not included with the data shown on the map unless they appear to be part of a signature related to mineralized rock.



MAPS SHOWING GEOCHEMICAL ANOMALIES AND DISTRIBUTION OF STREAM-SEDIMENT AND SOIL SAMPLES CONTAINING HIGH AND ANOMALOUS ABUNDANCES OF Au, Ag, As, Sb, Cu, Pb, Zn, AND Mo IN THE SOUTHWESTERN PART OF CORONADO NATIONAL FOREST AND ADJACENT AREAS, SOUTHEASTERN ARIZONA AND SOUTHWESTERN NEW MEXICO

Compiled by
Gary A. Nowlan
1996