


CORRELATION OF MAP UNITS AND MINERALIZATION





DESCRIPTION OF MAP UNITS


Gravel, sand and conglomerate (Holocene to Miocene)—Alluvium filling intermontane basins, on pediments, in alluvial aprons and stream terraces, and along water-courses.
Basalt (Pliocene to Miocene)—Lava flows and cinder cones.
Sedimentary and volcanic rocks, undivided (Miocene to Eocene)—Shale, to andesitic lava and tuff, and some interbedded conglomerate, sandstone, and shale.
Rhyolitic rocks (Miocene and Oligocene)—Includes lava flows, tuffs, and tuffaceous sandstone.
Rhyolitic tuffs (Miocene and Oligocene)—Aerial tuff, ash-flow tuff, tuff breccia, welded tuff, and some sedimentary rhyolite.
Rhyolitic lava flows (Miocene and Oligocene)—May include some intrusive bodies.
Intrusive rocks (Miocene and Oligocene)
Rhyolite (Miocene and Oligocene)—Dikes and plugs.
Granite (Oligocene)—Stocks.
Andesitic rocks (Oligocene)—Lava flow, breccia deposits, and interbedded sedimentary rocks.
Intrusive rocks (Eocene to Late Cretaceous)—Mainly Eocene to Late Cretaceous granite, monzonite, granodiorite, and diorite; some Oligocene to Late Cretaceous peraluminous two-mica and garnet-bearing granite. Includes Copper Creek Granodiorite.
Volcanic and sedimentary rocks (Eocene to Upper Cretaceous)—Andesitic lava flows and breccia sheets, rhyolitic tuff and welded tuff, and volcanoclastic sedimentary rocks.
Andesite (Eocene to Late Cretaceous)—Plugs, dikes, and stocks.
Rhyolite (Eocene to Late Cretaceous)—Plugs and dikes.
Sedimentary and volcanic rocks, undivided (Upper Cretaceous)—Volcanoclastic conglomerate, sandstone, lacustrine shale, and some andesitic and rhyolitic tuff.
Rhyolite (Upper Cretaceous)—Lava flows, tuffs, and interbedded conglomerate and sandstone.
Rhyolite tuff (Upper Cretaceous)—Includes aerial and ash-flow tuffs, tuff breccia, welded tuff, and sedimentary rocks.
Rhyolite lava flows (Upper Cretaceous)—Includes some tuff and sedimentary rocks.
Andesite (Upper Cretaceous)—Lava flows, breccia sheets, and interbedded conglomerate and sandstone.
Bishop Group (Lower Cretaceous)—Mainly gray shale and siltstone, and some sandstone, conglomerate, and limestone.
Basaltic andesite and andesite (Lower Cretaceous)—Lava flows, cinder deposits, and some dikes, sills, and plugs.
Dakota and Tensara Formations, undivided (Lower Cretaceous)—Andesitic to rhyolitic rocks, conglomeratic and sandstone.
Intrusive rocks (Jurassic)
Granite stocks
Rhyolite plugs
Volcanic and sedimentary rocks (Jurassic to Upper Triassic)—Rhyolitic welded tuff and lava flows, andesitic lava flows, and some sandstone, and redbeds. Includes Walnut Gap Formation, Canelo Hills Volcanics, and Garden Canyon and Mount Washburn Formations.
Metamorphic rocks (Triassic to Middle Proterozoic)—Metagranite, hornfels, and calc-silicate carbonate rocks.
Naco Group (Lower Permian and Pennsylvanian)—Mainly limestone and dolomite, some siltstone, sandstone, and marlstone.

Mi Sedimentary rocks (Mississippian)—Generally only Escabrosa Limestone; to the east unit also includes Paradise Formation, mostly shale.
Pl Lower Paleozoic formations, undivided (Upper Devonian to Middle Cambrian)—Mainly limestone and dolomite; some sandstone, shale, and conglomerate. Includes Paria Shale, Fortal, Sunnithin, Harra, El Paso, and Albigro Formations, Coronado Sandstone, and Bobsa Quartzite.
Diabase (Middle Proterozoic)—Includes some metadiorite, in sills, dikes, and plugs; line shows more acidic rock.
Apache Group (Middle Proterozoic)—Sandstone, shale, argillite, some conglomerate, and possibly some limestone.
Intrusive rocks (Middle Proterozoic)—Granite, granodiorite, and some dioritic sills, and lamprophyre.
Gneissic rocks (Middle and Early Proterozoic)
Metamorphosed granite and older schist or gneiss.
Pinal Schist (Early Proterozoic)—Schist, phyllite, metagranite, metagabbro, and meta-igneous rocks.
Contact—Dotted where concealed, queried where uncertain.
Fault—Showing slip; dotted where concealed or intruded, queried where uncertain. Where solid line becomes dotted line within a map unit, that unit is a composite of several formations, of which a younger one conceals faulting in an older one.
Normal fault—Ball and bar on downthrown side; dotted where concealed, queried where uncertain.
Thrust fault—Sawtooth on upper plate.
Glide fault—Open sawtooth on glide plate.
Complex fault—Earlier thrust fault on which later glide (grab) faulting took place.
Strike-slip fault—Arrow-sawtooth shows relative movement, queried where uncertain.
Oblique-slip fault—Composite of strike-slip and normal movement likely, but other type of movement may have occurred without the other.
Fold axis—Dotted where concealed; arrow shows direction of plunge.
Anticline
Overtured anticline—Side of closure of arrow ends is side of fold crest relative to fold axis.
Syncline
Syncline in foliation
Overtured syncline—Side of closure of arrow ends is side of fold trough relative to fold axis.
Strike and dip of beds
Horizontal
Inclined
Vertical
Overtured
Strike and dip of foliation
Inclined
Vertical
Cinder cone—Queried where uncertain.

Width of bar shows relative importance; the wider the bar, the more important, as based on a balance between total production and frequency of occurrence.



INDEX MAP SHOWING LOCATION OF CORONADO NATIONAL FOREST (GRAY AREAS).
Forest units shown on this plate are dark gray.



EXPLANATION OF TERRAIN-CLEARANCE DATA
Terrain clearance plotted as deflection off flight path—Base value 120 m in Atascosa-Cobre-Coches-Pajarito-San Luis-Tumacacori Forest unit, 140 m elsewhere; scale factor 50 m/minute of latitude or longitude.
INTRODUCTION
Residual NURE terrain-clearance data (observed clearance minus base value) were plotted as a deflection off the flight path. For north-south flight lines, deflections right and left of flight lines are positive and negative anomalies, respectively. For east-west lines, deflections above and below flight lines are positive and negative anomalies, respectively. A value near the mean anomaly for the data set was chosen as the base value, that is, the anomaly value that would plot on the flight path, and the scale factor was usually chosen to be about twice the standard deviation of the data. The scale factor is the number of anomaly units (m, meters) per minute of latitude or longitude on the map. Scale factors were chosen such that plotted profiles are easily read and display as much detail in the data as possible. For the 5-km-spaced NURE data and the 1:126,720 scale maps, a scale factor equivalent to two standard deviations proved satisfactory.

Santa Rita and Atascosa-Cobre-Coches-Pajarito-San Luis-Tumacacori Forest units
Whetstone and Patagonia-Huachuca-Canelo Forest units

TERRAIN-CLEARANCE MAPS FOR NURE PROFILES FOR THE SOUTHWESTERN PART OF CORONADO NATIONAL FOREST AND ADJACENT AREAS, SOUTHEASTERN ARIZONA AND SOUTHWESTERN NEW MEXICO

Compiled by
Mark E. Gettings
1996