

Table 3. Counts of planktic foraminifers from ODP 1018. Shaded samples were eliminated due to low counts or reworking.

Sample	Depth (rmcd)	<i>G. bulloides</i>	<i>Ga. glutinata</i>	<i>Gl. ruber</i>	<i>Gn. aequilateralis</i>	<i>Gr. inflata</i>	<i>Gr. scitula</i>	<i>N. "dupac"</i>	<i>N. dutertrei</i>	<i>N. pachyderma</i> (d)	<i>N. pachyderma</i> (s)	<i>Orbulina</i>	<i>T. quinqueloba</i>	Total
1018C 3H-4,47	22.60	23	0	0	0	0	0	0	2	0	18	0	0	43
1018C 3H-4,67	22.80	0	0	0	0	0	0	0	0	0	19	0	0	19
1018C 3H-4,77	22.90	0	0	0	0	0	0	0	0	0	23	0	0	23
1018C 3H-4,87	23.00	1	0	0	0	0	0	0	0	2	15	0	0	18
1018C 3H-4,97	23.10	8	0	2	0	0	0	11	8	10	75	0	1	115
1018C 3H-4,107	23.20	1	0	0	0	0	0	0	0	0	17	0	0	18
1018C 3H-4,117	23.30	1	0	0	0	0	0	0	0	3	22	0	0	26
1018C 3H-4,147	23.60	0	0	0	0	0	29	0	0	0	0	0	0	29
1018C 3H-5,17	23.80	3	0	0	1	0	0	0	0	2	85	0	0	91
1018C 3H-6,77	25.90	41	3	1	2	6	1	7	1	112	4	0	3	181
1018C 3H-6,82	25.95	3	0	0	0	0	0	2	6	6	0	0	1	18
1018C 3H-6,87	26.00	0	0	0	0	0	0	4	4	11	0	0	4	23
1018C 3H-6,92	26.05	3	2	0	0	2	0	1	7	2	2	0	1	20
1018C 3H-6,97	26.10	129	4	0	0	5	1	10	10	64	23	0	8	254
1018C 3H-6,102	26.15	154	7	1	0	7	3	14	8	68	3	2	3	270
1018C 3H-6,107	26.20	69	0	3	2	11	0	28	8	121	10	1	3	256
1018C 3H-6,112	26.25	63	4	2	0	2	5	21	27	128	3	1	1	257
1018C 3H-6,117	26.30	56	8	1	0	29	1	13	47	85	5	0	1	246
1018D 3H-4,142	26.33	119	2	3	2	16	0	15	17	46	0	2	2	224
1018C 3H-6,122	26.35	55	6	1	2	13	0	23	16	165	12	3	4	300
1018D 3H-5,2	26.43	2	0	0	1	1	0	1	4	4	0	0	0	13
1018D 3H-5,12	26.53	11	6	0	0	6	0	11	8	52	3	0	0	97
1018D 3H-5,22	26.63	4	0	0	0	1	1	1	0	2	1	1	1	12
1018D 3H-5,32	26.73	3	0	0	0	5	0	1	1	5	2	0	3	20
1018D 3H-5,42	26.83	4	4	0	0	2	1	1	3	6	11	1	2	35
1018D 3H-5,52	26.93	11	2	0	2	2	0	0	2	4	42	0	4	69
1018D 3H-5,62	27.03	6	5	0	0	2	0	2	1	5	88	1	1	111
1018D 3H-5,72	27.13	32	3	1	0	1	1	5	0	33	136	0	8	220