

Table 6. SST estimates for ODP Holes 1018C and 1018D. Dashes (-) indicate samples with unreliable SST estimates; \* insufficient information to calculate standard deviations; n = number of coretop sample matches with  $SCD \leq 0.15$ .

Sample	Depth <sup>a</sup> (rmcd)	age <sup>b</sup> (kyr)	n	SST <sub>cold</sub> (°C)	SST <sub>warm</sub> (°C)	2 $\sigma$ <sub>cold</sub> (°C)	2 $\sigma$ <sub>warm</sub> (°C)
1018C 3H-4,97	23.10	98.629	6	6.74	11.70	1.82	2.31
1018C 3H-5,17	23.80	102.759	24	0.48	4.75	1.33	1.71
1018C 3H-6,77	25.90	122.135	0	-	-	-	-
1018C 3H-6,97	26.10	124.082	0	-	-	-	-
1018C 3H-6,102	26.15	124.343	0	-	-	-	-
1018C 3H-6,107	26.20	124.605	1	13.03	16.51	*	*
1018C 3H-6,112	26.25	124.867	1	13.03	16.51	*	*
1018C 3H-6,117	26.30	125.129	0	-	-	-	-
1018D 3H-4,142	26.33	125.286	0	-	-	-	-
1018C 3H-6,122	26.35	125.390	1	13.03	16.51	*	*
1018D 3H-5,12	26.53	126.333	2	13.89	17.68	0.45	1.11
1018D 3H-5,52	26.93	128.427	5	3.19	7.23	0.43	0.69
1018D 3H-5,62	27.03	128.950	25	2.09	6.37	1.64	2.01
1018D 3H-5,72	27.13	129.474	10	6.72	12.31	1.83	1.92

<sup>a</sup> Lyle et al., in press.

<sup>b</sup> Poore et al., in prep.