

CORRELATION CHART SHOWING THE EAST-WEST LATERAL VARIATION IN THE INTERVALS BETWEEN COAL BEDS IN THE SOUTH LIMB OF THE MAHANOY BASIN

most of the major ridges in the mapped area. The formation is approximately 1200 feet thick in the western part of the area but thins to approximately 800 feet in the eastern part. The lower part of the Potsville formation extends southward across the Eisenhower Run anticline into the Southern anthracite field.

The rocks of the Potsville formation are mainly conglomerate and sandstone, but they also include small amounts of siliceous, claystone, and shale. Most of the rocks are gray, but brown, pink, and green beds are present near the bottom of the formation. The conglomerate is poorly bedded and contains well-sorted and well-sorted pebbles of quartz and scattered pebbles of chert embedded in a siliceous matrix. In diameter the quartz pebbles are as much as 4 inches and the chert pebbles are generally less than 2 inches. The sandstone in the formation is medium to coarse-grained and commonly grades laterally into conglomeratic sandstone and conglomerate.

A persistent coal bed known as the Lykens Valley (No. 2) coal bed is near the middle of the Potsville formation in all the basins in the area. Two nonpersistent coal beds—Buck Mountain (No. 3) and Little Buck Mountain (No. 4)—are present in the upper part of the formation.

**Allegheny formation.**—The Allegheny formation, the base of which was designated by White (1900, p. 624) as the Buck Mountain (No. 5) coal bed, overlies the Potsville formation and underlies the Conemaugh formation. The Allegheny formation ranges from 300 feet to 450 feet in thickness and averages about 400 feet. This formation is composed of conglomerate, sandstone, siliceous, claystone, and coal. The coal beds are the only laterally persistent lithologic units. In the lower part of the formation, sandstone and conglomerate are abundant; in the upper part, claystone and siliceous are more common, and the conglomerate is a minor constituent.

Five persistent coal beds—the Buck Mountain (No. 5), Seven-foot (No. 6) and No. 9—are present in the Allegheny formation. A nonpersistent coal bed, the Foraker (No. 8) bed, also is present near the top of the formation.

**Conemaugh formation.**—The Conemaugh formation overlies the Allegheny formation. The base of the Conemaugh is the Holmes (No. 10) coal bed according to Loban (1937, p. 46), and, although the lower 1,000 feet of the area is present in the central part of the Conemaugh has been removed by erosion. These rocks consist of sandstone, siliceous, claystone, and scattered lenses of conglomerate, interbedded with six persistent coal beds and several local beds (columns 1 and 2). The Holmes (No. 10), Primrose (No. 11), Orchard (No. 12), Little Orchard (No. 13), Diamond (No. 14), and Tracy (No. 15) are the persistent coal beds in the Conemaugh.

**OUTCROPS OF COAL**

Coal beds are rarely exposed in natural outcrops owing to the cover of soil. The outcrops shown on sheet 1 are in the position that they would occupy if the coal were projected through the soil to the surface of the ground. These positions may not agree with the locations of weathered coal, which, because of soil creep, are generally found lower on the slope than the original outcrop.

It was necessary to project some coal beds beyond their last known point of occurrence. This has been done to show the author's estimate of the extent of that particular coal horizon and does not mean that the coal is of suitable thickness throughout its indicated extent. Local or leader coal beds less than 18 inches thick were not mapped.

**THICKNESS OF COAL BEDS**

Listed below are the average thickness and range in thickness of coal beds in the area covered by this report. Figures on the right half of the table show the thickness of all coal in each bed; figures on the left half of the table show the thickness of all coal and all shale partings in each bed.

Most of the coal and bed thicknesses were obtained from mine company data and were chosen to show an average of many underground observations. Measurements of beds that may have been abnormally affected by deformation were not recorded. The maximum and minimum thicknesses shown in this table and in the columnar sections (sheet 2) are from all observed measurements along tunnels, mine coal measurements, drill cores, or sections measured by the Geological Survey personnel. Thickness figures have been omitted from the table where few reliable data are available.

Average thickness and range in thickness of coal beds in the eastern part of the Ashland quadrangle

Bed number	Bed		Coal		Percentage of refuse
	Average	Range	Average	Range	
	Feet	Feet	Feet	Feet	
17	0.9	5.0			
18	0.11	5.4			
14	8.0	3.6-11.0	6.4	2.0-9.4	20.8
13	6.4	4.4-12.2	4.7	1.0-10.0	29.5
12	8.4	3.6-13.1	6.6	1.2-9.8	22.0
11	1.8	3.6			
11T					
11	7.8	2.7-14.8	5.11	1.1-12.4	22.8
10S	1.2	6.1			
10	9.6	2.2-28.0	8.0	1.7-28.0	15.8
9S	7.4	2.8-13.1	6.5	1.8-10.0	12.5
9	2.0	14.7	7.1	5.1-12.8	22.7
8	18.0	12.2-24.4	16.0	9.0-22.2	10.6
8&9	26.0	13.0-44.0	20.7	9.0-34.8	23.3
7T	2.0	0.7-4.4	1.6	0.7-2.1	16.6
7	3.6	1.0-13.1	2.6	0.0-12.0	28.6
6	1.2	2.7			
6	3.0	1.7-11.0	2.7	0.0-6.7	32.6
5	0.9	3.1			
5	11.1	7.7-23.0	9.8	1.6-23.0	18.0
4	3.8	2.7-7.8	3.5	1.0-8.0	6.8
3					
3	0.7	6.2			

**COAL BEDS MINED IN THE AREA**

Eleven coal beds have been mined in the mapped area. Of these, the Buck Mountain (No. 5), Seven-foot (No. 6), Skidmore (No. 7), two splits of the Mammoth coal zone (Nos. 8 and 9), Holmes (No. 10), Orchard (No. 12), and the Diamond (No. 14) coal beds are economically the most important.

The Buck Mountain (No. 5) coal bed, basal unit of the Allegheny formation, overlies the more resistant rocks of the Potsville formation and underlies the remaining beds of the Allegheny, which are less resistant. As the result of differential resistance to erosion a topographic bench forms in some areas, making it possible to map the outcrop of the Buck Mountain coal bed rather accurately. In the western part of the Ashland quadrangle, on the west (see index map, sheet 2), the Buck Mountain consists of two or three splits, Nos. 5, 5M, and 5T, which are mined as separate coal beds. From the outcrop the splits merge eastward into the area of this report, and the combined No. 5 coal bed contains only discontinuous partings that are generally less than a foot thick. The Buck Mountain, which is extremely scarce, has an average thickness of nearly 12 feet in the mapped area.

The Seven-foot (No. 6) and Skidmore (No. 7) coal beds are approximately 160 and 130 feet, respectively, above the Buck Mountain (No. 5) coal bed in this area. The intervening rocks are chiefly sandstone, siliceous, claystone, and occasional lenses of conglomerate. Nos. 6 and 7 coal beds crop out in all of the basins in the mapped area. A split of the Skidmore coal bed, the Skidmore Top Split (No. 7T), crops out in the central part of the area in the North (William Penn) basin and in the southeastern part of the area in the Mahanoy basin. The Seven-foot coal bed has been mined at scattered localities throughout the area, and the Skidmore coal bed has been most extensively mined in the western part of the area in the East mine.

The Mammoth coal zone, consisting of the Mammoth Bottom Split (No. 8) and the Mammoth Top Split (No. 9) coal beds, has been mined as a single unit over most of the eastern half of the Ashland quadrangle. The Nos. 8 and 9 coal beds are separated by carbonaceous claystone and siliceous shale that may attain a thickness of 60 feet in some places. In parts of the Raven Run and Hammond mines, Nos. 8 and 9 were mined separately. Thus the Mammoth coal zone has been mined as a single unit, the average bed thickness is 26 feet, 10 inches. The persistence, quality, and thickness of the coal beds in the Mammoth coal zone make them the most economically important in the area.

The Holmes (No. 10) coal bed is important because of its greater average thickness, 9 feet, 6 inches, and its smaller than average percentage of refuse, 16 percent. Extensive mining and a rather uniform interval from the Mammoth coal zone are factors that make the Holmes coal bed useful as a stratigraphic marker.

The Orchard (No. 12) and Diamond (No. 14) coal beds, which are the youngest beds mined in this area, crop out in the more deeply folded North and Mahanoy basins. Both Nos. 12 and 14 coal beds have an average thickness of 8 feet. Parting of carbonaceous claystone or heavy coal generally constitute more than 20 percent of the bed thickness. These beds attain their maximum thickness in the west-central part of the area, where they have been mined extensively in the Hammond and East mines.

A number of coal beds listed in the preceding coalbeds table are not described in this report. These coals are generally of poor quality or of such local occurrence that they were of little economic importance at the time this report was written. This, however, does not preclude that they may not be of great economic importance as mining practices improve or as the market for coal increases.

**REFERENCES**

Ashburner, C. A., 1883, First report of progress in the anthracite coal region: Pennsylvania 2d Geol. Survey Rept. A4, 407 p.

Haley, B. R., Arndt, H. R., Rothrock, H. C., and Wagner, H. C., 1955, Geology of anthracite in the western part of the Ashland quadrangle, Pennsylvania: U. S. Geol. Survey Coal Inv. Map C-15.

Loban, S. R., 1937, Ground water in northeastern Pennsylvania: Pennsylvania Geol. Survey 4th ser. Bull. 74, 312 p.



Bedrock, H. R., Wagner, H. C., and Haley, B. R., 1950, Geology of anthracite in the west-central part of the Mount Carmel quadrangle, Pennsylvania: U. S. Geol. Survey Coal Inv. Map C-3.

Bedrock, H. R., Wagner, H. C., Haley, B. R., and Arndt, H. H., 1951, Geology of anthracite in the southeastern part of the Mount Carmel quadrangle, Pennsylvania: U. S. Geol. Survey Coal Inv. Map C-7.

Bedrock, H. R., Wagner, H. C., Haley, B. R., and Arndt, H. H., 1951, Geology of anthracite in the east-central part of the Mount Carmel quadrangle, Pennsylvania: U. S. Geol. Survey Coal Inv. Map C-10.

Bedrock, H. R., Wagner, H. C., Haley, B. R., and Arndt, H. H., 1953, Geology of anthracite in the southwestern part of the Mount Carmel quadrangle, Pennsylvania: U. S. Geol. Survey Coal Inv. Map C-12.


White, David, 1900, The stratigraphic succession of the fossiliferous beds of the Potsville formation in the southern anthracite coal field, Pennsylvania: U. S. Geol. Survey 23th Ann. Rept., pt. 1 (1899-99), p. 749-950.



STRUCTURAL CROSS SECTIONS



INDEX MAP OF PENNSYLVANIA SHOWING THE LOCATION OF THE ANTHRACITE FIELDS



INDEX MAP OF THE WESTERN MIDDLE ANTHRACITE FIELD SHOWING THE LOCATION OF MAPPED AREAS

**EXPLANATION**

Area described in this report

Area described in previous reports

**ROCKS OF MISSISSIPPIAN AGE**

The Mauch Chunk formation crops out in the northern part of the area on the Red Ridge and Centralia anticlines and in the southern part of the area on the Frackville anticline between Ashland Mountain and Broad Mountain. Only the upper 1,000 feet of the formation is exposed in the mapped area.

The rocks that comprise the Mauch Chunk formation are mainly sandstone, shale, siliceous, and fine-grained sandstone; but green siliceous and fine-grained sandstone, red or green medium to coarse-grained sandstone, and scattered lenses of gray or green conglomerate are also present. Most of the strata are quite lenticular. The Mauch Chunk appears to grade upward into the overlying Potsville formation.

**ROCKS OF PENNSYLVANIAN AGE**

Rocks of Pennsylvanian age consist of lenticular beds of conglomerate, sandstone, siliceous, claystone, and shale, interbedded with 12 persistent coal beds and several local coal beds. The lower part of the Pennsylvanian in the mapped area is predominantly conglomeratic, and the upper part is chiefly fine-grained. The coal beds, some of which extend throughout the Western Middle field, are the most persistent of the lithologic units, whereas the other strata exhibit so many variations that they are of minor value for an reference or key beds.

**Potsville formation.**—The Potsville formation, which includes the oldest rocks of Pennsylvanian age in the area, overlies the Mauch Chunk formation and underlies the Allegheny formation. The Potsville consists chiefly of resistant classic strata that form all the mountains and