

## Chapter 3

# **The GIS Project for the Geologic Assessment of In-Place Oil Shale Resources of the Eocene Green River Formation, Greater Green River Basin, Wyoming, Colorado, and Utah**



*Click here to return to*  
**Volume Title Page**

By Tracey J. Mercier, Gregory L. Gunther, and Christopher C. Skinner

## Chapter 3 of 6

### **Oil Shale Resources of the Eocene Green River Formation, Greater Green River Basin, Wyoming, Colorado, and Utah**

By U.S. Geological Survey Oil Shale Assessment Team

U.S. Geological Survey Digital Data Series DDS-69-DD

**U.S. Department of the Interior**  
KEN SALAZAR, Secretary

**U.S. Geological Survey**  
Marcia K. McNutt, Director

U.S. Geological Survey, Reston, Virginia: 2011

For more information on the USGS—the Federal source for science about the Earth, its natural and living resources, natural hazards, and the environment, visit <http://www.usgs.gov> or call 1-888-ASK-USGS

For an overview of USGS information products, including maps, imagery, and publications, visit <http://www.usgs.gov/pubprod>

To order this and other USGS information products, visit <http://store.usgs.gov>

Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Although this report is in the public domain, permission must be secured from the individual copyright owners to reproduce any copyrighted materials contained within this report.

Suggested citation:

Mercier, T.J., Gunther, G.L., and Skinner, C.C., 2011, The GIS project for the geologic assessment of in-place oil shale resources of the Eocene Green River Formation, Greater Green River Basin, Wyoming, Colorado, and Utah, *in* U.S. Geological Survey Oil Shale Assessment Team, ed., Oil shale resources of the Eocene Green River Formation, Greater Green River Basin, Wyoming, Colorado, and Utah: U.S. Geological Survey Digital Data Series DDS-69-DD, chap. 3, 2 p.

Contents

Introduction.....1

References Cited.....2

# The GIS Project for the Geologic Assessment of In-Place Oil Shale Resources in the Eocene Green River Formation, Greater Green River Basin, Wyoming, Colorado, and Utah

By Tracey J. Mercier, Gregory L. Gunther, and Christopher C. Skinner

## Introduction

A geographic information system (GIS) focusing on the Green River Formation, Greater Green River Basin of southern Wyoming, northern Colorado, and eastern Utah was developed as an analysis tool for the U.S. Geological Survey's (USGS) 2010 assessment of in-place oil shale resources in the Eocene Green River Formation. The Central Energy Resources Science Center of the USGS has also developed an internet mapping service (IMS) to deliver the GIS data to the public. This online mapping tool utilizes information from a database about the oil shale endowment of the western United States and converts the data into geospatial layers for analysis. The ability to view and query geologic features on an interactive map facilitates (1) an understanding of domestic oil shale resources for strategic planning, (2) formulating economic and energy policies, (3) evaluating lands under the purview of the Federal Government, and (4) developing sound environmental policies.

Several types of data are included in the GIS project, which contains oil shale zone boundaries, borehole locations, isopachs, oil shale Fischer assay results, and resource assessment results in tabular and geospatial data formats. The distribution and richness of oil shale for three assessment zones are illustrated in specialized maps created by the USGS Oil Shale Assessment Team; these are referred to as "raster maps or models." Models for the three oil shale zones are included in the Green River Formation, Greater Green River Basin GIS project. The IMS integrates these data with thematic geospatial layers for geographic reference using a web service-based architecture. The thematic layers or map services available through ArcGIS Online (ESRI, 2009) may include themes formally named Imagery, Streets, Shaded Relief, and Topography. Each formal theme can contain additional geospatial content, examples of which include:

- Highways
- Major and minor roads
- Railways
- Water features
- Administrative boundaries
- Cities, parks, and landmarks
- Public Land Survey System (PLSS)
- Surface ownership
- Surface and outcrop geology

To access the IMS and to download spatial data that formed the basis of the GIS for the Green River Formation, Greater Green River Basin, click on the following USGS Oil Shale Assessment website link: ([http://energy.cr.usgs.gov/other/oil\\_shale/](http://energy.cr.usgs.gov/other/oil_shale/)). Several data formats are provided, including Environmental Systems Research Institute (ESRI, 2006) shapefiles, cell-based GRIDs, and Triangular Irregular Network (TIN) models. These data are also exposed as web services based on a variety of industry-standard encodings, including Keyhole Markup Language (KML), Web Mapping Service (WMS), and ArcGIS Server REST. Additional encodings may be added based on technological need. Information for accessing these services is also available at the website.

All content is documented, based on standard metadata requirements. Access to the metadata is provided in a variety of formats that are incorporated into the downloadable datasets and can be viewed using the IMS.

## References Cited

Environmental Systems Research Institute (ESRI), 2006,  
ArcGIS, ver. 9.2: ESRI, Redlands, California, accessed on  
August 20, 2009, at [http://www.esri.com/software/arcgis/  
index.html](http://www.esri.com/software/arcgis/index.html).

Environmental Systems Research Institute (ESRI), 2009,  
ArcGIS Online. ESRI Website: ESRI, Redlands, California,  
accessed on August 20, 2009, at [http://resources.esri.com/  
arcgisonline/services/](http://resources.esri.com/arcgisonline/services/).



***Click here to return to  
Volume Title Page***