

A more complete description of the study area and the geologic and hydrologic characteristics of the San Antonio segment of the Edwards aquifer that make it vulnerable to contamination is provided in previous reports (Musgrove and others, 2010, 2016, 2019; Opsahl and others, 2018, 2020).

Hydrologic and water-quality data were collected during 2017–19 at two sets of paired surface-water and groundwater sites (fig. 1). The first site pair included Salado Creek (site 1) and the Shavano well (site 3), both in areas minimally influenced by urban development. The second site pair included West Elm Creek (site 2) and the Encino Rio well (site 4), where considerable urban development is taking place. Stages of the two creeks, water-level altitudes (WLAs) in the two wells, and the temperature and specific conductance of water in the wells were monitored continuously. Discrete water samples were collected from the wells and streams at approximately 8-week intervals and in response to storm events. These samples were analyzed for concentrations of chemical constituents and for a large suite of pesticides and pesticide degradates.

Temporal and Spatial Variability in Hydrology and Water Quality

During the first 11 months of the study, rainfall was below average, flow at West Elm Creek was infrequent, and WLAs at the Shavano well were uniformly declining (fig. 2). Following a period of above-average rainfall and resulting surface runoff in September 2018, flow in West Elm Creek was more frequent and stream stages were higher, and the WLA at the Encino Rio well increased nearly 70 feet (fig. 2). The rapid rise in WLA at the Encino Rio well represents groundwater recharge resulting from the direct hydrologic connectivity of SW and unconfined GW in the Edwards aquifer recharge zone. The Edwards aquifer is a limestone (karst) aquifer and such rapid responses to rainfall and runoff at both site pairs is to be expected, regardless of the degree of urbanization. Both continuous water-quality monitoring data and the chemical analysis of discrete water-quality samples showed corresponding changes in GW quality during periods of recharge (Opsahl and others, 2020).

The occurrence of pesticides in samples of stormwater runoff and groundwater collected during 2017–19 provides insight into the transport of water and any contained anthropogenic (human-created) contaminants into and within the Edwards aquifer. There were large differences in the number of different pesticides or pesticide degradates detected at the two SW sites: 7 different compounds were detected at the Salado Creek site, whereas 25 different compounds were detected at the more urbanized West Elm Creek site (table 1). The total number of pesticide detections per site and the average number of detections per sample were both five

times higher in samples collected from West Elm Creek than in samples collected from Salado Creek. Although the number of different pesticides detected (13) was the same at the two GW sites, the number of pesticide detections was markedly different. Consistent with the large differences in the number of pesticide detections and in the number of detections per site between the more urbanized and less urbanized SW sites, a total of 42 pesticide detections were measured in samples collected from the Shavano well, which represents the lesser urbanized area, whereas 74 pesticide detections were measured in samples collected from the more urbanized Encino Rio well site. Overall, a diverse assortment of pesticides was detected in the samples collected from both wells, but the more frequent occurrences at the Encino Rio well are consistent with a larger urban stormwater runoff component in recharge source water than at the Shavano well.




Figure 2. Time series of A, daily rainfall measured at the San Antonio International Airport (National Oceanic and Atmospheric Administration, 2020), B, water-level altitude and specific conductance from the Encino Rio well, and C, water-level altitude and stream stage data from the Encino Rio well and West Elm Creek, 2017–19.

Table 1. Summary of pesticide and pesticide degradate detections in surface-water and groundwater samples collected at sites in the San Antonio segment of the Edwards aquifer, 2017–19.

| | Surface-water sites (fig. 1) | | Groundwater sites (fig. 1) | |
|--|------------------------------|----------------|----------------------------|-----------------|
| | Salado Creek | West Elm Creek | Shavano well | Encino Rio well |
| Number of samples | 4 | 4 | 14 | 14 |
| Number of different pesticides detected | 7 | 25 | 13 | 13 |
| Total number of detections per site | 10 | 50 | 42 | 74 |
| Average number of detections per sample ¹ | 2.5 | 12.5 | 3.0 | 5.3 |

¹Calculated for each site as the total number of detections of all pesticides divided by the number of samples.

Implications for Edwards Aquifer Water Quality

Among all measured constituents, pesticides detected in discrete stormwater-runoff samples provided the clearest indication that urbanization is adversely affecting the quality of water in the Edwards aquifer. More specifically, a greater number of pesticides and a greater variety of pesticides were detected in samples collected from the more urbanized surface-water site. Not all pesticides have regulatory standards, but for those that do, there were no exceedances of individual pesticide concentrations relative to current (2020) drinking water standards (U.S. Environmental Protection Agency, 2019b). The observed hydrologic responses to rainfall and resulting surface runoff, and the corresponding changes in water quality in wells, are thought to result from the direct hydrologic connectivity of SW and unconfined GW. The patterns of GW-quality change, however, indicate mixing of water from multiple sources with the ambient water of the Edwards aquifer,

including recent surface-derived recharge and possibly inflow from other aquifers, such as the underlying Trinity aquifer. Therefore, the effects of urbanization on GW quality cannot be inferred solely from the input of stormwater runoff, as changes related to local and regional hydrologic conditions also must be considered. The deeper, confined zone of the Edwards aquifer, which is downgradient of the unconfined zone, is still susceptible to contamination from urban sources, though changes in water quality in this confined zone would likely occur over longer timescales (Musgrove and others, 2019). It is recognized and should be noted that a single study comparing the results of the analyses of data from two site pairs cannot support definitive conclusions about the full effect of urbanization on surface-water/groundwater quality; however, this study does provide useful insights about the spatial and temporal variability of stormwater runoff and unconfined groundwater that are consistent with a conceptual model that depicts the Edwards aquifer surface-water/groundwater system as a single water resource.



Pump hoist truck at the Shavano Park at Fawn Drive site.



U.S. Geological Survey hydrographers collecting a groundwater sample at the Encino Rio site.

References Cited

- City of San Antonio, 2020, Parks and facilities—Edwards aquifer: San Antonio Parks and Recreation web page, accessed April 22, 2020, at <https://www.sanantonio.gov/ParksAndRec/Parks-Facilities/All-Parks-Facilities/Gardens-Natural-Areas/Edwards-Aquifer>.
- Edwards Aquifer Authority, 2020, Edwards Aquifer Protection Program (EAPP)—Conservation easements: Edwards Aquifer Authority web page, accessed April 22, 2020, at <https://www.edwardsaquifer.org/aquifer-protection/groundwater-protection/conservation-easements/>.
- Greater Edwards Aquifer Alliance, 2020, Unique resource: Greater Edwards Aquifer Alliance web page, accessed April 22, 2020, at <http://www.aquiferalliance.net/unique-resource/>.
- Musgrove, M., Fahlquist, L., Houston, N.A., Lindgren, R.J., and Ging, P.B., 2010, Geochemical evolution processes and water-quality observations based on results of the National Water-Quality Assessment Program in the San Antonio segment of the Edwards aquifer, Texas, 1996–2006: U.S. Geological Survey Scientific Investigations Report 2010–5129, 93 p., accessed April 22, 2020, at <https://doi.org/10.3133/sir20105129>.
- Musgrove, M., Fahlquist, L., Stanton, G.P., Houston, N.A., and Lindgren, R.J., 2011, Hydrogeology, chemical characteristics, and water sources and pathways in the zone of contribution of a public-supply well in San Antonio, Texas: U.S. Geological Survey Scientific Investigations Report 2011–5146, 194 p., accessed April 22, 2020, at <https://doi.org/10.3133/sir20115146>.
- Musgrove, M., Opsahl, S.P., Mahler, B.J., Herrington, C., Sample, T.L., and Banta, J.R., 2016, Source, variability, and transformation of nitrate in a regional karst aquifer—Edwards aquifer, central Texas: *Science of the Total Environment*, v. 568, p. 457–469, accessed April 22, 2020, at <https://doi.org/10.1016/j.scitotenv.2016.05.201>.
- Musgrove, M., Solder, J.E., Opsahl, S.P., and Wilson, J.T., 2019, Timescales of water-quality change in a karst aquifer, south-central Texas: *Journal of Hydrology X*, v. 4, p. 1–16, accessed April 22, 2020, at <https://doi.org/10.1016/j.hydrox.2019.100041>.
- National Oceanic and Atmospheric Administration, 2020, Global Historical Climatology Network (GHCN): National Oceanic and Atmospheric Administration National Centers for Environmental Information database, accessed April 22, 2020, at <https://www.ncdc.noaa.gov/data-access/land-based-station-data/land-based-datasets/global-historical-climatology-network-ghcn>.
- Opsahl, S.P., 2012, Quality of surface-water runoff in selected streams in the San Antonio segment of the Edwards aquifer recharge zone, Bexar County, Texas, 1997–2012: U.S. Geological Survey Data Series 740, 20 p., 2 apps., accessed April 22, 2020, at <https://doi.org/10.3133/ds740>.
- Opsahl, S.P., Musgrove, M., Mahler, B.J., and Lambert, R.B., 2018, Water-quality observations of the San Antonio segment of the Edwards aquifer, Texas, with an emphasis on processes influencing nutrient and pesticide geochemistry and factors affecting aquifer vulnerability, 2010–16: U.S. Geological Survey Scientific Investigations Report 2018–5060, 67 p., accessed April 22, 2020, at <https://doi.org/10.3133/sir20185060>.
- Opsahl, S.P., Musgrove, M., and Mecum, K.E., 2020, Temporal and spatial variability of water quality in the San Antonio segment of the Edwards aquifer recharge zone, Texas, with an emphasis on periods of groundwater recharge, September 2017–July 2019: U.S. Geological Survey Scientific Investigations Report 2020–5033, 37 p., <https://doi.org/10.3133/sir20205033>.
- San Antonio Water System, 2020, Aquifer protection and evaluation: San Antonio Water System Protection and Compliance web page, accessed April 22, 2020, at https://www.saws.org/protecting-our-environment/water-resource-compliance-protection/aquifer_protection/.
- Schultz, A.L., 1994, 1994 review and update of the position of the Edwards aquifer freshwater/saline-water interface from Uvalde to Kyle, Texas: Edwards Underground Water District Report 94–05, 31 p., accessed April 22, 2020, at https://www.edwardsaquifer.org/science_docs/1994-review-and-update-of-the-position-of-the-edwards-aquifer-freshwater-saline-water-interface-from-uvalde-to-kyle-texas-plates/.
- Tremallo, R.L., Johnson, S., Hamilton, J.M., Winterle, J., Eason, S., and Hernandez, J.C., 2015, Edwards Aquifer Authority hydrologic data report for 2014: San Antonio, Tex., Edwards Aquifer Authority Report 15–01, 79 p., accessed April 22, 2020, at <https://www.edwardsaquifer.org/science-maps/research-scientific-reports/hydrologic-data-reports/>.
- U.S. Environmental Protection Agency, 2019a, Sole source aquifers for drinking water: U.S. Environmental Protection Agency web page, accessed April 22, 2020, at <https://www.epa.gov/dwssa>.
- U.S. Environmental Protection Agency, 2019b, Ground water and drinking water—National Primary Drinking Water Regulations: U.S. Environmental Protection Agency web page, accessed April 22, 2020, at <https://www.epa.gov/ground-water-and-drinking-water/national-primary-drinking-water-regulations>.
- White, W.B., 1988, Geomorphology and hydrology of karst terrains: New York, Oxford University Press, 464 p.

This fact sheet is based on the following report:

Opsahl, S.P., Musgrove, M., and Mecum, K.E., 2020, Temporal and spatial variability of water quality in the San Antonio segment of the Edwards aquifer recharge zone, Texas, with an emphasis on periods of groundwater recharge, September 2017–July 2019: U.S. Geological Survey Scientific Investigations Report 2020–5033, 37 p., <https://doi.org/10.3133/sir20205033>.

By Stephen P. Opsahl, MaryLynn Musgrove, and Keith E. Mecum

For more information, please contact

Director, [Oklahoma-Texas Water Science Center](https://www.usgs.gov/centers/tx-water)
U.S. Geological Survey
1505 Ferguson Lane
Austin, TX 78754
gs-w-txpublicinfo@usgs.gov
or visit <https://www.usgs.gov/centers/tx-water>

Publishing support provided by
Lafayette Publishing Service Center