

Prepared in collaboration with the National Park Service

Projected Sea-Level Rise and High Tide Flooding at Big Cypress National Preserve, Florida



Flooded wetlands in Big Cypress National Preserve (BICY), Florida, in August 2017. Photograph by National Park Service staff, BICY.


Overview

Objective

National parks and preserves in the South Atlantic-Gulf Region (<https://www.nps.gov/aboutus/index.htm>) contain valuable coastal habitats such as tidal wetlands and mangrove forests, as well as irreplaceable historic buildings and archeological sites located in low-lying areas. These natural and cultural resources are vulnerable to accelerated sea-level rise (SLR) and escalating high tide flooding (HTF) events. Through a Natural Resources Preservation Program-funded project during 2021–23, the U.S. Geological Survey (USGS), in collaboration with the National Park Service (NPS), estimated the probability of inundation at Big Cypress National Preserve (BICY) and several other parks under various SLR scenarios and contemporary HTF thresholds. The maps produced for this effort can be used to assess potential habitat change and explore how infrastructure and cultural resources within the park may be exposed to future flooding-related hazards. Though the study areas are different, the organization and wording of this publication are largely based on Thurman and others (2024a).

Big Cypress National Preserve

Located north of The Everglades on the western side of the Florida Peninsula, BICY covers 2,934.52 square kilometers of land (that is, area excluding water and aquatic bed wetlands from the National Oceanic and Atmospheric Administration [NOAA] Coastal Change Analysis Program [NOAA, 2019]) (fig. 1). Freshwater wetlands in BICY are essential to the health



Base from U.S. Geological Survey digital data
 Park boundaries from National Park Service (2024)
 Shaded bathymetry from U.S. Geological Survey, The National Map
 Orthoimagery from U.S. Department of Agriculture
 National Agriculture Imagery Program, 2017, 1.2 meters
 Universal Transverse Mercator, zone 17 north
 North American Datum of 1983

Figure 1. Location of Big Cypress National Preserve (BICY) within the State of Florida.

of The Everglades and the rich marine estuaries along Florida’s southwest coast (NPS, 2023). Because of SLR, the coastal habitats within BICY and other national parks and preserves in the region are at the forefront of concern. Specifically, these habitats are threatened unless they have the ability to adapt in place (for example, through vertical adjustment; Morris and others, 2002) or migrate to higher ground (for example, through tidal marsh transgression; Enwright and others, 2016). Similarly, SLR exposes important historic, cultural, and recreational sites to risk of damage or loss.

Relative Sea-Level Rise

SLR driven by global climate change is an ongoing and intensifying issue affecting low-lying coastal areas around the world (Sweet and others, 2022). Rates of SLR vary regionally because of factors such as the circulation and density of ocean water, rates of vertical land motion, and the location and volume of ice-mass loss from ice sheets and glaciers (Sweet and others, 2022). These localized rates, referred to as “relative SLR,” are differentiated from the global mean rate.

A recent multiagency report generated five potential global SLR scenarios based on the amount of greenhouse gases emitted into the atmosphere and the rate of ice-mass loss (Sweet and others, 2022). The five scenarios, termed “low,” “intermediate-low,” “intermediate,” “intermediate-high,” and “high,” each predict a different magnitude of future global mean SLR. Along with these global SLR scenarios, we used regional and local projections associated with each scenario to reflect relative SLR rates (fig. 2). Additionally, since there is uncertainty in how various natural processes will affect SLR, rates may vary within a given scenario. Whereas the median SLR projection can be

represented by a single value, the “likely” SLR can be represented by a range of values (that is, 17th to 83d percentile) (fig. 2).

High Tide Flooding

HTF occurs when spring tides (that is, maximum tides occurring during a full or new moon when the Earth, Moon, and Sun are aligned), high winds, or storms cause extreme water levels in coastal areas. For this effort, we used HTF thresholds defined by NOAA, which include thresholds designated “minor,” “moderate,” and “major” that vary on the basis of location (that is, latitude and longitude) (Sweet and others, 2022). Minor HTF events can cause disruptions such as flooding along trails and other infrastructure (fig. 3), moderate HTF events can cause greater disruptions and possibly damage, and major HTF events can cause destruction and lead to evacuations (NOAA, 2023).

Today, minor HTF events typically occur only a few days each year along the eastern Gulf of Mexico coast, and moderate and major HTF events typically occur less than once per year (Sweet and others, 2022). However, these events are exacerbated by relative SLR, and rapid increases in HTF frequency are expected within the next few decades (Thompson and others, 2021). By 2100, minor HTF is projected to occur more often than every other day under the intermediate-low SLR scenario, meaning that the threshold for a flood today could become the high tide level in the future (Sweet and others, 2018).

The increasing frequency and severity of HTF mean that even minor flooding events are no longer just a nuisance but are becoming a serious issue (Thompson and others, 2021). Cumulatively, the effects of many flooding events can be destructive to infrastructure and assets in low-lying areas (Sweet and others, 2018). Tidal flooding represents a contemporary effect of SLR and may be an indicator of challenges to come (Sweet and others, 2022).




Figure 2. Relative sea-level rise (SLR) scenarios for Big Cypress National Preserve (BICY), Florida, associated with the five global SLR scenarios from Sweet and others (2022). Sea-level change projected in this figure (modified from Thurman and others [2024b]) is relative to the year 1992.



Figure 3. Flooding at maintenance complex behind the headquarters of Big Cypress National Preserve (BICY), Florida, on September 29, 2022, the day after Hurricane Ian's landfall. BICY headquarters is the western visitor center in figures 4 and 5. Photograph by National Park Service staff, BICY.

Key Findings

Relative Sea-Level Rise

Inundation probabilities were mapped for BICY under the intermediate-low and intermediate-high SLR scenarios for the years 2050 and 2100 by using 1992 as the base year (fig. 4). For these scenarios and timesteps, the median SLR projection and likely range of values (that is, 17th to 83d percentile) are 33 centimeters (cm) for the intermediate-low scenario in 2050 (range: 27–38 cm), 67 cm for the intermediate-low scenario in 2100 (range: 58–77 cm), 43 cm for the intermediate-high scenario in 2050 (range: 33–57 cm), and 162 cm for the intermediate-high scenario in 2100 (range: 124–180 cm) (fig. 2) (Thurman and others, 2024b).

Under the intermediate-low scenario in 2050, about 10 percent of the park's nonwater area was classified as likely to be inundated (that is, probability greater than 0.66); under the same scenario, this percentage increased to about 16 percent by 2100 (fig. 4). For the intermediate-high scenario, about 12 percent of the park's nonwater area was classified as likely to be inundated by 2050 and about 32 percent by 2100 (Thurman and others, 2024b).

High Tide Flooding

Inundation probabilities were mapped for BICY under NOAA's minor, moderate, and major HTF thresholds (fig. 5). The minor threshold is 54 cm (plus or minus $[\pm]$ 19 cm, the root mean square error of linear regression, which is the square root of the average of the squared difference between projected values and actual values) above mean higher high water, and the moderate and major thresholds are 83 cm (\pm 25 cm) and 121 cm (\pm 39 cm) above mean higher high water, respectively (Sweet and

others, 2018, 2022). Under all three HTF thresholds, most of the park's nonwater area was classified as unlikely to be inundated. The likely inundation area increased from about 13 percent under the minor threshold to about 22 percent under the major threshold (Thurman and others, 2024b).

Applications of Maps

These maps of projected inundation from SLR and HTF can be used to assess current and future impacts of rising water levels to natural habitats, historic and cultural resources, and infrastructure within the park and can provide a resource for management decisions such as implementation of the resist-accept-direct (RAD) framework (Schuurman and others, 2020). Additionally, this mapping approach can be utilized to quantify and depict future habitat change driven by SLR, such as tidal saline wetland migration.

Methods

Data associated with this effort, along with details on the methods for the approximation of park areas that may be inundated from SLR and HTF, can be found in a USGS data release (Thurman and others, 2024b).

Acknowledgments

This Fact Sheet includes the results of research funded by the USGS and the NPS Natural Resources Preservation Program. We thank the NPS South Atlantic-Gulf Region and the staff of BICY, especially Thomas Forsyth and David Birdsall, for their input on SLR scenario choices.






Figure 4. Sea-level rise (SLR) inundation probabilities for Big Cypress National Preserve (BICY), Florida, under the SLR scenarios (Sweet and others, 2022) intermediate-low for the years *A*, 2050 and *B*, 2100 and intermediate-high for the years *C*, 2050 and *D*, 2100 (Thurman and others, 2024b). Sea-level change projected in this figure is relative to the year 1992. Probability is shown using three bins developed for the Intergovernmental Panel on Climate Change (2022): unlikely, probability less than or equal to 0.33; likely as not, probability greater than 0.33 and less than or equal to 0.66; and likely, probability greater than 0.66. Existing water areas were excluded from the analyses and area calculations.


A. Minor HTF threshold



B. Moderate HTF threshold



C. Major HTF threshold



- EXPLANATION**
- Inundation probability
 - Unlikely
 - Likely as not
 - Likely
 - Existing water area in 2015–17 (National Oceanic and Atmospheric Administration, 2019)
 - BICY (2,934.52 square kilometers land)
 - Visitor center

Base from U.S. Geological Survey digital data
 Park boundary and sites from National Park Service (2024)
 Universal Transverse Mercator, zone 17 north
 North American Datum of 1983

0 10 20 KILOMETERS
 0 10 20 MILES

Figure 5. High tide flooding (HTF) inundation probabilities for Big Cypress National Preserve (BICY), Florida, under the National Oceanic and Atmospheric Administration (Sweet and others, 2022) *A*, minor, *B*, moderate, and *C*, major HTF thresholds (Thurman and others, 2024b). Probability is shown using three bins developed for the Intergovernmental Panel on Climate Change (2022): unlikely, probability less than or equal to 0.33; likely as not, probability greater than 0.33 and less than or equal to 0.66; and likely, probability greater than 0.66. Existing water areas were excluded from the analyses and area calculations.

References Cited

- Enwright, N.M., Griffith, K.T., and Osland, M.J., 2016, Barriers to and opportunities for landward migration of coastal wetlands with sea-level rise: *Frontiers in Ecology and the Environment*, v. 14, no. 6, p. 307–316.
- Intergovernmental Panel on Climate Change, 2022, Climate change 2022—Mitigation of climate change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change: Cambridge, U.K., and New York, Cambridge University Press, 2,029 p., accessed February 2, 2024, at <https://doi.org/10.1017/9781009157926>.
- Morris, J.T., Sundareshwar, P.V., Nietch, C.T., Kjerfve, B., and Cahoon, D.R., 2002, Responses of coastal wetlands to rising sea level: *Ecology*, v. 83, no. 10, p. 2869–2877.
- National Oceanic and Atmospheric Administration [NOAA], 2019, C-CAP derived land cover—BETA [view 6 datasets—2015–2017 C-CAP derived 10 meter land cover—BETA]: National Oceanic and Atmospheric Administration dataset, accessed April 2022 at <https://coast.noaa.gov/digitalcoast/data/ccapderived.html>.
- National Oceanic and Atmospheric Administration [NOAA], 2023, What is high tide flooding?: National Oceanic and Atmospheric Administration web page, accessed February 2024 at <https://oceanservice.noaa.gov/facts/high-tide-flooding.html>.
- National Park Service [NPS], 2023, Big Cypress National Preserve, Florida: National Park Service web page, accessed May 2023 at <https://www.nps.gov/bicy/index.htm>.
- National Park Service [NPS], 2024, Administrative boundaries of National Park System units—National Geospatial Data Asset (NGDA) NPS national parks dataset: National Park Service dataset, accessed April 2024 at <https://irma.nps.gov/DataStore/Reference/Profile/2224545?Inv=True>.
- Schuurman, G.W., Hawkins-Hoffman, C., Cole, D.N., Lawrence, D.J., Morton, J.M., Magness, D.R., Cravens, A.E., Covington, S., O'Malley, R., and Fisichelli, N.A., 2020, Resist-accept-direct (RAD)—A framework for the 21st-century natural resource manager: Fort Collins, Colorado, National Park Service, Natural Resource Report NPS/NRSS/CCRP/NRR—2020/ 2213, 20 p., accessed July 14, 2023, at <https://doi.org/10.36967/nrr-2283597>.
- Sweet, W.V., Dusek, G., Obeysekera, J., and Marra, J.J., 2018, Patterns and projections of high tide flooding along the U.S. coastline using a common impact threshold: Silver Spring, Maryland, National Oceanic and Atmospheric Administration, National Ocean Service, NOAA Technical Report NOS CO-OPS 086, 56 p., accessed July 14, 2023, at https://tidesandcurrents.noaa.gov/publications/techrpt86_PaP_of_HTFlooding.pdf.
- Sweet, W.V., Hamlington, B.D., Kopp, R.E., Weaver, C.P., Barnard, P.L., Bekaert, D., Brooks, W., Craghan, M., Dusek, G., Frederikse, T., Garner, G., Genz, A.S., Krasting, J.P., Larour, E., Marcy, D., Marra, J.J., Obeysekera, J., Osler, M., Pendleton, M., Roman, D., Schmied, L., Veatch, W., White, K.D., and Zuzak, C., 2022, Global and regional sea level rise scenarios for the United States—Updated mean projections and extreme water level probabilities along U.S. coastlines: Silver Spring, Maryland, National Oceanic and Atmospheric Administration, National Ocean Service, NOAA Technical Report NOS 01, 111 p., accessed July 14, 2023, at <https://aambpublicoceanservice.blob.core.windows.net/ocean-serviceprod/hazards/sealevelrise/noaa-nos-techrpt01-global-regional-SLR-scenarios-US.pdf>.
- Thompson, P.R., Widlansky, M.J., Hamlington, B.D., Merrifield, M.A., Marra, J.J., Mitchum, G.T., and Sweet, W., 2021, Rapid increases and extreme months in projections of United States high-tide flooding: *Nature Climate Change*, v. 11, no. 7, p. 584–590, accessed July 14, 2023, at <https://doi.org/10.1038/s41558-021-01077-8>.
- Thurman, H.R., Enwright, N.M., Osland, M.J., Passeri, D.L., Day, R.H., and Simons, B.M., 2024a, Projected sea-level rise and high tide flooding at Timucuan Ecological and Historic Preserve, Florida: U.S. Geological Survey Fact Sheet 2024–3008, 6 p., accessed June 4, 2024, at <https://doi.org/10.3133/fs20243008>.
- Thurman, H.R., Enwright, N.M., Osland, M.J., Passeri, D.L., Day, R.H., Simons, B.M., Danielson, J.J., and Cushing, W.M., 2024b, Sea-level rise and high tide flooding inundation probability and depth statistics at Big Cypress National Preserve, Florida: U.S. Geological Survey data release, <https://doi.org/10.5066/P9JH8KQN>.
- By Hana R. Thurman,¹ Nicholas M. Enwright,² Michael J. Osland,² Davina L. Passeri,² Richard H. Day,² Bethanie M. Simons,¹ Jeffrey J. Danielson,² and William M. Cushing²
- ¹Cherokee Nation System Solutions, under contract to the U.S. Geological Survey
- ²U.S. Geological Survey

For more information about this publication, contact

Director, [Wetland and Aquatic Research Center](#)

U.S. Geological Survey

700 Cajundome Blvd.

Lafayette, LA 70506–3152

For additional information, visit

<https://www.usgs.gov/centers/wetland-and-aquatic-research-center-war-c>

Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Although this information product, for the most part, is in the public domain, it also may contain copyrighted materials as noted in the text. Permission to reproduce copyrighted items must be secured from the copyright owner.

Publishing support provided by

Lafayette Publishing Service Center

ISSN 2327-6932 (online)

<https://doi.org/10.3133/fs20243022>