

# DESCRIPTION OF THE LONDON QUADRANGLE.

## GEOGRAPHY.

*General relations.*—The territory represented by the London atlas sheet embraces an area of 950.4 square miles, extending from latitude 37° on the south to 37° 30' on the north, and from longitude 84° on the east to 84° 30' on the west. It includes, wholly or in part, the counties of Pulaski, Laurel, Rockcastle, Jackson, Lincoln, and Garrard, and it is named from London, the principal town within its borders. The adjacent quadrangles, so far as surveyed, are Richmond on the north, Beattyville on the northeast, Manchester on the east, Cumberland Gap on the southeast, and Williamsburg on the south.

In its geographic and geologic relations this quadrangle forms a part of the Appalachian province, which extends from the Atlantic coastal plain on the east to the Mississippi lowlands on the west, and from central Alabama to southern New York.

*Subdivisions of the Appalachian province.*—Respecting the attitude of the rocks, the Appalachian province may be divided into two nearly equal portions by a line which follows the northwestern side of the Appalachian Valley along the Allegheny front and the eastern escarpment of the Cumberland table-land. East of this line the rocks are greatly disturbed by innumerable folds and faults, and in many places they are so metamorphosed that their original form and composition can not now be determined. West of the division line the rocks are almost wholly sedimentary and the strata lie nearly flat, in the attitude in which they were deposited on the bottom of the sea. Since the western division lies almost wholly within the drainage basin of the Ohio River, it will be referred to in this description as the Ohio Basin.

*Ohio Basin.*—This portion of the province embraces the Cumberland Plateau and the Allegheny Mountains and the lowlands of western Tennessee, Kentucky, and Ohio. Its northwestern boundary is indefinite, but it may be regarded as coinciding with the Mississippi River as far up the stream as Cairo, and thence extending northeastward across the States of Illinois, Indiana, and Ohio to the western end of Lake Erie. Contrasted with the intensely folded strata of the Appalachian Valley the rocks of this region may be classed as horizontal, but, strictly speaking, they are rarely in this position, being gently inclined in various directions in different portions of the field. These slight undulations of the rocks have been produced by gentle uplifts which, though small by comparison, are pronounced geologic features of the region in which they occur.

The most prominent structural feature of the Ohio Basin is an arch in the strata, which has been styled the Cincinnati arch or anticline. The main portion of the fold enters the basin, as it is here outlined, from the direction of Chicago; it curves southward through Cincinnati and Lexington, Kentucky, and continues southwestward to Nashville, Tennessee. Originally the principal arch was supposed to extend northeastward from Cincinnati to Toledo, but evidence afforded by numerous oil and gas wells in that region has proved that the Toledo fold is only a small branch of the principal uplift. Stratigraphically the maximum development of this fold occurs in the vicinity of Lexington, where the Trenton limestone is exposed at an altitude of 1000 feet above sea level.

Geologically this arch separates the Ohio drainage basin into two parts, or structural basins, each of which contains coal-bearing rocks. The basin on the eastern side of the Cincinnati arch is generally known as the Appalachian coal field, and that on the western side as the coal field of western Kentucky or the central coal field of the United States. Besides these main structural features, the rocks of the Ohio Basin have been disturbed by a few small folds, and in places they have been broken by small faults.

*Topography of the Ohio Basin.*—The altitude of this division is greatest along the southeastern margin, where some of the ridges attain the

dignity of mountains. They are not continuous, and they do not form a system. At the north they constitute the Allegheny ranges, in the center they form a group of ranges limited on the northwest by Pine Mountain and on the southeast by Stone Mountain, and in the south the so-called mountains are only the escarpments of the Cumberland Plateau. The altitude of the mountainous belt varies from 500 feet in central Alabama to 2000 feet at Chattanooga, 3500 feet in the vicinity of Cumberland Gap, and from 2000 to 4000 feet throughout the northern portion of the Allegheny ranges.

From its extreme altitude on the southeastern margin the surface descends to less than 500 feet on the western border along the Mississippi River. This descent is not regular, but is accomplished by a number of steps or escarpments which mark the present extent of particularly hard beds and also the stages in the erosion of the surface to its present position. The highest and most pronounced escarpment is along the western margin of the Appalachian coal field, separating, in Kentucky, the great interior plain from the higher and more hilly region of the coal field, and, in Tennessee, marking the line between the eastern highlands and the Cumberland Plateau. In the latter State the escarpment is steep and regular and the plateau is very perfectly preserved, but in the former the rocks were not hard enough to protect the plain after it was uplifted, and as a consequence it was completely dissected by the numerous streams which drain its surface, leaving a hilly region in place of the plateau, and an irregular margin instead of an escarpment.

The great interior plain of Kentucky is continuous with both the eastern and the western highlands of Tennessee, and also probably with much of the highest land of Ohio and Indiana. Its general elevation along the western margin of the Appalachian coal field is about 1000 feet above sea level, but toward the west it probably descends to somewhat lower levels. The principal streams draining the Ohio Basin have cut deep channels below the surface of this plain, producing rugged topographic features in place of the gently undulating surface of the plain. In central Tennessee the drainage was especially active, and since the rocks exposed to the action of the streams were soft, the highland surface was entirely removed and a second plain was formed at a lower level. This is particularly well developed in the vicinity of Nashville, and it is known as the central basin of Tennessee.

Since the formation of the central basin, the land has been elevated again and the streams have cut deep gorges in its surface and deepened their old valleys in the region outside of the central basin.

*Topography of the London quadrangle.*—The London quadrangle is located upon the western margin of the Appalachian coal field, which at this particular point is destitute of its characteristic features. The hardest rocks do not reach outward to the margin of the field, and consequently the upper plain and the escarpment are lacking.

The major portion of the territory is drained by the Rockcastle River, which unites with the Cumberland River a few miles beyond the southern margin of the quadrangle. The Rockcastle River, throughout its entire course, flows in a narrow valley which is generally bordered on either side by perpendicular walls of sandstone or conglomerate. Below Line Creek its valley is extremely rugged and abounds in wild and picturesque scenery.

The southeastern portion of the territory is drained by Laurel River, another tributary of the Cumberland River. Near its headwaters this stream flows in a high, broad valley which is only slightly below the general level, but as it approaches the Cumberland it cuts deeper and deeper into the heavy conglomerate, until it also is confined by rocky walls from 100 to 400 feet in height.

Buck Creek, another tributary of the Cumberland, drains considerable territory in Pulaski County. It has cut through the thin capping of Coal Measure rocks, and flows in a deep gorge

cut wholly in the heavy-bedded Carboniferous limestone.

The only drainage not belonging to the Cumberland system is that of Dick River, which heads in Rockcastle County and flows northwestward into the Kentucky River just above Highbridge, Kentucky. In the London quadrangle this stream flows in a broad, flat valley, but its lower course is marked by a canyon like the canyons of the tributaries of the Cumberland River.

The general altitude of the surface of this quadrangle is not far from 1200 feet above sea level; above this altitude many knobs rise to a height of 300 feet, and below it the valleys are cut to depths sometimes as great as 550 feet. The topographic features found in this quadrangle, if considered apart from the general features of the region, can not be interpreted easily. When taken in connection with similar features in the surrounding territory, it is evident that a high-level peneplain once extended across this territory, the surface of which is now shown only in a few isolated hills in the northern portion of the quadrangle and in two or three similar eminences in the neighborhood of London. These stand at an altitude of 1500 feet, and they are presumably the sole representatives of the plateau which once existed here, and was continuous with the Cumberland table-land of Tennessee. Over most of this area the rocks then at the surface were too soft to preserve the peneplain, except in the northern portion, where the heavy conglomerates came to the surface and assisted in its preservation. In this portion of the area the limestone is much higher than it is farther south, and erosion would have had a more appreciable effect had it not been for the protecting cap of conglomerate.

In the vicinity of London the same peneplain was doubtless formed, but the rocks were not of adequate hardness to prevent its removal. In the epoch of active erosion which followed the formation of this peneplain the soft rocks were removed and a second regular surface was formed, but this surface is a structural plain, formed by a hard bed of conglomerate, which has not permitted the streams to cut lower.

Along the western side the presence of a large body of limestone above drainage level afforded opportunity for the streams to remove the surface to a second peneplain. Only a small area of this is seen in the quadrangle, but that is continuous with the great plain of central Kentucky. Below this plain, whose altitude is approximately 1000 feet above sea level, the gorges of the principal streams have been cut; hence they are necessarily a later feature than the plain itself.

Little direct evidence can be found concerning the geologic periods in which these various peneplains were formed. The uppermost plain is usually regarded as of Cretaceous and the lower as of post-Cretaceous age. Some evidence has been adduced which seems to limit the lower plain to the Eocene period, but at present its date can be regarded only as one of those unsettled questions which can not be definitely answered until more evidence is available.

## GEOLOGY.

### GENERAL SEDIMENTARY RECORD.

All of the rocks appearing at the surface within the limits of the London quadrangle are of sedimentary origin—that is, they were deposited by water. They consist of sandstones, shales, and limestones, having a total average thickness of 2000 feet. The materials of which they are composed were originally gravel, sand, and mud, derived from the waste of the older rocks and from the remains of plants and animals that lived while the strata were being laid down.

These rocks afford a more or less complete record of sedimentation from the upper part of the Silurian period to near the close of Carboniferous time. They also contain a record of the conditions of the land area which furnished the material for their formation. By knowing the conditions under which certain classes of rocks are formed, we can gain a fairly accurate idea of the distribution of land and

water and of the physical aspects of the land during the deposition of the rocks of this quadrangle.

The sea in which the Paleozoic sediments were laid down covered most of the Appalachian province and the Mississippi Basin. In the early stages of this era the eastern shore line of this ocean was probably along the Blue Ridge and the Smoky Mountains, but it migrated westward at intervals as the movements occurred which folded the rocks of the Appalachian Valley. Geologists do not agree concerning the westward migration of this shore line; many believe that the Coal Measure rocks were deposited entirely across the Appalachian Valley, and that they were uplifted as a whole at the close of the Carboniferous period. The original westward extent of these rocks is also an unsettled problem. Some believe that they were connected with the rocks of the same age in western Kentucky, while others are of the opinion that the Cincinnati arch formed an island in the Carboniferous sea over which the Coal Measures were never laid down.

The history of the continental area from which were derived the sediments now forming the rocks of the London quadrangle may at some future time be determined with considerable certainty, but at present our knowledge of the conditions of deposition will permit only the broadest generalizations.

From the outcrop of the lower Silurian limestones a few miles northwest of the London quadrangle, and from their presence in the Appalachian Valley southeast of this region, it is inferred that they are continuous from one outcrop to the other, and hence that they underlie this territory. The character of these rocks indicates that the territory represented by the London atlas sheet was deep sea during much of the early Paleozoic era.

The first indication of the rise of land areas in this body of water is found at the top of the Silurian sediments, but these formations are so greatly eroded in central Kentucky that it is impossible to locate the exact margin of the sea. In the closing stages of this era there may not have been any large body of land in this region, but there must have been at least islands upon which no deposition was taking place.

The conditions which prevailed during the deposition of the Devonian black shale have not yet been determined satisfactorily. In Pennsylvania and New York this epoch is characterized by immense deposits which have no representatives in the southern and western portions of the Appalachian province, except possibly in the fine-grained, black, carbonaceous shale of the Chattanooga formation. In this district the shale is 150 feet in thickness, but over much of the territory farther south it has a thickness of only a fractional part of the Kentucky measure.

Several theories have been advanced regarding the conditions which would permit of the deposition of only a few feet of carbonaceous shale in one locality while thousands of feet of sand and mud accumulated in another portion of the same province, but none has been accepted as entirely adequate. It has been argued that there are traces of shore formations in this shale on the flanks of the Cincinnati arch, and therefore that land probably existed in that locality during the deposition of the black shale, but no evidence, aside from the presence of fossil plants in this shale, could be found. The most satisfactory explanation assumes that the Devonian sea in the southern Appalachians was shallow, and received little or no sediment from the surrounding land areas and from the islands which existed in the area, because the land had previously been reduced to base-level. It was thus too low to afford fall by which streams might carry material to the sea.

The Waverly sea, which succeeded that of the Devonian period, also probably extended over much of the same district, but the conditions of the land were such that a liberal supply of waste was furnished for the formation of the Waverly shales. This sea deepened until almost the entire Appalachian province west of the Smoky Mountains and south of Pennsylvania was beneath its

Extent and counties.

General relations of topographic features and their significance.

Distinction between eastern and western divisions of the Appalachian province.

The Cincinnati arch.

Coal fields adjoining the Cincinnati arch.

Relief of the surface.

Ancient peneplain.

Second peneplain.

Ages of the peneplains.

Theories relating to the deposition of black shale.

Interpretation of the record in the rocks.

River systems.



surface, and limestone deposition was taking place over the entire area. This was followed by an interval in which muddy sediments were laid down, but the extent of the interval and the original thickness of the deposits are unknown. In the uplift which followed, the material just deposited was largely eroded; deep channels were cut in the land, extending in some cases through the shale and into the limestone to a depth of 100 feet.

This erosion interval represents, without doubt, the earlier portion of Coal Measure time, when the Cincinnati arch was dry land separating the two coal basins of Kentucky. At the beginning of the interval the sea occupied a basin much farther to the east than this region, and coal swamps flourished along its marshy borders, while the London quadrangle was dry land. Gradually the land subsided and the sea encroached toward the northwest, until finally it engulfed the land of this region. The advancing shore line was marked by accumulations of sand and gravel which filled the inequalities of the land and which have since been consolidated into sandstone and conglomerate. Whether this sea transgressed sufficiently to submerge the island of central Kentucky and connect with the western basin can not now be determined, but it is possible that it did, and that the sediments then laid down have been removed by erosion since the land was finally raised above the ocean level.

At the close of the Carboniferous period the strata were raised above the water and exposed to the action of the atmosphere. During the long periods which have since elapsed no marine sediments have been deposited on the surface, but the events of the passing ages are recorded in the forms sculptured from the land and in the river deposits. They have been suggested under the heading "Topography."

#### STRATIGRAPHY.

The strata exposed in the London quadrangle have a thickness of about 2000 feet. The thickness of the formations, the order of succession, and their general character are shown in the columnar sections, but a more detailed description of the individual beds and an indication of their probable equivalents in other fields are given in the following paragraphs.

#### SILURO-DEVONIAN STRATA.

In searching for oil and gas the drill has revealed the presence of a large body of rocks of Silurian age, but only the uppermost member shows at the surface within the limits of this quadrangle.

*Panola formation.*—This formation is named from the station of Panola, in Madison County, Kentucky. In its complete development it is a complex formation consisting of three members: coarse yellow sandstone at the base, fine blue shale in the middle, and a brown massive limestone at the top. The sandstone and shale are generally regarded as of Silurian, and the limestone as of Devonian age.

Only a small area of this formation outcrops in this quadrangle, and it is formed entirely of the uppermost member of the series. This member is the most constant, but occasionally it thins and disappears, leaving the black Devonian shale in contact with the blue calcareous shale of the Richmond formation. When fresh this limestone is of a bluish cast, but upon weathering it soon becomes brown. In places it is very cherty, but the chert is an irregular feature and frequently disappears in a distance of less than a mile. The outcrop on Dick River is the only exposure of this formation in the London quadrangle.

#### DEVONIAN STRATA.

Throughout the southern portion of the Appalachian province, and as far north as central Kentucky, the upper portion of the Devonian rocks is of unvarying lithologic character; it consists of black carbonaceous shale, which grows thinner and thinner toward the south, and which in places lies unconformably upon the rocks underneath. Northward from central Kentucky the Devonian increases rapidly in thickness, and many beds of coarse material appear in the mass of shale.

*Chattanooga shale.*—The Chattanooga forma-

tion is named from the city of the same name in eastern Tennessee, where it shows in typical form. It immediately overlies the Panola formation in a belt which crosses this territory from northeast to southwest. It outcrops on gentle slopes, and consequently its thickness is difficult of determination. In a well recently drilled at Broadhead it was found to be 150 feet in thickness, a measure which is probably more nearly correct than any that has been obtained at the surface. The line of separation between this shale and the limestone beneath is usually sharp and distinct, but in some localities in central Kentucky it is found interbedded with the limestone through a distance of about 20 feet.

The water carried by this shale becomes highly charged with mineral matter in solution, so that sulphur, chalybeate, and alum springs abound near it. It weathers rapidly into a white soil which is extremely poor—almost unfit for agricultural purposes. The excessive blackness of the fresh shale, its well-known bituminous character, and the presence of occasional thin seams of coaly matter have led many persons to search in this formation for coal, but no seams of consequence have ever been found.

#### CARBONIFEROUS STRATA.

Rocks of this age occupy almost the entire area of the London quadrangle. Although belonging to one geologic period, they have been separated into two series, Mississippian and Pennsylvanian.

#### MISSISSIPPIAN SERIES.

The rocks of this series are mainly of marine origin, and distinctly underlie the lowest coal-bearing stratum.

*Waverly shale.*—This formation overlies the Chattanooga shale, and is easily separated from the latter on account of its color. At its base the Waverly is a light-blue clay shale, which passes upward into sandy shale and argillaceous sandstone. The shale at the base abounds with light-blue or drab ironstone concretions, which on weathering change to a dark reddish brown. In many cases they have been mistaken for volcanic rocks on account of their dark color and their extreme toughness.

Over most of this quadrangle the Waverly is characterized by a great number of siliceous concretions, which ordinarily are only a few inches in diameter, but which occasionally attain a diameter of two feet. So numerous and so large are they in the northern portion of this area that they have given the name to Roundstone Creek, one of the principal tributaries of Rockcastle River. In the vicinity of this creek they occur at the top of the formation, but toward the south they are more abundant at other horizons, having been replaced at the top of the formation by a coarse yellow sandstone which is well shown at the crossing of Buck Creek west of Dallas.

The name Waverly is derived from Ohio, where it was used in designating this formation in the early surveys of that State. It makes a large portion of the hilly district in the northwestern corner of the London quadrangle, and it extends along the bottom of the valleys toward the southeast, far within the limits of the area where the limestone prevails. It has an average thickness of about 350 feet.

*Newman limestone.*—This formation is named from Newman Ridge, Hancock County, Tennessee, a type locality on the eastern side of the Appalachian coal field. The limestone in the London quadrangle is the representative, though possibly not the equivalent, of the Newman limestone in the type locality. In the type locality it is nearly 1500 feet thick, while in this quadrangle it is only about 250 feet thick.

It is a compact blue limestone with cherty beds at the base, and it varies in thickness from 100 to 250 feet. This great range in thickness is due to the fact that at the beginning of Pennsylvanian time a part of the area now occupied by the Newman limestone was land, and that much of the underlying rock was removed by erosion. The deep channels cut in this formation were, upon the return of the sea, filled with sand and gravel, so that at present they are apparent only when numerous measures of the limestone are compared.

The limestone outcrops in a wide band across the quadrangle from northeast to southwest. Where this rock forms the entire surface it makes a gently rolling country, but where it forms the

slopes of the valleys it frequently produces perpendicular cliffs. Its solubility leads to the formation of many caves, sinks, and underground channels. Even Roundstone Creek finds its way to the river, during ordinary stages of water, through such a channel.

*Pennington shale.*—This shale was so named from a water gap through Stone Mountain in Lee County, Virginia. The formation consists principally of calcareous shale, but it also includes many thin beds of impure limestone. It is variously colored, but greens and reds predominate. It succeeds the Newman limestone by gradual transition, and it is unconformably overlain by the Lee formation. In many places in this quadrangle it was entirely removed by erosion before the sediments of the Lee formation were laid down. Its greatest exposed thickness is about 90 feet.

This formation does not cover, in its outcrop, any appreciable amount of territory, for it is exposed usually on the steep slopes of the stream valleys, and it is eroded from all of the uplands which were originally covered by it.

#### PENNSYLVANIAN SERIES.

From an economic standpoint these are the most important rocks which come to the surface within the limits of this quadrangle, for coal is the principal mineral resource of this section. The strata consist entirely of sandstone and shale, which rest unconformably upon the rocks beneath, and are limited in their upward extent only by the amount of erosion which has occurred since they were deposited. They show great variation in character and thickness, and are evidently the result of shore or shallow-water deposition.

*Lee formation.*—This formation is named from Lee County, Virginia. It includes the basal portion of the coal-bearing rocks which appear on the western side of the Appalachian Basin, and is equivalent to the middle portion of the Pottsville series of Pennsylvania. The formation consists of sandy shale and sandstone, in which occur lentils of coarse material—massive sandstone and conglomerate—the whole ranging from 200 to 600 feet in thickness. Two of these lentils are especially heavy, constituting, topographically, the most important members of the formation. The Lee formation rests upon the eroded surface of the Newman limestone and the Pennington shale. The extent of the interval which separates these formations, and which is represented only by the unconformity, has never been determined with much exactness, but the evidence of fossil plants indicates that probably one-quarter of the Pottsville series is wanting on the western side of the Appalachian Basin in central Kentucky. The erosion interval undoubtedly represents a period of time in which the limestone area was dry land and the Pennsylvanian sea was located some distance to the eastward. Later, subsidence of the land allowed this sea to transgress upon the land, forming successive overlaps toward the west; each position of the shore was marked by sand and gravel, which were sorted by the waves and finally laid down as sandstone and conglomerate, while sandy mud was being deposited in the waters off shore. The result of these conditions is the local development of heavy beds in narrow belts along the margin of that ancient sea, which grade off in either direction into materials of entirely different composition. There are also apparent unconformities in the Lee formation itself which are indicative of oscillations during the influx of water by which some formations were eroded and others were deposited in their stead. The generally arenaceous character of the Lee formation and the irregularities of the beds which compose it render extremely difficult the separation of individual beds which can be traced over any considerable distance and represented on the map.

*Rockcastle conglomerate lentil.*—This is a heavy bed of sandstone or conglomerate extending from central Tennessee to the middle of this quadrangle, and is named from Rockcastle River, along which it is particularly prominent. It is the most conspicuous member of the Pennsylvanian series in the southern portion of this quadrangle. It forms high cliffs along Rockcastle River and its numerous tributaries from Line Creek to Cumberland River and up that stream as far as the falls, where it passes beneath water

level. This conglomerate is terminated abruptly toward the north along a line which crosses Wood Creek a mile above its junction with Hazel Patch Creek, Hawk Creek in the middle of its course, and the mouth of Line Creek. Toward the west it does not extend as far as Buck Creek, although rocks of this same general horizon are found in the hills west of that stream. Along the margin of the field in the southwestern corner of the quadrangle there are local developments of conglomerate which lie apparently at lower horizons. They are generally thin and discontinuous, and therefore are not worthy of representation on the map.

In the northern portion of the quadrangle there are heavy deposits of conglomerate at various points which have been provisionally correlated with the Rockcastle. The greatest development occurs in a belt about 4 miles in width which extends northward from Livingston nearly along the main course of Roundstone Creek. It continues northward for a distance of 6 miles beyond the margin of this territory, where it is terminated by the northern escarpment of the coal field.

This conglomerate is one of the most interesting features of the region, for it occupies a channel which was eroded through the Pennington shale and at least 100 feet into the Newman limestone. The channel was filled with well-rounded pebbles, which now form a mass of conglomerate, the top of which is frequently lower than the top of the limestone. The conglomerate varies in thickness from 150 feet in the center of the channel to zero on the edges of the mass. The channel in which this occurs was certainly eroded in the limestone before the deposition of the Pennsylvanian series and when this area was dry land; upon submergence, the channel was filled with gravel, which has since been slightly cemented into the conglomerate of to-day. It descends southward with the general dip of the rocks in this region, and passes beneath drainage level in Rockcastle River a little above the mouth of Roundstone Creek. This isolated outcrop of conglomerate is here provisionally correlated with the Rockcastle of the southern portion of the quadrangle, but the connection is merely assumed from the fact that the Livingston mass lies where the known Rockcastle would be found if continued. The stratigraphic relations of the two beds are different, but it seems probable that the Livingston area is merely an outlier of the main body and that it overlaps upon the eroded surface of the limestone beneath.

There are one or two masses of conglomerate in the valley of Horse Creek which occupy similar positions with reference to the surrounding rocks. No connection could be found between these isolated exposures and the main mass, but they have been regarded as the probable extension of the Rockcastle conglomerate.

*Corbin conglomerate lentil.*—This is named from Corbin, Whitley County, Kentucky, and, like the Rockcastle, it occupies a marginal position with reference to the coal basin. It extends 40 or 50 miles into Tennessee, where it is thin and irregular, but it develops into a massive bed from 100 to 200 feet in thickness in this portion of Kentucky. In the southeastern corner of the quadrangle the Corbin conglomerate has an extensive areal development, but farther north it is limited to a narrow belt, the western margin of which is formed either by the limit to which erosion has removed the conglomerate or by the margin of the original deposit; the eastern margin is formed by the line along which this formation goes under cover, or where it changes in character from a coarse conglomerate to a sandstone and then to a sandy shale. South of London the formation passes below drainage level before its character changes, but north of that place it fades gradually into shale before its horizon takes cover.

This conglomerate can be recognized generally by its peculiar pink or flesh color and by the softness of the cement with which the individual grains are bound together. Owing to this characteristic, it seldom forms cliffs, but weathers into rounded bosses and domes, which sometimes surprise the observer, who finds himself upon one of these outcrops without being aware that he has crossed an exposure of conglomerate. It is not always conglomeratic, but it preserves the same general character whether the component grains are sand or gravel. In the southern portion of the quadrangle this bed is not favorably disposed



for erosion, and consequently it forms the foundation for an extensive area of nearly level land; in the northern portion the distance between it and the limestone is less, allowing the streams to cut deep channels in the limestone and leaving the remnants of the conglomerate as mere caps to the hills.

The Lee formation is nearly equivalent to the Pottsville series of Pennsylvania, but it does not contain either the superior or the inferior limits of the type. Probably as much as one-quarter of the lower portion of the series is wanting in this section, being represented by the erosion interval at the base of the Pennsylvanian series. According to the evidence afforded by fossil plants the top of the Pottsville series occurs in a thick bed of shale which overlies the Corbin conglomerate.

**Breathitt formation.**—This formation includes all of the Carboniferous rocks lying above the Corbin conglomerate, or the top of the Lee formation. It is composed of shale and sandstone with occasional coal seams, but no individual bed is of sufficient importance to be shown as an independent formation. In the highest hills in the vicinity of London this formation shows about 550 feet in thickness. It is named from Breathitt County, Kentucky, where the formation is present in great force.

#### STRUCTURE.

To the eye of the observer the rocks of this quadrangle appear to lie horizontal, but when they are examined in detail and the altitude of one outcrop is compared with that of another, it is evident that the strata are seldom, if ever, in that position. The rocks were formed at the bottom of the sea, and since the sea bottom has generally less diversity of altitude than the present rock strata, it is evident that their present position is due to movement in the crust of the earth.

**Definition of terms.**—The strata when compared with a horizontal plane are found to be inclined. The inclination is known as the *dip* of the rocks. In the process of deformation the rocks have been thrown into arches and troughs. In describing these folds the term *syncline* is applied to the downward-bending trough, and the term *anticline* to the upward-bending arch. A synclinal axis is a line running lengthwise in the synclinal trough and at every point occupying its lowest part, toward which the rocks dip from either side. An anticlinal axis is a line which occupies at every point the highest portion of the anticlinal arch, and from which the rocks dip to either side. The axis may be horizontal or inclined. Its departure from the horizontal is called the *pitch*, and is usually only a few degrees in amount.

As a result of the strains and stresses which have affected the crust of the earth, the strata in many places have broken along certain lines, and the rocks on one side of the break have been lifted or depressed with reference to those on the other side. Where the rocks have been intensely folded, as in the Appalachian Valley, the breaks have developed from the compressed and overturned folds; but in the Ohio Basin the faults are due to tension, or the stretching of the strata. Faults of the former type are sometimes of great linear extent and of enormous displacement, and those of the latter are in this district generally short and of very slight displacement.

In addition to the crustal movements which have perceptibly deformed the rocks of this region, the province has been affected by vertical movements which have repeatedly elevated and depressed the surface of the land, but by amounts which are insignificant compared with the magnitude of the folds. These slight movements were not continuous, but occurred now and then, the periods of greatest activity being separated by intervals of quiet in which the agents of erosion had time to record their action on the face of the land.

**Structure of the London quadrangle.**—From the nearly horizontal position of the rocks in this territory it is apparent that the structure is very simple. Since the quadrangle lies on the north-

western margin of the Appalachian coal basin, the dips of its strata are generally toward the southeast throughout the quadrangle. There are variations from this regular southeastward descent of the rocks, but such exceptions are local, and have no effect on the structure as a whole. There are also such irregularities as the channels in which the conglomerate is deposited north of Livingston, but these are due to erosion and unconformity between the beds and not to disturbance. The rate at which the rocks dip varies with the position in the field. The northwestern half of the area lies upon the flank of the Cincinnati arch, and consequently the dip is much more pronounced in that locality than in the southeastern half, which lies farther within the coal basin.

In drilling some deep wells in this district it was found that the limestone dips more steeply in the southeastern portion of the quadrangle than do the coal-bearing rocks of the Pennsylvanian series. This is explained by the overlap at the base of the latter series, which permitted sedimentation to go on in the center of the basin while the margin was land and received no deposits. It has long been known that the Pottsville series increases in thickness from a few hundred feet on the western side of the field to 1500 or 2000 feet on the eastern side, but no one could say positively how this change was accomplished. It has lately been demonstrated, by study of the fossil plants of this series, that a large portion of the base of the series is entirely absent from the western margin of the Appalachian coal basin. Hence, in passing from the western edge of the field toward the center, older beds appear below those which form the margin of the field. The limestone passes beneath the lowest Pottsville beds as they develop toward the center of the basin, and consequently dips more steeply.

**Structure sections.**—The sections upon the structure sheet represent the strata as they would appear in the sides of a deep trench cut across the country. The sections are located at the upper edges of the blank spaces, along the lines A A and B B. The vertical and horizontal scales are the same, so that the actual form and slope of the land and the dips of the strata are shown. Minute details of structure can not be shown on a map of so small scale; therefore the sections are somewhat generalized from the dips observed in a belt of country a few miles in width along the line of the section.

#### MINERAL RESOURCES.

Rocks and minerals of economic importance are not very plentiful in this quadrangle. Coal, building stone, limestone, road metal, clay, and iron ore have been found within its limits, but only the first two have been developed beyond the local needs of a farming community.

**Coal.**—This is by far the most important mineral resource of the quadrangle; it has been mined for a number of years at Pine Hill and Livingston, but these deposits are now practically exhausted, and the production of coal is limited to the neighborhood of Pittsburg, where eight or ten companies are actively engaged in mining steam and domestic fuel.

It is a popular belief that coal seams are coextensive with the strata of which they form a part, and consequently that the existence of a workable bed in any locality is evidence that it can be found of the same thickness and quality in all adjacent regions. In order to search intelligently for coal it is essential that the prospector and operator should understand the mode of deposition of coal, its limitation in distribution, and its variability in composition and thickness. Coal is the result of great vegetable accumulations in swamps, and since swamps are usually of limited extent, the coal deposits resulting therefrom will necessarily be also limited in their range, and will not be coextensive with the overlying and underlying strata. Again, swamp vegetation accumulates usually on a somewhat uneven floor, and the seam will show the effect of the

inequalities upon which it rests, being thickest in the center of the area, or in the deeper portions of the swamp, and thinning toward the margin. All of the conditions which affect the quality of the coal are not known, but, from the mode of origin, it is probable that physical conditions varied in different portions of a coal swamp, and hence it is hardly likely that the quality of the coal will be constant throughout any very extensive field. On the whole, variations in coal seams appear to be the rule, and continuity and regularity the exception.

In the London quadrangle the deposits of coal are apparently very irregular; they occur at many horizons in the coal-bearing rocks, but they are generally local, and it seldom happens that more than one seam of workable thickness is found in any locality.

Owing to the unconformity at the base of this series and to the irregularity of the various strata composing the productive beds, it is almost impossible to correlate the coal outcrops found in this quadrangle, for the reason that there is no datum from which to measure. Since this is the case, no effort will be made to determine equivalency over broad areas, but the outcrops will be described individually.

In a belt a few miles in width lying along the southern margin of this quadrangle there is a prominent coal seam immediately beneath the Rockcastle conglomerate of the Lee formation, which has been worked at intervals for twenty-five or thirty years along the Cumberland River. It was found in a number of ravines south of Mount Sterling knob, showing 3 or 4 feet of splinty coal, but its full thickness could not be determined. It was also observed near the mouth of Cane Creek, but it was only partially exposed to view. Several outcrops are reported on Rockcastle River, but they were not visited, and the extent of the seam is problematical.

There is a lower seam in this district, but it is not known to be of workable thickness in this quadrangle. On the bluffs fronting Cumberland River it was worked years ago, and its thickness at this point is reported to be 30 inches. The probable equivalent of one of these seams has been worked along the western border of this quadrangle, near the main road from Somerset to Dykes, but its thickness at this point is not known.

Northward from Dykes there is a poor showing of coal in this formation. Several small seams show in outcrop, but they are generally too thin to be of commercial importance. At Livingston there were formerly extensive works on two small seams only a few feet above the limestone, but they are now abandoned, as the deposits of workable coal are exhausted. The mines at Pine Hill, on presumably the same seam, are also abandoned, owing to the exhaustion of the pocket of coal upon which they were opened. Isolated outliers of what appears to be the same seam occur in the vicinity of Mount Vernon, but they are probably too limited in extent to be of much value.

In the territory north of the main Rockcastle River there are a number of small coals in the Lee formation. They have been described in detail in the State report on Jackson and Rockcastle counties. Five seams have been recognized, some of which may possibly have a workable thickness over a portion of this area, but none of them are of great promise. Their general thickness is less than 3 feet, and since they outcrop in an exceedingly rough section of the country, it will presumably be a long time before they are developed. The one of most promise seems to be the cannel seam of Horselick Creek. It was not seen by the writer, but if it has the thickness given in the State reports it will be a valuable deposit when transportation can be secured.

Decidedly the most important seam of coal in this quadrangle occurs within a few feet of the base of the Breathitt formation. It is known as the Pittsburg or Altamont coal. It is of workable thickness only over a small area which extends along the railroad from London to Altamont, and from near the line

of the railroad to Raccoon Creek. It was formerly opened and worked to some extent at Lily, but the seam was too thin for profitable mining and has been abandoned. This locality is evidently on the southern edge of the coal swamp, for in a cut south of town it shows only 12 inches in thickness. In its best development, near Pittsburg, it ranges in thickness from 36 to 41 inches. It is mined by eight or ten companies in and about Pittsburg, and finds a ready sale as a steam and domestic fuel.

A few small seams occur in the country east of Pittsburg, but it is probable that they are of no present commercial value.

**Building stone.**—Along the valley of Roundstone Creek the Waverly formation carries a bed of fine building stone, which has been quarried at Langford for a number of years. It consists of a fine-grained bluish sandstone, which is soft when freshly quarried but which soon hardens by the action of the elements into a very durable stone. Its principal defect is the thinness of the bed of workable stone, which renders quarrying unprofitable as soon as the cover becomes of moderate thickness.

Stone for rough work could be found at a number of places, both in the limestones of the Mississippian and in the sandstones of the Pennsylvanian series, but it has never been used except for local purposes.

**Limestone.**—This kind of rock is abundant in the western portion of the quadrangle. Much of it is of good quality for the manufacture of lime, and also for use as road metal, but it has been utilized only in a small way.

**Miscellaneous.**—Residual clay of good quality for the manufacture of brick could be obtained at many points on the limestone outcrop, but it has never been used. Fire or under clay accompanies many of the coal seams, but it has never received attention, and its value is not accurately known. Surface indications of iron ore are common, but no deposits of value are known in the territory. From time to time interest has been manifested in exploration for oil and gas, and several deep wells have been drilled in this region, but so far as known the results are unsatisfactory and the wells have been abandoned.

#### SOILS.

The surface of this quadrangle is too deeply cut by streams to make good farming lands. There are, however, two areas of low relief which, so far as the surface is concerned, are tolerably well adapted to agricultural pursuits, but which are very different in value on account of the difference in the soil which is found upon them. One of these areas is composed of the outcrop of the Newman limestone in the western portion of the area. The soil derived from this formation is fairly good, and the region has the appearance of a prosperous agricultural section, but this limestone appears to be almost if not quite destitute of phosphate, and consequently the soils can not compare with those of the "Bluegrass" region, which are derived from the Silurian limestones. The soil characterizing the Newman limestone is easily exhausted, and when once worn out will not renew itself, but requires artificial fertilizers.

The second area of moderately low relief is in the southeastern corner of the quadrangle, and its soils are formed from the decay of the sandstones and shales of the coal-bearing rocks. The soils derived from such rocks are poor and thin, and ill adapted to any sort of agricultural pursuit.

Along the Rockcastle River the valleys are generally floored by the Newman limestone or the Pennington shale, but the extreme narrowness of the valleys and the steepness of the slopes render them of little value to the farmer.








The Waverly formation produces hilly lands and a soil so poor that it is almost valueless for farming purposes.

M. R. CAMPBELL,  
Geologist.

February, 1898.

COLUMNAR SECTIONS

GENERALIZED SECTION FOR THE NORTHERN HALF OF THE LONDON QUADRANGLE. SCALE: 500 FEET = 1 INCH.						
PERIOD.	FORMATION NAME.	SYMBOL.	COLUMNAR SECTION.	THICKNESS IN FEET.	CHARACTER OF ROCKS.	CHARACTER OF TOPOGRAPHY AND SOIL.
CARBONIFEROUS	(Corbin conglomerate-lentil.)	(Clec)		0-150	Conglomerate or coarse pink sandstone.	Rounded ridges. Sandy soil.
	Lee formation.	Cle		300-600	Sandy shale and sandstone with a few seams of coal.	Hills and ridges. Generally poor soil.
	(Rockcastle conglomerate-lentil.)	Cler		0-150	Coarse conglomerate.	Cliffs. Sandy soil.
	Pennington shale.	Cpn		0-60	Red and green shales and thin limestone.	Clay soil.
	Newman limestone.	Cn		100-250	Blue limestone with a few nodules of chert. Cherty limestone.	Cliff and hill lands. Generally fertile soil where slopes are not too steep.
	Waverly formation.	Cwv		350	Greenish, calcareous and argillaceous sandstones. Light-blue clay shale with iron concretions.	Hilly lands. Very poor soil.
SIL. DEV.	Chattanooga shale.	Dc		150	Black carbonaceous shale.	Valleys. Poor soil.
	Panola formation.	SDp		20+	Brown limestone, generally cherty, at the top; light-blue clay shale below.	Valleys. The shale forms poor soil and bad roads.

GENERALIZED SECTION FOR THE SOUTHERN HALF OF THE LONDON QUADRANGLE.						
SCALE: 500 FEET = 1 INCH.						
PERIOD.	FORMATION NAME.	SYMBOL.	COLUMNAR SECTION.	THICKNESS IN FEET.	CHARACTER OF ROCKS.	CHARACTER OF TOPOGRAPHY AND SOIL.
CARBONIFEROUS	Breathitt formation.	Cbt		500	Sandy shale and coarse ferruginous sandstone.  Pittsburg, Ky., coal seam at the base.	Hilly country, with gentle slopes and rounded summits. Soil fair on shale outcrop; poor on sandstone.
	(Corbin conglomerate-lentil.)	Clec		0-150	Conglomerate or coarse pink sandstone.	Cliffs. Sandy soil.
	Lee formation.	Cle		500-1000	Sandy shale and sandstone with a few seams of coal.	Gently rolling uplands in the vicinity of London, and ridges near Rockcastle River.
	(Rockcastle conglomerate-lentil.)	(Cler)		(0-150)	(Coarse conglomerate.)	Generally poor soil. (Cliffs. Sandy soil.)
	Pennington shale.	Cpn		0-150	Red and green shales and thin beds of limestone.	Valleys or slopes. Clay soil, sometimes fertile.
	Newman limestone.	Cn		225	Blue limestone with a few nodules of chert. Cherty limestone.	Valleys. Generally fertile soil.
	Waverly formation.	Cwv		100+	Calcareous sandstone.	Rocky valleys.

NAMES OF FORMATIONS.

PERIOD.	NAMES AND SYMBOLS USED IN THIS FOLIO.		MARIUS R. CAMPBELL: RICHMOND FOLIO, U. S. GEOLOGICAL SURVEY, 1896.		GEOLOGICAL SURVEY OF KENTUCKY: REPORTS ON LINCOLN AND GARRARD COUNTIES, 1882, BY W. M. LINNEY; REPORT ON ROCKCASTLE COUNTY, 1891, BY GEO. M. SULLIVAN.	
CARBONIFEROUS	Breathitt formation.	Cbt				
	Corbin conglomerate-lentil.	Clec	Corbin conglomerate-lentil.		Conglomerate.	
	Lee formation.	Cle	Lee formation.			
	Rockcastle conglomerate-lentil.	Cler	Rockcastle conglomerate-lentil.			
	Pennington shale.	Cpn	Pennington shale.		Upper Sub-Carboniferous.	
	Newman limestone.	Cn	Newman limestone.			
SIL. DEV.	Waverly formation.	Cwv	Waverly formation.		Lower Sub-Carboniferous.	
	Chattanooga shale.	Dc	Chattanooga shale.		Black shale.	
	Panola formation.	SDp	Panola formation.		Corniferous limestone. Crab Orchard shale.	

MARIUS R. CAMPBELL,  
*Geologist.*