

# DESCRIPTION OF THE NORFOLK QUADRANGLE.

By N. H. Darton.

## GEOGRAPHY.

*General relations.*—The Norfolk quadrangle embraces the region lying between the parallels 36° 30' and 37° north latitude and the meridians 75° 30' and 76° 30' west longitude. It measures approximately 35 miles from north to south and 56 miles from east to west, and contains about 1913 square miles, of which about one-half is ocean and bay and the other half is land with inlets. In Virginia it comprises Norfolk, Princess Anne, and small portions of Nansemond, Warwick, and Elizabeth City counties. In North Carolina it includes the northern margins of Currituck and Camden counties. The land portion is a low plain extending to the Atlantic Ocean on the east and to Chesapeake Bay and James River on the north. It is deeply invaded by tide water in the channels of James, Nansemond, Elizabeth, Lynnhaven, North Landing, and Northwest rivers, and in Back Bay, North Bay, and some minor waters. Its fresh-water streams are small and reach tide water at no great distance from their sources. The quadrangle includes the greater part of the Dismal Swamp with its central body of water, Lake Drummond.

*Coastal Plain province.*—The eastern portion of the Atlantic slope of the United States embraces two distinct provinces: the Piedmont Plateau, a region of undulating plains, extending eastward from the Blue Ridge with gradual declivity and underlain by ancient crystalline rocks; and the Coastal Plain, a province bordering the ocean, deeply invaded by tide-water estuaries and underlain by gently east-dipping unconsolidated strata from early Cretaceous to latest Pleistocene in age. The Norfolk quadrangle lies entirely within the Coastal Plain province.

The Coastal Plain continues its slope eastward beyond the Atlantic shore line, its eastern margin lying at a moderate depth at the edge of the continental plateau, 100 miles offshore, where it terminates in the great continental slope, 3000 to 10,000 feet high. From the eastern portion of the Piedmont Plateau there extends across the Coastal Plain to the verge of this slope a very smooth and even surface, inclining gently southeastward and broken only by the broad, shallow valleys of the rivers, in the larger of which tide water reaches as far west as the eastern margin of the crystalline rocks of the Piedmont Plateau.

The highest altitudes in the Coastal Plain province are about 400 feet above sea level; its submarine margin is about 300 feet below sea level. So gentle is the inclination and so perfect the unity of the plain that if the land were elevated or depressed 100 or 200 feet the shore line would simply be shifted about the same number of miles. Thus the position of the coast may be considered an accident of the present slope and altitude of the land—indeed, between the mouth of Hudson River and Chesapeake Bay the present coast does not coincide with the trend of the province, but cuts obliquely across it, so that, while only about half the province is submerged in the latitude of Richmond, it is nearly all beneath the ocean in the latitude of New York. Below tide level the province is an even and nearly level sea bottom; above tide level it comprises lowlands of broad, flat terraces which skirt the coast and the estuaries, and extensive areas of higher plateau surfaces lying between the large valleys and rising gradually westward. The principal waterways are broad, shallow estuaries, flanked sometimes by tidal marshes, sometimes by low sea cliffs, and demarked from the higher plateau surface by moderately steep slopes. The lesser waterways are commonly estuarine in their lower reaches, but narrow and steep bluffed in their upper portions, frequently heading in nar-

row ravines cut sharply in the extensive plains of the divides.

## GEOGRAPHIC DIVISIONS.

*Low, level plain.*—The principal feature of the Norfolk quadrangle is a very level terrace elevated from 10 to 20 feet above sea level and intersected by extensive tide-water areas and a few shallow valleys of fresh-water streams. Upon this plain to the westward is the Dismal Swamp, and to the north and east, along the bay and ocean shores, there are sand dunes, which have a height of 70 feet at Cape Henry. The general terrace is part of a low plain which constitutes the surface of all the Eastern Shore of Virginia and Maryland. It also forms a moderately wide margin along the western shore of Chesapeake Bay, extends up James River in low terraces as far as Richmond, and, in the area of the Norfolk quadrangle and adjoining region, reaches as far as the western edge of Nansemond County. This plain is traversed by James River and Hampton Roads. It is crossed transversely by a shallow trough which is occupied on the north by the estuarine channel of Elizabeth River and Southern Branch, and on the south by North Landing River. In the middle of the depression, from North Landing to Great Bridge, there is a swamp, through which the Albemarle and Chesapeake Canal has been excavated. The plain is trenched by a number of valleys reaching tide water in their lower portions. Of these Nansemond River, Western Branch, Eastern Branch, Tanner Creek, Mason Creek, Little Creek, Lynnhaven River, and Broad Bay flow to the north into James River or Chesapeake Bay; and Northwest River and its little branches, together with North Landing River, flow southward into Currituck Sound. Knott Island is an outlying portion of the terrace level, cut off by tide-water inlets which join Currituck Sound. North of Knott Island lie Back Bay and North Bay, separated from the ocean eastward by a long spit of beach sand. Wide areas of the terrace level are almost perfectly smooth, without any undulation perceptible to the eye. It is thought probable that the entire plain area aside from the marsh and swamps was originally a forest of short-leaved pine with occasional trees of other varieties. This pine forest, together with many small wooded areas which were occupied by cypress swamp, has been in greater part cleared. In the vicinity of Virginia Beach and thence to Cape Henry stumps are now exposed, mainly of cypress, up to the very margin of the terrace, which is being eroded directly by the waves of the ocean. Portions of these plains in the eastern and northern parts of the quadrangle are fairly well drained, owing to the sandy nature of the soil, but a large area in the southwestern part, with imperfect drainage, is occupied by the Dismal Swamp. A small detached area of swamp is known as The Green Sea.

*Dismal Swamp.*—The larger portion of this great fresh-water morass lies in the Norfolk quadrangle. It is an area of moderately elevated, nearly level land with such imperfect drainage that it remains constantly inundated to a slight depth. The outlines of the swamp area are irregular, as shown on the map, and usually the limits are not well defined, the position of the edges of the wet portion varying with the rainfall and the presence of the swamp flora. Some marginal areas which are cleared have ceased to be swampy except in wet weather. The swamp is heavily wooded and contains extensive canebrakes (see Illustration sheet, figs. 1 to 4). The swamp, which slopes gradually upward to the southwest, varies in altitude from 12 to 22 feet above mean tide level. Near its center there is a picturesque body of open water, known as Lake Drummond (see Illustration sheet, figs. 5 and 6). This lake is nearly circular in outline, about

2½ miles in diameter, and until recently its surface was slightly over 22 feet above mean tide level, being the highest portion of the swamp. At one time its depth was about 15 feet, due in part to damming of the swamp by the banks of canals. Now, owing to the deepening of the canal feeder, the lake is only about 6 feet deep and the surface correspondingly lower, which is lower than it is known to have ever been before. Its floor is largely covered with white sand. The lake water is light brown in color, due to a considerable amount of finely divided vegetal matter in suspension. It is thought to be perfectly wholesome, and as it is famous for its keeping properties, it has been used extensively for supplying ships for long voyages. The lake is surrounded by woods, and at some points cypress trees are found growing in the water (see fig. 6 on Illustration sheet). The depth of the water decreases rapidly in the woods adjoining the lake, and over the swamp area in general it is rarely more than 1½ feet, except possibly in very wet weather. The average depth is from 1 to 3 inches, but in many portions the average depth is from 6 to 8 inches. In very dry seasons the amount diminishes all over the swamp area.

Some marginal portions of the swamp have been drained for farming land, for which the soil is admirably adapted. The swamp area known as The Green Sea was originally a portion of the main swamp, but the Dismal Swamp Canal, which traverses the eastern portion of the area from north to south, has in a measure drained the intervening region. This canal sustains the water level and the resulting swamp conditions to the west, but has reclaimed from inundation a zone of considerable width to the east, an area which is further drained by the branch ditch known as the Herring Canal. The swamp flora is characterized by the occurrence of bald cypress, juniper, black gum, and extensive canebrakes.

The swamps lie in shallow basins in the surface of the general terrace of the Norfolk region. The basins are now filled to the general level of the surrounding country with vegetal accumulations, which have a maximum thickness of about 20 feet. In recent excavations for a gate on the feeder about half a mile east of Lake Drummond there were exposed 10 feet of peat filled with roots and tree trunks, lying on 8 feet of clear peat which merged with the overlying beds, and this in turn was underlain by fossiliferous sand of late Neocene age. The thickness of the swamp deposits decreases toward the periphery of the present swamp area, but so few excavations have been made along the border zone that the conditions of thinning are not known. The upper beds of peaty materials merge gradually into the sands of the adjoining area, so that no boundary line can be given.

The basin of the Dismal Swamp owes its origin to an extensive depression in the surface of the Columbia formation. At first this hollow was probably a slough in the terrace surface. When the Columbia formation was deposited James River had essentially its present course, but emptied into open water some distance northwest of the swamp. Its main current appears to have built a bar or broad delta which extended eastward and thus built up the terrace plain that lies east and southeast of Norfolk. Between this delta and the steep slope at the edge of the highlands which lie a short distance west of the Norfolk quadrangle there remained an area of lowland, a slough which was not built up appreciably by the Columbia deposits. When the delta was uplifted it became a high terrace with good drainage conditions, while the slough became a swamp filled with luxuriant vegetation, and it has so continued ever since. At first, when the vegetation was young, relatively fine-grained peat accumulated, but as the forest grew older, roots and trunks were intermixed

with the finer materials, and finally the depression was filled up to the general level of the country by these accumulations. It is now so remote from the larger drainage ways and so choked with canebrakes that its drainage is still very imperfect and the swamp conditions continue over nearly all the original basin area. Lake Drummond is no doubt the remaining portion of an original center pond, probably greatly encroached on by the forests and canebrakes. It is probable also that during some periods the lake was dry for a short time. Its bottom has been raised somewhat by vegetal accumulations, but probably its water level has just about kept pace with the general rise of the swamp surface.

*Tide marsh.*—The widest areas of tide marsh are those around Back Bay and at the head of Currituck Sound. Tide marshes occur in all the tide-water inlets, but they are generally of limited extent. Tide marsh extends up North Landing River to North Landing, and up its branch, West Neck Creek. On North Landing River it merges into fresh-water marsh. Tide marsh owes its origin to the growth of marine vegetation and the deposition of fine silt, its materials being partly earthy and partly vegetal. Brackish water is most favorable to its development. It tends to fill up the tide-water inlets, and to grow out into sheltered bays, notwithstanding moderate wave action, and eventually to constrict the bays to tidal channels. It advances along lines where the waves are least violent and makes rapid progress behind sand spits and other similar protections which break the force of the wind. The plants in the marsh grow so close together that they favor the deposition of mud and sand, and their widely extended roots and stout stems bind the deposits together. The wide neck of marsh at the Causeway is an excellent illustration of marsh growth, it having accumulated here under the lee of Knott Island. It is probable that Back Bay and Currituck Sound were once connected by a strait which has been closed by this marsh growth. The marsh is now encroaching upon the bays on the west side of Back Bay, and it is making rapid progress in filling up the bays and inlets north of Back Bay.

*Sand dunes.*—Along the shores of the ocean and of Chesapeake Bay there are accumulations of dune sand of geologically recent origin. They attain their greatest prominence on Cape Henry. In this region the sand dunes are generally confined to a narrow zone near the shore, just out of reach of high tide, where they occur in groups of various sizes. The material is loose beach sand which has been blown back from the beach by the wind. At some localities the sand has largely remained in its present place for centuries, but at other places it is blown by the wind at the present time. At Cape Henry the dunes attain a height of 70 feet and extend along the beach as a high ridge to Lynnhaven Inlet. From this inlet to Little Creek, and again from Little Creek to beyond Ocean View, there is a nearly continuous line of dunes constituting a narrow ridge from 15 to 25 feet high. The finer material from the dunes is often carried some distance inland by the winds and merges into the sandy loam which is derived from the surface of the Columbia formation. On this account it is not possible to indicate on the map any precise boundary for the dune-sand areas. In the vicinity of Cape Henry the dune sand is advancing inland comparatively rapidly. The tops of a few old cypress trees project above the summits and slopes of some of the higher dunes near the cape, and in "The Desert," an ill-drained forested area behind the dunes, the sand is accumulating rapidly about the tree trunks. "The Desert" has a surface of low dunes, which occur mainly in faint ridges approximately concentric to the present curve of the coast line of the cape. This

Location of quadrangle.

Terrace extending along the Eastern Shore of Virginia and Maryland.

Reclaimed swamp land.

Submarine extension of Coastal Plain.

Dunes along Chesapeake Bay.

Origin of Dismal Swamp.

Origin of "The Desert."

Lake Drummond.

condition indicates a series of physical changes, beginning with the formation on the terrace level of low dunes of sand, which was followed by the advance of forest growth, and this by the reinvasion of sand which is gradually killing and burying the forest.

The sand dunes about Cape Henry were described a hundred years ago by B. H. Latrobe (Trans. Am. Philos. Soc., Vol. IV, 1799, pp. 439-444, 1 plate; see also Am. Jour. Sci., Vol. XL, 1865, pp. 261-264).

A comparison of his description with the present conditions shows how little change there has been in a century. At that time the dunes were apparently as high as at present and were rapidly encroaching upon the forests in "The Desert." The sand is blown up the steep front of the dune and much of it is carried down the gentler western slope far into the forest. The line of sand dunes, it is thought, has caused the imperfect drainage of the forested area by damming the water which flows down the slight slope of the inland terrace toward the ocean. During the first sixteen years after the establishment of the lighthouse at Cape Henry it was estimated that the dunes had risen about 20 feet in height, and at one definitely located point, a short distance from the lighthouse, had proceeded into the desert about 350 yards. It was predicted by Mr. Latrobe more than a hundred years ago that if the dunes should continue to advance at this rate for twenty or thirty years they would swallow up the whole swamp, but it is evident that the rate of accumulation has greatly diminished in later years, for the dunes do not appear to have progressed much farther than they were in Mr. Latrobe's time.

## GEOLOGY.

### GENERAL SEDIMENTARY RECORD.

The geologic deposits of the Norfolk quadrangle comprise sandy loams, sands, clays, marls, peat, and muck. They are in greater part of sedimentary origin, but some of the sands are eolian, and the marsh accumulations have been aided and augmented by plant growth. The general surface formation is a sheet of sandy loam of no great thickness. This is underlain by an extensive series of Coastal Plain deposits lying on a floor of the crystalline rocks which constitute the surface of the Piedmont region to the west, but which slope far below sea level in their extension eastward under the Coastal Plain.

The rocks of the Coastal Plain consist of broad sheets of sands, gravels, clays, diatomaceous earth, marls, and glauconitic sands comprised in a succession of formations which dip very gently to the southeast. They rise above sea level in regular succession westward and northward, and in the Norfolk region have an aggregate thickness over 2300 feet. They

outcrop in the Norfolk quadrangle are in heavy-faced type. The lower formations reach the surface farther west in Virginia, except the marine Cretaceous deposits, the outcrops of which begin in Maryland. The Pleistocene formations are the only ones seen at the surface in the Norfolk quadrangle, but the Neocene, Eocene, and Cretaceous formations have been explored considerably by well borings.

Classified according to origin, these deposits of the Coastal Plain may be divided into two groups: one corresponds more or less closely with those formations now in process of deposition in the estuaries and along the shores in the immediate vicinity; the other corresponds closely to off-shore sediments known from soundings to be in process of deposition over the more deeply submerged portions of the province. In general the shore and estuarine deposits overlie those of the second group, and are thus known to be the younger. They record certain modifications in geography due to changes in altitude of the land, and, moreover, display certain distinctive characteristics indicating the climate of the periods during which they were laid down. The older formations contain abundant remains of marine organisms, preserved as fossils, and thus these deposits are records of periods during which the land stood lower and the sea consequently extended farther inland than at present. The lowest and oldest formation of the Coastal Plain series, which does not come to the surface in the Norfolk district, appears to be devoid of marine fossils, but its beds contain impressions of leaves, together with lignitized wood and other vegetal fossils, as well as the bones and teeth of dinosaurs. In addition, portions of the deposit are coarse and irregularly bedded, so that this formation, like the younger deposits, appears to bear record of land conditions and thus indicates that the land had an altitude many feet higher than that at present.

From the character of the materials it is known that the deposits of the Coastal Plain province were derived from the rocks of the neighboring interior (Piedmont and Appalachian) provinces.

### ARCHEAN PERIOD.

The "bed rock" underlying the Coastal Plain sediments has been reached by the well at Fort Monroe at a depth of 2246 feet. It is a light-colored rhyolite of very compact texture and great hardness. In the drilling operation it was penetrated 8 feet and samples were secured by which it was identified. Doubtless the same rock underlies the entire Norfolk area, sloping gently eastward.

### CRETACEOUS PERIOD.

*Potomac formation.*—The deep borings at Fort Monroe and the Norfolk waterworks penetrated

amount to the east, under the Coastal Plain. It is supposed that the top of the Potomac formation was reached at a depth of 945 feet at Fort Monroe and 782 feet in the Norfolk waterworks boring. The deposits penetrated by the well borings are shown on the Columnar Section sheet. They are mainly sands of light-gray color and varying degrees of fineness. The material is quartz with occasional grains of other minerals and flakes of mica. Pebbles of quartz are of frequent occurrence. Sandy clays occur in scattered thin beds, and at various horizons there is a general admixture of a small amount of clay with the sand. The clay bed at 1218 feet in the Norfolk waterworks boring was a tough, variegated red, gray, and buff clay, strikingly like some of that exposed in Maryland. When this boring was at a depth of 1320 feet in the brown-red sand, there was obtained a specimen of *Exogyra*, but there is some uncertainty as to the authenticity of its occurrence in the formation at this depth. It is a marine Cretaceous fossil which would not be expected in the Potomac formation, as it belongs to a much higher horizon. From the evidence of leaves, wood, and the character of the sediments, as observed in the belt of outcrop, the Potomac formation is probably an estuarine deposit. There is considerable uncertainty as to the depth at which the Potomac formation was entered in the deep well at Fort Monroe, for the boring operations were conducted by a method which results in considerable mixing up of material, and the sands and clays between 800 and 1000 feet were not sufficiently distinctive in appearance to indicate their age. In the well sunk at the Chamberlain Hotel, Old Point Comfort, some years ago, marine Cretaceous fossils were found to a depth of 845 feet, with the Potomac formation apparently lying about 100 feet below. As the top of this formation is indicated with a fair degree of distinctness at 782 feet in the Norfolk waterworks boring, there is apparently a considerable westward slope of the top of the formation between the two localities. As there is evidence of considerable unconformity at the base of the marine Cretaceous sediments, this feature may be due to an excavation of a shallow channel on the surface of the Potomac formation in this vicinity.

*Marine Cretaceous formations.*—Overlying the Potomac formation in the Norfolk region are sands and sandy clays containing abundant fossil shells of Cretaceous age, specimens of which have been found in the deep borings. The fossils are the same as those which characterize the marine Cretaceous formations outcropping extensively in New Jersey and from North Carolina south. In Maryland these formations thin out a short distance south of Washington, and in eastern Virginia they do not reach the surface at all, but their extension underground is well established by the evidence of the deep borings about Norfolk. The evidence, however, is not sufficient to afford grounds for subdividing the deposits so as to apply formation names to them.

In the Norfolk waterworks well the Cretaceous shells began to appear at a depth of 715 feet, or possibly 700 feet, and they were found in great abundance down to a depth of 775 feet. Only one species was observed, which is a small *Exogyra*, precisely similar in general form to *Exogyra costata*, but having a smooth surface or showing only very faint crenulations. The shells vary in length from  $\frac{1}{4}$  inch to  $1\frac{1}{4}$  inches. The containing material is a dark sandy clay with sand streaks. It is somewhat micaceous throughout, and appears not unlike the Matawan formation in Maryland and Delaware. A single shell was thought to have been obtained at a depth of 1320 feet, but this was not positively determined. In the well at Lambert Point, Norfolk, borings at depths of 563 to 610 feet yielded marine Cretaceous fossils, including *Exogyra costata* Say? and the following additional species: *Astarte ocolirata* Gabb, *Ostrea plumosa* Morton, *Gouldia? decemmaria* Conrad, *Gryphaea vesicularis* Lam., *Liopistha (Cymella) bella* Conrad, *Corbula* sp., *Modiolus* sp., and *Baculites?* In the well at Chamberlain Hotel, Old Point Comfort, fragments of *Terebratula harlani* Morton were

obtained when the boring was at a depth of 845 feet, presumably from gravel extending from 840 to 845 feet. It is possible that this fossil came from a less depth. It was not expected that the marine Cretaceous would be found here so much deeper than in the wells at Lambert Point and at Norfolk waterworks, which are several miles southeastward. This would indicate a local northwesterly dip of the beds. The dark, sandy, micaceous clay reported from 850 to 920 feet in the Chamberlain well appears to be typical marine Cretaceous material.

By a comparison of the records of the deep borings at Fort Monroe, Norfolk, and Lambert Point, it is believed that the marine Cretaceous deposits have a thickness of at least 65 feet, and possibly much more. In the Norfolk waterworks well, as above shown, the marine fossils have a range of 80 feet, and a portion of the still lower micaceous gray sands may also be marine.

### EOCENE PERIOD.

*Pamunkey formation.*—In the Norfolk region this formation is deeply buried, but the deep boring at the Norfolk waterworks has afforded a fairly definite conception of its relations. It is a formation which outcrops extensively along Potomac, Rappahannock, Mattaponi, Pamunkey, and James rivers in Virginia, toward the western margin of the Coastal Plain belt. It there lies on an eroded surface of the Potomac sands, sandstones, and clays, and has a thickness of over 150 feet. In the Norfolk waterworks boring it is much thinner. The only fossils it has yielded in this boring are foraminifera, of which the following were recognized, at the depths stated:

#### Eocene foraminifera from boring at Norfolk waterworks.

685 feet; *Nodosaria obliqua* L.; *Dentalina confluens* Reuss; *Bulimina buchiana* d'O.; *Uvigerina pygmaea* d'O.; *U. tenuistriata* Reuss; *Rotalia orbicularis* d'O., and *R. propinqua* Reuss.  
695 feet: *Nodosaria* sp.; *Bulimina buchiana* d'O.; *Cristellaria cultrata* M.; *Truncatulina haidingeri* d'O., and *T. lobatula* W. & J.

From a depth of 665 feet there was a sample containing foraminifera, but they give no conclusive evidence as to the age of the formation. There was found an abundance of *Bulimina buchiana* d'O., which is regarded as typical Eocene, together with *Rotalia orbicularis* d'O., which is an Eocene form, although also found in later formations. At 695 feet *Bulimina buchiana* d'O. is associated with forms that are more abundant in Cretaceous rocks but that occur in both older and younger rocks. They are *Cristellaria cultrata* M. and *Truncatulina haidingeri* d'O., both of which are common in the New Jersey Cretaceous marls. On this account, and because of a marked change in the character of the deposits, it is thought probable that the Pamunkey formation extends only to a depth of 680 feet. It appears to begin at 640 feet, where the clays suddenly become dark colored, sandy, and glauconitic. The formation is thus limited to 40 feet of dark, glauconitic sands and sandy clays. The glauconite is a characteristic constituent in the surface outcrops to the west. It is a bottle-green mineral sprinkled in small grains through the deposit in sufficient amount to give a dark-green tint to the sand. The mineral is a silicate of iron and potash, a product of various organisms which live in a moderately deep sea.

The boring at Lambert Point passed through the Pamunkey formation, but the record was not sufficiently definite for its identification. It is nearer the surface than was expected, as marine Cretaceous fossils were obtained at 563 feet. In the deep well at Fort Monroe it is probable that the beds from about 610 feet to at least 710 feet, and possibly those from 710 to 840 feet also, are Pamunkey in age. The glauconite admixture is distinctive, especially when it is associated with foraminifera, as it is from 610 to 710 feet. Some shell fragments were found at a depth of 840 feet which appear to be Eocene species, but their identification is not established and they may have been derived from a higher level in the well. Sharks' teeth of Eocene type were found in the boring from the old well at Fort Monroe between 580 and 590 feet, but they probably do not represent the life existing at the time these beds were deposited. They are probably pebbles from

Formations, and associated unconformities, underlying the Coastal Plain region of eastern Virginia.

Period.	Formation.	Character.	Thickness in feet.
Pleistocene	Alluvium, etc.	River silt, peat, muck, beach sand, dune sand, etc.	0 to 60
	Unconformity.		
	Columbia formation	Sandy loams, sands, and clays.	10 to 30
Neocene	Unconformity.		
	Lafayette formation (Pliocene?)	Gravel, orange sands, and loams.	25 to 40
	Unconformity.		
	Pliocene strata	Marls and sands.	10 to 20
Eocene	Chesapeake formation (Miocene)	Fine sands, clays, and diatomaceous deposits.	35 to 565
	Unconformity.		
	Pamunkey formation	Glauconitic sands, marls, and clays.	30(?) to 300
Cretaceous	Marine deposits	Clays and sands.	0 to 500
	Unconformity.		
Archean	Potomac formation	Sands and clays.	200 to 1300
	Great unconformity.		
	Crystalline rocks	Granites, gneisses, etc.	

range in age from early Cretaceous to late Pleistocene. The successive formations are in most cases separated by unconformities, each representing a period during which the surface was uplifted above the sea and sculptured by waves and streams, so that when each succeeding deposit was laid down its strata were more or less discordant with the partially eroded strata of the preceding period. The formations which underlie the Coastal Plain region of eastern Virginia are shown in the accompanying table; those which

sands and clays undoubtedly belonging to the Potomac formation, which is so extensively exposed along the upper estuarine portions of James, Rappahannock, and Potomac rivers in Virginia. It constitutes the lower members of the Coastal Plain series, lying on the floor of granites, gneisses, and other crystalline rocks of the Piedmont region. In the outcrops in Virginia it exhibits a maximum thickness of about 600 feet along Potomac River near Washington, but it thickens to more than double this

Eocene rocks inclosed in the overlying Chesapeake formation, a feature often observed in surface outcrops of the basal beds of that formation. It is to be expected that the Pamunkey formation would be found to be thicker at Old Point Comfort than at Norfolk, for in its surface outcrops to the west it is over 150 feet thick.

NEOCENE PERIOD.

**Chesapeake formation (Miocene).**—Underlying the thin mantle of Pleistocene and Pliocene materials in the Norfolk region there is a thick mass of marls and clays of Miocene age, known as the Chesapeake formation. It does not outcrop at the surface in the quadrangle, but it is reached by several wells. The deep borings at the Norfolk waterworks, Lambert Point, and Fort Monroe pass entirely through the formation and afford complete sections of its beds. The materials are mainly shell marls and light-gray clays, with occasional darker portions and sandy beds. In the Norfolk waterworks boring the formation appears to have been entered at a depth of about 70 or 80 feet, and it extended to 640 feet, a thickness of 565 feet. Its uppermost beds are 70 feet of gray clays, sandy clays, and sands, then 110 feet of light-gray clay, 70 feet of darker sandy clays in part of greenish tint, below which there are light-gray clays extending from 330 to 640 feet in depth. Many of the strata yielded fossil shells, of which the following, found at the depths stated, were the most distinct.

Upper Miocene fossils from boring at Norfolk waterworks.

- 75 feet: Pecten jeffersonius Say, P. clintonius Say, Turritella alticoستا Conrad, Rangia cuneata Conrad.
- 85 feet: Rangia cuneata Conrad.
- 105 feet: Abra aequalis Say, Pecten jeffersonius Say, P. madisonius Say, Yoldia levis Say, Turritella alticoستا Conrad, Cadulus thallus Conrad, Solen.
- 115 feet: Rangia cuneata Conrad, Ostrea virginica Gmel., Yoldia levis Say, Drillia limatula, Turritella alticoستا Conrad, Rangia clathrodonta Conrad.
- 125 feet: Leda acuta Conrad, Turritella alticoستا Conrad, Cadulus thallus Conrad.
- 135 feet: Yoldia levis Say, Cadulus thallus Conrad.
- 175 feet: Pecten eboreus Say.
- 185 feet: Pecten clintonius Say.
- 325 feet: Cardium islandicum Chemnitz (?).
- 345 feet: Mulinia lateralis Say.
- 365 feet: Pecten eboreus Conrad.

The fossils listed above indicate upper Miocene age. *Rangia cuneata* has never before been reported from beds as old as these. From the lower borings less distinctive remains are obtained. Diatoms with sponge spicules and foraminifera occur from 580 to 625 feet, notably at 585 and 605 feet, where they are very abundant. The proportion of diatoms is hardly sufficient to entitle the material to be classed as an infusorial earth. They are the same species as those which occur in the lower Chesapeake beds outcropping at Richmond, Va., and other points along the western zone of outcrop of the formation. Sponge spicules occur in various strata between 355 and 505 feet. Foraminifera are present in considerable abundance and variety at a depth of 645 feet, including *Bulimina elongata* d'O., *Uvigerina pygmaea* d'O., *U. tenuistriata* Reuss, *Rotalia soldanii* d'O., *Nonionina scapha* F. & M., and *N. boueana* d'O.

The Lambert Point boring passed through the Chesapeake formation, affording the section given on the Columnar Section sheet. Unfortunately it was mainly the sandy constituents of the washings that were saved, so that the beds appear to be predominantly sandy and no close comparison can be made with the Norfolk waterworks boring. The only Chesapeake fossils obtained were between 235 and 264 feet. They included *Perna maillata* Lam. and *Pecten jeffersonius* Say. Foraminifera, echinus spines, and sponge spicules occurred from 17 to 563 feet. As Cretaceous fossils occur at 568 feet in this boring, and the Pamunkey formation is probably represented, the Chesapeake beds are here much thinner than in the other borings, the thick diatomaceous bed is absent, and, if the data are authentic, the base of the formation rises rapidly westward from the Norfolk waterworks boring, where it is at a depth of about 640 feet.

At Fort Monroe the Chesapeake formation extends from 40 to about 590 feet. *Dosinia acetabulum* Conrad and *Pecten madisonius* Say

were reported at a depth of 50 feet in the new well. In the old well other distinctive Miocene shells were obtained at intervals to 400 feet, and diatoms in abundance at 558 feet. Between 580 and 590 feet sharks' teeth were obtained, which were either deposited as pebbles in the basal Chesapeake bed or occurred in place in the underlying Pamunkey formation.

The following fossils have been reported from the Chesapeake formation in the new well at Fort Monroe at depths of 50 to 100 feet, the larger number coming from about 50 feet:

Miocene fossils from Fort Monroe well.

- |                              |                            |
|------------------------------|----------------------------|
| Columbella (Anachis) sp. ?   | Dosinia acetabulum Conrad. |
| Cadulus thallus Conrad.      | Pecten sp. ?               |
| Dentalium attenuatum Say.    | Pecten madisonius Say.     |
| Eulima sp. ?                 | Pecten jeffersonius Say.   |
| Olivella sp. ?               | Glycymeris subovatus.      |
| Scala multistriata Say.      | Tellina declivis Say.      |
| Turbonilla.                  | Ostrea subreflexa.         |
| Turritella alticoستا Conrad  | Lucina crenulata Conrad.   |
| Nassa acuta Say.             | Yoldia limatula Say.       |
| Ptychosalpinx atilis Conrad. | Crepidula sp. ?            |
| Venericardia granulata Say.  | Crepidula formicata Say.   |
| Cardium laqueatum Conrad.    | Crepidula costata Say.     |
| Corbula sp. ?                | Venus mercenaria Linne.    |
| Corbula inaequalis Say.      | Astarte undulata Say.      |
| Meretrix sayana Conrad.      | Balanus proteus.           |
| M. convexa Say.              |                            |

The clays are richly diatomaceous between the depths of 530 and 560 feet, and among the fossil forms are those which characterize the great bed of diatomaceous earth in the Chesapeake formation underlying the Coastal Plain through New Jersey, Delaware, Maryland, Virginia, and southward. With the base of the Chesapeake formation at 580 feet at Fort Monroe and at 640 feet at the Norfolk waterworks, 11 miles south-south-east, a dip of about 6½ feet per mile is indicated, and apparently about the same dip is indicated by the diatom beds.

**Pliocene strata.**—Just beneath the covering of Columbia and alluvial formations in the Norfolk region there is a thin layer of marls and sands of Pliocene age. They do not outcrop at the surface, but are reached by many wells and have been uncovered in places by the excavations for the canals. In most localities they contain large numbers of shells, many identical with living or relatively recent forms, with others of late Neocene age intermixed with them. These deposits were extensively exposed by the Dismal Swamp Canal excavations, notably at three points: one on the main canal, 4 miles south of Wallacetown, another on the feeder from Lake Drummond, about halfway between the lake and the main canal, and the other on the main canal in the vicinity of Lilly, N. C. According to L. Woolman (Proc. Phila. Acad. Nat. Sci. for 1898, p. 414), the fossils collected from these beds are as follows:

Fossils from the Pliocene strata at Lilly, N. C.

- |   |                                  |
|---|----------------------------------|
| Arca limula Conrad.                         | Cæcum cooperi Smith.             |
| A. plicatura Conrad var. sub-lineolata d'O. | Ostrea virginica Gmel.           |
| Corbula contracta Say (numerous.)           | Crepidula plana Say.             |
| Meretrix convexa Say.                       | Eupleura caudata Say.            |
| Mulinia lateralis Say.                      | Fulgur canaliculatum Say.        |
| Nucula proxima Say.                         | F. carica Gmel. (numerous).      |
| Venus mercenaria Linne.                     | Nassa trivittata Say (numerous). |
| Lucina crenulata Conrad.                    | Scala lineata Say.               |
| Olivella nitidula Dillw.                    | Turbonilla reticulata Ads.       |
| Polynices (Neverita) duplicata Say.         | Spisula solidissima Dillw.       |
| Tornatina canaliculata d'O.                 | Tellina tenera Say.              |
| Nassa obsoleta Say.                         | Ensis directus Conrad.           |
| Divaricella quadrisulcata d'O.              | Urosalpinx cinereus Say.         |
|   | Astrangia danae Agassiz.         |

In some of the clays containing these shells there are also diatoms in abundance, comprising many species ranging from Miocene to the present, a few never observed in Miocene beds and several which are supposed to belong to the Miocene exclusively. The latter were probably derived from the Chesapeake clays exposed in the outcrops to the west of this region. Similar diatoms are contained in the gray clays lying between 25 and 65 feet below the surface in the Norfolk waterworks boring. At a depth of 25 feet in this boring occurs *Rangia cuneata* Gray, a Recent to Pliocene form. At Great Bridge Pliocene beds were exposed by the excavations of the Albemarle and Chesapeake Canal, and yielded a few fossils, and on the Jericho Canal southeast of Suffolk the same beds yielded a large number of distinctive Pliocene forms. It is evident from the widespread occurrence of the fossils that the entire area of the Norfolk quad-

rangle is underlain by Pliocene deposits, except probably the deep channel extending down James River and out to the ocean through Chesapeake Bay. This channel undoubtedly cuts into the underlying Chesapeake formation. Judging by its fauna, the Pliocene is of about the same age as the Croatan beds of North Carolina. The Lafayette formation, of supposed Pliocene age, shown in the table of formation names, does not occur in the Norfolk area.

PLEISTOCENE PERIOD.

**Columbia formation.**—The entire land area of the Norfolk quadrangle is covered by a thin sheet of loams and sands, called the Columbia formation, the surface of which constitutes the wide, low plain so characteristic of the region. The formation was laid down along the coast in a belt which, in this region, extended back to the base of the highlands along the west side of the Dismal Swamp. The precise physical conditions under which it was deposited are not known, but it covered a flood plain bordering the sea. The thickness of the Columbia formation is usually from 20 to 55 feet, and in the greater part of the area its base is slightly below tide-water level. The surface on which it lies is known to be somewhat irregular or gently rolling, but the precise configuration of this surface is not everywhere known. The Columbia formation is cut through by the deeper valleys, notably those of James, Elizabeth, and Nansemond rivers, Hampton Roads, and Chesapeake Bay. It is being cut away along the ocean front, where the upper edge often presents a low cliff extending along the beach, ordinarily at the level of high tide. The beach sand is usually banked up against the cliff, and beach or dune sands completely bury the formation locally. The lower portion of the formation extends out under the ocean for a greater or less distance, and sometimes it is bare of sand in the zone between high and low tide levels.

Under the surface the Columbia formation presents a relatively uniform character throughout, but there are some local variations, and there is considerable range in thickness. Large boulders are occasionally inclosed in the finer material, and their occurrence can be explained only by the hypothesis that they were transported by floating ice. A very instructive series of borings was made sometime ago by the Norfolk City Water Department in the region east of the city, which passed through the Columbia deposits to the marl of the underlying Chesapeake formation. The results are given in the following sections:

Myers farm.

Fine sand	1 to 11
Loose sand	11 to 18
Sand and sandy clay	18 to 50
Gravel with water, lying on marl which was penetrated to a depth of 117 feet	50 to 52

Taylor farm.

Muck	1 to 19
Sand and sandy clay	19 to 51
Red sand	51 to 58
Blue clay	58 to 68
Coarse sand with water	68 to 85½
White sand lying on marl	85½ to 97½

Poorhouse tract swamp.

Muck	1 to 5
Fine sand	5 to 9
Sand and clay	9 to 32½
Blue clay	32½ to 35½
Sandy clay	35½ to 38½
Sand and clay	38½ to 45½
Gravel with water, lying on marl	45½ to 51½

Poorhouse tract turnpike.

Sand	1 to 18
Blue clay	18 to 31½
Sandy clay	31½ to 34½
Loose sand	34½ to 42½
Gravel with water, lying on marl	42½ to 50

City property east of city limits.

Fine sand	1 to 9½
Sand and sandy clay	9½ to 44
Small gravel with water, lying on marl	44 to 46½

Babcock farm.

Sand	1 to 11½
Loose sand with water	11½ to 25½
Sand	25½ to 30½
Sand, clay, and muck	30½ to 53

Lake Lawson.

Yellow clay with gravel streaks	1 to 10
Sand	10 to 24½
White sand	24½ to 31
Sandy clay	31 to 36½
Coarse white sand with water, lying on marl	36½ to 38

Taylor woods.

Sand	1 to 10½
Sand, clay, and marl	10½ to 40½
Gravel with water, lying on marl which was penetrated to a depth of 60 feet	40½ to 40½

Drummond woods.

Sand	1 to 4
White sand	4 to 13
Sand and clay	13 to 21
Clay and sandy clay	21 to 30½
Sandy clay	30½ to 36½
White sand	36½ to 38½
Gravel with water, lying on marl	38½ to 44

TOPOGRAPHIC DEVELOPMENT.

During the Cretaceous, Eocene, and Neocene periods this region was the center of deposition of widely extended sheets of sands, clays, and marls. The periods of deposition were separated by intervals during which the strata were brought to the surface and more or less eroded. The configuration of the land surfaces in these periods of uplift and degradation is not known, but during the final stages of the Neocene the uplifted area had been smoothed to a nearly level plain. In early Pleistocene time there was an uplift with perceptible tilting to the eastward, during which wide troughs were excavated in the James River and the other larger Coastal Plain valleys, and a coastal lowland was cut which extends westward to a line that passes northward from near Suffolk across the lower ends of the tide-water peninsulas of eastern Virginia lying west of Chesapeake Bay. In this process the Lafayette formation was removed and the underlying formations were leveled to an approximate plain which in greater part lies a few feet below the present tide-water level. There were next flood-plain conditions, during which the thin sheet of Columbia formation was laid down over this eroded area, the product being a smooth plain extending from a short distance west of Chesapeake Bay eastward across the bay and the Eastern Shore and along the James and other large tide-water valleys. Then followed uplift, during which the James and other rivers which have their sources in the higher land to the west, cut their channels through the Columbia deposits, and Susquehanna River with branches from the other Coastal Plain rivers excavated the trough which is now occupied by Chesapeake Bay. At this stage there were also initiated the small streams which now lie entirely in the Columbia plain, such as Elizabeth, Nansemond, Northwest, North Landing, and Lynnhaven rivers in the Norfolk region. Small local streams also isolated a fragment of terrace in Knott Island. It was in this stage of uplift that a channel was cut through the depression now occupied by the Albemarle and Chesapeake Canal. The Atlantic Ocean also cut a sea scarp on the slope of the Columbia terrace, probably several miles east of the present shore line. Next came the submergence in which the deeper depressions were flooded with tide water, the Atlantic Ocean advanced to the vicinity of the present shore line, and the minor drainage ways ceased their active cutting, in most cases having their mouths so drowned as greatly to decrease their grade. This subsidence is still in progress, but at a very gradual rate.

Probably in the later stages of the last uplift and during the early part of the subsidence dune sands began to accumulate on the beaches along the shores adjacent to the ocean. Some of the dunes have continued to rise until now they are over 70 feet high on Cape Henry. They occupy a narrow zone along the beach to Willoughby Spit and are found even on the shore of Elizabeth River. Spits also have been built from the main land, Willoughby Spit and the beach extending from the southern border of the quadrangle northward to Sand Bridge being conspicuous examples. Marsh growth has progressed in many parts of the

Coastal lowland produced during early Pleistocene uplift.

Columbia formation deposited during flood-plain conditions.

Excavation of river channels in Columbia terrace.

Submergence and flooding of the lower channels.

inclosed area behind the beach along the sea coast. These spits show that the prevailing shore currents are southward along the sea coast and westward along Chesapeake Bay.

Geologic processes which are now in progress and which have continued for sometime during the submergence are the augmentation of the sand dunes and spits, growth of the marshes, and deposition of alluvium, mainly under tide water.

## ECONOMIC PRODUCTS.

### SOILS.

The soils of the Norfolk quadrangle present considerable diversity in character, varying from pure sand through sandy loam and clay to swamp muck. The greater part of the district has a surface covering of Columbia formation, with a soil consisting mainly of sandy loam. This is the normal soil of the very fertile truck region which is extensively cultivated about Norfolk. The proportion of the organic content of the soil is not large, but with proper fertilization the soil has the physical condition most favorable to the growth of a number of vegetables and fruits. The local variations in the soils are gradual but frequent and the differences are perceived less readily in the soil than in the relative productiveness in garden truck. In the extreme eastern part of the area the soils are more sandy, owing in part to coarser materials in the Columbia formation in that direction and to admixture of blown sand from the beach. The sands on the beach and in the sand dunes are nearly barren of vegetation. The soils of the present swamp areas are often too carbonaceous for immediate use, but in some cases this feature has been corrected by exposure to the weather after the land has been cleared of trees, drained of water, and ploughed. These soils are usually rich in plant food and prove to be highly productive.

The following are mechanical analyses of typical truck soils from the vicinity of Norfolk made by the Department of Agriculture:

Mechanical analyses of truck soils from vicinity of Norfolk.

Constituent. (Size of grains indicated in millimeters.)	Five-halves west of Norfolk.		Four-halves west of Norfolk.		Three-halves east of Norfolk.		Two-halves east of Norfolk.	
	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	
Clay (.005 to .0001).....	7.15	8.40	8.88	14.35				
Fine silt (.01 to .005).....	5.90	7.33	4.79	8.88				
Silt (.05 to .01).....	15.14	11.54	20.20	31.45				
Very fine sand (.1 to .05).....	7.51	5.71	6.63	10.16				
Fine sand (.35 to .1).....	38.25	41.03	12.96	5.12				
Medium fine sand (.5 to .25).....	23.27	24.04	42.12	25.17				
Coarse sand (1 to .5).....	1.42	0.64	0.27	2.67				
Organic matter, water, loss.....	1.36	1.36	4.15	2.20				

The first three soils are characteristic of the lighter varieties. The last is much heavier and is suited to cabbage and spinach.

The soils of the swamps vary from pure peat to clayey loam. Two leading varieties are recognized, the "juniper" or "light" swamp, and the "black gum" or "dark" swamp. The first is nearly pure peat, consisting of a brown mass of vegetal fragments derived from the juniper or white cedar, which is the characteristic tree of "light" swampy areas. The thickness of the deposit is often 8 to 10 feet. From 75 to 95 per cent of the material is organic. When such land is cleared and drained the peat cakes and hardens so that it resembles charred wood. Land of this sort is practically worthless. The black gum swamp deposits which have been laid down in various portions of the Dismal and other swamps and which bear a forest of cypress, black gum, and red maple, are well adapted to agriculture in most cases. This soil contains a large amount of organic matter, which is mainly in its upper portion. When it is properly drained and cultivated the amount of organic matter gradually diminishes, but it has been found in the drained areas that after being under cultivation for fifty years the soil still retains enough organic matter to remain black in color. The organic matter furnishes nitrogenous materials to plants, so that the soil is a rich one, but its disposition to retain moisture renders it rather slow for the raising of early

vegetables. The soils are also notably acid, which has to be neutralized by repeated applications of lime. The percentage of clay in the swamp soils is large, for the sluggish drainage in the swampy areas does not bring much sand, and the principal inorganic sediments are very fine flocculent clayey materials. This character greatly retards artificial drainage of the region, so that in reclaiming swamp lands numerous ditches and extensive tiling are necessary. The recent deepening of the Dismal Swamp Canal has reduced the general water level through most portions of the large swamp, and in the future this condition will greatly facilitate reclamation. There are extensive areas of the swamp which can be economically drained and which have rich and lasting soils, and the region has good prospects of being valuable agriculturally in the future. It is not expected that the soils will be available for truck farming to the same extent as the dry plains of the surrounding region, but they will yield crops of many important staples.

### UNDERGROUND WATERS.

Under the greater portion of the area of the Norfolk quadrangle the coarser basal beds of the Columbia formation contain considerable water. This is the source of supply of hundreds of shallow private wells scattered over the region and of a portion of the city supplies for Norfolk and Portsmouth. The water is low in mineral constituents, but contains a moderate amount of organic matter. Unfortunately it is subject to surface contamination, owing to the imperfect protection afforded by the relatively permeable sandy loam under which it lies. Where wells are near sources of pollution they are soon contaminated, and probably throughout the region there is some seepage of surface water containing malarial germs, drainage of manured fields, etc. In sinking wells in this region it is very desirable to locate them as far as possible from stables, cess-pools, or ponds of stagnant water.

The present supply for Norfolk is obtained mainly from ponds east of the city, and, although a surface water, its quality is fair. The supply is supplemented by groups of wells driven into the Columbia formation. Portsmouth is supplied by a series of shallow driven wells. In a series of test borings made by Norfolk east of the city the following results were obtained.

#### Results of test borings made east of Norfolk.

Location.	Depth to water.	Rise of water. (+ above surface; - below surface.)	
		Feet.	Ft. In.
Myers farm.....	50 to 52	- 7	
Taylor farm.....	68 to 85½	+ 20½	
Poorhouse tract, swamp.....	45½ to 51	+ 14	
Poorhouse tract, on turnpike.....	42 to 50	- 2½	
City property.....	44 to 46½	- 3	
Babcock farm.....	38½ to 53	- 6	2½
Lake Lawson.....	36½ to 38	- 22	
Taylor woods.....	40½	+ 6	
Drummond woods.....	38½ to 44	- 7	

Deep wells in the Norfolk region so far have not proved successful in obtaining pure water. Water is found in greater volume in some of the deep borings, but it has been rather too salty for domestic use. The first horizon of this saline water is at a depth of 738 feet in the deep well at the waterworks east of Norfolk, and at 608 feet at Lambert Point. Other deeper horizons of saline waters are at 785, 805, 950, 975, 984, 1038, and 1190 feet in the well east of Norfolk and at 945 feet in the well at the Chamberlain Hotel, Old Point Comfort.

The boring east of Norfolk was made by the City Water Department for the purpose of testing the water contents of all the Coastal Plain formations down to the crystalline rock, or "bed rock." It was made at the waterworks at Moore's bridges, 5 miles east-northeast of the City Hall. The boring was begun April 7, 1896, and continued at intervals, through 1896, 1897, and 1898, to a depth of 1760 feet without obtaining a flow of fresh water. The record of this boring is given on the Columnar Section sheet. The flows were as follows:

At 738 feet, 10 gallons a minute; 785 feet, 35 gallons; 805 feet, 25 gallons; 950 feet, 75 gallons; 975 feet, 10 gallons; 984 feet, 15 gallons; 1038

feet, 25 gallons; and 1072 feet, 150 gallons. At 1190 feet there was water that did not quite overflow, at 1220 and 1227 feet and at several lower levels small volumes of salt water were found; at 1480 feet there was a large flow of very salt water, and at the bottom there appeared to be a strong flow of water which carried so much sand that it choked the casing. The well was cased with 12-inch pipe to a depth of 777 feet 7 inches, with 10-inch pipe to 1207 feet 9 inches, with 8-inch pipe to 1538 feet 11 inches, and with 6-inch pipe to 1730 feet.

Boring was discontinued on account of a string of tools having dropped into the well; otherwise there would have been no difficulty in sinking the casing much deeper. It is probable that within the next 600 feet the bed rock would have been penetrated and that water not too saline for use would have been found in the coarse sand immediately above it. The boring was dynamited at 1070 feet, with the expectation of opening the pipe to the 150-gallon flow at that depth, but the operation appears to have failed. The chemical character of the principal flows from the Norfolk boring is as follows:

Chemical character of principal flows from Norfolk boring.  
[In grains per United States gallon.]

Constituent.	738 feet.	1,088 feet.	1,072 feet.
Chloride of sodium.....	112.33	158.83	164.10
Carbonate of soda.....	30.21		
Carbonate of lime.....	4.06		
Total mineral matter.....	150.36	195.12	213.03
Organic matter.....		.40	.41
Ammonia, free.....	.18	.18	.17
Ammonia, albuminoid.....	.006	.005	.002

The boring at Lambert Point was made several years ago for the Norfolk and Western Railroad Company. Its depth is 616 feet and its casing, which is a 6-inch pipe, extends to a depth of 606 feet. The pipe stands about 4 feet above the ground and the water rises 19 feet above the outlet. The flow is stated to be 65 gallons a minute. Temperature, about 70°. The water is bright and clear, but it contains the following large amounts of mineral matters, which give it a saline taste:

Mineral matter in water from boring at Lambert Point.

	Grains per U. S. gallon.
Chloride of sodium.....	28.92
Sulphate of potash.....	1.39
Sulphate of soda.....	2.77
Bicarbonate of lime.....	.93
Bicarbonate of magnesia.....	.78
Bicarbonate of soda.....	28.09
Oxide of iron and alumina.....	.08
Silica.....	.50

This flow appears to be from the same bed of the marine Cretaceous as the one yielding the flow at 738 feet at the Norfolk waterworks boring. The well record is given on the Columnar Section sheet.

At Fort Monroe three deep wells have been bored. The first, in 1864, was sunk in the fort by the Government to a depth of 907 feet, but yielded only a small quantity of saline water at 599 feet. The second, sunk in 1896 at the Chamberlain Hotel, reached a depth of 945 feet, where a flow was obtained which is estimated at about 50 gallons a minute and which rises over 17 feet above the surface, or about 22 feet above tide level. The water is reported as being slightly saline. The third well was sunk at the fort by the Government in the early part of 1902 to a depth of 2254 feet. It passed entirely through the Potomac formation and entered the crystalline rocks 8 feet. At depths of 1320 and 2131 feet water was struck and found to be salty and of small flow. The water at the lower horizon rose in the pipe to within 20 feet of the orifice, or to a height of 17 feet below tide. Other horizons of water not tested were struck at depths of 285, 630, 985, 1520, 1630, 1915, and 1945. The well is cased to the 2131-foot flow and an unsuccessful effort has been made since its completion to cut the casing at the untested water levels. The record of these wells is given on the Columnar Section sheet. It is thought that the water horizon at 945 feet in the Chamberlain Hotel well is the same as that which yields saline water at 1038 feet at the Norfolk waterworks well, 11 miles south-southeast. This would indicate a dip in that direction of about 9 feet to the mile, but it is probable that the maximum dip of the water-

bearing beds is due east, which would make the dip approximately 20 feet per mile. On an east-west line passing through Fort Monroe the crystalline rock is now known to descend at a rate of about 34 feet per mile.

At Newport News a boring was made several years ago to a depth of 600 feet and no water was obtained. This fact and the absence of water at a similar depth in the Fort Monroe borings indicate that the higher water horizons at Lambert Point and Norfolk waterworks do not extend far west. At Money Point, 5 miles south of Norfolk, a 5-inch well was sunk to a depth of 562 feet, and furnished a good supply of slightly ferruginous water which rose 10 feet above tide level. Its source of supply is low in the Chesapeake formation, in sand under an 8-foot bed of rock. Another well, about half a mile southwest, was sunk 450 feet and yielded salt water.

At Virginia Beach various attempts have been made to obtain artesian water, but the borings have not been sufficiently deep. One well is stated to have been sunk about 600 feet without finding water in appreciable amount. A well having a depth of 70 feet yielded a moderate supply of somewhat ferruginous water.

A well bored to a depth of 147 feet in Norfolk formerly yielded a fair supply of hard water, but after the introduction of the city waterworks it fell into disuse and was finally abandoned.

In the vicinity of Jacksondale and Lynnhaven wells are reported to be from 12 to 14 feet deep, through stiff clay and red sandy clay into white sand, and have plenty of excellent water. Near London Bridge, on the ridge east of Lynnhaven River, the depths vary from 8 to 10 feet. About Kempsville, in the region at the head of Eastern Branch of Elizabeth River, the water is generally brackish or hard in shallow wells. Fairly good water is obtained from driven wells at a depth of 80 feet. Owing to some hard stratum at that depth at Kempsville no deeper driven wells have been practicable. Around Thalia an abundant supply of good water is obtained from wells 12 to 15 feet deep. One at the post-office, 12 feet deep, furnishes water for 190 head of stock and for dairy use. The water is in sand under a 2-foot layer of clay. In the low ridge passing through Nimmo good water occurs in sand at depths of 10 to 15 feet. At Sigma, nearer the ocean, the wells are deeper, varying from 20 to 25 feet, but the water is slightly brackish.

About North Landing wells vary from 10 to 14 feet in depth. On the wide, flat area about Hickory there are numerous wells from 10 to 15 feet deep, all of which obtain good supplies of excellent water in sand and gravel, in places overlain by blue sandy clay. In the region about Fentress wells from 10 to 15 feet in depth obtain satisfactory water supplies. In the vicinity of Cornland wells are from 8 to 12 feet deep, the deeper ones usually furnishing a satisfactory volume of water. They pass through about a foot of dark soil, 3 or 4 feet of clay, and 6 to 8 feet of sand. There are several driven wells 18 to 45 feet deep. Below 12 feet these usually pass through black quicksand, which contains very bad water. The beds from 25 to 30 feet contain but little water; from 36 to 45 feet there is sand, with clear, cool water in abundance, but slightly mineralized. In the settlement of Benefit, east of Cornland, plenty of water is obtained in wells 10 to 12 feet deep. Farther north, at Grassfield, the wells are 8 to 10 feet deep on the lower lands, but yield poor water. On the higher slopes to the south wells 12 to 14 feet deep furnish good water. About Gilmerton wells average 8 to 10 feet deep, but a few are sunk from 20 to 30 feet. On the peninsula north of Norfolk, about Tanner Creek, the wells are 8 to 10 feet deep.

It is reported that in the North Carolina portion of the Norfolk quadrangle the wells average from 10 to 15 feet in depth and obtain plenty of water for local use. In the vicinity of Lilly, which is on the reclaimed portion of the Dismal Swamp area, the water is not of satisfactory quality. A well at the post-office 20 feet deep yielded water which had such a brackish taste that it could not be used for household purposes. The Lake Drummond water is taken from the canal and used at this place.

June, 1901.