

Ride the Rockies

Frisco to Frisco Loop

2007 Geology Highlights

START

Day 1 Sunday, June 17
Frisco to Steamboat Springs **98 MILES**

44 MILES Monday, June 18 **Day 2**
Steamboat Springs to Craig

89 MILES Tuesday, June 19 **Day 3**
Craig to Rifle

Day 4 Wednesday, June 20
Rifle to Glenwood Springs **36 MILES**

Day 5 Thursday, June 21
Glenwood Springs to Aspen **43 MILES**


61 MILES Friday, June 22 **Day 6**
Aspen to Leadville

Day 7 Saturday, June 23
Leadville to Frisco **51 MILES**

FINISH

422 Miles

Geology of the Region



Landsat Image of the Region



Geology



Quaternary—0 to 1.8 million years ago
This is the geologic time period during which the present landscape formed. Glaciation peaked and waned several times, sculpting cirques (semicircular-shaped bowls at the heads of mountain valleys) and U-shaped valleys. Last major glaciers retreated about 12,000 years ago.

- Includes alluvium (sand, gravel, and silt deposited by rivers and streams), eolian (windblown) deposits, glacial deposits, landslide deposits, and young volcanic rocks (basalt flows).

Tertiary—1.8 to 66 million years ago
A major mountain-building episode, the Laramide orogeny, occurred during this period—70 to 45 million years ago. Erosion then exposed basement rocks and created a flat surface. Erosion of this surface during regional uplift—beginning 10 to 5 million years ago—shaped the present mountain landscape. Rifting (faulting) began about 30 million years ago, creating the Arkansas and San Luis Valleys.

Sedimentary rocks of Tertiary age
Includes sandstone, siltstone, shale, claystone, and conglomerate (rounded rock fragments in a fine-grained matrix).

Igneous rocks of Tertiary age
Includes volcanic rocks, such as basalt, rhyolite, and ash-flow tuffs (especially in the San Juan Mountains), and intrusive rocks with compositions similar to granite.

Cretaceous—66 to 144 million years ago
A seaway flooded Colorado, depositing shallow marine, shoreline, and swamp sediments. Dinosaurs became extinct by the end of this period.

- Includes primarily shale, sandstone, and coal, and minor limestone and conglomerate (rounded rock fragments in a fine-grained matrix).

Lakes

Jurassic and Triassic—144 to 245 million years ago (Includes some rocks as old as 320 million years)
The Ancestral Rockies were eroded during this time of deserts, intermittent streams, salt flats, coastal plains, dunes, and deltas. Dinosaur fossils and footprints are found in deposits of ancient river channels.

- Includes sandstone, siltstone, and claystone, and minor limestone, gypsum, and conglomerate (rounded rock fragments in a fine-grained matrix).

Permian and Pennsylvanian—245 to 320 million years ago
During this time, rocks were uplifted to form the Ancestral Rocky Mountains, which were just as high and rugged as our present mountains. Erosion of older sediments resulted in deposition along mountain flanks and in basins.

- Includes sandstone, siltstone, shale, conglomerate (rounded rock fragments in a fine-grained matrix), gypsum, and limestone.

Mississippian to Cambrian—320 to 540 million years ago
This was a time of widespread marine deposition when Colorado was intermittently below sea level.

- Represented mostly by limestone, but also includes quartzite, sandstone, shale, and dolomite.

Precambrian—older than 540 million years ago (Includes rocks as old as about 1.8 billion years in Colorado)
The Precambrian accounts for more than 85 percent of geologic time. These rocks are referred to as the basement rocks; they are exposed in the cores of major mountain ranges and in some of the deeper canyons. They are the products of metamorphism (changes in the chemistry and fabric resulting from heat and/or pressure) and igneous intrusion (emplacement of molten rock).

- Includes intrusive rocks, chiefly granite, and metamorphic rocks such as gneiss, schist, and quartzite.

About This Image




This satellite image is a portion of a mosaic of sixteen Landsat 7 scenes that cover parts of the State of Colorado. The scenes were acquired between October 1999 and October 2000.

Landsat 7 is a medium-resolution, polar-orbiting satellite managed by the United States Geological Survey (USGS). The instrument is an eight-band spectrometer. This image is a false-color composite, which combines bands 5 (SWIR), 4 (near-IR), and 3 (Red) and displays them as red, green, and blue, respectively.

The appearance of shaded relief was created using the National Elevation Dataset, which is based on digital elevation models that have a resolution of 30 meters. The 2007 Ride The Rockies route was superimposed using the ESRI Data and Maps roads database in ArcInfo.

Landsat 7 data are received, processed, and distributed by the USGS EROS Data Center in Sioux Falls, South Dakota. The EROS Data Center is the national archive for land remote sensing data, which includes aerial photography of the U.S., elevation data, declassified satellite photography, and Landsat images. Information on the archive holdings may be obtained by contacting the EROS Data Center, Customer Services Department at 1-800-252-4547, email custserv@usgs.gov or browse <http://earthexplorer.usgs.gov>

The image on this poster may be downloaded for free from the USGS. Go to <http://www.eros.usgs.gov/rtr.htm> to link to the download page.



Text by Janet L. Slate
Edited by Jon W. Raese
GIS and digital cartography by Paco VanSistine
Layout and design by Amber Hess