


CLIMATE AND WATER YIELD



ing.
on the

EXPLANATION		
<p>Adequate yields Easy construction of wells Dissolved solids content is generally less than 1000 ppm</p> <hr/> <p>Small well outlet Easily censored Iron content and hardness of water is high for domestic use</p>	Advantages	<div style="display: flex; align-items: center;"> <div style="width: 10px; height: 10px; background-color: #007bff; margin-right: 5px;"></div> Good </div> <div style="display: flex; align-items: center;"> <div style="width: 10px; height: 10px; background-color: #6c757d; margin-right: 5px;"></div> Fair </div> <div style="display: flex; align-items: center;"> <div style="width: 10px; height: 10px; background-color: #dc3545; margin-right: 5px;"></div> Poor </div>
	Disadvantages	

2. The annual snow runoff of the drainage basin is 100 mm. The snow cover has an half an inch, most of which drains in the early spring. The snow cover is not uniform, and the snowmelt is uneven. The snowmelt is usually in late May to early June. The snowmelt is usually in late May to early June.
3. The snowmelt is usually in late May to early June. The snowmelt is usually in late May to early June.
4. The snowmelt is usually in late May to early June. The snowmelt is usually in late May to early June.
5. The snowmelt is usually in late May to early June. The snowmelt is usually in late May to early June.
6. The snowmelt is usually in late May to early June. The snowmelt is usually in late May to early June.
7. The snowmelt is usually in late May to early June. The snowmelt is usually in late May to early June.
8. The snowmelt is usually in late May to early June. The snowmelt is usually in late May to early June.
9. The snowmelt is usually in late May to early June. The snowmelt is usually in late May to early June.
10. The snowmelt is usually in late May to early June. The snowmelt is usually in late May to early June.
11. The snowmelt is usually in late May to early June. The snowmelt is usually in late May to early June.
12. The snowmelt is usually in late May to early June. The snowmelt is usually in late May to early June.
13. The snowmelt is usually in late May to early June. The snowmelt is usually in late May to early June.
14. The snowmelt is usually in late May to early June. The snowmelt is usually in late May to early June.

We express our appreciation to the well owners and well drillers in the area for their excellent cooperation in providing basic data for this study. We also thank the water superintendents of the municipalities for permission and cooperation in test pumping municipal wells.


Minnesota Division of Waters, 1959, Hydrologic Atlas of Minnesota: Division of Waters Bulletin 10, 182 p.

Nikiforoff, C. C. and others, 1939, Soil Survey (Reconnaissance) of the Red River Valley area, Minnesota: U.S. Dept. Agriculture Series 1939: No. 25.

Prior, C. H. and Hess, J. H., 1961, Floods in Minnesota, Magnitude and Frequency: Minnesota Division of Waters Bulletin 12, 142 p.

Thornthwaite, C. W. and Mather, J. R., 1955, Instructions and tables for computing potential evapotranspiration and the water balance: Drexel Institute of Technology, Publications in Climatology, Vol. X, no. 3.

OF THE MUSTINKA AND



For sale by U. S. Geological Survey, price \$2.00 per set.