# GROUND WATER

## SELECTED REFERENCES

The following list of reports is not a complete bibliography of the hydrology of the Great Basin Region. It includes only the most recent areal reports of the more heavily developed areas and selected technical reports of local and regional interest. Many published and unpublished basic-data reports and some technical reports have been omitted. More complete information on published and unpublished hydrologic data in all or specific parts of the region can be obtained from the district offices of the U.S. Geological Survey in Carson City, Nev.; Salt Lake City, Utah; Boise, Idaho; and Cheyenne, Wyo.

Bagley, J.M., Jeppson, R.W., and Milligan, C.H., 1964, Water yields in Utah: Utah State Univ. Agr. Expt. Sta. Spec. Rept. 18.

Baker, C.H., Jr., 1970, Water resources of the Heber-Kamas-Park City area, north-central Utah: Utah Dept. Nat. Resources Tech. Pub. 27.

Bjorklund, L.J., and Robinson, G.B., Jr., 1968, Ground-water resources of the Sevier River basin between Yuba Dam and Leamington Canyon, Utah: U.S. Geol. Survey Water-Supply Paper 1848.

Bolke, E.L., and Price, Don, 1969, Hydrologic reconnaissance of Curlew Valley, Utah and Idaho: Utah Dept. Nat. Resources Tech. Pub. 25.

Burnham, W.L., Harder, A.H., and Dion, N.P., 1969, Availability of ground water for large-scale use in the Malad Valley-Bear River areas of southeastern Idaho—an initial assessment: U.S. Geol. Survey open-file rept.

Busby, M.W., 1966, Annual runoff in the conterminous United States: U.S. Geol. Survey Hydrol. Inv. Atlas HA-212.

Carpenter, C.H., Robinson, G.B., Jr., and Bjorklund, L.J., 1967, Ground-water conditions and geologic reconnaissance of the upper Sevier River basin, Utah: U.S. Geol. Survey Water-Supply Paper 1836.

Carpenter, Everett, 1913, Ground water in Box Elder and Tooele Counties, Utah: U.S. Geol. Survey Water-Supply Paper 333.

Cohen, Philip, 1963, An evaluation of the water resources of the Humboldt River valley near Winemucca, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources Bull. 24.

———1964, A brief appraisal of the ground-water resources of the Grass Valley area, Humboldt and Pershing Counties, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 29.

Cohen, Philip, and Everett, D.E., 1963, A brief appraisal of the ground-water hydrology of the Dixie-Fairview Valley area, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 23.

Cohen, Philip, and Loeltz, O.J., 1964, Evaluation of hydrogeology and hydrogeochemistry of Truckee Meadows area, Washoe County, Nevada: U.S. Geol Survey Water-Supply Paper 1779-S.

Cordova, R.M., 1970, Ground-water conditions in southern Utah Valley and Goshen Valley, Utah: Utah Dept. Nat. Resources Tech. Pub. 28.

Cordova, R.M., and Subitzky, Seymour, 1965, Ground water in northern Utah Valley, Utah; A progress report for the period 1948–63: Utah State Engineer Tech. Pub. 11.

Crosthwaite, E.G., 1963, Ground-water appraisal of Antelope and middle Reese River valleys, Lander County, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 19.

Dion, N.P., 1969, Hydrologic reconnaissance of the Bear River basin in southeastern Idaho: Idaho Dept. Reclamation Water Inf. Bull. 13.

Eakin, T.E., 1962, Ground-water appraisal of the Imlay area Humboldt River basin, Pershing County, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 5.

Eakin, T.E., and Hughes, J.L., 1967, Water-resources appraisal of Steptoe Valley, White Pine and Elko Counties, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 42.

Eakin, T.E., and Lamke, R.D., 1966, Hydrologic reconnaissance of the Humboldt River basin, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources Bull. 32.

Eakin, T.E., Maxey, G.B., Robinson, T.W., Fredericks, J.C., and Loeltz, O.J., 1951, Contributions to the hydrology of eastern Nevada: Nevada State Engineer Water-Resources Bull. 12.

Eakin, T.E., Moore, D.O., and Everett, D.E., 1965, Water-resources appraisal of the upper Reese River valley, Lander and Nye Counties, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 31.

Everett, D.E., and Rush, F.E., 1965, Water-resources appraisal of Lovelock Valley, Pershing County, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 32.

———1967, A brief appraisal of the water resources of Walker Lake area, Mineral, Lyon, and Churchill Counties, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 40.

Fenneman, N.M., 1931, Physiography of the western United States: New York, McGraw-Hill Book Co., 534 p.

Feray, D.E., Oetking, P., and Renfro, H.B., 1968, Geologic highway map of the Pacific Southwest region, California and Nevada: Am. Assoc. Petroleum Geologists U.S. Geol. Highway Map Ser., Map 3.

Feth, J.H., Barker, D.A., Moore, L.G., Brown, R.J., and Veirs, C.E., 1966, Lake Bonneville: Geology and hydrology of the Weber Delta district, including Ogden, Utah: U.S. Geol. Survey Prof. Paper 518.

Gates, J.S., 1965, Reevaluation of the ground-water resources of Tooele Valley, Utah: Utah State Engineer Tech. Pub. 12.

Glancy, P.A., 1968, Water-resources appraisal of Butte Valley, Elko, and White Pine Counties, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 49.

Hahl, D.C., and Langford, R.H., 1964, Dissolved mineral inflow to Great Salt Lake and chemical characteristics of the salt lake brine, part II, technical report: Utah Geol. and Mineralog. Survey Water-Resources Bull. 3.

Hahl, D.C., and Mundorff, J.C., 1968, An appraisal of the quality of surface water in the Sevier Lake basin, Utah, 1964: Utah Dept. Nat. Resources Tech. Pub. 19.

Hardman, George, 1965, Nevada precipitation map: Nevada Univ. Agr. Expt. Sta.

Harrill, J.R., 1968, Hydrologic response to irrigation pumping in Diamond Valley, Eureka and Elko Counties, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources Bull. 35.

———1969, Hydrologic response to irrigation pumping in Hualapai Flat, Washoe, Pershing, and Humboldt Counties, Nevada 1966–67: Nevada Dept. Conserv. and Nat. Resources Water Resources Bull. 37.

———1970, Effects of ground-water development on the water regimen of Paradise Valley, Humboldt County, Nevada 1964–68, and hydrologic reconnaissance of the tributary areas: Nevada Dept. Conserv. and Nat. Resources Water Resources Bull. 39.

Harrill, J.R., and Worts, G.F., Jr., 1968, Estimated water use in Nevada 1950–65: Nevada Dept. Conserv. and Nat. Resources Water Resources-Inf. Ser. Rept. 7.

Hely, A.G., Mower, R.W., and Harr, C.A., 1970, Water resources of Salt Lake County, Utah: Utah Dept. Nat. Resources Tech. Pub. 31.

Hood, J.W., 1971, Hydrologic reconnaissance of the Park Valley area, Box Elder County, Utah: Utah Dept. Nat. Resources Tech. Pub. 30.

Hood, J.W., and Price, Don, 1970, Hydrologic reconnaissance in Grouse Creek valley, Box Elder County, Utah: Utah Dept. Nat. Resources Tech. Pub. 29.

Hood, J.W., Price, Don, and Waddell, K.M., 1969, Hydrologic reconnaissance of Rush Valley, Tooele County, Utah: Utah Dept. Nat. Resources Tech. Pub. 23.

Hood, J.W., and Rush, F.E., 1965, Water-resources appraisal of the Snake Valley area, Utah and Nevada: Utah State Engineer Tech. Pub. 14 (also Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 34).

Hood, J.W., and Waddell, K.M., 1968, Hydrologic reconnaissance of Skull Valley, Tooele County, Utah: Utah Dept. Nat. Resources Tech. Pub. 18.

———1969, Hydrologic reconnaissance of Deep Creek valley, Tooele and Juab Counties, Utah and Elko and White Pine Counties, Nevada: Utah Dept. Nat. Resources Tech. Pub. 24.

Huxel, C.J., Jr., 1966, Effects of irrigation development on the water supply of Quinn River valley area, Nevada and Oregon, 1950–64: Nevada Dept. Conserv. and Nat. Resources Water Resources Bull. 34.

———1969, Water resources and development in Mason Valley, Lyon and Mineral Counties, Nevada, 1948–65: Nevada Dept. Conserv. and Nat. Resources Water Resources Bull. 38.

Kohler, M.A., Nordenson, T.J., and Baker, D.R., 1959, Evaporation maps for the United States: U.S. Weather Bureau Tech. Paper 37.

Lamke, R.D., and Moore, D.O., 1965, Interim inventory of surface-water resources of Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources Bull. 30.

Loeltz, O.J., and Eakin, T.E., 1953, Geology and water resources of Smith Valley, Lyon and Douglas Counties, Nevada: U.S. Geol. Survey Water-Supply Paper 1228.

Malmberg, G.T., 1967, Hydrology of the valley-fill and carbonate-rock reservoirs, Pahrump Valley, Nevada-California: U.S. Geol. Survey Water-Supply Paper 1832.

Malmberg, G.T., and Worts, G.F., Jr., 1966, The effects of pumping on the hydrology of Kings River valley, Humboldt County, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources Bull. 31.

Marine, I.W., and Price, Don, 1964, Geology and ground-water resources of the Jordan Valley, Utah: Utah Geol. and Mineralog. Survey Water Resources Bull. 21.

Maxey, G.B., 1968, Hydrogeology of desert basins: Ground Water, v. 6. no. 5, p. 10–22.

Meinzer, O.E., 1911, Ground water in Juab, Millard, and Iron Counties, Utah: U.S. Geol. Survey Water-Supply Paper 277.

———1917, Geology and water resources of Big Smoky, Clayton, and Alkali Spring Valleys, Nevada: U.S. Geol. Survey Water-Supply Paper 423.

Mower, R.W., 1965, Ground-water resources of Pavant Valley, Utah: U.S. Geol. Survey Water-Supply Paper 1794.

Mower, R.W., and Nace, R.L., 1957, Water consumption by water-loving plants in the Malad Valley, Oneida County, Idaho: U.S. Geol. Survey Water-Supply Paper 1412.

Price, Don, and Bolke, E.L., 1970, Hydrologic reconnaissance of the Sink Valley area, Tooele and Box Elder Counties, Utah: Utah Dept. Nat. Resources Tech. Pub. 26.

Robinove, C.J., and Berry, D.W., 1963, Availability of ground water in the Bear River valley, Wyoming: U.S. Geol. Survey Water-Supply Paper 1539-V.

Robinson, G.B., Jr., 1971, Ground-water hydrology of the San Pitch River drainage basin, Sanpete County, Utah: U.S. Geol. Survey Water-Supply Paper 1896.

Rosa, J.M., 1968, Water yield maps for Idaho: Agr. Research Service Rept. 41-1410.

Rush, F.E., 1967, Water-resources appraisal of Washoe Valley, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 41.

———1968a, Water-resources appraisal of Clayton Valley-Stonewall Flat area, Nevada and California: Nevada Dept. Conser. and Nat. Resources Water Resources-Reconn. Ser. Rept. 45.

———1968b, Water-resources appraisal of Thousand Springs valley, Elko County, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 47.

Rush, F.E., and Everett, D.E., 1964, Ground-water appraisal of Monitor, Antelope, and Kobeh Valleys, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 30.

———1966, Water-resources appraisal of Little Fish Lake, Hot Creek, and Little Smoky Valleys, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 38.

Rush, F.E., and Kazmi, S.A.T., 1965, Water-resources appraisal of Spring Valley, White Pine and Lincoln Counties, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 33.

Sandberg, G.W., 1966, Ground-water resources of selected basins in southwestern Utah: Utah State Engineer Tech. Pub. 13.

Sinclair, W.C., 1962a, Ground-water resources of Pine Forest Valley, Humboldt County, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 4.

———1962b, Ground-water resources of Hualapai Flat, Washoe, Pershing, and Humboldt Counties, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 11.

———1963a, Ground-water appraisal of the Black Rock Desert area, northwestern Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 20.

———1963b, Ground-water appraisal of the Pueblo Valley-Continental Lake region, Humboldt County, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 22.

Smith, R.E., and Gates, J.S., 1963, Ground-water conditions in the southern and central parts of the East Shore area, Utah, 1953–1961: Utah Geol. and Mineralog. Survey Water Resources Bull. 21.

Stokes, W.L., ed., 1964, Geologic map of Utah: Utah Univ.

U.S. Public Health Service, 1962, Drinking water standards: Public Health Service Pub. 956.

U.S. Weather Bureau, 1960, Climate of the United States - Wyoming: Climatography of the United States, No. 60–48.

———1963, Normal annual and May–September precipitation (1931–60) for the State of Utah, map of Utah. scale 1:500,000.

———1965, Mean annual precipitation, 1930–57, State of Idaho: U.S. Soil Conserv. Service in cooperation with the U.S. Weather Bureau.

Utah Water and Power Board and Utah State University, 1963, Developing a State water plan, Utah's water resources—problems and needs—a challenge: Utah Water and Power Board-Utah State Univ.

Waddell, K.M., and Price, Don, 1972, Quality of surface water in the Bear River basin, Utah, Wyoming, and Idaho: U.S. Geol. Survey Hydrol. Inv. Atlas HA-417.

Walker, G.E., and Eakin, T.E., 1963, Geology and ground water of Amargosa Desert, Nevada-California: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 14.

Worts, G.F., Jr., and Malmberg, G.T., 1966, Hydrologic appraisal of Eagle Valley, Ormsby County, Nevada: Nevada Dept. Conserv. and Nat. Resources Water Resources-Reconn. Ser. Rept. 39.

Young, R.A., and Carpenter, C.H., 1965, Ground-water conditions and storage in the central Sevier Valley, Utah: U.S. Geol. Survey Water-Supply Paper 1787.

## VALLEY GROUND-WATER RESERVOIRS

Valley ground-water reservoirs, which consist of the saturated valley-fill deposits, are the chief developed source of ground water in the region. The occurrence and movement of water in the consolidated rocks surrounding the valleys are controlled largely by geologic structures such as faults, joints, and folds, and the resulting complex aquifer systems are difficult to delineate or exploit. Water from these aquifer systems discharges in springs and seeps and sustains the base flow of streams, or moves into the valley ground-water reservoirs as subsurface recharge. Most of the water that moves through the consolidated rocks ultimately discharges within the drainage basin in which it originates as precipitation, but some water may travel long distances in interbasin flow systems and discharge in a basin far removed from its place of origin. Development of ground water in some of the valley reservoirs has resulted in overdraft and long-term declines of water levels. Such areas in Nevada and Utah have been designated "critical ground-water areas" and further development is restricted or controlled by the State water administrators.

## GROUND-WATER DEVELOPMENT

Most of the valley ground-water reservoirs have undergone some degree of development by wells. At least 23 areas of ground-water development yielded more than 10,000 acre-feet to wells annually, and withdrawal by wells from all developed valley ground-water reservoirs totaled about 1 million acre-feet in 1965. Natural discharge from the valley ground-water reservoirs—exclusive of those in areas of upstream depletion (see runoff map, sheet 2) averages about 2.4 million acre-feet annually. The estimate (about 3 million acre-feet) for upstream depletions in the nominal water supply (sheet 1) includes natural ground-water discharge in the areas of upstream depletions.

*Estimated natural ground-water discharge, exclusive of areas of upstream depletion, and withdrawal by wells*

| Subregion | Discharge (Acre-feet per year) | |
|---|---|---|
| | Natural discharge (Average) | Withdrawal by wells (1965) |
| Bear River | 40,000 | 175,000 |
| Great Salt Lake | 630,000 | 300,000 |
| Sevier Lake | 329,000 | 280,000 |
| Humboldt | 480,000 | 140,000 |
| Central Lahontan | 205,000 | 42,000 |
| Tonopah | 680,000 | 98,000 |
| Total (rounded) | 2,400,000 | 1,000,000 |

## GROUND-WATER STORAGE

The estimated 190 million acre-feet of water stored in the upper 100 feet of saturated material in the valley ground-water reservoirs represents only a small fraction of the total ground-water storage in the region. The thickness of the saturated fill in most of the valleys exceeds 100 feet, and in some valleys it exceeds 1,000 feet. Moreover, large volumes of water are stored in the consolidated rocks of the region. Therefore, the 190 million acre-feet is the minimum volume of water that is economically recoverable from storage. Not all the ground water in storage is fresh, however; most of the valley reservoirs contain some saline water, especially those reservoirs near the major terminal lakes and sinks. Hence, full development of the valley ground-water reservoirs would require desalinization of some of the water.

*Estimated ground-water storage in the upper 100 feet of saturated valley-fill deposits*

| State | Assumed average specific yield (percent) | Area (acres) | Storage in upper 100 feet of saturation (1,000 acre-feet) |
|---|---|---|---|
| **BEAR RIVER SUBREGION** | | | |
| Idaho | 10 | 538,000 | 5,400 |
| Utah | 10 | 537,000 | 5,400 |
| Wyoming | 10 | 123,000 | 1,200 |
| Total | | | 12,000 |
| **GREAT SALT LAKE SUBREGION** | | | |
| Idaho | 10 | 108,000 | 1,100 |
| Nevada | 10 | 404,000 | 4,000 |
| Utah | 10 | 1,490,000 | 14,900 |
| | 5 | 5,980,000 | 29,900 |
| Total | | | 49,900 |
| **SEVIER LAKE SUBREGION** | | | |
| Utah | 10 | 1,670,000 | 16,700 |
| | 5 | 990,000 | 5,000 |
| Total | | | 21,700 |
| **HUMBOLDT SUBREGION** | | | |
| Nevada | 10 | 3,910,000 | 39,100 |
| | 5 | 908,000 | 4,500 |
| Total | | | 43,600 |
| **CENTRAL LAHONTAN SUBREGION** | | | |
| Nevada | 10 | 1,530,000 | 15,300 |
| | 5 | 166,000 | 830 |
| Total | | | 16,130 |
| **TONOPAH SUBREGION** | | | |
| Nevada | 10 | 5,060,000 | 50,600 |
| **GREAT BASIN REGION** | | | |
| Idaho | | 646,000 | 6,500 |
| Nevada | | 11,980,000 | 114,300 |
| Utah | | 10,700,000 | 71,900 |
| Wyoming | | 123,000 | 1,200 |
| Total (rounded) | | | 190,000 |

## QUALITY OF GROUND WATER

The general suitability of water for most uses is closely related to the dissolved-solids content of the water, although concentrations of individual materials in the water can limit its usefulness for some purposes. For domestic use, the U.S. Public Health Service (1962) recommends water with a maximum dissolved-solids content of 500 mg/l (milligrams per liter), but waters containing up to 1,000 mg/l are generally used for domestic supplies where better water is not available. The combination of dissolved-solids content and SAR (sodium-adsorption ratio) is often used to evaluate the suitability of the water for irrigation use. The larger the dissolved-solids content and SAR of the water, the less suitable the water is likely to be for irrigation. However, these factors must be considered in conjunction with soil types and drainage conditions, as well as the presence of substances, such as boron, that are toxic to certain plants.




GRAPH SHOWING RELATION BETWEEN DEPTHS AND DISSOLVED SOLIDS IN WATER FROM SELECTED WELLS.

## QUALITY VARIATION WITH DEPTH

In most places, there is an increase in dissolved solids in ground water with increasing depth. In some parts of the region, however, shallow ground water is more highly mineralized than deeper ground water. This condition commonly occurs in and near terminal sinks of topographically closed valleys where deep, relatively fresh water moves upward under artesian pressure into shallower aquifers and dissolves minerals along its upward course. In these terminal areas, the mineral content of shallow ground water is concentrated by evapotranspiration.




MAP SHOWING GENERAL DISTRIBUTION OF VALLEY GROUND-WATER RESERVOIRS AND CRITICAL GROUND-WATER AREAS (1965 LEVEL OF DEVELOPMENT).




MAP SHOWING DISSOLVED SOLIDS AND LOCATIONS OF PRINCIPAL SALINE SPRINGS




| Map number | Areas of principal withdrawal | 1965 withdrawal (nearest thousand acre-feet) |
|---|---|---|
| 1 | Kings River Valley (Rio King area) | 32,000 |
| 2 | Quinn River Valley (Orovada District) | 45,000 |
| 3 | Humboldt River | 35,000 |
| 4 | Pahrump Valley | 35,000 |
| 5 | Gem Valley | >10,000 |
| 6 | Gentile Valley | >10,000 |
| 7 | Upper Malad Valley | >10,000 |
| 8 | Cache Valley | 28,000 |
| 9 | East Shore area | 59,000 |
| 10 | Tooele Valley | 20,000 |
| 11 | Jordan Valley | 102,000 |
| 12 | Salt Lake Valley, Utah | 67,000 |
| 13 | Goshen Valley | 14,000 |
| 14 | Southern Utah Valley | 17,000 |
| 15 | Juab Valley | 18,000 |
| 16 | Sevier Desert | 27,000 |
| 17 | Sanpete Valley | 12,000 |
| 18 | Pavant Valley | 69,000 |
| 19 | Escalante Valley | 44,000 |
| 20 | Milford district | 44,000 |
| 21 | Beryl-Enterprise district | 70,000 |
| 22 | Cedar City Valley | 16,000 |
| 23 | Parowan Valley | 15,000 |

MAP SHOWING PRINCIPAL AREAS OF NATURAL GROUND-WATER DISCHARGE AND WITHDRAWALS BY WELLS.




MAP SHOWING SODIUM-ADSORPTION RATIO

# WATER IN THE GREAT BASIN REGION; IDAHO, NEVADA, UTAH, AND WYOMING

By
Don Price, T. E. Eakin, and others
1974

INTERIOR—GEOLOGICAL SURVEY, RESTON, VIRGINIA—1974