TABLE. 1—*Generalized correlation chart showing time-stratigraphic, rock-stratigraphic and regional hydrogeologic units in the southeastern United States Coastal Plain*

ERA-THEM	SYSTEM	Series	Gulf Coast Stage	Mississippi (N–S)	Alabama (W–E)	Georgia (W–E)	South Carolina (W–NE)	North Carolina (SE–Mid-central)	Northern Florida (Panhandle–Northern peninsula)
CENOZOIC	QUATERNARY	Holocene and Pleistocene		Alluvium and terrace deposits	Alluvium and terrace deposits	Alluvium and terrace deposits	Alluvium and terrace deposits; Pamlico Formation; Talbot Formation; Penholoway Formation; Wicomico Formation; Okefenokee Formation; Waccamaw Formation; Bear Bluff Formation; Socastee Formation	Alluvium and terrace deposits; Canepatch Formation; Waccamaw Formation; Bear Bluff Formation; Yorktown Formation	Terrace deposits
		Pliocene		Citronelle Formation; Graham Ferry Formation	Citronelle Formation				Citronelle Formation
	TERTIARY	Miocene		Pascagoula Formation; Hattiesburg Formation; Catahoula Sandstone	Undifferentiated; Catahoula Sandstone	Hawthorn Formation	Hawthorn Formation; Ashley Formation; Edisto Formation; Chandler Bridge Formation; Cooper Formation; Ashley Member	Hawthorn Formation; Pungo River Formation; Belgrade Formation; River Bend Formation	Alum Bluff Group; Hawthorn Formation; Tampa Limestone and equivalents; Suwannee Limestone
		Oligocene	Chickasawhayan	Paynes Hammock Formation; Chickasawhay Limestone	Paynes Hammock Formation; Chickasawhay Limestone	Cooper Marl; Suwannee Limestone			Chickasawhay Limestone; Ocala Limestone; Suwannee Limestone
			Vicksburgian	Bucatunna Formation; Byram Formation; Glendon Formation; Marianna Formation; Mint Spring Formation	Bucatunna Formation; Byram Formation; Glendon Formation; Marianna Formation; Mint Spring Formation; Red Bluff Formation				Bucatunna Clay Member of Byram Formation; Byram Formation; Glendon Limestone; Marianna Limestone; Bumpnose Limestone
		Eocene	Jacksonian	Forest Hill Formation; Red Bluff Formation; Yazoo Formation	Yazoo Formation; Ocala Limestone	Yazoo Clay; Ocala Limestone; Moodys Branch Formation; Barnwell Formation (updip); Santee Limestone; Gosport Sand	Barnwell Formation (updip); McBean Formation (updip); Santee Limestone; Gosport Sand	Castle Hayne Formation	Ocala Limestone; Moodys Branch Formation
			Claibornian	Moodys Branch Formation; Cocoa Sand Member of Moodys Branch Formation; Cocaina Creek Member of Moodys Branch Formation; Cocktield Formation; Cook Mountain Formation; Sparta Sand; Zilpha Clay; Winona Sand; Neshoba Sand Member; Basic City Shale Member; Meridian Sand Member; Tallahatta Formation	Gosport Sand; Lisbon Formation; Tallahatta Formation	McBean Formation (updip); Lisbon Formation; Tallahatta Formation; Warley Hill Marl (updip); Congaree Formation (updip); Fishburne Formation (subsurface); Upper Confining Unit	Beaufort Formation	Lisbon Formation; Tallahatta Formation; Undifferentiated lower Eocene rocks; Oldsmar Limestone; Cedar Keys Formation
			Sabinian	Bashi Formation; Hatchetigbee Formation; Tuscahoma Formation; Nanafalia Formation	Bashi Formation; Hatchetigbee Formation; Tuscahoma Formation; Nanafalia Formation	Bashi Formation; Hatchetigbee Formation; Tuscahoma Formation; Nanafalia Formation; Cedar Keys Formation; Black Mingo Formation			
		Paleocene	Midwayan	Naheola Formation; Porters Creek Clay; Tippah Sand Lentil; Clayton Formation	Naheola Formation; Porters Creek Formation; Clayton Formation	Clayton Formation	Clayton Formation; Williamsburg Formation; Snapp Formation; Rhems Formation; Beaufort (?) Formation (subsurface)		
MESOZOIC	CRETACEOUS	Upper Cretaceous	Navarroan	Owl Creek Formation; Prairie Bluff Formation; Ripley Formation; Clayton Formation; Coon Creek Tongue; McNairy Sand Member; Mcshan Formation; Coon Creek Tongue	Prairie Bluff Formation; Providence Sand; Ripley Formation	Providence Sand; Ripley Formation	Peedee Formation	Peedee Formation	Rocks of Navarroan age
			Taylorian	Demopolis Chalk; Coffee Sand; Arcola Limestone Member; Mooreville Chalk	Demopolis Chalk; Cusseta Sand Member; Mooreville Chalk	Cusseta Sand	Black Creek Formation	Black Creek Formation	Rocks of Taylorian age
			Austinian	Tombigbee Sand Member; Eutaw Formation	Tombigbee Sand Member; Eutaw Formation; Blufftown Formation	Blufftown Formation; Eutaw Formation	Unnamed sediments in coastal areas; Middendorf Formation	Unnamed sediments in coastal areas; Middendorf Formation	Rocks of Austinian age
			Gulfian	McShan Formation	McShan Formation	Atkinson Formation (upper part–subsurface)	Cape Fear Formation	Cape Fear Formation	
			Eaglefordian	Tuscaloosa Group; Gordo Formation; Coker Formation; Unnamed upper member; Mantee sandy member; Massive sand; Marine shale	Gordo Formation; Coker Formation; Tuscaloosa Group; Unconformity; Absence of Eagleford and Woodbine; Marine shale	Tuscaloosa Formation; Atkinson Formation (lower part–subsurface); Marine shale	Unnamed rocks; Marine shale	Unnamed rocks; Marine shale	Unnamed rocks; Unnamed rocks; Atkinson Formation
			Woodbinian						
		Lower Cretaceous (Comanchean)	Washitan and Fredericksburgian	Undifferentiated	Undifferentiated	Undifferentiated	Undifferentiated	Undifferentiated	Undifferentiated; ?
PALEOZOIC	JURASSIC			Undifferentiated	Undifferentiated	Undifferentiated	Undifferentiated	Undifferentiated	Undifferentiated
	TRASSIC			Undifferentiated	Undifferentiated	Undifferentiated	Undifferentiated	Undifferentiated	Undifferentiated

EXPLANATION

SURFICIAL AQUIFER

UPPER CONFINING UNIT

FLORIDAN AQUIFER SYSTEM

SOUTHEASTERN COASTAL PLAIN AQUIFER SYSTEM

A1, aquifer unit

C1, confining unit

A2, aquifer unit

C2, confining unit

A3, aquifer unit

C3, confining unit

A4, aquifer unit

BASE OF SYSTEM

ABSENT

LOCAL UNIT NOT CONSIDERED PART OF SYSTEM

[1] Mancini and others (1985) considered the Marine Shale of Southern Alabama to be part age equivalent to a portion of the Woodbine Formation and/or Eagle Ford Group of Texas.

[2] Since the report was prepared the Cooper Formation in South Carolina has been raised to group and its members to formation rank. (Weems and Lemon, 1984).

[3] Brown and others, 1972; Brown and others, 1979.

A4 REGIONAL AQUIFER

The A4 aquifer is the most regionally extensive clastic aquifer of the Southeastern United States Coastal Plain (plate 3). Correlative sand beds that are part of this aquifer extend in the subsurface or outcrop from Tennessee into central North Carolina. The upper surface of the A4 aquifer slopes gently seaward at a gradient of 15 to 30 feet per mile in Georgia, northern Florida, South Carolina, and adjacent counties of North Carolina but slopes more steeply in Alabama and Mississippi (30 to 50 feet per mile or greater). Where this aquifer unit crops out in western Georgia, Alabama, and Mississippi, its landward limit marks the inner margin of Coastal Plain sediments. Cropping out as a 40- to 50-mile wide band in eastern Mississippi–western Alabama, its width of outcrop decreases gradually in eastern Alabama and western Georgia; the A4 unit is completely covered in eastern and central Georgia by cropping rocks of Late Cretaceous and Tertiary age. Where the updip limit of this aquifer is shown to occur well south of the Fall Line such as in eastern Georgia and South Carolina, its location is due to erosion or nondeposition of equivalent strata. Although the A4 aquifer does not crop out in South Carolina, equivalent strata are exposed locally along the Cape Fear River in North Carolina. The A4 aquifer is absent along the axis of the Peninsular arch in northeastern Florida and southeastern Georgia.

Poorly sorted, nonmarine, mottled red to reddish-brown, fine to coarse grained, feldspathic quartz sand considered, in part, equivalent to the outcropping Cape Fear Formation and in part, equivalent to Unit F (Brown and others, 1972) forms the top of the A4 aquifer in South Carolina and adjacent counties of North Carolina. Interbedded noncalcareous clay, siderite, hematite, pyrite, shell fragments and a few microfossils are common minor constituents associated with these water-bearing rocks.

Subsurface strata previously described as the upper part of the Atkinson Formation (Applin and Applin, 1967) outline the upper surface of the A4 aquifer in Georgia, in northern Florida, and in southeastern Alabama. In these areas, rock that comprises the upper surface of the aquifer consist largely of marine deposits but includes some nonmarine beds. The aquifer typically consists in these localities of glauconitic quartz sand and sandstone that are interbedded with silt, shale and minor limestone; the sands are feldspathic in shallow updip regions.

In most of Mississippi and Alabama, and in the northwestern Georgia Coastal Plain, strata that are equivalent to the Eutaw or Eutaw and McShan Formation(s) make up the top of the A4 aquifer. In those localities, the aquifer is comprised of sparsely fossiliferous, greenish gray to yellowish brown, fine to coarse grained, glauconitic, calcareous sand that is interbedded with gray micaceous shale. Minor constituents include volcanic ash (bentonite), siderite, pyrite and lignite.

In downdip areas of southwestern Mississippi, the permeability of the Eutaw and McShan sand decreases as these rocks grade into dense marine chalk and argillaceous shale that are part of the C3 confining unit (fig. 5). As a result, the top of the A4 aquifer here consists of underlying sand, sandstone, and gravel that are considered equivalent to the upper part of the Tuscaloosa Group. Important water-bearing units within the A4 aquifer that are part of the Tuscaloosa Group or Formation include: (1) deltaic, highly cross-bedded, fine to coarse grained, ferruginous quartz sand and cherty gravel deposits ascribed to the Gordo Formation; (2) fluvial and nearshore marine, glauconitic quartz sand deposits of the Coker Formation; and (3) nonmarine, massively bedded, medium to coarse grained quartz sand of the so-called "massive sand" section. Taken as a whole, strata of the Tuscaloosa Group or Formation tend to be interbedded with dark, micaceous marine shale as they grade coastward.

Outcropping and subsurface strata that are part of the Coffee Sand form the top of the A4 aquifer in some of the northeastern counties of Mississippi. These strata consist of massively bedded, marine, fine to medium grained, loosely consolidated, glauconitic quartz sand that is often laminated with silt and clay; minor constituents include mica, siderite, and lignite.

Permeable clastic rocks of Early Cretaceous age (Washitan and Fredericksburgian Stages) form the base of the A4 aquifer in downdip areas of Georgia, Alabama, Mississippi, and northern Florida but are not known to crop out in the Southeastern Coastal Plain. These rocks consist of red to reddish brown sand, sandstone, and gravel that were deposited under nonmarine conditions; interlaminated marine and nonmarine shale, siltstone and minor limestone beds are also present.

Deeply buried sedimentary rocks of Jurassic age that are found in parts of Georgia, Alabama, Mississippi, and Florida underlie the basal Cretaceous sedimentary rocks. However, they are not considered to be part of the Southeastern Coastal Plain regional aquifer system. Jurassic rocks were excluded from the system because they are not known to contain ground water with a dissolved-solids content of less than 10,000 mg/L (milligrams per liter). It is assumed that little or no movement of ground water occurs in these deeply buried saline aquifers.

RELATION OF STRATIGRAPHY AND PERMEABILITY

The extent and thickness of the major aquifer and confining units that comprise the Southeastern Coastal Plain aquifer system are primarily a function of gradational changes in permeability, and secondarily, a result of differential movement within the rocks that comprise the Coastal Plain and its floor. The following discussion demonstrates the control of these factors on the distribution, geometry, and character of the major aquifers and confining units of the Southeastern Coastal Plain aquifer system.

The relation between an aquifer's permeability and lithology is an important criterion in delineating the boundaries of water-bearing units. For example, the highly permeable Floridan aquifer system overlies and is hydraulically interconnected with the less-permeable A2 aquifer in Georgia, southwestern South Carolina, and southern Alabama. The boundary between the two aquifers represents a transitional, or facies, boundary separating carbonate rocks of the Floridan from the underlying clastic rocks. This boundary generally represents a contrast in permeability. From a regional perspective, the permeability of the Floridan is one or more orders of magnitude higher than that of the underlying A2 aquifer. Consequently, the configuration of the top of the A2 aquifer is largely coincident with the base of the Floridan aquifer system (Miller, 1982a) where the two systems are juxtaposed. The "transgression" of this boundary across time-stratigraphic lines, shown on several of the hydrogeologic sections presented in this report (figs. 4, 6, 8, 9, and 10), explains the diversity of rock units that comprise the top of the A2 aquifer.

The wavy, undulating contour pattern that typifies the surface of the A2, C2, A3, and C3 regional units generally reflects local changes in permeability. This relationship is shown by the configuration of the basal surface of the A3 aquifer (plate 2B). Hydrogeologic sections F-F' and G-G' (figs. 10 and 11) show the A3 aquifer grading coastward from porous sand, sandstone, and minor limestone beds found in updip localities into low-permeability chalk and calcareous shale that lie in the deep subsurface of southwestern Georgia and southeastern Alabama. Consequently, the permeable parts of the A3 aquifer thin seaward greatly. These sections also show that permeable rocks of Tertiary age extend further downdip than the underlying permeable Cretaceous rocks that comprise the A3 aquifer. This is in contrast to southeastern Georgia where clastic Cretaceous strata of the A3 aquifer grade seaward to permeable limestone (Lawson Limestone) that is part of the Floridan aquifer system (fig. 9). Such major permeability variations overshadow the influence of minor structural features and account for many of the local highs and lows drawn on the maps showing the top or base of the different regional aquifer units. For example, the apparent structural low that is mapped on the base of the A2 aquifer in Colquitt and Grady Counties in southwest Georgia and its alignment with the position of the postulated Gulf Trough (compare plates 1A, 2B, and fig. 13) is considered to be coincidental.

In spite of the generally poor correspondence between hydrogeologic an stratigraphic units of the Southeastern United States Coastal Plain, several large and small-scale structural features are subtly reflected on the top of the major hydrogeologic units. Major embayments that appear on maps of some of these units include the Southeast Georgia, Southwest Georgia, and Mississippi embayments. The Southeast Georgia embayment (Maher, 1971) is centered and most prominent in coastal counties of southeastern Georgia and is evident on the map showing the surface of the A4 aquifer (plate 3). The Southwest Georgia embayment forms the dominant structural element in the southern tier counties of Georgia and the central Florida panhandle; it is strongly reflected by the contour pattern shown on the A4 aquifer map. The Mississippi embayment, with its north-south trending axis nearly coincident with the Mississippi River (fig. 13), strongly controls the western and southwestern dip of beds in western Mississippi.

The Cape Fear and the Peninsular arches form two major salients of the Southeastern United States Coastal Plain (fig. 13). They have resulted in the nondeposition or erosion of Tertiary clastic rocks in eastern South Carolina as well as the absence of the basal Cretaceous sands in the southeast Georgia-northeast Florida region (see plans 1A, 1B, and 3). The effect of the Wiggins anticline in Southeastern Mississippi is limited to the deep subsurface strata that comprise the surface of the A4 aquifer (plate 3). Shallower overlying clastic aquifers (A2, A3) do not extend as far south and were not influenced by this structural feature. The Jackson dome, near Jackson, MS, forms a positive feature and strongly controls the configuration of the A2 and A4 aquifers (plates 1A, and 3).

Small- and large-scale faults associated with Pickens-Gilbertown fault system (fig. 13) bound graben-type structures that displace sediments downward in varying degrees across southwestern Alabama and south-central Mississippi. Detailed mapping of these complex features is beyond the scope of this report but the position of the system as previously mapped (Gulf Coastal Association of Geological Societies, 1972) is shown. Downwarping or faulting associated with the postulated Gulf trough (Herrick and Vorhis, 1963, and Gelbaum, 1978) in south Georgia has been inferred on the basis of limited well control and is mapped on the upper surface of the A2 aquifer. However, this feature is not discernable at the base of the A2 aquifer and its depositional and structural influence may have been limited to post-Sabine time. Likewise, the postulated Miller fault of Faye and Prowell (1982) is not recognizable at the top of any of the aquifers or confining units mapped herein. The Miller fault, which supposedly displaced upper Cretaceous rocks several hundred feet and affected the potentiometric surface of two aquifers in Burke County, GA and Allendale and Barnwell Counties, S.C., has been shown by subsequent drilling (Bechtel Corporation, 1982) not to exist.

An unnamed northwest-trending structural lineament or possible fault centered in Marion and Dillon Counties, S.C., and Robeson County, N.C., is inferred on the basis of an anomalous change in dip on the upper surface of the A4 aquifer (plate 3). There is subparallel alignment of this postulated structural component with the so-called "Florence basin" of Triassic age (Popenoe and Zietz, 1977; Daniels and others, 1983), possibly indicating that border faulting associated with the Florence basin continued from Triassic into Late Cretaceous time. However, information is not available to confirm or deny this feature or ascertain its affect on the regional ground-water flow system.



PLATE 3.—Geology and configuration of the A4 regional aquifer.



FIGURE 11.—Generalized hydrogeologic section from eastern Alabama to northern Mississippi.



From Applin and Applin, 1967; Vernon and Puri, 1955; Copeland, 1968; Williams, 1969; Cederstrom and others, 1979; Gelbaum and Howell, 1982

FIGURE 13.—Principal structural features of the Southeastern Coastal Plain of the United States.



FIGURE 12.—Generalized hydrogeologic section in central Alabama.

SELECTED REFERENCES

Anderson, E.G., 1980, Regional stratigraphy of the Woodbine-Tuscaloosa (northern coastal region): Gulf Coast Association of Geological Societies Transactions, v. 30, p. 35–39.

Applin, E.R., 1955, A biotacies of Woodbine age in the southeast Gulf Coast region: U.S. Geological Survey Professional Paper 264–I, p. 187–197.

Applin, E.R., and Applin, P.L., 1964, Logs of selected wells in the Coastal Plain of Georgia: Georgia Geological Survey Bulletin 74, 229 p.

Applin, P.L., 1951, Preliminary report on buried Pre-Mesozoic rocks in Florida and adjacent states: U.S. Geological Survey Circular 91, 28 p.

Applin, P.L., and Applin, E.R., 1944, Regional subsurface stratigraphy and structure of Florida and southern Georgia: American Association of Petroleum Geologists Bulletin, v. 28, p. 1673–1753.

_____ 1947, Regional subsurface stratigraphy and correlation of middle and early Upper Cretaceous rocks in Alabama, Georgia and north Florida: U.S. Geological Survey Oil and Gas Investigations (Preliminary), Chart 26.

_____ 1965, The Comanche Series and associated rocks in the subsurface in central and south Florida: U.S. Geological Survey Professional Paper 447, 84 p.

_____ 1967, The Gulf Series in the subsurface in northern Florida and southern Georgia: U.S. Geological Survey Professional Paper 524–G, 35 p.

Babcock, Clarence, 1969, Geology of the Upper Cretaceous clastic section, northern peninsular Florida: Florida Geological Survey Information Circular 60, 44 p.

Bechtel Corporation, 1982, Studies of the postulated Millet Fault - Report prepared for Georgia Power Company with Southern Company Services, v. 1 and 2.

Berg, R.R., and Cook, B.C., 1968 Petrography and origin of Lower Tuscaloosa sandstones, Mallalieu Field, Lincoln County, Mississippi: Gulf Coast Association of Geological Societies Transactions, v. 18, p. 242–255.

Bicker, Jr., A.R., 1969, Geologic map of Mississippi: Mississippi Geological Survey, scale 1:500,000, 1 sheet.

Blackwelder, B.W., and Ward, L.W., 1979, Stratigraphic revision of the Pliocene deposits of North and South Carolina: South Carolina Geological Survey Geologic Notes, v. 23, no. 1, p. 33–49.

Blackwelder, B.W., Ward, L.W., Gohn, G.S., and Poore, R.Z., 1979, Stratigraphic revision of Eocene, Oligocene, and Lower Miocene Formations of South Carolina: South Carolina Geological Survey Geologic Notes, v. 23, no. 1, p. 2–32.

Boswell, E.H., 1963, Cretaceous aquifers of northeastern Mississippi: Mississippi Board of Water Commissioners Bulletin 63–10, 202 p.

_____ 1976a, The lower Wilcox aquifer in Mississippi: U.S. Geological Survey Water-Resources Investigations 60–75, 3 sheets.

_____ 1976b, The Meridian-upper Wilcox aquifer in Mississippi: U.S. Geological Survey Water-Resources Investigations 76–79, 3 sheets.

_____ 1977, The Eutaw-McShan aquifer in Mississippi: U.S. Geological Survey Water-Resources Investigations 76–134, 2 sheets.

_____ 1978a, The Tuscaloosa aquifer system in Mississippi: U.S. Geological Survey Water-Resources Investigations 78–98, 3 sheets.

_____ 1978b, The Coffee Sand and Ripley aquifers in Mississippi: U.S. Geological Survey Water-Resources Investigations 78–114, 1 sheet.

_____ 1979, The Citronelle aquifer in Mississippi: U.S. Geological Survey Water-Resources Investigations 78–131, 1 sheet.

Boswell, E.H., Moore, G.K., MacCary, L.M., and others, 1965, Cretaceous aquifers in the Mississippi embayment, with a discussion on the quality of the water by H. G. Jeffery: U.S. Geological Survey Professional Paper 448–C, 37 p.

Brown, P.M., Brown, D. L., Reid, M. S., and Lloyd, O. B., Jr., 1979, Evaluation of the geologic and hydrologic factors related to the waste-storage potential of Mesozoic aquifers in the southern part of the Atlantic Coastal Plain, South Carolina, and Georgia: U.S. Geological Survey Professional Paper 1088, 37 p.

Brown, P.M., Miller, J.A., and Swain, F.M., 1972, Structural and stratigraphic framework, and spatial distribution of permeability of the Atlantic Coastal Plain, North Carolina to New York: U.S. Geological Survey Professional Paper 796, 79 p.

Cagle, J.W., 1963, Geologic map of Escambia County, Alabama: Alabama Geological Survey Map 26, scale 1:60,000, 1 sheet.

Cederstrom, D.J., Boswell, E.H., and Tarver, G.R., 1979, Summary appraisals of the nation's ground-water resources - South Atlantic - Gulf Region: U.S. Geological Survey Professional Paper 813–O, 35 p.

Cepek, P., Hay, W.M., Masters, B.A., and Worsley, T.R., 1968, Calcereous plankton in the Upper Cretaceous of Alabama, in Scott, J.C., Chairman, Facies changes in the Selma Group in central and eastern Alabama: Alabama Geological Society Guidebook for the Sixth Annual Field Trip, p. 27–32.

Christopher, R.A., 1982a, Palynostratigraphy of the basal Cretaceous units of the eastern Gulf and southern Atlantic Coastal Plains, in Arden, D.D., Beck, B.F., and Morrow, E., eds., Symposium on the Geology of the Southeastern Coastal Plain: Georgia Geological Survey Information Circular 53, p. 18–23.

_____ 1982b, The occurrence of the Complexopollis-Aritanopollis Zone (Palynomorphs) in the Eagle Ford Group (Upper Cretaceous) of Texas: Journal of Paleontology, v. 56, no. 2, p. 525–541.

Christopher, R.A., Owens, J.P., and Sohl, N.F., 1979, Late Cretaceous palynomorphs from the Cape Fear Formation of North Carolina: Southeastern Geology, v. 20, no. 3, p. 145–159.

Cleaves, A.W., 1980, Depositional systems and lignite prospecting models: Wilcox Group and Meridian sandstone of northern Mississippi: Gulf Coast Association of Geological Societies Transactions, v. 30, p. 283–307.

Colquhoun, D.J., Heron, S.D., Jr., Johnson, H.S., Jr., Pooser, W.K., and Siple, G.E., 1969, Up-dip Paleocene-Eocene stratigraphy reviewed: South Carolina Geological Survey Geologic Notes, v. 13, no. 1, p. 1–25.

Colquhoun, D.J., Oldham, R.W., Bishop, J.W., and Howell, P.D., 1982, Updip delineation of the Tertiary limestone aquifer, South Carolina: Final Technical Completion Report to Office of Water Policy, Department of the Interior, 93 p.

Colquhoun, D.J., Woolen, I.D., Van Nieuwenhuise, D.S., Padgett, G.G., Oldham, R.W., Boylan, D.C., Bishop, J.W., and Howell, P.D., 1983, Surface and subsurface stratigraphy, structure, and aquifers of the South Carolina Coastal Plain, Report to Department of Health and Environmental Control, Ground Water Protection Division, 78 p.

Cooke, C.W., 1936, Geology of the Coastal Plain of South Carolina: U.S. Geological Survey Bulletin 867, 196 p.

_____ 1943, Geology of the Coastal Plain of Georgia: U.S. Geological Survey Bulletin 941, 121 p.

Copeland, C.W., ed., 1966, Facies changes in the Alabama Tertiary: A Guidebook for the Fourth Annual Field Trip of the Alabama Geological Society, 103 p.

_____ 1968a, Geology of the Alabama Coastal Plain - a guidebook: Alabama Geological Survey Circular 47, 97 p.

_____ 1968b, Facies changes in the Selma Group in central and eastern Alabama, in Scott, J.C., Chairman, Facies changes in the Selma Group in central and eastern Alabama: Alabama Geological Society Guidebook for the Sixth Annual Field Trip, p. 2–26.

Copeland, C.W., Newton, J.G., and Sell, D.M., eds., 1976, Cretaceous and Tertiary faults in southwestern Alabama: Alabama Geological Society Guidebook for the Fourteenth Annual Field Trip, 114 p.

Cramer, H.R., 1969, Structural features of the Coastal Plain of Georgia: Southeastern Geology, v. 10, no. 2, p. 111–123.

_____ 1974, Isopach and lithofacies analysis of the Cretaceous and Cenozoic rocks of the Coastal Plain of Georgia, in Stafford, L.P., ed., Symposium on the Petroleum Geology of the Georgia Coastal Plain: Georgia Geological Survey Bulletin 87, p. 21–43.

Cramer, H.R., and Arden, D.D., 1980, Subsurface Cretaceous and Paleogene geology of the Coastal Plain of Georgia: Georgia Geologic Survey Open-File Report 80-8, 184 p.

Cushing, E.M., 1966, Map showing altitude of the base of freshwater in coastal plain aquifers of the Mississippi embayment: U.S. Geological Survey Hydrologic Investigations Atlas HA–221, 1 sheet.

Cushing, E.M., Boswell, E.H., and Hosman, R.L., 1964, General geology of the Mississippi embayment: U.S. Geological Survey Professional Paper 448-B, 28 p.

Cushing, E.M., Boswell, E.H., Speer, P.R., Hosman, R.L., and others, 1970, Availability of water in the Mississippi embayment: U.S. Geological Survey Professional Paper 448–A, 13 p.

Daniels, David L., Zietz, Isidore, and Popenoe, Peter, 1983, Distribution of subsurface Lower Mesozoic rocks in the southeastern United States, as interpreted from regional aeromagnetic and gravity maps, in Gohn, Gregory, S., ed., Studies related to the Charleston, South Carolina earthquake of 1886 - tectonics and seismicity: U. S. Geological Survey Professional Paper 1313, p. R1–R24.

Davis, M.E., 1980, Generalized geologic map and southern extent of major use of ground water from principal aquifers in Alabama: Alabama Geological Survey Circular 105, scale 1:1,000,000, 1 sheet.

Davis, M.E., Sparkes, A.K., and Peacock, B.S., 1983, Results of a test well in the Nanafalia Formation near Melvin, Alabama: U.S. Geological Survey Water-Resources Investigations 82–4108, 17 p.

Devery, D.M., 1982, Subsurface Cretaceous strata of Mississippi: Mississippi Bureau of Geology Information Series 82–1, 24 p.

Drennen, C.W., 1953, Reclassification of outcropping Tuscaloosa Group in Alabama: American Association of Petroleum Geologists Bulletin, vol. 37, no. 3, p. 522–538.

Eargle, H.D., 1955, Stratigraphy of the outcropping Cretaceous rocks of Georgia: U.S. Geological Survey Bulletin 1014, 101 p.

Faye, R.F., and Prowell, D.C., 1982, Effects of Late Cretaceous and Cenozoic faulting on the geology and hydrology of the Coastal Plain near the Savannah River, Georgia and South Carolina: U.S. Geological Survey Open-File Report 82–156, 73 p.

Fenneman, N.M., 1938, Physiography of the eastern United States: New York, McGraw Hill, 714 p.

Fenneman, N.M., and Johnson, D.W., 1946, Map of physical division of the United States: U.S. Geological Survey Map 7–C.

Fisher, V.N., 1974, Northeast-southwest stratigraphic cross section of the Jurassic sediments, Newton County to Jasper County, Mississippi: Mississippi Geological Survey, 1 sheet.

Frederiksen, N.O., Gibson, T.G., and Bybell, L.M., 1982, Paleocene-Eocene boundary in the eastern Gulf Coast: Gulf Coast Association of Geological Societies Transactions, vol. 32, p. 289–294.

Gandle, L.A., 1979, The Oligocene-Eocene aquifer system in Mississippi: U.S. Geological Survey Water-Resources Investigations 79–28, 2 sheets.

Gelbaum, C.S., 1978, The geology and ground water of the Gulf Trough: Georgia Geologic Survey Bulletin 93, p. 38–47.

Gelbaum, C.S., and Howell, Julian, 1982, The geography of the Gulf Trough, in Arden, D.D., Beck, B.F., and Morrow, E., eds., Second Symposium on the geology of the Southeastern Coastal Plain: Georgia Geologic Survey Information Circular 53, p. 140–153.

Georgia Geological Survey, 1976, Geologic map of Georgia, no. 7, scale 1:500,000, 1 sheet.

Gibson, T.G., Mancini, E.A., and Bybell, L.M., 1982, Paleocene to Middle Eocene stratigraphy of Alabama: Gulf Coast Association of Geological Societies, vol. 32, p. 449–458.

Gibson, T.G., 1979, Paleocene to Middle Eocene depositional cycles in eastern Alabama and western Georgia, in Arden, D.D., Beck, B.F., and Morrow, Eleanore, eds., Symposium on the geology of the Southeastern Coastal Plain: Georgia Geological Survey Information Circular 53, p. 53–63.

Gohn, G.S., Bybell, L.M., Christopher, R.A., Owens, J.P., and Smith, C.C., 1979, A stratigraphic framework for Cretaceous and Paleogene margins along the South Carolina and Georgia coastal sediments, in Arden, D.D., Beck, B.F. and Morrow, E., eds., Symposium on the geology of the Southeastern Coastal Plain: Georgia Geological Survey Information Circular 53, p. 64–74.

Gohn, G.S., Bybell, L.M., Smith, C.C., and Owens, J.P., 1978, Preliminary stratigraphic cross sections of Atlantic Coastal Plain sediments of the Southeastern United States—Cenozoic sediments along the South Carolina coastal margin: U.S. Geological Survey Miscellaneous Field Studies Map MF–1015-B, 2 sheets.

Gohn, G.S., Christopher, R.A., Smith, C.C., and Owens, J.P., 1978, Preliminary cross sections of Atlantic Coastal Plain sediments of the southeastern U.S.—Cretaceous sediments along the South Carolina Coastal Margin: U.S. Geological Survey Miscellaneous Field Studies Map MF–1015–A, 2 sheets.

Gohn, G.S., Hazel, J.E., Bybell, L.M., and Edwards, L.E., 1983, The Fishburne Formation (Lower Eocene), a newly defined subsurface unit in the South Carolina Coastal Plain: U.S. Geological Survey Bulletin 1537–C, 16 p.

Gohn, G.S., Higgins, B.B., Smith, C.C., and Owens, J.P., 1977, Lithostratigraphy of the deep corehole (Clubhouse Crossroads corehole 1) near Charleston, South Carolina, in Rankin, D.W., ed., Studies related to the Charleston, South Carolina, earthquake of 1886—A preliminary report: U.S. Geological Survey Professional Paper 1028, p. 59–70.

Gohn, G.S., Smith, C.C., Christopher, R.A., and Owens, J.P., 1980, Preliminary stratigraphic cross sections of Atlantic Coastal Plain sediments of the southeastern United States—Cretaceous sediments along the Georgia coastal margin: U.S. Geological Survey Miscellaneous Field Studies Map MF–1015–C, 2 sheets.

Gulf Coast Association of Geological Societies, 1972, Tectonic map of the Gulf Coast Region U.S.A.: American Association of Petroleum Geologists, scale 1:1,000,000, 1 sheet.

Grubb, H.F., 1984, Planning report for the Gulf Coast reg