DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY

MISCELLANEOUS GEOLOGIC INVESTIGATIONS
MAP I-587



INDEX TO GEOLOGIC MAPPING

NOTE: Numbers refer to serial numbering of published sources of information. Limits of areas are not shown where they follow state boundaries or major rivers

SOURCES OF PUBLISHED DATA

1. Bingham, J. W., and Grolier, M. J., 1965, Geologic map and sections of parts of Grant, Adams, and Franklin Counties, Washington: U.S. Geol. Survey open-file rept. Map, scale 1:60,000.
2. Bond, J. G., 1963, Geology of the Clearwater embayment: Idaho Bur. Mines and Geology Pamph. 128, 83 p., 30 illus., map, scale 1:250,000.
3. Brown, C. E., and Thayer, T. P., 1966, Geologic map of the Canyon City quadrangle, northeastern Oregon: U.S. Geol. Survey Misc. Geol. Inv. Map I-447.
4. Calkins, F. C., 1905, Geology and water resources of a portion of east-central Washington: U.S. Geol. Survey Water-Supply Paper 118, 96 p., 17 figs.
5. Capps, S. R., 1941, Faulting in western Idaho and its relation to the high placer deposits: Idaho Bur. Mines and Geology Pamph. 56, 20 p.
6. Collier, A. J., 1914, The geology and mineral resources of the John Day region: Oregon Bur. Mines and Geology, v. 1, no. 3, 47 p., map, scale 1:500,000.
7. Fisher, R. V., 1967, Early Tertiary deformation in north-central Oregon: Am. Assoc. Petroleum Geologists Bull., v. 51, no. 1, p. 111–125, 6 figs.
8. Fitzsimmons, T. P., 1949, Petrology of the southwest quarter of the Pine quadrangle, Oregon: Univ. of Washington (Seattle), Ph.D. thesis.
9. Foxworthy, B. L., 1962, Geology and ground-water resources of the Ahtanum Valley, Yakima County, Washington: U.S. Geol. Survey Water-Supply Paper 1598, 97 p., 11 figs., map, scale 1:62,500.
10. Gilluly, James, 1937, Geology and mineral resources of the Baker quadrangle, Oregon: U.S. Geol. Survey Bull. 879.
11. Gilluly, James, Reed, J. C., and Park, C. F., Jr., 1933, Some mining districts of eastern Oregon: U.S. Geol. Survey Bull. 846–A, 140 p., 29 figs.
12. Graham, C. E., 1949, Structure of the western portion of the Lewiston downwarp in southeastern Washington: Wash. State Univ. (Pullman), Master's thesis, 56 p., 4 pls., 7 figs.
13. Griggs, A. B., 1966, Reconnaissance geologic map of the west half of the Spokane quadrangle, Washington and Idaho: U.S. Geol. Survey Misc. Geol. Inv. Map I-464, scale 1:125,000.
14. Hamilton, Warren, 1962, Late Cenozoic structure of west-central Idaho: Geol. Soc. America Bull., v. 73, p. 511–516, 1 fig.
15. ———— 1963, Metamorphism in the Riggins region, western Idaho: U.S. Geol. Survey Prof. Paper 436, 95 p., 3 pls., 79 figs.
16. ———— 1963, Columbia River basalt in the Riggins quadrangle, western Idaho: U.S. Geol. Survey Bull. 1141–L, 37 p., 9 figs., map.
17. Hampton, E. R., and Brown, S. G., 1964, Geology and ground-water resources of the upper Grande Ronde River basin, Union County, Oregon: U.S. Geol. Survey Water-Supply Paper 1597.
18. Hodge, E. T., 1931, Geologic map, Madras quadrangle, Oregon: Oregon State Univ. (Corvallis), scale 1:125,000.
19. ———— 1932, Geologic map, north-central Oregon: Univ. of Oregon (Eugene), Pub. Geology Ser., v. 1, no. 5, scale 1:250,000.
20. Hogenson, G. M., 1964, Geology and ground water of the Umatilla River basin, Oregon: U.S. Geol. Survey Water-Supply Paper 1620.
21. Huntting, M. T., and others, 1961, Geologic map of Washington: Washington Div. Mines and Geology, scale 1:500,000.
22. Kinnison, H. B., and Seeva, J. E., 1963, Effects of hydraulic and geologic factors on streamflow of the Yakima River basin, Washington: U.S. Geol. Survey Water-Supply Paper 1595.
23. Kirkham, V. R. D., 1927, Orofino and Lapwai area: Idaho Bur. Mines and Geology Pamph. 24.
24. Laval, W. N., 1956, Stratigraphy and structural geology of portions of south-central Washington: Univ. of Washington (Seattle), Ph.D. thesis, 208 p.
25. Libbey, F. W., 1943, Some mineral deposits in the area surrounding the junction of the Snake and Imnaha Rivers in Oregon: Oregon Dept. Geology and Mineral Industries GMI Short Paper 11, 17 p., 5 figs.
26. Lukanuski, J. N., 1963, Geology of part of the Mitchell quadrangle, Jefferson and Crook Counties, Oregon: Oregon State Univ. (Corvallis), Master's thesis, 90 p., 22 figs.
27. Lupher, R. L., and Warren, W. C., 1942, The Ascitn stage of the Snake River canyon near Lewiston, Idaho: Jour. Geology, v. 50, no. 7, p. 866–881, 4 figs.
28. Lystrom, D. J., Ness, W. L., and Hampton, E. R., 1967, Ground water of Baker Valley, Baker County, Oregon: U.S. Geol. Survey Hydrol. Inv. Atlas HA–242.
29. Maide, H. E., 1965, Guidebook for field conference E, northern and middle Rocky Mountains: Internat. Assoc. Quaternary Research, 7th Cong., map.
30. Newcomb, R. C., 1965, Geology and ground-water resources of the Walla Walla River basin, Washington-Oregon: Washington Div. Water Resources Water-Supply Bull. 21, 151 p., 19 figs., map.
31. ———— 1967, The Dalles-Umatilla syncline, Oregon and Washington: U.S. Geol. Survey Prof. Paper 575–B, p. B88–B93, 3 figs.
32. ———— 1969, Effect of tectonic structure on the occurrence of ground water in the basalt of the Columbia River Group of the Dalles area, Oregon and Washington: U.S.Geol.Survey Prof. Paper 383–C, 33 p., 18 figs., map.
33. Newton, V. C., and Corcoran, R. E., 1963, Petroleum geology of the western Snake River Basin, Oregon-Idaho: Oregon Dept. Geology and Mineral Industries, Oil and Gas Inv., no. 1, 67 p., 4 pls., 2 figs.
34. Pardee, J. T., 1941, Preliminary geologic map of the Sumpter quadrangle, Oregon: Oregon Dept. Geology and Mineral Industries.
35. Patterson, P. V., 1966, Geologic map, in watershed work plan, Wolf Creek, Union County, Oregon: U.S. Dept. of Agriculture, Soil Conserv. Service.
36. Pebbles, J. J., 1962, Engineering geology of the Cartwright Canyon quadrangle: Idaho Bur. Mines and Geology Pamph. 127, 69 p., 31 figs.
37. Peck, D. L., 1964, Geologic reconnaissance of the Antelope-Ashwood area, north-central Oregon: U.S. Geol. Survey Bull. 1161–D, 26 p., 9 figs., map.
38. Pigg, J. H., Jr., 1961, The lower Tertiary sedimentary rocks in the Pilot Rock and Heppner areas, Oregon: Univ. of Oregon (Eugene), Master's thesis, 67 p., 22 figs.
39. Prostka, H. J., 1962, Geology of the Sparta quadrangle, Oregon: Oregon Dept. Geology and Mineral Industries Geol. Map Ser. 1, scale 1:62,500.
40. ———— 1967, Preliminary geologic map of the Durkee quadrangle, Oregon: Oregon Dept. Geology and Mineral Industries Geol. Map Ser. GM–33, scale 1:62,500, 4 sht.
41. Ptacek, A. D., 1965, Cenozoic geology of the Seven Devils Mountains, Idaho: Univ. of Washington (Seattle), Ph.D. thesis, 96 p., 2 pls., 38 figs.
42. Robinson, J. W., and Price, Don, 1963, Ground water in the Prineville area, Crook County, Oregon: U.S. Geol. Survey Water-Supply Paper 1619–P.
43. Ross, C. P., 1938, The geology of the Wallowa Mountains: Oregon Dept. Mines and Geology Bull. 3.
44. Ross, C. P., and Forrester, J. D., 1959, Geologic map of the State of Idaho: U.S. Geol. Survey and Idaho Bur. Mines and Geology, scale 1:500,000.
45. Russell, I. C., 1897, A reconnaissance in southeastern Washington: U.S. Geol. Survey Water-Supply Paper 4.
46. ———— 1901, Geology and water resources of Nez Perce County, Idaho, Part I: U.S. Geol. Survey Water-Supply Paper 53.
47. ———— 1901, Geology and water resources of Nez Perce County, Idaho, Part II: U.S. Geol. Survey Water-Supply Paper 54.
48. Savage, C. N., 1961, Geology and mineral resources of Gem and Payette Counties (Idaho): Idaho Bur. Mines and Geology County Rept. 4, 49 p., 13 figs.
49. ———— 1965, Economic geology of carbonate rocks adjacent to the Snake River south of Lewiston, Idaho: Idaho Bur. Mines and Geology Mineral Resources Rept. 10, 26 p., 2 pls., 8 figs.
50. Schmidt, D. L., 1964, Reconnaissance petrographic cross section of the Idaho batholith in Adams and Valley Counties, Idaho: U.S. Geol. Survey Bull. 1181–G.
51. Sheppard, R. A., 1967, Geology of the Simcoe Mountains volcanic area, Washington: Washington Div. Mines and Geology Geol. Map GM–3, scale 1:250,000.
52. Smeden, H. W., 1959, Geology of part of the northern Wallowa Mountains: Univ. of Washington (Seattle), Ph.D. thesis.
53. Smith, G. O., 1901, Geology and water resources of a portion of Yakima County, Washington: U.S. Geol. Survey Water-Supply Paper 55.
54. ———— 1903, Geologic folio of the Ellensburg quadrangle, Washington: U.S. Geol. Survey Geol. Folio 86, 7 p., 3 maps, scale 1:125,000.
55. Smith, W. D., and Allen, J. E., 1941, Geology and physiography of the northern Wallowa Mountains, Oregon: Oregon Dept. Geology and Mineral Industries Bull. 12, 64 p., 16 figs.
56. Stearns, H. T., and Anderson, A. L., 1966, Geology of the Oxbow and Snake River near Homestead, Oregon: Idaho Bur. Mines and Geology Pamph. 136, 23 p., 10 figs.
57. Swanson, D. A., 1966, Tieton volcano, a Miocene eruptive center in the southern Cascade Mountains, Washington: Geol. Soc. America Bull., v. 77, no. 11, p. 1293–1314, 7 figs.
58. ———— 1967, Yakima basalt of the Tieton River area, south-central Washington: Geol. Soc. America Bull. 78, no. 9, p. 1077–1110, 1 pl., 3 figs.
59. Taylor, E. M., 1960, Geology of the Clarno Basin, Mitchell quadrangle, Oregon: Oregon State Univ. (Corvallis), Master's thesis, 173 p., 2 pls., 24 figs.
60. Thayer, T. P., 1937, Some relations of later Tertiary volcanology and structure in eastern Oregon: Internat. Geol. Cong., 20th, Mexico, 1956, Vulcanologia del Cenozoico, sec. 1, v. 1, p. 231–245.
61. Trauger, F. D., 1950, Ground-water resources of Baker Valley, Baker County, Oregon: U.S. Geol. Survey open-file rept., 100 p., 12 figs., 3 maps, scale 1:62,500.
62. Valller, T. L., 1967, The geology of part of the Snake River canyon and adjacent areas in northeastern Oregon and western Idaho: Oregon State Univ. (Corvallis), Ph.D. thesis, 267 p., 3 pls., 37 figs.
63. Wagner, N. S., 1958, Important rock units of northeastern Oregon: Oregon Dept. Geology and Mineral Industries, The Ore Bin, v. 20, no. 7, p. 63–68, map.
64. Wagner, W. R., 1945, A geologic reconnaissance between the Snake and Salmon Rivers north of Riggins, Idaho: Idaho Bur. Mines and Geology Pamph. 74.
65. Walcott, W. E., and Neff, G. E., 1950, Report on the flow structure of the basalt in the Coulee City-Long Lake-Soap Lake area, Columbia Basin project: U.S. Bur. Reclamation Rept. duplicated, 11 p., 21 illus., map, scale 1:36,000.
66. Walters, K. L., and Grolier, M. J., 1960, Geology and ground-water resources of the Columbia Basin project area, Washington, Volume I: Washington Div. Water Resources Water-Supply Bull. 8, 518 p., 28 figs.
67. Waring, G. A., 1913, Geology and water resources of a portion of south-central Washington: U.S. Geol. Survey Water-Supply Paper 316.
68. Waters, A. C., 1965, Geomorphology of south-central Washington, illustrated by the Yakima East quadrangle: Geol. Soc. America Bull., v. 66, p. 663–684, 2 pls., 3 figs., map, scale 1:62,500.
69. Wells, F. G., 1961, Geologic map of Oregon west of the 121st meridian: Oregon Dept. Geology and Mineral Industries, scale 1:500,000.
70. Williams, Howel, 1957, A geologic map of the Bend quadrangle, Oregon (1:125,000) and a reconnaissance geologic map of the central portion of the high Cascade Mountains (1:250,000): Oregon Dept. Geology and Mineral Industries.

SOURCES OF UNPUBLISHED DATA

Bishop, Donald T., Northwestern Idaho.
Brooks, Howard C., Mineral quadrangle, Oregon-Idaho.
Deacon, Robert J., Central Washington.
Hamilton, Warren, West-central Idaho.
Hampton, E. R., North-central Oregon.
Hogenson, G. M., Yakima Valley.
Neff, George E., Central Washington.
Norvitch, Ralph F., North-central Idaho.
Powers, Howard A., West-central Idaho.
Robinson, Paul T., Ochoco Mountains area.
Robison, James H., North-central Oregon.
Ross, Sylvia H., Northwestern Idaho.
Swanson, Donald A., Central Oregon.
Wagner, Norman S., Northeastern Oregon.

STRUCTURES IN RELATION TO GROUND WATER

This map was compiled to show the areas in which geologic structure favors the accumulation and storage of ground water in the basalt of the Columbia River Group of Miocene and Pliocene age. Published and unpublished sources of data were used, supplemented in places by field reconnaissance. In addition to folds and faults in the basalt, the map includes a few structures in rocks older than the basalt, where those structures affect the position and water-bearing capabilities of the nearby basalt.

Most of the known folds that show structural relief of more than 100 feet and faults that have more than 50 feet displacement are included. In zones of close and complex deformation, minor structures have been omitted for lack of space or have been generalized. Except in the area south of the Ochoco anticline, where faults and fractures were plotted from aerial photographs without ground checking, nondisplacement breaks of the extension-fracture type, sometimes called regional joints, have been omitted.

The structures that displace the basalt are geologically youthful, Pliocene and Quaternary in age. Most of the anticlines are tectonic mountains, and the synclines are tectonic valleys.

Many of the folds, especially where they are distant from other structures, are smoothly symmetrical in cross section, but within areas of major deformation some of the folds contain all degrees of asymmetry. Within the folds of greatest asymmetry, the shorter limb is turned up steeply, or even overturned and thrust faulted in places. The direction of the overasteepening is consistent in some long asymmetrical folds, such as northward in the Horse Heaven and Aldrich Mountain anticlines and southward in the Cleman Mountain and Tygh Ridge anticlines; but it changes from side to side along other folds, such as the Columbia Hills and the Umtanum anticlines.

Most of the long, prominent faults are normal and dip 60 to 85 degrees. A few high-angle reverse faults have been identified, and low-angle thrust faults are common locally in and near the steeply folded limbs of some of the anticlines. Because of the rock slump and talus cover and the shattered condition of the basalt along those zones of rupture, thrust faults have been observed in only a few places. Strike-slip faults have been observed only along lateral boundaries of large blocks that end in an overstepened limb of a large anticline.

The vertical displacement of the basalt in the folds ranges approximately from the 100-foot selected minimum to about 2,000 feet. Many of the larger folds have amplitudes of 500 to 2,000 feet. The displacement on faults shown on the map ranges from 50 feet to about 3,000 feet. This greatest displacement occurs along some of the larger faults bounding the Grande Ronde valley, the John Day fault, and the long faults in central Idaho. Most of the faults shown have displacements of less than 500 feet. The horizontal displacement along the few thrust faults identified and the few strike-slip faults shown is less than a few hundred feet.

The structures fall within five main groupings: (1) Basin-and-range type of block faulting, most of which fades northward into longer open-folded structures (Thayer, 1957) or dies out in the southern limb of the (2) Blue Mountain anticline; (3) the long north-south tilt-block faults along the west side of the Idaho batholith; (4) the Cascadian structures, of which the Yakima Ridges are the main elements; and (5) the band of almost horizontal basalt extending along the west side of the northern Rocky Mountains from Grangeville to the Okanogan Highlands (interrupted only by the Lewiston downwarp).

EXPLANATION

Pliocene, Pleistocene, and Holocene	QTs	Sedimentary deposits — *Younger than basalt of the Columbia River Group. Includes continental sedimentary deposits principally the Dalles Formation; Ellensburg Formation and Mascall Formation except for the lowest layers which are interstratified with the basalt; Madras Formation; Rattlesnake Formation; Idaho Group; and Ringold Formation, as well as some glacial, glaciofluvial eolian, alluvial and collurial deposits. Small amounts of volcanic rocks are intercalated in some of these sedimentary deposits. Deposits that are predominantly of eolian origin are omitted*
	QTv	Volcanic rocks — *Younger than basalt of the Columbia River Group. Basaltic; andesitic and dacitic lavas and cinder cone deposits largely belonging to the Cascade lavas and extensions therefrom but includes small amounts of lava distant from the Cascade Range*
Pliocene or Pleistocene	QTi	Intrusive rocks — *Andesitic porphyry intruded into the basalt of the Columbia River Group in the Columbia Gorge. Equivalent to the feeder conduits for some of the Cascade lavas*
Miocene and Pliocene	Tcr	Basalt of the Columbia River Group — *Accordantly layered flows of dense basaltic lava. Total thickness ranges from featheredge to more than 5,000 feet; individual flows range generally from 10 to 200 feet thick and average 80 feet. Includes some layers of the lower part of the Ellensburg Formation in the Yakima Valley, some interlayered parts of the Latah Formation at the northeastern edge, the Pogetta Formation at the southeastern edge, and small parts of the Mascall Formation at the southern edge of the main area of the basalt. Includes small areas of some younger lavas such as the platy andesites of the Baker and La Grande Valleys, (Gilluly, 1937; Hampton and Brown, 1964) not yet differentiated*
	Ts	Strawberry Volcanics — *Basalt, andesite, and rhyolite flows and breccia in part contemporaneous with, and interfingering with, basalt of the Columbia River Group in the John Day River basin. Includes some volcanic rocks younger than the basalt of the Columbia River Group proper near the Pasalina basin. Extends beyond boundaries of this mapping*
	Tpcr	Rocks older than basalt of the Columbia River Group — *Rocks of various types older than the basalt. Includes small areas of sedimentary deposits, such as the Latah Formation, which are partly contemporaneous with the basalt. Shown only where in proximity to the basalt of the Columbia River Group*

TERTIARY AND QUATERNARY
TERTIARY OR QUATERNARY
TERTIARY
PRECAMBRIAN TO TERTIARY

GEOLOGIC SYMBOLS

- Contact — *Dashed where approximately located*
- Fault — *Dashed where approximately located, dotted where concealed. Parallel arrows show direction of strike slip. Bar and ball on downthrown side*
- Thrust fault — *Sawteeth on upper plate*
- Axis of anticline showing direction of plunge — *Dashed where approximately located, dotted where concealed*
- Dome
- Axis of syncline showing direction of plunge — *Dashed where approximately located, dotted where concealed*
- Axis of monocline and direction of dip from axis

SCALE 1:500 000
10 0 10 20 30 MILES
10 0 10 20 30 KILOMETERS

Base from U.S. Geological Survey
State base maps, 1961–65

TECTONIC STRUCTURE OF THE MAIN PART OF THE BASALT OF THE COLUMBIA RIVER GROUP WASHINGTON, OREGON, AND IDAHO

By
R. C. Newcomb

1970

United States (Columbia River basin). Structure, 1:500,000. 1970.
cop. 2.

M(200)
I-587
cop. 2

USGS LIBRARY
JUL 20 1970
RESTON, VIRGINIA

G3700
.S9W3
1970
N38
I-587

USGS LIBRARY - RESTON
3 1818 00175211 0

INTERIOR—GEOLOGICAL SURVEY, WASHINGTON, D.C.—1970—80063
For sale by U. S. Geological Survey, price $1.00