



CORRELATION OF MAP UNITS

Qa	Quaternary and Tertiary	Quaternary and Tertiary
OTg	Older Tertiary and Cretaceous	Tertiary and Cretaceous
INTRO	INTRO	INTRO
SEDI	SEDI	SEDI
INTR	INTR	INTR
MTA	MTA	MTA
MTM	MTM	MTM
MTN	MTN	MTN
MTX	MTX	MTX
MTY	MTY	MTY
MTZ	MTZ	MTZ
MTA	MTA	MTA
MTB	MTB	MTB
MTC	MTC	MTC
MTD	MTD	MTD
MTX	MTX	MTX
MTY	MTY	MTY
MTZ	MTZ	MTZ

DESCRIPTION OF MAP UNITS

- Qa** Alluvium, colluvium, and glacial deposits (Holocene and late Pleistocene)—In northwest part of area (Stromboli Lake quadrangle), locally includes Troublesome Formation (Miocene and Oligocene) adjacent to Proterozoic rock units.
- OTg** Gravels (Quaternary and Tertiary)—Rounded to subangular pebbles, and boulders derived from metamorphic and igneous rocks mostly to the west of the map area. Patterned areas consist of debris derived almost entirely from Precambrian rocks and found on higher slopes above modern drainage channels.
- INTRO** Intrusive rocks—May include minor amounts of mafic rocks, such as porphyrite.
- SEDI** Sedimentary rocks—May include the Fountain Formation (Permian and Pennsylvanian) (Triassic and Permian) and Lyons Sandstone (Permian). Fountain Formation, predominantly arkosic conglomerate and moderately coarse grained sandstone. Lyons Formation, red and light-green calcareous sandstone; Lyons Sandstone, red and pink fine-grained to very fine grained sandstone.
- INTR** Intrusive rocks—May include minor amounts of mafic rocks, such as porphyrite.
- MTA** Monzonite, granodiorite, and syenite—May include minor amounts of mafic rocks, such as porphyrite.
- MTB** Quartz monzonite and monzonite—Mostly massive, gray to buff, fine to medium-grained quartz monzonite or monzonite displaying aligned subparallel foliation and aligned biotite laths, especially along borders of plutons. U-Pb zircon age 1,400 m.y. (Alenikoff and others, 1990).
- MTC** Monzonite to granodiorite facies—Medium to coarse-grained, grayish-white and black, foliated monzonite to granodiorite. Exposed locally in Mount Evans batholith. Contains 6-8 percent magnetite. U-Pb zircon age of 1,443 m.y. (Alenikoff and others, 1990).
- MTD** Inclusions—Pretise inclusions mostly of biotite gneiss in Silver Plume Granite.
- MTX** Quartz monzonite (Early Proterozoic)—Light gray to light tan, leucocratic, and fine to medium grained. Includes Twin Spring Quartz Monzonite of the Eldorado Springs area. Field evidence suggests Twin Spring is both younger and the same age as the Boulder Creek Granodiorite. Patterned where unit contains numerous biotite gneisses.
- MTY** Granodiorite (Early Proterozoic)—Gray to pinkish-gray, medium to coarse-grained gneissic granodiorite; weathers darker pinkish gray. Exposed locally at Mt. Morrison and in Clear Creek and Turkey Creek canyons. Logically fits the 1,700 m.y. isochron of the Boulder Creek Granodiorite (C.E. Hedge, oral communication, 1972).
- MTZ** Boulder Creek Granodiorite and associated rocks of the Roubidoux Pluton (Early Proterozoic)—Medium grayish-white and black, medium to very coarse grained, locally porphyritic, predominantly granodiorite but includes lenses and layers of gabbro, hornblende diorite, hornblende, and quartz diorite. Rb-Sr determinations suggest the Boulder Creek is about 1,700 m.y. old (Peterson and others, 1968; Reed and others, 1987).
- Xg** Gabbro and related rocks (Early Proterozoic)—Dark gray to pinkish-gray, black and white, massive, coarse-grained to very coarse grained rock ranging in composition from melaphyre to quartz diorite; contains intermediate plagioclase and orthopyroxene and clinopyroxene.
- Xgd** Hornblende diorite, quartz-bearing hornblende diorite, quartz diorite, and hornblende (Early Proterozoic)—Gray to black, medium to fine-grained, gray to black hornblende-bearing rocks. Exposed in small plutons, lenses, pods, and dikes.
- EARLY PROTEROZOIC METAMORPHIC ROCKS** (The nomenclature and modal chemical and quantitative analyses of the metamorphic rocks are given in Gable (1996)).
- Amphibolite**—Dark greenish-gray to black, fine to medium-grained, nonlayered to poorly layered, usually to strongly foliated rock, composed mostly of hornblende and plagioclase.
- EARLY PROTEROZOIC METASEDIMENTARY AND METAVOLCANIC ROCKS**
- Xq** Quartzite and quartz gneiss—Quartzite is white, gray, or pinkish or purplish gray, medium to coarse grained, and locally conglomeratic. Interbedded with quartzite schist and some calc-silicate gneiss lenses. Quartzite along Coal Creek and in Eldorado Springs area is interbedded with conglomeratic quartzite. At some localities, foliated quartz gneiss is conspicuously layered and contains variable amounts of garnet, magnetite-ilmenite, and epidote.
- Zones of closely spaced shear zones**
- Boundary of Idaho Springs-Robson shear zone**
- Folds**—Showing approximate trace of axial plane and direction of plunge. Dotted where concealed; quartered where inferred.
- Antiform**
- Overturned antiform**
- Synform**
- Overturned synform**
- Small-scale linear and planar features**
- Bearing and plunge of axes of minor folds**—Symbols show type of minor folds
- Strike and dip of foliation**
- Inclined**
- Vertical**
- Bearing and plunge of lineation**—Lineations defined by aligned minerals, mineral streaks, and fold axes or intersection of planar elements. Symbols may be combined with foliation symbol.

- Xqs** Quartz-biotite-muscovite schist—Silver gray to dark gray, fine to medium-grained, micaceous schist locally containing north-south trending cordierite, cordierite, and garnet and small amounts of staurolite and sillimanite. Mapped along mountain front in Rabbit Butte area.
- Xf** Feldspar-rich gneiss—Light gray, leucocratic, fine to medium-grained, microcline-plagioclase-quartz-biotite gneiss. Contains cordierite, foliated, granitic in appearance. Locally garniferous and albitically interbedded with conformable thin to thick layers and lenses of hornblende gneiss, amphibolite, biotite gneiss, and locally calc-silicate rock.
- Xh** Interlayered feldspar-rich gneiss and hornblende gneiss—Feldspar-rich gneiss may contain hornblende and is generally darker than the feldspar-rich gneiss unit (Xf). Interlayered hornblende gneiss is black, white, and dark gray or greenish gray and similar in composition to the hornblende gneiss in hornblende gneiss unit (Xgd). Layers and lenses of biotite gneiss, amphibolite, calc-silicate rock, and biotite-quartz-plagioclase gneiss make up 10-30 percent of unit.
- Xgh** Hornblende gneiss and amphibolite—Black, black and white, and dark gray or greenish gray, fine to medium grained, and locally weak to strong foliation. Consists of hornblende-plagioclase-quartz gneiss or hornblende-clinopyroxene-quartz-biotite gneiss in places interbedded with amphibolite, calc-silicate gneiss, biotite-quartz-plagioclase gneiss, and minor quartzite.
- Xhs** Interlayered hornblende gneiss, calc-silicate gneiss, and amphibolite—Black and white, light gray, greenish gray, and silty gray, fine to medium grained, layered; has moderate to good compositional layering and moderate to weak foliation. More leucocratic than hornblende gneiss (Xgh) because individual gneiss layers are thicker and contain more silts, calcite, and quartz-bearing layers. Contains minor cordierite-biotite gneiss and biotite-quartz-plagioclase gneiss.
- Xhgt** Biotite gneiss—Dark gray, fine to medium-grained, foliated, biotite-quartz-plagioclase gneiss, generally leucocratic, has weak to strong foliation, and interlayered with hornblende gneiss, calc-silicate gneiss, and sillimanite-biotite-quartz gneiss. Patterned areas consist of fine lenses and layers of granite gneiss.
- Xma** Biotite gneiss and amphibolite—Black and white, light gray, greenish gray, and silty gray, fine to medium grained, biotite-quartz-plagioclase gneiss interlayered and intergraded; includes protomylonitic lenses and layers of amphibolite. Locally contains lenses and layers of hornblende-biotite-plagioclase gneiss, calc-silicate gneiss, quartzite, and pegmatite and layers of granite gneiss and pegmatite.
- Xmb** Biotite-plagioclase gneiss—Interbedded and interlayered biotite-quartz-plagioclase gneiss and sillimanite-biotite-quartz-plagioclase gneiss, has protomylonitic lenses and layers of granite gneiss and pegmatite, and at some localities contains small lenses and layers of calc-silicate gneiss, amphibolite, and quartzite. Contains much more amphibole than the biotite gneiss and amphibolite unit (Xma).
- Xng** Garnet-biotite gneiss—Exists as individual layers that grade into and are interlayered with minor units of biotite-quartz-plagioclase gneiss, garnet-sillimanite-biotite gneiss, and cordierite-garnet-sillimanite-biotite gneiss.
- Xgn** Quartz-plagioclase gneiss—Leucocratic, poorly foliated, and contains thin layers of biotite gneiss, feldspar-rich gneiss, quartzite, hornblende gneiss, and amphibolite. Only mapped along mountain front south of Morrison.
- Xgnc** Cordierite-garnet-gneiss-biotite gneiss—Dark gray to black, medium to coarse grained, foliated to massive. Interlayered with thinner layers of cordierite-biotite gneiss, garnet-biotite gneiss, and cordierite-garnet-sillimanite-biotite gneiss. K-feldspar is not present or is rare in these rocks.
- Xgnc** Cordierite-garnet-sillimanite-biotite gneiss ± K-feldspar ± plagioclase—Generally gray to very dark gray, fine to medium grained, foliated to massive. May be finely grained with one or more of the following types of gneiss: garnet-sillimanite-biotite, cordierite-biotite, garnet-biotite, cordierite-garnet-biotite, cordierite-sillimanite-biotite, magnetite-sillimanite-biotite, and cordierite-magnetite-sillimanite-biotite.
- Xgnc** Sillimanite-biotite gneiss—Banded light gray, very dark gray, or black at some localities, interbedded with biotite-quartz-plagioclase gneiss and layers and lenses of amphibolite, biotite gneiss, calc-silicate gneiss, and garnet-biotite gneiss. South of the Idaho Springs-Robson shear zone, primary muscovite and trace amounts of cordierite present in unit. Pattern shows area that contains protomylonitic lenses and layers of Silver Plume Granite (Ysp).
- Xgnc** Cordierite-garnet-sillimanite-biotite gneiss and minor magnetite-sillimanite-biotite gneiss—Light gray to nearly black, fine to medium grained, compositionally layered. In places layers discontinuous due to stringers and clots of pegmatite and knots of cordierite, sillimanite, biotite, and magnetite. Both rock types gradational into cordierite-garnet-sillimanite-biotite gneiss (Xgnc) and sillimanite-biotite gneiss (Xgnc).
- Xcb** Cordierite-sillimanite-biotite gneiss—Light gray to nearly black, fine to medium grained, compositionally layered. Mapped only in Evergreen-Squaw Pass area.
- Xncs** Calc-silicate gneiss—Variable gray, green, white, or black and fine to coarse grained. Color depends on the type and quantity of minerals present, such as hornblende, diopside, biotite, calcic amphibole, scapolite, and quartz. Foliation poor but compositional layering good.
- Xi** Rutile-bearing, sillimanite-quartz gneiss and biotite-bearing, sillimanite-quartz gneiss and biotite-bearing, sillimanite-quartz gneiss—Exists as thin lenses in the Evergreen-Squaw Pass area.
- Xsu** Sillimanite-quartz gneiss containing accessory rutile lenses—Light gray to gray, fine to medium grained, and foliated. Exists only as thin lenses 15 cm to 30 m wide in the Evergreen and Squaw Pass quadrangles; most are of limited extent but one is 6 km long.
- Xsf** Augen gneiss—Fine-grained, sheeted gneiss containing small, pink feldspar xenocrysts in a well-foliated biotite matrix. Mapped only in Rabbit Butte area.

REFERENCES CITED

Alenikoff, J.N., DeWitt, Ed., Reed, J.C., Jr., and Walker, Marianne, 1990, The Mount Evans batholith—an anomalous 1.4 Ga pluton in the Front Range, Colorado [abs.]. Geological Society of America Abstracts with Programs, v. 22, no. 6, p. 1.

Braddock, W.A., 1969, Geology of the Empire quadrangle, Grand, Gilpin, and Clear Creek Counties, Colorado. U.S. Geological Survey Professional Paper 616, 56 p.

Bryant, B.H., 1974a, Reconnaissance geologic map of the Conifer quadrangle, Jefferson County, Colorado. U.S. Geological Survey Miscellaneous Field Studies Map MF-597, scale 1:24,000.

—, 1974b, Reconnaissance geologic map of the Pine quadrangle, Jefferson County, Colorado. U.S. Geological Survey Miscellaneous Field Studies Map MF-598, scale 1:24,000.

—, 1976, Reconnaissance geologic map of the Bailey quadrangle, Jefferson and Park Counties, Colorado. U.S. Geological Survey Miscellaneous Field Studies Map MF-616, scale 1:24,000.

Bryant, B.H., Miller, R.D., and Scott, G.R., 1973, Geologic map of the Indian Hills quadrangle, Jefferson County, Colorado. U.S. Geological Survey Geologic Quadrangle Map GQ-1073, scale 1:24,000.

Cable, D.J., 1968, Geology of the crystalline rocks in the western part of the Morrison quadrangle, Jefferson County, Colorado. U.S. Geological Survey Bulletin 1221-E, 45 p.

—, 1969, Geologic map of the Nederland quadrangle, Boulder and Gilpin Counties, Colorado. U.S. Geological Survey Geologic Quadrangle Map GQ-853, scale 1:24,000.

—, 1972, Geologic map of the Turagan quadrangle, Boulder, Gilpin, and Jefferson Counties, Colorado. U.S. Geological Survey Geologic Quadrangle Map GQ-978, scale 1:24,000.

—, 1980, Geologic map of the Gold Hill quadrangle, Boulder County, Colorado. U.S. Geological Survey Geologic Quadrangle Map GQ-1265, scale 1:24,000.

—, 1996, Metamorphism, metamorphism, and provenance of the early Proterozoic metamorphic rocks of the central Front Range, Colorado. U.S. Geological Survey Open-File Report 96-222, 2 sheets, map scale 1:100,000.

Cable, D.J., and Madole, R.F., 1976, Geologic map of the Ward quadrangle, Boulder County, Colorado. U.S. Geological Survey Geologic Quadrangle Map GQ-1277, scale 1:24,000.

Cable, D.J., and Sims, P.K., 1969, Geology and regional metamorphism of some high-grade cordierite gneisses, Front Range, Colorado. Geological Society of America Special Paper 128, 97 p.

Hawley, C.C., and Moore, F.B., 1967, Geology and ore deposits of the Lawson-Dumont-Fall River district, Clear Creek County, Colorado. U.S. Geological Survey Bulletin 1231, 92 p.

Hedge, C.E., 1970, Whole-rock Rb-Sr age of the Pike's Peak batholith, Colorado. U.S. Geological Survey Professional Paper 700-B, p. B86-B89.

Lovings, T.S., and Goodland, E.N., 1950 (1951), Geology and ore deposits of the Front Range, Colorado. U.S. Geological Survey Professional Paper 222, 317 p.

Marsh, S.P., and Sheridan, D.M., 1976, Rutile in Precambrian sillimanite-quartz gneiss and related rocks, east-central Front Range, Colorado. U.S. Geological Survey Professional Paper 959-G, 17 p.

Moench, R.H., 1964, Geology of Precambrian rocks, Idaho Springs district, Colorado. U.S. Geological Survey Bulletin 1182-A, 70 p.

Moench, R.H., Harrison, J.E., and Sims, P.K., 1962, Precambrian folding in the Idaho Springs-Central City area, Front Range, Colorado. Geological Society of America Bulletin, v. 73, p. 35-58.

Panmon, R.C., 1980, Mineral resources of the Indian Peaks study area, Boulder and Grand Counties, Colorado, with a section on interpretation of aeromagnetic data, by Gordon Johnson. U.S. Geological Survey Bulletin 1463, 109 p.

Peterson, Z.E., Hedge, C.E., and Braddock, W.A., 1968, Age of Precambrian events in the northeastern Front Range, Colorado. Journal of Geophysical Research, v. 73, no. 6, p. 2277-2296.

Peterson, W.L., 1964, Geology of the Flatts Canyon quadrangle, Colorado. U.S. Geological Survey Bulletin 1181-C, 23 p.

Reed, J.C., Jr., Beckert, M.E., Pardo, W.R., Palterer, J.S., and Alenikoff, J.N., 1987, Evolution of the Early Proterozoic Colorado Province—constraints from U-Pb geochronology. Geology, v. 15, no. 9, p. 861-865.

Schroeder, D.A., 1995, Geologic map of the Strawberry Lake quadrangle, Grand County, Colorado. U.S. Geological Survey Geologic Quadrangle Map GQ-1764, scale 1:24,000.

Scott, G.R., 1963, Bedrock geology of the Kessler quadrangle, Colorado. U.S. Geological Survey Professional Paper 421-B, p. 71-125.

Sheridan, D.M., and Marsh, S.P., 1976, Geologic map of the Squaw Pass quadrangle, Clear Creek, Jefferson, and Gilpin Counties, Colorado. U.S. Geological Survey Geologic Quadrangle Map GQ-1337, scale 1:24,000.

Sheridan, D.M., Maxwell, C.H., and Abbot, A.L., 1967, Geology and uranium deposits of the Rabbit Butte district, Jefferson County, Colorado, with a section on Paleozoic and younger sedimentary rocks, by Richard Van Horn. U.S. Geological Survey Professional Paper 520, 121 p.

Sheridan, D.M., Reed, J.C., Jr., and Bryant, B.H., 1972 (1973), Geologic map of the Empire quadrangle, Jefferson County, Colorado. U.S. Geological Survey Miscellaneous Geologic Investigations Series Map I-786-A, scale 1:24,000.

Sims, P.K., and Cable, D.J., 1967, Petrology and structure of Precambrian rocks, Central City quadrangle, Colorado. U.S. Geological Survey Professional Paper 564-E, 56 p.

Taylor, R.B., 1976, Geologic map of the Black Hawk quadrangle, Gilpin, Jefferson, and Clear Creek Counties, Colorado. U.S. Geological Survey Geologic Quadrangle Map GQ-1248, scale 1:24,000.

Twesto, Ogden, 1987, Rock units in the Precambrian basement in Colorado. U.S. Geological Survey Professional Paper 1321-A, p. A1-A54.

Van Horn, Richard, 1972, Surficial and bedrock geologic map of the Golden quadrangle, Jefferson County, Colorado. U.S. Geological Survey Folio of the Golden quadrangle, Colorado. Map I-761-A, scale 1:24,000.

Wells, J.D., 1967, Geology of the Eldorado Springs quadrangle, Boulder and Jefferson Counties, Colorado. U.S. Geological Survey Bulletin 1221-D, 85 p.

Wrucke, C.T., and Wilson, R.F., 1967, Geologic map of the Boulder quadrangle, Boulder County, Colorado. U.S. Geological Survey Open-File report, scale 1:24,000.

Young, E.J., 1991, Geologic map of the East Portal quadrangle, Boulder, Gilpin, and Grand Counties, Colorado. U.S. Geological Survey Miscellaneous Investigations Series Map I-2212, scale 1:24,000.

Base from U.S. Geological Survey, Denver West, 1983, Estes Park, 1:50,000.

Universal Transverse Mercator projection

1983 North American datum

APPROXIMATE MAIN SECTION LINE

SCALE 1:100,000

0 1 2 3 4 5 MILES

0 1 2 3 4 5 KILOMETERS

CONTOUR INTERVAL 50 METERS

NATIONAL GEODETIC VERTICAL DATUM OF 1989

MAP LOCATION

CONVERSION FACTORS

Multiply	By	To obtain
centimeters (cm)	0.3937	inches (in.)
meters (m)	3.281	feet (ft)
kilometers (km)	0.6214	miles (mi)

Geology compiled 1976-84

Revised in 1995 with assistance of William G. Johnson, Jr. Map digitized by Springfield and Springfield, Digital cartography by Dennis Wiley and David Dumencous. Edited by F.C. Brunsten.

GEOLOGIC MAP OF THE PROTEROZOIC ROCKS OF THE CENTRAL FRONT RANGE, COLORADO

Compiled by
Dolores J. Gable
2000

- SOURCES OF GEOLOGIC DATA**
- | | |
|--------------------------------|--------------------------------|
| 1. Panmon (1980) | 17. Braddock (1969) |
| 2. Panmon (1980) | 18. Marsh and others (1962) |
| 3. Panmon (1980) | 19. Marsh and Moore (1967) |
| 4. Cable (1980) | 20. Sheridan and Marsh |
| 5. Cable and Madole (1976) | 21. Sheridan and others |
| 6. Panmon (1980) | 22. Sheridan and others (1973) |
| 7. Panmon (1980) | 23. Bryant and others (1974) |
| 8. Schneider (1995) | 24. Bryant (1976) |
| 9. Young (1991) | 25. Bryant (1976) |
| 10. Cable (1969) | 26. Bryant (1976) |
| 11. Gable (1972) | 27. Petersen (1964) |
| 12. Wells (1967) | 28. Scott (1963) |
| 13. Van Horn (1972) | |
| 14. Sheridan and others (1967) | |
| 15. Taylor (1976) | |
| 16. Sims and Gable (1967) | |
- Any use of trade names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Geological Survey.
- For sale by U.S. Geological Survey Information Services, Reston, Virginia 20192. Catalog # I-2465. Manuscript accepted for publication April 24, 1997.
- This report is available as PDF file at <http://www.gsa.gov>
- ISBN 0-607-08372-3