


Mercator projection
Geodetic Reference System 1980; North American Datum 1983
Longitude of central meridian 70°19' W; latitude of true scale 41°39' N.
False easting 0 m; false northing 0 m
This map is not intended for navigational purposes.



Shoreline data were extracted from the National Oceanic and Atmospheric Administration Medium Resolution Digital Vector Shoreline (see text).

DISCUSSION
Introduction

This map shows the sea floor in shaded relief view at a scale of 1:25,000, with sea floor depth as topographic contours overlaid in blue. It is based on multibeam echo-sounder data collected during four cruises conducted between the fall of 1994 and the fall of 1998. The map is part of a 3-quadrangle map series showing the area offshore of Boston, Mass., that is companion to the Stellwagen Bank National Marine Sanctuary map series (Valentine and others, 2001, 2003a-c; also see location map). Other maps of Quadrangle 1 depict topographic contours (Butman and others, 2003a), and backscatter intensity over shaded relief and topographic contours (Butman and others, 2003b).

The multibeam echo-sounder surveys were conducted aboard the vessel *Frederick G. Creed*, a SWATH (Small Waterplane Area Twin Hull) ship that surveys at speeds up to 15 knots. A Simrad Subsea EM 1000 Multibeam Echo Sounder (95 kHz), mounted on the starboard pontoon of the *Creed*, was used to acquire these data. The bathymetric soundings were gridded at 6 m/pixel resolution and smoothed using a 9-cell by 9-cell median filter; contours having a 5-meter interval were generated from the resulting grid. The shaded relief image was created by vertically exaggerating the topography four times and then artificially illuminating the relief by a light source positioned 45 degrees above the horizon from an azimuth of 350 degrees. In the resulting image, topographic features are enhanced by strong illumination on the northward-facing slopes and by shadows cast on south-facing slopes. The image accentuates small features that could not be effectively shown by contours alone at this scale. Blank areas in the image represent places where no data exists; data coverage begins 6 to 7 km offshore. The shoreline was extracted from the National Oceanic and Atmospheric Administration Medium Resolution Digital Vector Shoreline (data downloaded from <http://seaserver.nos.noaa.gov/projects/shoreline/shoreline.html>, Segment EC30_05, digitized from Chart 13267 (National Ocean Service, 1997)). Topographic lows are identified by hachured contours (hachures face deeper water).

Some features in the image are artifacts of data collection and environmental conditions. They include small highs and lows and unnatural-looking features, and patterns oriented parallel or perpendicular to survey tracklines (tracklines run northwest-southeast). For example, the wrinkle-like features in the northeastern part of the quadrangle, which are about 100 to 150 m long and are oriented northeast-southwest and perpendicular to the ship's track, are a result of heave of the vessel during data collection caused by large surface waves. The northwest-southeast-trending raised linear features, for example noticeable near 42°19' N, 70°43' W, are a result of a Simrad hardware calibration problem with the acoustic beams aimed directly below the ship. Other northwest-southeast-trending features, especially noticeable along the shallow southwestern part of the mapped area of the quadrangle

(for example, see the linear feature extending between 42°16.37' N, 70°40.79' W and 42°15.63' N, 70°39.80' W), are a result of marginal data overlap where survey lines were spaced too far apart.

Geologic history

The major topographic features shown in this map series were formed by glacial processes that occurred in several stages. Ice containing rock debris moved across the region, sculpting its surface and depositing sediment, forming the ridges and valleys that characterize the region. Other features are the result of processes at work when much of the area was covered with rotting ice, and when at the same time small valley glaciers and ice falls were active. Ice retreat and marine submergence occurred between 18 and 14 ka, resulting in a highstand of sea level approximately 33 m above modern sea level about 14 ka (Oldale and others, 1993). A lowstand of sea level approximately 45 m below modern sea level occurred about 12 ka as the earth's crust rebounded from ice loading. Thus, the sea floor of Massachusetts Bay in water depths shallower than about 45 m was reworked during the marine transgression between 18 and 14 ka, again during a relatively rapid sea-level regression between 14 and 12 ka, and finally during the transgression between 12 ka and the present. Today, the surficial sediments and features are reworked and shaped by tidal and storm-generated currents, which erode and transport sediments from the shallow areas into the deeper basins. Over time, the shallow areas affected by these processes have become coarser as sand and mud are removed and gravel remains, and the deeper basins have been built up to as they receive the sand and mud. Knebel and Côté (1995) have identified areas of erosion, sediment reworking, and deposition in this region.

Quadrangle 1 features

One of the most striking aspects of the sea floor shown by this survey is the variability in bottom morphology and texture over scales of a few kilometers or less, caused by both natural and anthropogenic processes. The topography, surface features, and surficial sediment texture are the result of glacial processes, reworking during the last rise in sea level, reworking by modern processes, and the disposal of dredged and other material in this region over the last century.

The topographic contours in this region of western Massachusetts Bay run approximately northwest-southeast. Northeastward of about the 45-meter contour, the sea floor slopes steadily toward Stellwagen Basin (see location map). In water depths between 35 and 45 m, in the area between 42°16' N, 70°37' W and 42°18' N, 70°39' W, rock pinnacles rise 5 to 10 m above the surrounding sea floor. These rugged features are 100 to 300 m in horizontal extent. A series of depressions less than 1 m deep and characterized by extremely sharp boundaries occurs between 42°15' N, 70°36' W and 42°18' N, 70°40' W. Examples of these features are

centered near (1) 42°17.26' N, 70°38.29' W; (2) 42°17.0' N, 70°37.80' W; (3) 42°16.75' N, 70°38.40' W; (4) 42°16.40' N, 70°37.17' W; and (5) 42°15.5' N, 70°36.27' W, and they suggest sediment movement and sorting of bottom sediments. These features are floored with high-backscatter material composed of gravel and sand (Butman and others, 2003b); bottom photographs show large bedforms (Gutierrez and others, 2001). Between 35 and 30 m water depth, there is a shoreward transition from smooth to rough sea floor. However, some areas in this depth range still appear smooth (that is, little texture is evident) in the shaded relief image; examples are the regions centered near 42°14.35' N, 70°37.5' W and 42°17.1' N, 70°41.4' W. At depths shallower than about 30 m the sea floor is rough, and there is a suggestion of northeast-southwest-trending ridges having steep faces toward the southeast; the ridges are particularly noticeable in the area between 42°14.6' N, 70°38.6' W and 42°15.5' N, 70°39.5' W.

Ribbon-like features (for example, near 42°16.88' N, 70°40.46' W and 42°14.57' N, 70°37.91' W) typically are 0.5 to 1 km long and less than 40 m wide, and are suggestive of active sediment transport. The relief across these features is about 25 cm or less. Bottom photographs show that the northern feature is floored with coarse sand shaped into large bedforms (Gutierrez and others, 2001). No photographs are available of the southern feature.

Between 70°43' W and 70°44' W, northward from 42°19' N, an unnatural-appearing roughness is caused by individual dumps of anthropogenic material. In the backscatter intensity image (Butman and others, 2003b), individual dumps are clearly visible as features having high backscatter intensity.

ACKNOWLEDGMENTS

The multibeam echo-sounder surveys were conducted with support from the Canadian Hydrographic Service and the University of New Brunswick. For their skillful work at sea, we thank G. Costello, N. Doucet, J. Gagne, R. Sanfalcon, and B. Tessier of the Canadian Hydrographic Service, E. Radford of the National Oceanic and Atmospheric Administration, and the officers and crew of the Canadian Hydrographic Service survey vessel *Frederick G. Creed*. We also thank J.H. Clark of the University of New Brunswick for his assistance in data acquisition, processing, and interpretation.

REFERENCES CITED

Butman, Bradford, Hayes, Laura, Danforth, W.W., and Valentine, P.C., 2003a, Sea floor topography of Quadrangle 1 in western Massachusetts Bay offshore of Boston, Massachusetts: U.S. Geological Survey Geologic Investigations Series Map I-2731-A, scale 1:25,000.

—, 2003b, Backscatter intensity, shaded relief, and sea floor topography of Quadrangle 1 in western Massachusetts Bay offshore of Boston, Massachusetts:

U.S. Geological Survey Geologic Investigations Series Map I-2731-C, scale 1:25,000.

—, 2003c, Shaded relief and sea floor topography of Quadrangle 2 in western Massachusetts Bay offshore of Boston, Massachusetts: U.S. Geological Survey Geologic Investigations Series Map I-2732-B, scale 1:25,000.

Gutierrez, B.T., Butman, Bradford, and Blackwood, D.S., 2001, Photographs of the sea floor of western Massachusetts Bay, offshore of Boston, Massachusetts: July 1999: U.S. Geological Survey Open-File Report 00-427, 1 CD-ROM.

Knebel, H.J., and Côté, R.C., 1995, Seafloor environments within the Boston Harbor-Massachusetts Bay sedimentary system: a regional synthesis: *Journal of Coastal Research*, v. 11, no. 1, p. 230-251.

National Ocean Service, 1997, Massachusetts Bay: National Oceanic and Atmospheric Administration, National Ocean Service, Chart 13267, scale 1:80,000.


Oldale, R.N., Colman, S.M., and Jones, G.A., 1993, Radiocarbon ages from two submerged strandline features in the western Gulf of Maine and a sea-level curve for the northeastern Massachusetts coastal region: *Quaternary Research*, v. 40, no. 1, p. 38-45.

Valentine, P.C., Middleton, T.J., and Fuller, S.J., 2001, Maps showing sea floor topography, sun-illuminated sea floor topography, and backscatter intensity of the Stellwagen Bank National Marine Sanctuary region off Boston, Massachusetts: U.S. Geological Survey Open-File Report 00-410, 1 CD-ROM, scale 1:60,000.

Valentine, P.C., Baker, J.L., and Unger, T.S., 2003a, Sea floor topography of the Stellwagen Bank National Marine Sanctuary off Boston, Massachusetts: U.S. Geological Survey Geologic Investigations Series Map I-2676-A, scale 1:60,000.

Valentine, P.C., Unger, T.S., and Baker, J.L., 2003b, Sun-illuminated sea floor topography of the Stellwagen Bank National Marine Sanctuary off Boston, Massachusetts: U.S. Geological Survey Geologic Investigations Series Map I-2676-B, scale 1:60,000.

—, 2003c, Backscatter intensity and sun-illuminated sea floor topography of the Stellwagen Bank National Marine Sanctuary off Boston, Massachusetts: U.S. Geological Survey Geologic Investigations Series Map I-2676-C, scale 1:60,000.



Location map showing mapped Quadrangle 1 shaded in blue. The western Massachusetts Bay map series comprises quadrangles 1-3 (outlined in black); the maps for Quadrangle 1 include this map and Butman and others (2003a,b). The shaded relief and sea floor topography of Quadrangle 2 are shown at scale 1:25,000 in Butman and others (2003c). Quadrangles 1-18 (outlined in gray) compose the companion Stellwagen Bank National Marine Sanctuary (SRNMS) map series. The shaded relief and sea floor topography of the entire area of quadrangles 1-18 is shown at scale 1:60,000 in Valentine and others (2001, 2003b); it is also shown by quadrangle at scale 1:25,000 as U.S. Geological Survey Geologic Investigations Series Maps I-2701 (for quadrangle 1) through I-2718 (for quadrangle 18). The SRNMS boundary is shown as a dashed line. Selected bathymetric contours are labeled in meters.

**SHADED RELIEF AND SEA FLOOR TOPOGRAPHY OF QUADRANGLE 1
IN WESTERN MASSACHUSETTS BAY OFFSHORE OF BOSTON, MASSACHUSETTS**

By

Bradford Butman, Laura Hayes, William W. Danforth, and Page C. Valentine

2003