U.S. DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY

GEOLOGIC INVESTIGATIONS SERIES
MAP I-2802

Table 1.—Summary of drill-hole data in the Van Buren North quadrangle, Missouri.

[There are no logs for the upper parts of most of the drill holes. Og, Gasconade Dolomite; €e, Eminence Dolomite; €p, Potosi Dolomite; €dd, Derby-Doerun Dolomite (usage of Missouri Geological Survey, 1979); €dt, Davis Formation; €b, Bonneterre Formation; €l, Lamotte Sandstone; Yrm, Rhyolite of Russell Mountain; Yg, Mesoproterozoic granite. nd, no data available. Datum is mean sea level]

Drill-hole number	Location	Collar elevation, in feet	Contact elevation, in feet	Bottom elevation, in feet
Drill-hole data on file with Missouri Department of Natural Resources, Geological Survey and Resource Assessment Division				
19649	SW1/4SE1/4 sec. 7, T. 28 N., R. 1 W.	530	No data to 492 492–Og-Ke 583–Kp-Ke 690–Ke-Kd 740–Kd-Ke 760–Ke-Kl 765–Kl-Cr 767–Cr-g	895
19669	SE1/4SW1/4 sec. 9, T. 28 N., R. 1 W.	510	No data to 495 495–Kp-Kdd 605–Kdd-Ke 710–Ke-Kd 730–Kd-Ke 745–Ke-Cr 760–Cr-g	404
19737	SW1/4SE1/4 sec. 12, T. 28 N., R. 1 W.	850	No data to 558 558–Og-Ke 590–Ke-Kp 640–Kp-Kdd 665–Kdd-Ke 690–Ke-Kd 710–Kd-Ke 715–Ke-Cr 730–Cr-g	435
20118	NW1/4SE1/4 sec. 8, T. 28 N., R. 1 W.	850	No data to 623 623–Og-Ke 650–Ke-Kp 698–Kp-Kdd 705–Kdd-Ke 805–Ke-Kl 810–Kl-Cr 830–Cr-g	405
20189	NW1/4SE1/4 sec. 4, T. 28 N., R. 1 W.	580	No data to 550 550–Ke-Kp 610–Kp-Kdd 615–Kdd-Ke 660–Ke-Kl 665–Kl-Cr 680–Cr-g	445
20241	SE1/4NE1/4 sec. 13, T. 28 N., R. 2 W.	600	No data to 585 585–Ke-Kp 610–Kp-Kdd 633–Kdd-Ke 640–Ke-Kl 642–Kl-Cr 660–Cr-g	455
21589	NW1/4NW1/4 sec. 12, T. 27 N., R. 1 W.	649	No data to 524 524–Og-Ke 569–Ke-Kp 621–Kp-Kdd 631–Kdd-Ke 636–Ke-Cr 1096–Cr-g	-1096
21668	NW1/4SE1/4 sec. 15, T. 28 N., R. 1 W.	760	No data to 681 681–Og-Ke 710–Ke-Kp 750–Kp-Kdd 754–Kdd-Ke 759–Ke-Kl 762–Kl-Cr 1060–Cr-g	458
22002	NE1/4NW1/4 sec. 12, T. 27 N., R. 1 W.	660	No data to 590 590–Og-Ke 630–Ke-Kp 719–Kp-Kdd 740–Kd-Ke 760–Ke-Cr 1061–Cr-g	-1001
23113	SW1/4NW1/4 sec. 27, T. 28 N., R. 1 W.	500	No data to 473 225–Og-Ke 450–Ke-Kp 515–Kp-Kdd 520–Kdd-Ke 654–Ke-Kl 656–Kl-Cr 1090–Cr-g	-1001
20977	NW1/4SE1/4 sec. 17, T. 27 N., R. 1 W.	700	No data to 681 681–Og-Ke 715–Ke-Kp 755–Kp-Kdd 771–Kdd-Ke 776–Ke-Cr 1096–Cr-g	-1137
Drill-hole data on file with U.S. Bureau of Land Management				
CT-5	SE1/4NW1/4 sec. 33, T. 28 N., R. 1 W.	510	No data to 255 255–Ke-Cr 1160–Cr-Cdd 1240–Cdd-Cp 1260–Cp-Cdd 1310–Cdd-Cb 1390–Cb-Cr	806
PD-15-1	NE1/4SE1/4 sec. 15, T. 27 N., R. 1 W.	706	No data to 126 126–Ke-Cdd 1240–Cdd-Cp 1430–Cp-Cb 1530–Cb-Cr 1564–Cr-Yg	-858
SH-4	NE1/4NW1/4 sec. 21, T. 28 N., R. 1 W.	520	No data to 458 458–Ke-Cr g	-860
SH-12	SE1/4NW1/4 sec. 7, T. 28 N., R. 1 W.	560	No data to 498 498–Ke-Cr g	-915
SH-35	SW1/4NW1/4 sec. 14, T. 28 N., R. 1 W.	580	No data to 400 400–Ke-Cr g	-804



Figure 2.—Columnar section of exposed Paleozoic units in the Van Buren North quadrangle.

CORRELATION OF MAP UNITS
Holocene and Pleistocene
QUATERNARY
TERTIARY
Unconformity
Lower Ordovician
ORDOVICIAN
Upper Cambrian
CAMBRIAN
Unconformity
MESOPROTEROZOIC
* In cross section only
DESCRIPTION OF MAP UNITS
Alluvium (Holocene and Pleistocene?)—Gravel, sand, and clay along the bed and active flood plains of modern stream valleys. Material consists mostly of subrounded to subangular, fine- to coarse-grained, and quartzite clasts in a matrix of sand containing silt and clay. Thickness ranges from 0 to 20 ft
Landslide deposits (Holocene and Pleistocene?)—Chaotic jumble of dolomite, chert, sandstone, and quartzite clasts in a clayey and silty matrix. Includes large slump blocks and jumbles of chert. Locally with hummocky topography. 20 to 100 ft thick
Muck and clayey silt (Holocene and Pleistocene)—Silt, clay, and muck filling sinkholes. Commonly contains water. Thickness unknown
Terrace deposits (Pleistocene?)—Relatively flat areas along floors and flanks of modern valleys that underlie some upland seasonal floors of present streams. Material consists of large cobble-size to sand-size, subrounded to subangular chert, sandstone, and quartzite clasts within a matrix of sand, silt, and clay. In some locations a significant fraction of the matrix is of subangular to subrounded loess. Terraces abutting the Current River and major streams generally have a 6- to 10-ft-high cutbank descending to the water or a pronounced slope descending to streams. Near area of Qa. Thickness at least 10 ft and possibly more along larger stream courses
Residuum derived from the Roubidoux Formation (Ordovician) and Tertiary(?)–Cretaceous(?) sand, chert, sandstone slabs and sandy chert rubble and cobbles and clayey, commonly reddish, soil. Siliceous, minor knob, and ridge caps mostly derived from the Roubidoux Formation and some carbonates from the underlying upper part of the Gasconade Dolomite. Commonly mixed with alluvium. Contact with the Gasconade Dolomite is placed where identifiable clasts of dolomite are found in the residuum. Near contacts, or where residuum thickness approaches zero. Unit thickness may exceed 40 ft on some knobs and tapers down to zero on flanking hillsides
Upland gravel deposit (Quaternary and Tertiary)—Gravel deposits, as high as 300 ft above modern streams. Consists of chert, containing mostly subrounded to rounded clasts of chert, sandstone, and quartzite in a matrix of silt, sand, and clay. Commonly contains reddish clay. Near contacts, or where residuum thickness approaches zero. Unit thickness may exceed 40 ft on some knobs and tapers down to zero on flanking hillsides
Roubidoux Formation (Lower Ordovician)—Sandstone, sandstone, and chert. Dolomite, very light gray to medium-gray, pale-orange, or pink-ish-gray, very fine grained to medium-grained, commonly thin- to thick-laminated; interbedded with quartz sandstone, white to pale orange, weathering to light-gray, fine- to medium-grained, sand grains coated by coarse-grained, poorly sorted, thin- to thick-bedded, commonly crossbedded or cone-in-cone; well-rounded quartz grains commonly have angular overgrowths; most beds dolomite-cemented; some silica-cemented. Dolomite commonly replaced by white, gray, or brown chert, commonly stromatolitic and occasionally oolitic or fossiliferous. Heavy mineral fraction dominated by tourmaline, zircon, and garnet. Basal 30 ft contains abundant 1- to 5-ft-thick layers of sandstone; overlying beds are dolomitic and medium- to thick-bedded, but rare except for occasional impressions of the snail Lecanospira. Bedding surfaces of sandstone beds commonly have ripple marks formed by currents. Lower contact of unit mapped at lowest orthoquartzite or sandstone. Unit thickness ranges from 0 to 350 ft. Thickest in southeast corner of the map area. Regionally the Roubidoux ranges from 100 to 350 ft thick; thickness determined from wells in the Ozark region (Bretz and others, 1965)
Gasconade Dolomite (Lower Ordovician)—Dolomite, chert, sandstone, and orthoquartzite. Dolomite is the dominant rock type and is described in figure 2 but not shown separately on this map. Dolomite of upper unit, light-gray, medium- to thick-bedded, commonly thick-bedded, vuggy; weathers to a pitted surface. Dolomite of upper part of middle unit, light-gray, fine- to coarse-grained, medium- to thick-bedded with white to light-gray chert nodules and lenses. The uppermost bed of upper part of middle unit commonly contains abundant chert, up to 5 ft thick. Dolomite of middle part of middle unit, light-gray to very light gray, medium- to coarse-grained, commonly contains chert as nodules, stringers, and beds; chert varieties are porcelaneous, oolitic, and fossiliferous; some beds cross-laminated. Dolomite of lower part of middle unit, formerly called the Van Buren Formation, dolomite, light-gray to yellowish-gray, fine- to medium-grained, thin- to medium-bedded. The lower unit is the Gunter Sandstone Member, light-gray to white sandstone, sandy dolomite, or orthoquartzite interbedded with light-gray to tan, fine- to coarse-grained, thin-bedded dolomite. Contact with underlying Eminence Dolomite is placed at base of lowest sandstone or sandy dolomite of the Gunter Sandstone Member. Thickness of the Gunter ranges from 15 to 44 ft. Maximum thickness of exposed Gasconade is about 280 ft
Eminence Dolomite (Upper Cambrian)—Dolomite, light- to medium-gray, occasionally mottled red, pink, or green; medium- to coarse-grained, medium- to thick-bedded, commonly massive; minor amounts of chert, locally white or drusy, in nodules and irregular fragments, mostly in upper half of formation; minor quartz sandstone, medium- to thick-bedded. Contact with underlying Potosi Dolomite is gradational and placed above the highest bed of brown or drusy dolomite. Unit is variably thick near and over the Proterozoic knobs in the region. Exposed thicknesses range from 140 ft to as much as 270 ft; in well logs the thickness is as much as 363 ft (table 1)
Potosi Dolomite (Upper Cambrian)—Dolomite, light-brownish-gray to pale-yellowish-brown, fine- to medium-grained, thick-bedded, vuggy and porous, with drusy chert; locally stromatolitic; bituminous odor typical of freshly broken rock. Thickness ranges from 336 to 480 ft on the basis of drill hole data (table 1)
Derby-Doerun Dolomite (usage of Missouri Geological Survey, 1979) (Upper Cambrian)—Shown in cross section only. Dolomite, siltstone, and shale. Dolomite, buff to brown or light-gray, fine to medium-grained, thin- to medium-bedded, argillaceous, silty, with minor amounts of chert and sparse sulfide minerals; vuggy and porous. Siltstone and shale, thin-bedded, interbedded with dolomite. Thickness ranges from 100 to 209 ft on the basis of drill hole data (table 1)
Davis Formation (Upper Cambrian)—Shown in cross section only. Shale, siltstone, sandstone, dolomite, and limestone conglomerate. Shale, dark-green, fissile, thin to thick-bedded; as much as 50 percent of formation. Limestone, light-gray, fine-grained to coquinoid, dense, locally glauconitic. Contains intraformational sedimentary breccias and pseudonodules. Ranges from 131 to 185 ft thick on the basis of drill hole data (table 1)
Bonneterre Formation (Upper Cambrian)—Shown in cross section only. Dolomite, limestone, siltstone, and shale. Dolomite, light-gray, fine- to medium-grained, medium-bedded; contains algal structures; locally glauconitic and shaly; sparse to abundant sulfide minerals. Limestone, brownish-gray to pink, fine-grained, thin-bedded, fossiliferous, locally oolitic; more common in lower part of formation. Siltstone, quartzose, light to dark-gray, laminated. Shale, dark-green, thin-bedded; occurs as sparse thin beds and partings. Where the underlying Lamotte Sandstone is missing, basal 2 to 20 ft of Bonneterre consists of dolomite-cemented porphyry pebbles and cobbles. Major lead-bearing unit in the Ozark region. Thickness in the Van Buren North quadrangle ranges from less than 95 ft to 515 ft on the basis of drill hole data (table 1)
Lamotte Sandstone (Upper Cambrian)—Shown in cross section only. Sandstone, quartzose, light-gray, yellow, brown, or red; dominantly medium-grained, moderately to well-sorted, well-indurated; locally contains interbedded red to purple silty shale and, in upper part, scattered lenses of arenaceous dolomite. Felsite pebble or boulder conglomerate commonly present at base. Thickness ranges from 2 to 224 ft on the basis of drill hole data (table 1)
Rhyolite of Russell Mountain (Mesoproterozoic)—Ash-flow tuff, porphyritic, trachytic to rhyolitic, purple to dark-purplish-red. Probably equivalent to either the Rhyolite of Russell Mountain (McDowell and Harrison, 2000) or the Tuff of Little Thorny Mountain (Harrison and others, 2002)
Granite (Mesoproterozoic)—Shown in cross section only
EXPLANATION OF MAP SYMBOLS
PLANAR FEATURES
OTHER FEATURES
DISCUSSION
INTRODUCTION
The Van Buren North 7.5-minute quadrangle is located in south-central Missouri on the Salem Plateau of the Ozark Plateaus physiographic province (Fenneman, 1938; Bretz, 1965; Harrison and others, 1994; Weary and Doctor, 2014). Most of the area within the quadrangle is publicly owned, and much of the area along the Current River is administered by the Ozark National Scenic Riverways unit of the National Park Service. Most land along the western quadrangle margin is part of the Peck Ranch State Conservation Area, and much of the eastern quadrangle is within the Mark Twain National Forest and the Cardarova, Beal, and Paint Rock State Conservation Areas. Privately owned areas are used largely for farming, timber production, and recreation. The northern part of the town of Van Buren occupies the extreme southeast corner of the map area, but most of the quadrangle is unpopulated. The only major highways in the area are U.S. Highway 60 and Business 60 in the southeast corner of the quadrangle. Missouri Highway D along the eastern Current River arm and Rte. BB in the northeast are central area. The map area has topographic relief of about 550 ft, with elevations ranging from about 630 ft along the Current River to about 1,180 ft on hilltops about 1,000 ft in the west-central area.
Fieldwork was conducted during the winter months of 1997 and 1998. Bedding and joint attitudes were recorded from bedrock exposures wherever observed. Many contact elevations were determined by interpolation along a point of known elevation.
MESOPROTEROZOIC ROCKS
Proterozoic igneous rocks exposed in the Van Buren North quadrangle are a southwestern extension of the rocks in the St. Francois Mountains of eastern Missouri (fig. 1). Uranium-lead isotopic analyses established that granite at 1,511.5±80.4 Ma for granite from a corehole in the Winona quadrangle to the west (Van Schmus and Bickford, 1981) and at 1,470±10 Ma for granite from a corehole in the Ellington quadrangle to the northeast (Bickford and others, 1981), and (b) 1,470.4±2.7 Ma for the rhyolite of Shutin Mountain in the Segall Mountain quadrangle to the west (Harrison and others, 2000).
One small outcrop of rhyolite (unit Yrm) occurs in the map area in the center of sec. 24, T. 28 N., R. 1 W. in the southwest corner. The rhyolite is porphyritic and purple to dark purplish red. Whole-rock major-oxide analyses yielded the following composition: SiO_2, 72.64 percent; Al_2O_3, 13.23 percent; FeO, 0.21 percent; Fe_2O_3, 2.21 percent; MnO, 0.06 percent; CaO, 0.52 percent; TiO_2, 0.24 percent; Na_2O, 1.91 percent; K_2O, 7.31 percent; and P_2O_5, 0.03 percent, and a loss on ignition of 1.55 percent. Generally, however, Mesoproterozoic rocks are encountered only in deep (1,000–2,000 ft) exploration drill holes (table 1). All but one of the drill holes in the quadrangle encountered rocks that were described as granites. The only nearby granite outcrop is in the Big Spring quadrangle about 2 mi to the southeast of the southeast corner of the map area. The granite and its surrounding volcanic deposits were deformed, tilted, and subjected to uplift and erosion throughout Late Proterozoic time. The regional relief on this buried erosional surface has been reported to be as much as 2,000 ft (Hawkesworth and Ethridge, 1978).
PALEOZOIC STRATIGRAPHY
The Van Buren North quadrangle is underlain by lower Paleozoic marine strata ranging from about 1,200 to 2,300 ft in thickness; the rocks consist chiefly of dolomite, chert, sandstone, and orthoquartzite. From oldest to youngest, these units are the Upper Cambrian Lamotte Sandstone, Bonneterre Formation, Davis Formation, Derby-Doerun Dolomite (usage of Missouri Geological Survey, 1979), Potosi Dolomite, and Eminence Dolomite; and the Lower Ordovician Gasconade Dolomite and Roubidoux Formation. Only the last four units are exposed in the quadrangle. The stratigraphic column, figure 2, summarizes the general lithologies and thickness of the Paleozoic strata that crop out in the Van Buren North quadrangle.
In Late Cambrian time, the Ozark region was flooded by an epicontinental sea that covered most of the Midwestern part of the United States. Because of strong subsidence in the Reelfoot rift, southeastern Missouri was one of the areas were submerged, and high areas remained above water as islands. In the map area, the basal Cambrian unit, the Lamotte Sandstone, onlaps the flanks of these Mesoproterozoic bedrock hills.
During deposition of the Bonneterre Formation, algal reefs developed along the flanks of the islands of Proterozoic rocks in the Late Cambrian sea (Lyle, 1977). These reefs, especially to the northwest along a mining district known as the Viburnum Trend (about 30 mi northwest of the map area), later became the focus of precipitation of important lead deposits (Thacker and Anderson, 1977). Most of the island hills in the map area were entirely buried by the end of Bonneterre deposition, but subsequently these topographic features have affected depositional patterns through differential compaction of sediments deposited on and around their flanks.
Deposition, mostly of carbonate sediments, continued through the rest of the Cambrian and into the beginning of the Ordovician, sequentially accumulating the Davis Formation, Derby-Doerun Dolomite, Potosi Dolomite, and Eminence Dolomite.
The Davis, which is exposed in outcrops in the Ozark region (Lyle, 1977), the St. Francois County, Mo., by Buehler (1957), The Davis Formation conformably overlies the Bonneterre Formation and comprises shale, siltstone, sandstone, dolomite, and limestone conglomerate. The Davis, along with the overlying Derby-Doerun Dolomite, has low primary permeability and acts as an aquitard, part of the St. Francois confining unit, between the water St. Francois aquifer and the upper Ozark aquifer (Kleeschulte and Seeger, 2000).

Figure 1.—Map showing location of the Van Buren North 7.5-minute quadrangle and the geomorphic subdivisions of Missouri. The St. Francois Mountains region is underlain mostly by Cambrian and Proterozoic rocks, the Salem Plateau by Cambrian and Ordovician rocks, the Springfield Plateau by Mississippian rocks, the Central Lowland province by Ordovician to Pennsylvanian rocks and Quaternary periglacial sediments, and the Mississippi embayment by Cretaceous to Quaternary sediments (adapted from Imes and Emmett, 1994).

Figure 3.—Compass-rose diagram showing the dominant joint trends in the Paleozoic rocks of the Van Buren North quadrangle. Joint sets were weighted by counting closely spaced joint sets three times, moderately spaced sets two times, and widely spaced sets one time. This system approximates the importance of each set relative to the volume of the rocks. Interval is 10 degrees. Numbers on diagram are percent of total. n, number of joints measured.
The Derby-Doerun Dolomite was originally two separate formations described from outcrops near mines operated by the Derby Lead Co. and the Doe Run Lead Co. in St. Francois County, Mo. (Thompson, 1995). The formation comprises thin- to medium-bedded dolomites intercalated with thin-bedded siltstone and shale.
The Potosi Dolomite of Late Cambrian age is the oldest Paleozoic unit exposed in the map area. Here mapped, the Potosi crops out in two small areas at the base of the bluffs in sec. 5, T. 28 N., R. 1 W., along the eastern margin of the Current River flood plain at the southeastern edge of the map area. The name Potosi was originally used by Winslow (1894) in a much broader sense for rocks exposed near the town of Potosi in Washington County, Mo. Dake (1930) redefined the Potosi Dolomite as used presently. The Potosi is a fine- to medium-grained, light-brownish-gray to pale-yellowish-brown dolostone, locally dolomitic, that has siliceous drusy as a permineralization of the dolomite or as vug linings, and a field smell on freshly broken surfaces. In outcrop in the map area, and in the Van Buren South and Big Spring quadrangles to the south and east, the Potosi characteristically is more intensely jointed than the overlying units, with subvertical fractures. Pronounced cavern development exists at or near the contact between the Potosi and the overlying Eminence Dolomite and is marked by bedding unconformities and chert breccias, suggesting a permeability contrast across that boundary. The Potosi is characteristically unfossiliferous. Its contact with the overlying Eminence Dolomite is gradational through a few tens of feet and is interfingering, with alternating beds of light-brown dolomite of the Potosi and massive, cliff-forming, pale-red dolomite of the Eminence formations, especially near the Mesoproterozoic knobs in the region (Ordoñez and others, 1999).
The Eminence Dolomite was described by Ulrich (1911) from exposures in the vicinity of the town of Eminence, Mo. Most of the exposures occur in the floor and along the walls of the valley of the Current River and its main tributaries. The Eminence comprises medium- to thick-bedded and commonly massive, medium- to coarse-grained, light to medium-gray dolomite and cherty dolomite with a few minor sandstone beds and white cherts. Many of the beds are stromatolitic, with pits and vugs developed between the laminae, and others are crossbedded. The thick, nonCherty dolomite beds commonly weather into irregular pinnacles; especially beds lying subjacent to sandstone of the Eminence or of the overlying Gunter Sandstone Member of the Gasconade Dolomite.
A conspicuous interval of sandstone and dolomite about 10 ft thick occurs within the Eminence between 30 and 90 ft below the top of the formation. The base of the interval is a bedding unit of sandstone bed, 18 to 24 in thick, and the top is a thin interval of fine- to medium-bedded dolomite and an upper sandstone, 4 to 12 in thick, although ledge forming, that commonly has ripple marks and can be localized kilometers. This sandstone-dolomite interval was observed in outcrops scattered throughout the map area, has been reported in the Stegall Mountain 7.5-minute quadrangle to the west (Harrison and others, 2002), and has been observed in the Van Buren South and Big Spring 7.5-minute quadrangles to the south and southeast. Assuming that this interval is correlative across this rugged part of the Ozarks, it may mark a regressive event similar to, but less widespread than, the regression indicated by the unconformity beneath the basal Gunter Sandstone Member of the overlying Gasconade Dolomite. The sandstone in this interval is easily mistaken for the Gunter in some localities. The variation of thickness of the Eminence above the sandstone-dolomite interval is probably due to erosion down into the Eminence prior to deposition of the overlying Gunter.
Bridge (1930) reported the Eminence to be abundantly fossiliferous, especially with fossils and gastropods in residual cherts. No such fossils were observed during the course of this study. Conodonts recovered from two samples about 80 ft below the top of the Eminence in sec. 13, T. 28 N., R. 2 W. (USGS Nos. 11902-CO and 11903-CO), indicate Late Cambrian to very Early Ordovician ages (J.E. Repetski, USGS, written commun., 2001). Previous maps produced by this USGS project placed the Cambrian-Ordovician boundary in the upper part of the Eminence Dolomite (McDowell, 1998; Orndorff and others, 1999; McDowell and Harrison, 2000; Orndorff and Harrison, 2001; Harrison and others, 2002; Weems, 2002; Harrison and McDowell, 2003). However, recent international agreement on redefinition of this boundary requires that it be placed at the contact between the Eminence and Gasconade Dolomites (Cooper and others, 2001). Bridge (1930) speculated that an important unconformity exists at the base of the Gasconade. Conodont biostratigraphy suggests that there may be multiple unconformities within the basal Gasconade interval and that the basal Gasconade is significantly diachronous across the area between Van Buren and the town of Eminence, about 22 mi to the northwest (Repetski and others, 2000).
The thickness of the Eminence ranges from about 140 ft to as much as 270 ft (Harrison and others, 2002) and up to 363 ft reported from drill holes (table 1). Variability in thickness is the result of the non-planar transition from the underlying Potosi Dolomite and was leveling by the unconformity at the base of the overlying Gasconade Dolomite.
The Eminence Dolomite is the main cave-bearing formation in the Van Buren North quadrangle, and caves are concentrated just below the Gunter Sandstone Member of the overlying Gasconade Dolomite and to a lesser extent below the sandstone-dolomite interval within the Eminence.
In earliest Ordovician time a regression of the sea produced an unconformity on top of the Eminence Dolomite. A subsequent marine transgression in the Early Ordovician resulted in deposition of the Gasconade Dolomite. The Gasconade was named by Nason (1892) for exposures along the Gasconade River in Laclede, Pulaski, and Phelps Counties, central Missouri. The basal 15 to 44 ft consists of interbedded sandstone, orthoquartzite, and thin-bedded dolomite and sandy dolomite named the Gunter Sandstone Member by Ball and Smith (1903) for exposures along the Niangua River at Gunter near Hahatonka Springs, Camden County, central Missouri. The lower part of the Gunter is composed of relatively thin quartz sandstone and dolomite beds that commonly have ripple marks on the bedding surfaces. The upper part is typically thick-bedded or massive, light-gray to white quartz sandstone as much as 15 ft thick, producing a prominent topographic bench on many of the hillsides.
Bridge (1930) subdivided the Gasconade into upper, middle, and lower units of the Gunter in the Van Buren North quadrangle is different from that in outcrops observed in other areas to the northwest (Orndorff and others, 1999), where the total thickness rarely exceeds 15 ft and the lower sandstone is more prominent than the upper. Bridge (1930) used the name Van Buren Formation for the Gunter Member and superjacent medium-bedded non-cherty and cherty dolomite below an oolite bed in the middle part of the Gasconade. The Van Buren was defined by its faunal content, and the name Van Buren Formation is no longer used. Pratt and others (1998) subdivided lower part of the Gasconade in the Van Buren quadrangle into a lower unit, the Gunter, and an upper unit, the Ozark Township Formation. The Ozark Township Formation is characterized by a persistent cryptozoan chert. The Cryptozoan chert is distinctive in outcrop and in drill holes (Pratt and others, 1998). Conodonts recovered from the chert bed were obtained in the Van Buren North quadrangle. However, Cryptozoan chert bodies indicating this horizon were observed in numerous locations at higher elevations within the area, especially in the area of sec. 26, T. 28 N., R. 2 W. As shown in figure 2, the lower part of the Gasconade is the Gunter Sandstone Member. The middle part of the Gasconade includes the upper, cherty part of Bridge's Van Buren Formation and dolomite above the oolite bed, up to and including the Cryptozoan chert. The upper part of the formation above the chert is thick-bedded dolomite.
Fossils are rare in the Gasconade within the map area, though the presence of planispiral gastropods and cephalopods from cherts within this unit have been reported at Fort Leonard Wood, about 75 mi to the southwest, and at a locality just northeast of the town of Van Buren (Harrison and others, 1996; J.E. Repetski, oral commun., 2000). Conodonts collected from just above the Cryptozoan chert in the NW1/4 sec. 27, T. 28 N., R. 1 W. (USGS No. 11901-CO), indicate Early Ordovician (early Ibexian) age (J.E. Repetski, written commun., 2001). Conodonts from the Gasconade in the Low Wassie quadrangle to the southwest also are indicative of an early Early Ordovician (early Ibexian) age (J.E. Repetski, written commun., 1997).
The Roubidoux Formation was named for exposures in the area of Roubidoux Creek in Pulaski County, Mo. (Nason, 1892). The contact between the Roubidoux and the underlying Gasconade is probably conformable and is marked at the base of the first significant orthoquartzite or sandstone in the Roubidoux. In this area the first sandstone is commonly thin bedded and is overlain by dolomite and cherty dolomite. The Roubidoux consists of interbedded dolomite, cherty dolomite, sandy dolomite, dolomitic sandstone, quartz sandstone, orthoquartzite, and chert. The carbonate in the formation is generally removed or obscured by residuum, so the sandstones are the most prominent lithology in outcrops and residuum. The Roubidoux probably accumulated in shallow water, because dolomite-rich intervals contain common algal stromatolites and oolites, while sandstone-rich intervals contain abundant desiccation cracks and ripple marks suggestive of periodic exposure and widely variable current directions.
No outcrops of Roubidoux were observed in the Van Buren North quadrangle, but the formation was identified in the southwest corner of the quadrangle on the basis of (1) stratigraphic thickness above the base of the Gasconade Dolomite or above the Gasconade Cryptozoan chert, (2) abundance of sandstone residuum, and (3) changes in local physiography, such as a break in slope.
Fossils generally are rare in the Roubidoux formation, though impressions of snails (mostly Lecanospira) locally are common in chert and sandstone. A few fossiliferous beds were reported by Bridge (1930) but none was observed in the course of this study. The only fossils from the Roubidoux in the quadrangle are from the upper part of the formation elsewhere within the Salem Plateau, and Harrison and others (1996) reported the occurrence of conodonts indicative of early Early Ordovician (early to middle Ibexian) age. Carver (1961), on the basis of petrographic studies, concluded that most of the gastropod and brachiopod fossils in this unit have been obscured by dolomitization.
The top of the Roubidoux has been removed by erosion in the map area. The Roubidoux is reported to be as much as 250 ft thick in other parts of Missouri (Thompson, 1995). As much as 165 ft of Roubidoux may be present in the southwestern part of the Van Buren North quadrangle.
After deposition of the Roubidoux, a number of younger Early Ordovician units accumulated which are preserved elsewhere in and around the Ozark Plateaus. These probably were present at one time in the map area.
Local residual concentrations of fossiliferous chert (Mississippian (Osagean) chert that probably accumulated in paleo-sinkholes were found in two locations in the southwest corner of the map area (Harrison and others, 1996; J.E. Repetski, oral commun., 2000), marked by triangles on the map. Similar deposits have also been found to the southwest in the Birch Tree quadrangle, to the west in the Bartlett and Winona quadrangles, and to the north in the Exchange quadrangle (Bridge, 1930). The parent unit of these residual rocks probably once extended across the Van Buren North quadrangle, but subsequently was stripped away by erosion. Pennsylvanian deposits, although unknown from the quadrangle, are known from the northern Ozark region (Unklesbay and Vineyard, 1992). This suggests that a regional transgression may have occurred across the Ozarks in Pennsylvanian time. No evidence has been found to suggest Permian deposition in the Ozark region.
POST-PALEOZOIC HISTORY
The Mesozoic history of the Ozark region is virtually unknown. Deposits of this age, if they once were present, have been eroded from the area. Based on depositional patterns in Mesozoic rocks throughout North America, the Ozark region probably was emergent and undergoing erosion in Triassic, Jurassic, and Early Cretaceous time. During the Late Cretaceous, however, the Ozark region probably was covered by an inland sea that covered most of the United States west of the present Mississippi River (Smith and others, 1994). It has been suggested that glauconitic Cretaceous sandstone, now stripped away, may have been the source of the iron contained in the "primary limonites" that occur in the southeastern Ozark region (Potter, 1955, p. 1251. Although missing today from the Ozark Plateaus, Upper Cretaceous deposits still are preserved to the southeast in the Mississippi embayment.
The Ozark region may also have been covered by the sea in early Paleocene time (Smith and others, 1994), but throughout the remainder of the Tertiary Period the Ozarks were emergent and have been above sea level and undergoing weathering and erosion. This implies that the existing stream patterns were established no earlier than the Tertiary. The widespread occurrence of thick residual soils indicates that the Ozark area has undergone prolonged weathering, at least since middle Tertiary time (McKeown and others, 1988). The existing rolling upland topography, with its deeply incised stream valleys, probably has resulted from regional Quaternary uplift, which was accompanied by the capture of much of the paleo-Gasconade River drainage basin by tributaries of the White River (Ellrich and Siemens, 1977). The development of several Pliocene to Pleistocene terrace deposits at successively lower elevations indicates that uplift and downcutting have been episodic and not continuous (Flint, 1941). It remains controversial whether the existence of uplift, with concomitant cave development, have resulted in the formation of regional flat erosional surfaces called peneplains (Bretz, 1956, 1962, 1965), or whether the existing landforms represent a dynamic equilibrium between bedrock hardness and erosion rates (Hack, 1960; Reams, 1968).
SURFICIAL GEOLOGY
Several levels of surficial terrace deposits occur along the Current River. Detailed differentiation and mapping of these terraces across the map area is beyond the scope of the present project. In several upland areas of the map, thin patchy deposits of subrounded to rounded cobbles and pebbles of chert, sandstone, and quartzite were observed (QTg). These upland gravels may be as high as 300 ft above the present river level and possibly are fluvial deposits from the major streams prior to downcutting of the present steep valleys. It is also possible that some isolated deposits of QTg, on high, conical knobs, actually may be sinkhole fills, which have subsequently been exhumed.
Valley floors are underlain by alluvial deposits of Holocene to possibly late Pleistocene age (Flores, 1985; Alkemade and others, 1995) that were derived mostly by the washing of clay, silt, and chert fractions from the adjacent upland residual soils and to a lesser extent by the reworking of loess.
Pleistocene loess is thin or absent over most of the Salem Plateau (Ebens and Connor, 1980). However, the Van Buren North quadrangle includes extensive areas