




This map shows the configuration of the bedrock surface if all unconsolidated earth materials were removed.



CONTOURS ON THE BEDROCK SURFACE,
IN FEET ABOVE OR BELOW (-)
MEAN SEA LEVEL

Contour interval 50 feet; hachures
indicate closed depression


REFERENCES

- Flint, R.F., 1964, The surficial geology of the Branford quadrangle, with map: Connecticut Geol. and Nat. History Survey Quad. Rept. 14, 45 p.
- Mazzaferro, D.L., 1973, Hydrogeologic data for the Quinnipiac River basin, Connecticut: Connecticut Water Resources Bulletin 75, 54 p.
- Sanders, F.E., Guidotti, C.V., and Wilde, Pat., 1964, Foxon fault and Gaillard graben in the Triassic of southern Connecticut: Connecticut Geol. and Nat. History Survey Rept. Inv. 2, 16 p.
- Ward, Freeman, 1920, The Quaternary geology of the New Haven region, Connecticut: Connecticut Geol. and Nat. History Survey Bull. 29, 78 p.

Base from U.S. Geological Survey, 1967;
photorevised 1972

10,000-foot grid based on Connecticut
coordinate system

1,000-meter Universal Transverse
Mercator grid ticks, zone 18



CONTOUR MAP OF THE BEDROCK SURFACE, BRANFORD QUADRANGLE, CONNECTICUT

By
Charles E. Brown