

MISCELLANEOUS FIELD STUDIES

MAP MF-606 A


EXPLANATION

This map is intended to serve as an aid in reconnaissance evaluation of unconsolidated materials and can be used to identify areas of potential interest.

The units on this map indicate the first material of substantial thickness (generally greater than 3 feet (1 m) encountered beneath the soil layer. The soil layer (commonly a foot or two (about 0.5 m) thick) is not mapped. Other materials, different in composition, may underlie each map unit (see block diagram) or may occur as minor lenses within each map unit.

THIS MAP SHOULD NOT BE USED AS A SUBSTITUTE FOR ONSITE INVESTIGATION.

Most unconsolidated materials are mixtures of the three particlesize classes defined in the diagram below. This diagram also relates these three size classifications to the Wentworth classification (Wentworth, 1922) which can be compared with other classifications used in engineering and soil science.



Coarse particles (stones) include granules, pebbles, cobbles, and boulders. Medium particles include very coarse, coarse, medium, and fine sand-sized particles. Fine particles include very fine sand-, silt-, and clay-sized particles. Very fine sand is included in this class because it commonly occurs with finer materials, and because very fine sand, silt, and clay behave similarly when watersoaked and under stress.

Materials mapping involves a visual estimate of particle-size distribution in a deposit by the field geologist. Percentages of particle sizes therefore may vary somewhat from place to place beyond the limits defined in the map units below.



SAND AND GRAVEL

Particle sizes range from 100 percent coarse to 25 percent coarse and 75 percent medium. Material may occur

as thin layers of well-sorted sand interbedded with thin layers of well-sorted gravel; as poorlysorted layers of mixed sand and gravel; or as distinct pockets of well- to poorly-sorted sand and gravel. May contain minor amounts of fine particles




Particle sizes range from 25 percent coarse and 75 percent medium through 100 percent medium to 50 percent medium and 50 percent fine. Material commonly occurs as well- to poorlysorted sand in layers of variable thickness



VERY FINE SAND, SILT, AND CLAY

Particle sizes range from 50 percent fine and 50 percent medium to 100 percent fine. Material may occur as layers of poorly sorted very fine sand, silt, and clay; as regularly alternating layers of silt and very fine sand interbedded with clay; or as thick, massive beds of clay with only minor amounts of very fine sand and silt. Locally contains scattered coarser particles






Poorly sorted nonlayered mixture of coarse, medium, and fine particles in varying proportions. Some till, averaging less than 10 feet (3 m) thick, is sandy, loose, and very stony; other till, commonly more than 10 feet (3 m) thick, is less sandy, less stony, and very compact. Where these tills occur together, the loose, sandy till is always on top. The compact till forms the N. bulk of many smooth elongate hills (drumlins) even where the sandy till is exposed at the surface



SWAMP DEPOSITS

Dark, decomposed organic material intermixed with varying amounts of sand, silt, and clay. Locally contains scattered stones



Shown only for roads, highways, dams, solid-waste disposal, and other major construction. In addition, in urban centers and other areas of dense development, fill of variable thickness and extent may overlie the natural materials shown on the map.

af, predominantly earth fill Gaft, predominantly trash fill

	_
	-
	-
	-

BEDROCK (LEDGE) OUTCROP Bedrock exposed at ground surface; may be partly covered by thin soil. Ruled pattern shows areas of small, closely-spaced outcrops



In general, lakes and ponds greater than 5 acres, or streams wider than 200 feet (60 m)

SOURCES OF DATA

00.1974

Schnabel, R.W., 1971, Surficial geologic map of the Southwick quadrangle, Massachusetts and Connecticut: U.S. Geol. Survey Geol. Quad. Map GQ-891. Wentworth, C.K., 1922, A Scale of grade and class terms for clastic sediments: Jour. Geology, v. 30, p. 377-392.

M(200) MF606-

01

MAP SHOWING UNCONSOLIDATED MATERIALS, SOUTHWICK QUADRANGLE, MASSACHUSETTS-CONNECTICUT By Robert W. Schnabel Connecticut (Southwick guad.). Su For sale by U.S. Geological Survey, price 50 cer

3 1818 00180351 7

1974