

GEOLOGY GENERALIZED FROM FOSTER (1970)

CORRELATION OF MAP UNITS

UNCONSOLIDATED DEPOSITS

Q Su QUATERNARY

IGNEOUS AND METAMORPHIC ROCKS

TM29 TM296 TERTIARY OR MESOZOIC

Kr] CRETACEOUS(?)

KJm CRETACEOUS OR JURASSIC

SEDIMENTARY ROCKS

DESCRIPTION OF MAP UNITS

PALEOZOIC(?)

UNCONSOLIDATED DEPOSITS

Q54 UNCONSOLIDATED SEDIMENTARY DEPOSITS SEDIMENTARY ROCKS

kr DETRITAL ROCKS (CRETACEOUS?) KJm MENTASTA ARGILLITE OF RICHTER (1967) (JURASSIC OR CRETACEOUS)

IGNEOUS AND METAMORPHIC ROCKS QTb BASALT

Tm MAFIC VOLCANIC ROCKS TF FELSIC TUFF, WELDED TUFF, LAVA, AND HYPABYSSAL INTRUSIVE ROCKS

TM29 GRANITIC ROCKS, UNDIVIDED

MER ULTRAMAFIC ROCKS

R4 DIORITE

Ep€m METAMORPHIC ROCKS, UNDIVIDED

GEOLOGIC SYMBOLS

CONTACT, APPROXIMATELY LOCATED U, UPTHROWN SIDE; D, DOWNTHROWN SIDE

---- FAULT OR LINEAMENT FROM AERIAL PHOTOGRAPHS LINE SEPARATES NORTHERN (YUKON-TANANA UPLAND) POPULATION OF GEOCHEMICAL SAMPLES FROM SOUTHERN (ALASKA RANGE) POPULATION

X BASE METAL PROSPECTS NORTH OF THE TANANA RIVER

GEOCHEMICAL SYMBOLS O BACKGROUND VALUES WEAKLY ANOMALOUS VALUES

STRONGLY ANOMALOUS VALUES

This series of geochemical maps shows the distribution of copper in four sample media: (A) the oxide residue (the oxalic-acid-leachable fraction) of the stream sediment, (B) the minus-80-mesh stream sediment, (C) the ash of streambank sod (mixed organic and inorganic material) collected beneath the water level, and (D) the ash of aquatic bryophytes (mosses). The copper data are plotted on base maps showing generalized geology and the drainage pattern. The map symbols show the sample sites and ranges of values in the following manner: (1) open symbols denote background, (2) small black symbols represent weakly anomalous values, and (3) large black symbols denote strongly anomalous values. Because the small black symbols represent weakly anomalous values, they are considered to be significant only where they are closely associated with strongly anomalous metal values either in the same sample medium or with anomalous values in other sample media. The ranges of values represented by the symbols are shown on the histograms which accompany the geochemical maps. An explanation of sampling, preparation, and analytical procedures is given in Circular 734, which accompanies this folio. Complete analytical data for geochemical samples collected by the U.S. Geological Survey in the Tanacross quadrangle are available in a U.S. Geological Survey open-file report (O'Leary and others, 1976).

Of the four sample media, the oxide residue (mainly secondary iron-manganese oxides) of stream sediment and the aquatic bryophytes act as scavenging agents of ions in solution in the stream waters. The copper content of these media, therefore, is indicative of the amounts of copper migrating in solution from bedrock and colluvium. The copper content of the streambank sod represents both copper scavenged from solution by the organic material and the copper content of the detrital material in the sod. The copper content of the minus-80-mesh stream sediment, on the other hand, mainly represents the amount of copper within the detrital material of the stream sediment.

Copper values in the ash of streambank sod show a relatively high positive correlation with the organic content of the sample. This high correlation suggests that the amount of organic material noticeably influences the copper content of the sod. A regression analysis—log copper vs. organic content—was used to determine the influence of organic content on the variation of copper values in the ash of the sod. This type of analysis allows separation of those high copper values that reflect the concentration of background amounts of copper by organic material from those high values that are derived from a mineralized source. Values from the regression analysis--shown as residuals--were used on the geochemical map (fig. C). The distribution of the residuals is shown on the upper of the two accompanying histograms. The lower histogram shows the distribution of original copper concentrations in the ash of the streambank sod.

The copper values in the ash of aquatic bryophytes were not adjusted on the basis of percent of organic material because the organic content of the bryophytes shows little variation. The histograms and other statistical data for copper in the oxide residue of stream sediment (fig. A) and in the minus-80-mesh stream sediment (fig. B) show two populations. One population (generally lower values) represents the copper content of the samples collected in the maturely dissected, forested terrain of the Yukon-Tanana Upland--that part of the quadrangle north of the Tanana River. The other population of generally higher copper values represents samples collected in the rugged, mountainous terrain of the Alaska Range--south and west of the heavy black line on the map. In the maturely dissected terrain, chemical weathering is probably the main factor controlling the mobility of copper. This type of weathering may be characterized by the solution of sulfide and other minerals and a general dispersion and impoverishment of copper and other base metals in the weathering zone. In the rugged mountainous terrain, on the other hand, mechanical weathering is the primary process controlling element dispersion. In this environment, impoverishment of metals in the weathering zone due to chemical processes is a minor factor.

In the terrain north of the Tanana River, anomalous copper values in the oxide residue of the stream sediment (fig. A), and to a lesser degree in the other three sample media, delineate known mineralized zones and several potential areas of mineralized rock. The most prominent anomalous area is near the east border of the quadrangle (fig. A). Several of the anomalous values in the northern part of the zone represent samples collected in streams draining an area where there is at least one porphyry copper prospect—in T. 22 N., R. 21 E. In this area, scattered anomalous copper values in the minus—80—mesh stream sediment (fig. B), and in the ash of streambank sod (fig. C) and of aquatic bryophytes (fig. D), roughly correlate with the anomalous values in the oxide residue of the stream sediment. The scattered anomalous copper values in the oxide residue south of this area suggest the presence of additional zones of mineralized

A cluster of high copper values in the ash of streambank sod (fig. C) indicates an anomalous zone in the west-central part of the quadrangle. Scattered anomalous copper values in the other sample media in the vicinity of the high copper values in sod ash suggest a rather extensive area that should be explored further. Anomalous molybdenum, lead, zinc, and arsenic values in this same general area (Curtin, Day, Carten, Marsh, and Tripp, 1976; Curtin and others, 1976b, C; Curtin, O'Leary, and Carten, 1976) give further evidence of mineralization in this part of the quadrangle.

In that part of the quadrangle north of the Tanana River, six base metal prospects were not defined by high copper values in the four sample media. These prospects are located in T. 24 N., R. 10 E.; T. 21 N., R. 14 E.; T. 21 N., R. 16 E.; T. 18 N., R. 15 E.; T. 16 N., R. 18 E.; and in T. 24 N., R 20 E. The surface features of at least two of the prospects are similar to those of two occurrences of porphyry copper mineralization that were outlined by high copper values in one or more of the sample media. The absence of high copper values around these prospects, however, indicates either that the copper content of the altered and mineralized rock is low or that the amount of mineralized rock is too small to produce copper-bearing dis-

The results demonstrate that copper occurrences are more completely defined by the use of a combination of sample media than by any one of the sample media when used alone.

persion trains that could be detected at the sampling density used in this study.

Patterns defining areas of copper potential are shown on the composite geochemical map of copper and molybdenum distribution (Curtin and others, 1976a), which is included in this folio.

REFERENCES CITED


Curtin, G. C., Day, G. W., Carten, R. B., Marsh, S. P., and Tripp, R. B., 1976, Geochemical maps showing the distribution and abundance of molybdenum in the Tanacross quadrangle, Alaska: U.S. Geol. Survey Misc. Field Studies Map MF-767G, Curtin, G. C., Day, G. W., O'Leary, R. M., Marsh, S. P., and Tripp, R. B., 1976a, Composite geochemical map of anomalous copper and molybdenum distribution in the Tanacross quadrangle, Alaska: U.S. Geol. Survey Misc. Field Studies Map MF-767M, 1 sheet, scale 1:250,000.

______1976b, Geochemical maps showing the distribution and abundance of lead in the Tanacross quadrangle, Alaska: U.S. Geol. Survey Misc. Field Studies Map MF-767H, 1 sheet, scale 1:500,000.


_____1976c, Geochemical maps showing the distribution and abundance of zinc in the Tanacross quadrangle, Alaska: U.S. Geol. Survey Misc. Field Studies Map MF-767I, 1 sheet, scale 1:500,000. Curtin, G. C., O'Leary, R. M., and Carten, R. B., 1976, Geochemical maps showing the distribution and abundance of arsenic in the Tanacross quadrangle, Alaska: U.S. Geol. Survey Misc. Field Studies Map MF-767J, 1 sheet, scale 1:500,000.

Foster, H. L., 1970, Reconnaissance geologic map of the Tanacross quadrangle, Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-593, scale 1:250,000. O'Leary, R. M., McDanal, S. K., McDougal, C. M., Day, G. W., Curtin, G. C., and Foster, H. L., 1976, Spectrographic and chemical analyses of geochemical samples and related data from the Tanacross quadrangle, Alaska: U.S. Geol. Survey

OF TOTAL SAMPLES COPPER, IN RESIDUALS (log Cu vs. % organic) Standard deviation = .175 Correlation coefficient
Cu vs. % organic = .5350 Number of samples = 594



A. Copper in the oxide residue of stream sediment



SYMBOL AND PERCENTAGES OF TOTAL SAMPLES

15 20 30 50 70 70 100 100 700 700 700 8000 COPPER, IN PARTS PER MILLION

YUKON-TANANA UPLAND

Arithmetic mean = 188

Standard deviation = 259

Geometric mean = 150 Geometric deviation = 1.81

Number of samples = 519

SYMBOL AND PERCENTAGE OF TOTAL SAMPLES

15 20 30 50 70 70 70 500 500 500 500


COPPER, IN PARTS PER MILLION

ALASKA RANGE Arithmetic mean = 398

Standard deviation = 284 Geometric mean = 312


Geometric deviation = 2.04 Number of samples = 81

 <10
 10
 20
 30
 40
 40
 50
 60
 60
 90
 90
 110
 120
 130 COPPER, IN PARTS PER MILLION YUKON-TANANA UPLAND Arithmetic mean = 13.9 Standard deviation = 8.4 Geometric mean = 12.2 Geometric deviation = 1.64 Number of samples =499 SYMBOL AND PERCENTAGE OF TOTAL SAMPLES 7 10 10 10 20 20 20 20 70 70 70 00 COPPER, IN PARTS PER MILLION ALASKA RANGE Arithmetic mean = 32 Standard deviation = 16 Geometric mean = 28.4 Geometric deviation = 1.67



D. Copper in the ash of aquatic bryophytes (mosses)

C. Copper in the ash of streambank sod



B. Copper in the minus-80-mesh stream sediment


BASE FROM U. S. GEOLOGICAL SURVEY, 1:250,000, TANACROSS QUADRANGLE, 1964

QUADRANGLE LOCATION

Scale 1:500,000 1 inch equals approximately 8 miles

Number of samples = 80





GEOCHEMICAL MAPS SHOWING THE DISTRIBUTION AND ABUNDANCE OF COPPER IN THE TANACROSS QUADRANGLE, ALASKA

G. C. CURTIN, G. W. DAY, R. M. O'LEARY, S. P. MARSH, AND R. B. TRIPP

CHARGE FROM THE U.S. GEOLOGICAL SURVEY, RESTON, VA. 22092

laska (Tanacross Guid.). Copper



1976

For sale by U. S. Geological Survey, price \$.50

BACKGROUND INFORMATION RELATING TO THIS MAP IS PUBLISHED

AS U.S. GEOLOGICAL SURVEY CIRCULAR 734, AVAILABLE FREE OF