

MAP B.-ZINC IN THE MAGNETIC FRACTION

CORRELATION OF MAP UNITS

Qtg	Qtb	Tcv	QUATERNARY
Tf	Tfm	Tt	TERTIARY
Tkt			
Ksv			CRETACEOUS
Pps			PERMIAN AND PENNSYLVANIAN
Mcs			MESSIPIAN TO CAMBRIAN
Yd			MIDDLE PROTEROZOIC
Yf			
Ym			EARLY PROTEROZOIC

[illegible]

STRIKE AND DIP OF BEDS

$\swarrow 5$ Inclined
+ Vertical

STRIKE AND DIP OF FOLIATION--includes primary flow
foliation of volcanic rocks and secondary
metamorphic foliation of metamorphic rocks

$\swarrow 50$ Inclined
+ Vertical

mineral concentrations, thereby providing clues to regional patterns of mineralization; (2) showing new areas containing enriched metals, not previously known from mining activity; and (3) providing supportive data to the interpretive and summary maps (Watts and Hassemer, 1986) and the mineral resource assessment (Richter and others, 1983). Background information that pertains to published components of the Silver City folio is also available (Richter and others, 1983).

The quadrangle lies almost entirely within the southern Basin and Range province, an area of north- to northwest-trending mountain ranges and intervening basins (fig. 1). The small area in the northeastern part of the quadrangle, characterized by horizontal bedrock and alluvium, is in the transition zone between the Basin and Range and Colorado Plateaus provinces. The westerly flowing Gila River, and

consist of diverse rocks ranging in age from Proterozoic through Cenozoic. The four main groups of rocks exposed the ranges are, from oldest to youngest: (1) Proterozoic metamorphic and plutonic rocks; (2) Paleozoic and Mesozoic sedimentary rocks; (3) Cretaceous and early Tertiary volcanic and plutonic rocks; and (4) middle Tertiary volcanic and plutonic rocks. The mountain ranges are separated by structural and topographic basins filled with late Tertiary to Holocene sedimentary rocks and

A map of the Arizona Plateau Province, showing the location of the study area. The map includes labels for 'ARIZONA', 'PLATEAUS PROVINCE', and 'TRANS'. A dashed line indicates the boundary of the study area.

Figure 1.--Index map of Arizona and New Mexico showing the location of the study area (stippled) and some physiographic features.

**GEOCHEMICAL MAPS
ZINC IN TWO FRACT
SILVER CITY 1° X 2**

In most cases the material analyzed consisted of two fractions (nonmagnetic and magnetic) of heavy (specific gravity greater than 2.8) minerals separated from the bulk alluvial sediments by density contrast (gold pan and heavy liquids) and released by dissolution of the heavy residue. The nonmagnetic fraction, however, the nonmagnetic fraction contained most of the detectable or enriched concentrations and therefore is the only fraction shown for these elements. Methods of collecting and processing samples are discussed in detail by others (1979; Watts and Hassner, 1986; Watts and others, 1984). The samples were examined microscopically after initial preparation, and were then pulverized to a fine powder and analyzed for 30-31 elements by emission spectrography (Grimes and Mazzancino, 1968; McNeal and others, 1983).

Microscopic scans of the samples prior to their treatment with HF revealed the presence of the nonmagnetic fraction, regardless of tectonic terrane. The zircon grains in the samples from the volcanic terrane were characterized by a wide range of sizes (from 10 to 100 μm) and shapes (from rounded to elongated) and were associated with magneite and/or containing black inclusions, often in both volcanic and preterrestrial terrane. The zircon grains in the samples from the volcanic aggregate strata (intergrowths) of various minerals such as quartz, feldspar, and calcite were characterized by a wide range of sizes (from 10 to 100 μm) and shapes (from rounded to elongated). Concentrates from regionally metamorphosed rocks (metagabbro, amphibolites, and metabasites) contained abundant shandite and muscovite and lesser amounts of zircon. The zircon grains in the samples from the tectonized zone (Isken) in all types of pre-terrestrial terrane usually contained abundant shandite and muscovite and lesser amounts of zircon. The zircon grains in the samples from the tectonized zone (Isken) in all types of pre-terrestrial terrane usually contained abundant shandite and muscovite and lesser amounts of zircon. The zircon grains in the samples from the tectonized zone (Isken) in all types of pre-terrestrial terrane usually contained abundant shandite and muscovite and lesser amounts of zircon.

INTERPRETIVE IMPLICATIONS OF THE HEAVY-MINERAL FRACTIONS

In the magnetic fraction, high element contents are mostly associated with the various oxides and oxyhydroxides of iron and manganese. The sources of these element-enriched detrital grains are diverse but undoubtedly

The statistical distribution of the data is shown on the histogram (fig. 2). The frequency distributions of the two fractions of heavy-mineral concentrates are superimposed on each other on the histogram to facilitate their comparison, which is useful for interpretations. The symbols used on the maps correspond to class intervals of

Drewes, Harald, Houser, B. B., Hedlund, D. C., Richter, D. H., Thorman, C. H., and Finnell, T. L., 1985, Geology map of the Silver City 1° x 2° quadrangle, New Mexico and Arizona, U.S. Geological Survey Miscellaneous Publication 85-1.

Figure 1

CONTOUR INTERVAL 200 FEET
NATIONAL GEODETIC VERTICAL DATUM OF 1929

Hassmer, J. R., Ficklin, W. H., Motoko, J. M., and Watts, K. C., with contributions from Preston, R. L., Seaple, S. M., and Ward, F. N., 1983, Analytical results for 325 water samples from the Silver City 1° x 2° quadrangle, Arizona and New Mexico: U.S. Geological Survey Open-File Report 83-10.

McDonald, J. R., Fourn, C. E., Hassmer, J. R., and Watts, K. C., 1983, Analytical results for stream-sediment concentrates, sieved stream-sediment, rock, and water samples from the Silver City 1° x 2° quadrangle, New Mexico, Arizona, and New Mexico: U.S. Geological Survey Open-File Report 83-10, 220 pgs. [Computer tape available from U.S. Department of Commerce, National Technical Information Service, Springfield, VA 22161.]

Richter, J. D., Houser, C. K., Kline, D. C., Kline, D. P., Sharp, W. M., Drewes, Harold, Hedlund, D. C., Raines, G. L., and Hassmer, J. R., in press, The Continental

Richter, D. H., Sharp, W. N., Watts, K. C., Reines, G. L., Mouser, B. R., and Klein, D. P., 1983, Mineral resource assessment of the Silver City 1° x 2° quadrangle, New Mexico and Arizona: U.S. Geological Survey Open-File Report 83-924, 77 p.

VanTrump, George, Jr., and Miesch, A. T., 1977, The U.S. Geological Survey RASS-STATPAC system for management and statistical reduction of geochemical data:

Investigations Series Map 1-1310-B, 2 sheets.
Watts, K. C., Hassner, J. R., Forn, C. L., and Stens, D. F., 1986a, Geochemical maps showing distribution and abundance of lead in two fractions of stream-sediment concentrates, Silver City 1" x 2" quadrangle, New Mexico and Arizona: U.S. Geological Survey Miscellaneous Field Studies Map MF-1183-A, scale 1:250,000.
1986b, Geochemical maps showing distribution and

Miscellaneous Field Studies Map MF-1183-E, scale 1:250,000.
 1986d, Geochemical maps showing distribution and abundance of silver in two fractions of stream-sediment concentrates, Silver City 1" x 2" quadrangle, New Mexico and Arizona: U.S. Geological Survey
 Miscellaneous Field Studies Map MF-1183-F, scale 1:250,000.
 1986e, Geochemical maps showing distribution and

Geological Survey Miscellaneous Field Studies Map MF-1183-H, scale 1:250,000.
 1966g, Geochemical maps showing distribution and abundance of tin in two fractions of stream-sediment concentrates, Silver City 1° x 2° quadrangle, New Mexico and Arizona: U.S. Geological Survey Miscellaneous Field Studies Map MF-1183-I, scale 1:250,000.
 1966h, Geochemical maps showing distribution and

Watts, K. C., Hessemer, J. R., and Stems, D. F., 1984, Geochemical halos in the Silver City mining region and adjacent areas, Grant County, New Mexico: U.S. Geological Survey Bulletin 1534, 85 p.

CLASS INTERVAL

[illegible]

rogram showing distribution of zinc. Symbols shown within each interval of reported element concentrations corresponds with symbol map. Nonmagnetic fraction is shaded; magnetic fraction is unshaded. Number below class interval symbol is number of samples in nonmagnetic fraction; lower number is number of samples in the magnetic fraction.


LIBRARY

ANCE OF
STRATES,

U.S. LIBRARY - WESTON
3 1818 00210916 1

643027

M
M



By _____