

**EXPLANATION**

Mineral resource tracts -- Numbers refer to descriptions in table 2


Sub-tracts -- Letters refer to descriptions in table 1

Epithermal gold veins

Porphyry copper

Polymetallic vein

Base from U.S. Geological Survey  
Port Moller, 1953 (revised 1988);  
Simeonof Island, 1953; Steповak Bay,  
1963 (revised 1981)  
Universal Transverse Mercator projection



**DESCRIPTION OF MAP UNITS**

Geology by Frederic H. Wilson, Robert L. Detterman, and Ellen E. Harris. Stratigraphic nomenclature and age assignment from Detterman and others. In press.

**SURFICIAL DEPOSITS AND SEDIMENTARY ROCKS**

**Qs Surficial deposits (Holocene and Pleistocene)**—Unconsolidated, poorly to well-sorted, poorly to moderately well stratified sand, gravel, and silt. Also includes silvial, colluvial, glacial, marine, lacustrine, and eolian deposits. Locally includes reworked volcanic ash and debris-flow deposits.

**Tm1 Milky River Formation (Pliocene)**—Volcanogenic near-marine sedimentary rocks and interlayered flows and sills. Upper part of unit contains numerous porphyritic andesitic lava flows, lahar deposits, and tuff beds interlayered with sedimentary rocks. Lower part consists almost entirely of coarse-grained, highly crossbedded and channelled, fluvial volcanic sandstone and cobble-bearing conglomerate that has clastic composition mainly of volcanic debris.

**Tb1 Bear Lake Formation (late Miocene)**—finer, territic marine and nonmarine (Wright, 1971; Niles, 1984) sandstone, conglomerate, siltstone, and shale. Sandstone is moderately well sorted and grains are moderately well rounded. Conglomerate horizons are made up of well rounded clasts, of which about 40 to 50 percent are quartz and chert, 20 to 30 percent are volcanic fragments, 10 to 15 percent are granitic clasts, and remainder is lithic sedimentary clasts. Unit is abundantly fossiliferous.

**Tu Unga Formation (middle Miocene to late Oligocene)**—Volcaniclastic sedimentary dominant in upper part, whereas carbonaceous shale and coal are confined to debris and are poorly consolidated. Many conglomeratic beds are lahar deposits and are poorly consolidated. Many conglomeratic beds are lahar deposits and are poorly consolidated. Many conglomeratic beds are lahar deposits and are poorly consolidated.

**Tbc Bekolka Formation (middle and early Miocene? and late Oligocene?)**—Mainly tuffaceous, volcaniclastic sandstone, siltstone, and conglomerate and interbeds of tuff and volcanic breccia (McLean, 1979). Rocks are dominantly red, pink, and purple and very well indurated. Potassium-argon dates from volcanic breccia on Dagle Island is 11.72±0.41 Ma (sample 85AP9; 50k; table 2, pamphlet); if mapping is correct, then some of the Bekolka is of Miocene age.

**Ts Steповak Formation (early Eocene)**—Upper part is olive-gray and yellowish-brown sandstone rich in unsorted volcanic debris and silt and lower part is dark, commonly laminated siltstone and shale deposited as a deep-water turbidite and Megalaura distributed throughout upper part is characteristic of water depths no greater than 30 to 50 m (Louie Marincovich, Jr., written commun., 1983-1986).

**Tl1 Tledet Formation (middle Eocene to late Paleocene)**—Pale-yellowish-brown to gray-green interbedded sandstone, conglomerate, siltstone, and shale. Lithic clasts in conglomerates are dominantly andesitic and volcanic debris that contains 20 to 30 percent volcanic clasts. Most volcanic clasts are not fresh Mesozoic source rather than contemporaneous magmatic activity. In sharp contrast to most overlying units (Detterman and others, in press). In type area just east of Pavlof Bay, characteristic lithologies include shallow marine mainly of brachiopod-strewn type, that are typical for the major part of Tolsta Formation.

**Kc Chiglik Formation (late Cretaceous)**—Dominantly a light-olive gray to olive gray sandstone and interbedded olive gray to olive-black siltstone and conglomerate of multicolored chert, white quartzite, and minor volcanic rocks. Unit is cyclic nearshore marine, tidal-flat, nonmarine flood-plain, and fluvial deposits (Parchild, 1977; Detterman, 1978). Nonmarine parts may contain coal beds as much as 2 m thick. Marine fossils, mainly pelecypods, indicate a late Campanian to early Maestrichtian age (J.W. Miller, written commun., 1983-88).

**Kh Hoodoo Formation (late Cretaceous)**—Typically dark-gray to black, rhythmically sandstone. Becomes more sandy upward, and at Hoodoo Mountain contains quartz. Sandstone beds range from 0.5 to 1 m thick and siltstone and shale beds range from 1 to 2 m thick and have individual layers as thin as 1 cm (Detterman and others, 1981). Sparse megafossils indicate an age of late Campanian to early Maestrichtian (J.W. Miller, written commun., 1983-88). Depositional environment is characteristic of lower to upper slope of a submarine fan; structures imply submarine slumping and turbidly current flow.

**Volcanic rocks (late Miocene)**—Andesite, basalt, and dacite lava flows, sills, and plugs. Intrusive rocks typically cap ridges and include massive flows, agglomerate, and lahar deposits. Minor porphyritic alteration is characteristic, except near San Diego Bay where sericitic and argillite alteration is pervasive. Hypabyssal rocks are of similar composition and were very shallowly emplaced. Potassium-argon ages range from 10.40±0.49 to 6.07±0.23 Ma (Wilson and others, 1994).

**Tm Meshik Volcanics (early Oligocene and late Eocene)**—Well exposed in mountains north of Iliof Bay and consists of basalt and andesite lava flows, coarse volcanic rubble, lahar deposits, tuff, and minor volcaniclastic sedimentary rocks. Volcanic sedimentary rocks are temporally and lithologically equivalent to the Steповak Formation.

**Tp Popof volcanic rocks (early Oligocene and late Eocene)**—Lava flows, lahar deposits, debris flows, ash-flow tuff, and tuff on Popof, Unga, Korovin, and Andronia Islands. Lithologically and temporally equivalent to the Meshik Volcanics, but units crop out spatially separate from the Meshik Volcanics and is indicative of local volcanic sources; hence, its separate informal designation. Gradationally interfingers with the Steповak Formation.

**INTRUSIVE ROCKS**

**Qh1 Younger intrusive rocks, undivided (Holocene and Pleistocene)**—Hypabyssal dacite plugs and domes at Quaternary volcanic centers, particularly Trader Mountain and Mount Dana. Porphyritic dacite at Trader Mountain has a K-Ar age of 0.98±0.06 Ma (see table 2, pamphlet).

**T1 Older intrusive rocks, undivided (Pliocene and late Miocene)**—Medium to coarse-grained, equigranular granodiorite to quartz diorite plutons and stocks containing hornblende, biotite, and pyroxene. Typically display well-developed hornblende zones and sporadic hydrothermal alteration in surrounding country rocks. Potassium-argon ages range from 9.43±0.26 to 3.21±0.14 Ma (see table 2, pamphlet).

**Tm1 Intrusive rocks, undivided (Tertiary)**—Small intrusive bodies of quartz diorite or diorite, which are typically hypabyssal and contain phenocrysts of pyroxene or hornblende in a fine-grained groundmass.

**Tg Granitic rocks (Paleocene–Miocene)**—Medium-grained biotite granodiorite and quartz monzonite plutons that have hydrothermal granular textures and local development of potassic feldspar phenocrysts as much as 1 cm in size. Crop out on Nagai, Big and Little Konuih, Simeonof, Chernaubura and Bird Islands. Potassium-argon ages (Bark, 1985; Moore, 1974; Kettle and Turner, 1976) range from 6.5±3.5 to 57.9±1.8 Ma (recalculated using constants of Steiger and Jager, 1977).

**METAMORPHIC ROCKS**

**QTC Contact-metamorphosed rocks (early Quaternary or late Tertiary)**—Contact metamorphosed and hydrothermally altered rocks in mountain east of Mount Dana. Rocks probably consist of the Hoodoo Formation (R1), undivided Tertiary volcanic rocks (Tm), and Chiglik Formation (Kc). Unit is well indurated, very fine grained, and intensely fractured; sulfide mineralization is common, and resultant iron staining is ubiquitous on weathered surfaces.

**— Fault**—Dashed where approximately located; dotted where concealed; queried where uncertain.

**— Thrust or high-angle reverse fault**—Dashed where approximately located; dotted where concealed; queried where uncertain. Southwest on upper plate.

**Volcanic crater**

See pamphlet for references cited.

**MAPS SHOWING THE RESOURCE ASSESSMENT OF THE PORT MOLLER, STEPOVAK BAY, AND SIMEONOF ISLAND QUADRANGLES, ALASKA PENINSULA**

By  
**Frederic H. Wilson, Willis H. White, Robert L. Detterman, and James E. Case**

1996

Geology mapped by L.M. Angelson, 1983-85; J.E. Case, 1977-78, 1983-85; C.L. Connor, 1982; John Decker, 1985; R.L. Detterman, 1977-78, 1982-86; G.D. DuBois, 1985; B.M. Gamble, 1983-84; Stephanie Zuranski, 1986; Louie Marincovich, Jr., 1982, 1984; J.W. Miller, 1982-85; M.A. Pomonis, 1983; Nora Shew, 1982; F.R. Weber, 1983-85; F.H. Wilson, 1977-78, 1982-86; M.E. Yeart, 1977-78, 1982-85

Edited by C.L. Ostergren; prepared by Tom Chase  
Manuscript approved for publication February 9, 1994