

U.S. DEPARTMENT OF THE INTERIOR

U.S. GEOLOGICAL SURVEY

GEOLOGIC MAP OF THE CRAIG, DIXON ENTRANCE,  
AND PARTS OF THE KETCHIKAN AND PRINCE RUPERT QUADRANGLES,  
SOUTHEASTERN ALASKA

Compiled by

DAVID A. BREW

TO ACCOMPANY MISCELLANEOUS FIELD STUDIES MAP MF-2319

1996

## CONTENTS

	Page
Introduction	3
Descriptions of lithotectonic terranes	7
Mineral resources	10
Description of map units	10
References	31
Index to map units, by name	44

## ACCOMPANYING ILLUSTRATIONS

Map sheet: Geologic map of the Craig, Dixon Entrance, and parts of the Ketchikan and Prince Rupert quadrangles, southeastern Alaska; includes:

Correlation of Map Units

Brief Description of Map Units

Figure 1. Index map showing location of study area and physiographic provinces and sections (from Wahrhaftig, 1965).

Figure 2. Sources of geologic information, 1983-1995.

Figure 3. Lithotectonic terrane and major fault map of southeastern Alaska and adjacent regions. Dashed line is boundary of the Craig, Dixon Entrance, and parts of the Ketchikan and Prince Rupert quadrangles map area (after Brew and others, 1991a, b, 1992a). Major faults are labeled as follows: BR, Border Ranges; CHS, Chatham Strait; CLS, Clarence Strait; CRML, Coast Range megalineament; FQ, Fairweather-Queen Charlotte; NA, Nahlin, NS, Neva Strait; THL, Tally Ho-Llewellyn; and TR, Transitional.

# **GEOLOGIC MAP OF THE CRAIG, DIXON ENTRANCE, AND PARTS OF THE KETCHIKAN AND PRINCE RUPERT QUADRANGLES, SOUTHEASTERN ALASKA**

**Compiled By**

**David A. Brew**

## INTRODUCTION

### BACKGROUND AND PURPOSE

This map and report covers the southwestern part of southeast Alaska (fig. 1). The geology of this area is critical to the understanding of the whole region because here are found the oldest isotopically dated rocks and the region's most complete, most fossiliferous, most varied, and relatively unmetamorphosed and undeformed Paleozoic stratigraphic section. The rocks in the eastern part of the area include features transitional to the tectonic, magmatic, and metamorphic character of the Coast Mountains geology of southeastern Alaska.

This compilation was stimulated (1) by the need for a coherent geologic map of the area for use in the study of the undiscovered mineral resources of the Tongass National Forest and adjacent lands (Brew and others, 1991a; Brew and Drinkwater, 1991; Brew and others, 1992a); (2) by the related requirements that (a) correlative units mapped by different recent workers be examined and summarized, and (b) newly available age information from fossil determinations and isotopic dating be integrated into the descriptions of the map units; and (3) by the need for a geologic base map for use in interpretation of the geochemical study of the Craig study area (Cathrall and others, 1993a, b; Cathrall, 1994). It brings together several regional geologic studies which differ somewhat in their treatment of the geology of the Craig, Dixon Entrance, and the western parts of the Ketchikan and Prince Rupert quadrangles. The compilation was made to reconcile these differences.

In regard to point 2b, above and to stratigraphic nomenclature in general: As part of this compilation process I have examined both the fossil and isotopic evidence used by the geologists who produced the maps listed in the following paragraph and the new fossil information that has been published in journals since some of those maps were produced. From this synthesis came age assignments for some formal geologic units that differ in part from previous assignments. The evidence for these changes is cited in every case. Inasmuch as the age of a unit is not a criterion in the definition of lithostratigraphic units (North American Commission on Stratigraphic Nomenclature, 1983, Article 22(e), p. 856), any such changes in age are not a formal stratigraphic problem. Nor is the extension of such an age assignment from the area where to rocks have been dated into areas where the same lithostratigraphic unit is present, but not dated. In some cases I have extended the use of formal names beyond where they have been previously applied, into areas where informal unit names have been used in the past. In all such cases I followed the published suggestions of recent mappers in the area. No rock units with previously applied formal names have been assigned to different units with different names. Informal names are applied to both metamorphic and plutonic lithodemic units to facilitate organization and presentation.

The compilation is based mainly on: (1) the 1:250,000-scale geologic map of the Petersburg area based on 1978 to 1983 mapping by Brew and others (1984); (2) the 1:250,000-scale geologic map of the Craig quadrangle based on 1950 to 1971(?) mapping by Eberlein and others (1983); (3) the 1:250,000-scale geologic map of the Ketchikan and Prince Rupert quadrangles based on 1970(?) to 1977 mapping by Berg and others (1988); (4) the several different reports and maps by G.E. Gehrels and J.B. Saleeby that are cited below; (5) the 1:200,000-scale geologic map of Annette Island by Karl (1992) and Horton and others (1992); and on unpublished mapping by J.F. Baichtal of the U.S. Forest Service. The areas covered by these different reports are shown on figure 2.

## LOCATION AND PHYSIOGRAPHY

The map area covers the southwesternmost part of the Alexander Archipelago (fig. 1). It includes most of Prince of Wales Island, the third largest island (after Kodiak and Hawaii) of the United States, and an abundance of smaller islands of greatly differing size. Steep, forested mountains; deep, broad valleys; lakes and streams; and many tidewater bays and coves dominate the landscape. Most of the lower parts of the area are covered by heavy forest and brush, with minor muskegs and meadows. It rains a lot.

Physiographically, the map area (fig. 1) consists of parts of the Coastal Trough and Pacific Border Ranges provinces of Wahrhaftig (1965). The Coastal Trough province is a series of lowlands that extend the length of the Pacific Mountain System, interrupted along its length by mountain groups. The Pacific Border Ranges province consists mainly of several mountain ranges bordering the Pacific Coast. Within the map area, the Coastal Trough province is represented by part of the Kupreanof Lowland and the Pacific Border Ranges province is represented by the Prince of Wales Mountains.

The Kupreanof Lowland consists of islands, channels, and adjacent areas. Most of it is rounded heavily glaciated terrain with a local relief of 300-500 feet (90-160 m) and a maximum relief of 1,000-1,500 feet (310-470 m) separated by complex waterways. Some blocklike mountains that rise above the general level have summits at 2,000-3,000 feet (620-930 m). The Lowland is characterized by many lakes.

The Prince of Wales Mountains consists of moderately rugged glaciated mountains with rounded summits 2,000- 3,500 feet (620-1,090 m) high. The high point of the map area is Pin Peak, 3,806 feet (1,185 m) high, about 10 miles (16 km) east of Klawock. The mountains are dissected by steep-walled U-shaped glacial valleys and by several fiords 600-1,000 feet (190-310 m) deep. Several passes less than 500 feet (160 m) high cross the range. Lakes are abundant and karst topography and locally extensive cave systems occur in localities underlain by carbonate rock. The east and northeast front of the range is steep and abrupt against Clarence Strait, but the north and northwest front against the Kupreanof Lowland is not as distinct (Wahrhaftig, 1965). The karst topography of this province has been the focus of intense cave exploration in recent years and one cave has been found to contain very significant bear remains (Heaton and Grady, 1992a,b, 1993; Heaton, 1995).

There are five larger communities and several smaller ones in the map area. The main ones are the ferry terminal community of Hollis, the logging village of Thorne Bay; the fishing town of Craig; the fishing village of Klawock, which is populated mostly by Tlingit Indians; and the fishing village of Hydaburg, which is populated mostly by Haida Indians. There are also numerous abandoned logging camps, mining and quarry camps, and Indian village sites. Logging, fishing, mineral exploration, and tourism are the major economic activities in the area. The Alaska Marine Highway System connects Hollis by ferry with Ketchikan to the east of the map area. An extensive network of U.S. Forest Service roads that are now part of the State of Alaska highway system cover the central part of Prince of Wales Island (U.S. Forest Service, 1987).

Most of the map area is part of the Tongass National Forest, but significant areas are Alaska Native Regional and Village Corporation, State of Alaska, and private lands, and scattered patented mining claims (U.S. Forest Service, 1987). Five Wilderness Areas within the Tongass Forest are administered by the U.S. Forest Service (1987) and one National Wildlife Refuge on Forrester Island is administered by the U.S. Fish and Wildlife Service (Brew and Drinkwater, 1991).

## HISTORY OF GEOLOGIC AND RELATED STUDIES

Geologic investigations in the map area fall into several roughly defined periods. The first was during the latter part of the 1800's when Prince of Wales Island and adjacent areas received early attention in the form of prospecting and sporadic reconnaissance studies. The first systematic geologic mapping and mineral-deposit studies were those of Theodore Chapin in the early 1900's which are reported in Buddington and Chapin (1929), the Wright brothers' studies (Wright and Wright, 1908; Wright, 1915), and the extensive mapping of Buddington himself (Buddington and Chapin, 1929).

Mineral exploration, mining, and the quarrying of marble (Burchard, 1913, 1920) and limestone (for cement) continued in the 1800's and 1900's up to World War II, with some interruption during World War I (See references to local U.S.G.S studies in the compilations of Cobb). Immediately following World War II, U.S. Geological Survey parties visited and studied several metallic-mineral-deposit localities, some in detail (Robinson and Twenhofel, 1953; Kennedy, 1953; Warner and others, 1961).

Somewhat later U.S.G.S. studies (Rossmann and others, 1956; Sainsbury, 1961; MacKevett, 1963) were concerned with geophysics, areal mapping, and mineral deposits. In the late 1950's through the 1970's E.H. Cobb (Cobb, 1972a, b, c, d; 1978a, b; Cobb and others, 1968; Cobb and Elliott, 1980) compiled mineral-deposit information from all available sources. Condon (1961) compiled a reconnaissance geologic map of the Craig quadrangle using both available previous mapping and aerial photographs. This same period saw the first intense work on the Alaskan-type mafic-ultramafic plutons in and close to the map area (Ruckmick and Noble, 1959; Taylor, 1967; Irvine, 1974)

The next roughly defined period of U.S.G.S. and other studies was in the late 1960's, but the effort really started with the 1950's work of Eberlein in the western Prince of Wales Island area; this series of studies emphasized regional geologic, geophysical, and geochemical mapping (Barnes and others, 1972a, b; 1975; Berg, 1972, 1973, 1978, 1980, 1982; Berg and others, 1976, 1978a, b, 1988; Clark and others, 1971; Clark and Greenwood, 1972; U.S. Geological Survey, 1977, 1979). During this same general period State of Alaska parties were doing detailed studies of selected mineral deposits (Terreid and others, 1978) and an extended series of paleontologic studies was started (Armstrong, 1970; and the numerous references of Churkin, Savage, and Soja) that continues to the present. The original work of Eberlein, together with that of several colleagues, was released in Eberlein and others (1983).

One result of these geologic studies was the inclusion of some new information in the regional-scale geologic map compilations of Beikman (1974, 1975). Other regional-scale results were the regional mineral-resource-oriented compilations of Berg and others (1981) and Berg (1984) and the aeromagnetic compilations of Decker (1979) and Decker and others (1981).

In the late 1970's and 1980's the U.S.G.S, through G.D. Eberlein and H.C. Berg, started supporting J.B. Saleeby of the California Institute of Technology in geologic and geochronologic studies of southern Prince of Wales Island. This led to the involvement and support of Saleeby's students G.E. Gehrels and C.M. Rubin; to further support by the National Science Foundation; and to the eventual publication of a series of important topical and regional studies by the California Institute of Technology group (Gehrels, 1980, 1991; Gehrels and others, 1983, 1987; Gehrels and Saleeby, 1986, 1987; Rubin and Saleeby, 1987, 1991, 1992; Saleeby, 1991, 1992). In addition, at least one thesis (Sullivan, 1991) of local significance has been produced.

In 1978, the U.S.G.S started a regional geologic mapping and mineral-resource assessment study in what was then the almost totally unmapped Petersburg project area, just north of the Craig quadrangle. This project required examining and extending as appropriate the stratigraphic units described by Ovenshine and Webster (1970), Eberlein and Churkin (1970a), Eberlein and others (1978), and Eberlein and others (1983) from the Craig map area. The major Petersburg project maps and reports to date are Brew and others (1984, 1989b), Grybeck and others (1984), Barnes and others (1989), and Karl and Koch (1990).

Following the completion of the Petersburg project, the U.S.G.S. initiated a geochemical, geophysical, and mineral-resource assessment study in the Craig and Dixon Entrance quadrangles. To date the results of the aeromagnetic survey (U.S. Geological Survey, 1984), a mineral-resource study (Gehrels and others, 1983), and the geochemical studies have been reported (McDanal and others, 1991; Detra and others, 1992; Cathrall and others, 1993a, b; Cathrall, 1994); and mineral-resource-assessment information provided (D.J. Grybeck, written commun., 1990) for the regional mineral-resource study of Brew and others (1991a). During the U.S.G.S. study, the U.S. Bureau of Mines studied some of the alkalic igneous rocks (Warner and Barker, 1989; Barker and Mardock, 1988, 1990) and also started an ongoing study of the identified mineral resources of the Ketchikan Mining District, which includes the study area (Maas and others, 1991; Clautice and others, 1994). Several mineral-exploration companies have been active in the Prince of Wales Island part of the study area for the past several years and are continuing their efforts. The results of a cooperative U.S. Geological Survey-Bureau of Indian Affairs geological and geophysical study of the mineral resources of Annette Island during this same period are reported by Karl (1992) and Horton and others (1992). Drinkwater and Celzina (1994) described the geochemistry of the alkalic plutons in the area and Himmelberg and Loney (1995) described several of the ultramafic-mafic intrusions in the area. More recently, J.F. Baichtal of the U.S. Forest Service has done detailed geologic mapping in selected areas, mostly on Prince of Wales Island.

Other regional-scale studies concerned in part with the map area are the geothermal compilation of Motyka and Moorman (1987); the U.S. Geological Survey (1989) Mineral Resources Data System compilation of mineral-deposit information; and the mineral-deposit-locality compilations of Nokleberg and others (1987, 1994a, b).

Most of the information mentioned above was evaluated and used in the U.S.G.S. study of the undiscovered mineral resources of the Tongass National Forest and adjacent lands (Brew and others, 1991a; Brew and Drinkwater, 1991; Brew and others, 1992a) and much of it was used by the U.S. Bureau of Mines in their study of the identified resources (Coldwell, 1990). These studies were available to the U.S. Forest Service for their land planning process and mineral-resource considerations are contained in the various land-use classifications proposed for the Craig, Dixon Entrance and adjacent areas (U.S. Forest Service, 1990, 1991).

Finally, as the compiler of this map, I have had some, but not extensive, field experience in the area. It consists of a reconnaissance trip to eastern Prince of Wales Island in 1967, geologic mapping in the Dall Island and Craig areas in 1969, lengthy reconnaissance and familiarization trips to Dall Island and western Prince of Wales Island in 1978 and 1979, geologic mapping in the northernmost part of the Craig quadrangle in 1978 to 1981 as part of the Petersburg project (Brew and others, 1984, 1989b), and geologic mapping in the vicinity of Ketchikan and Gravina Islands (Brew and Karl, 1988a, b, c; Ford and Brew, 1988, 1994). As part of the study of the undiscovered mineral resources of the Tongass National Forest and adjacent lands (Brew and others, 1991a, 1992a; Brew and Drinkwater, 1991; Brew and others, 1992a), I examined almost all of the available geologic, geochemical, and geophysical information in some detail.

### SOME FEATURES OF THIS REPORT

Three specific aspects of this report should be emphasized: one is the origin of the unit descriptions, and the second is the alphabetical index of map units at the end of the text, and the third is a *caveat* regarding the geology of the inland areas. These aspects are described briefly here, together with a note on the geologic-time divisions used in this report for the Silurian Period.

(1) As explained in the section of this report titled "Description of Map Units", those descriptions have been adapted directly from the citations given for each unit. In most cases the descriptions have been shortened and altered. In every case, however, the descriptions are those of the original workers.

(2) The "Brief Description of Map Units" section on the map sheet that includes the page numbers for the complete descriptions of the units and the alphabetically-arranged index (by map unit title) at the end of this text are intended to allow the reader to go back and forth between the text and the map with a minimum of page turning and searching.

(3) Finally, there were very few logging or other roads in the interior of any of the islands in the study area at the time that most of the geologic mapping was done. Contacts were located on the map using a high degree of intuition and photogeologic interpretation. Detailed follow-up mapping along the now-existent roads (U.S. Forest Service, 1987) would certainly result in many changes. Thus this map cannot be used to locate contacts between map units precisely in the previously inaccessible areas.

The geologic time series divisions of the Silurian used in this report are based on those proposed by Berry and Boucot (1970) for North America, and they are not the divisions in previous common use by the U.S. Geological Survey. However, some Geological Survey reports have already used the Berry and Boucot divisions (Hansen, 1991, p. 61). Those divisions are, from youngest to oldest, Pridoli, Ludlow, Wenlock, and Llandovery. In this report, rocks assigned to Pridoli and Ludlow are considered to be the Upper Silurian and those assigned to Wenlock and Llandovery to be the Lower Silurian. There is no Middle Silurian in this scheme.

#### ACKNOWLEDGMENTS

First of all, thanks are due to the geologists who did most of the mapping that I have compiled here: Hank Berg, Mike Churkin, Allan Clark, Don Eberlein, George Gehrels, Don Grybeck, Ed MacKevett, Tom Ovenshine, Pete Sainsbury, Jason Saleeby, and Charlie Rubin. Next, thanks are due to the many paleontologists who have determined the fossils and established the ages of most of the rock units, but especially to Claire Carter, Mike Churkin, Norm Savage, and Connie Soja; the last two have helped a great deal in providing data and checking my correlations for this report. Next, thanks are due to George Gehrels, Marv Lanphere, Jason Saleeby, and Don Turner for their dating of hitherto poorly bracketed granitic units and their country rocks. Next, I acknowledge the encouragement and gentle prodding of John Cathrall during the compilation, discussions with Sue Karl about the Descon Formation and related rocks, Don Grybeck for some corrections to the previous geologic compilation in the Trocadero Bay and Noyes Island areas, and Jim Baichtal of the Forest Service for sharing his unpublished mapping in the Tuxekan Passage area. Finally, thanks to the technical reviewers, Sue Karl and Don Grybeck; to the Geologic Names Unit reviewer, Ron Le Compte; and to the map editor, Carol Ostergren.

#### DESCRIPTIONS OF LITHOTECTONIC TERRANES

Southeastern Alaska and adjacent British Columbia is where the tectonostratigraphic/lithotectonic terrane concept was first conceived and developed (Monger and Ross, 1971; Jones and others, 1972, Berg and others, 1972, 1978; Monger and others, 1982; Brew and Ford, 1983, 1984; Monger and Berg, 1987) and both the tectonic evolution and the evolution of tectonic understanding of the region are therefore of special interest. Brew and others (1991a, b, 1992a) and Brew (in Nokleberg and others, 1994c) have summarized a current interpretation of the lithotectonic terrane situation in southeastern Alaska. Some of that summary applies to this map area and selected parts are abstracted here.

Southeastern Alaska is comprised of several types of lithotectonic elements. The fundamental elements are accreted terranes, meaning terranes that originated elsewhere and were transported by plate-tectonic movements and joined to previously accreted terranes or to proto-North America in approximately their present locations. The boundaries between terranes are commonly obscured by later geologic events, but vestiges of some remain. After accretion, the terranes and their boundaries were covered in part by overlap assemblages made up of sedimentary rocks derived from the underlying juxtaposed terranes. All of these elements were faulted and rearranged by later movements. Finally, post-accretion units of sedimentary and volcanic rocks were deposited after the last major movements.

Brew and others (1991a, b, 1992a) and Brew (unpub. data, 1996) summarized the lithotectonics of northern southeastern Alaska and adjacent parts of British Columbia, where all of the elements are present. The terranes and overlap assemblage rocks are shown on figure 3. From west to east, the generally north-south-trending elements (including the boundaries) are the (1) Pacific plate, (2) Transitional fault contact with the Yakutat terrane, (3) Yakutat terrane, (4) Fairweather-Queen Charlotte Islands fault, (5) Chugach terrane, (6) Border Ranges fault, expressed as the western limit of the Tarr Inlet suture zone, (7) Wrangellia terrane, (8) obscured, probably stratigraphic, contact of Wrangellia terrane with Alexander terrane, (9) Alexander terrane, (10) Alexander terrane with stratigraphically overlying Wrangellia terrane, (11) stratigraphic contact of Wrangellia terrane with the Gravina overlap assemblage, (12) Gravina overlap assemblage, (13) fault contact of Gravina overlap assemblage with the Behm Canal structural zone, (14) Behm Canal structural zone, (15) fault contact of the Behm Canal structural zone with the Nisling terrane, (16) Nisling terrane, (17) Meade fault and (or) Tally Ho-Llewellyn fault contact of Nisling terrane with Stikine terrane, (18) Stikine terrane, (19) Laberge overlap assemblage on Stikine and Cache Creek terranes. (20) Nahlin fault contact of Stikine terrane with Cache Creek terrane, and (21) Cache Creek terrane. The Coast Range megalineament (Brew and Ford, 1978) is an important young north-south element that cuts above-listed elements 10, 11, 12, 13, and 14.

The westward elements (1 through 4, above) of this array are still active, but the other parts were assembled more or less as follows: Chugach terrane to Wrangellia terrane in mid- to Late Cretaceous time; Wrangellia terrane to Alexander terrane before Carboniferous time (presuming that they were ever apart); Alexander terrane to Yukon prong and Stikine terranes in Middle Jurassic to Late Cretaceous time; Yukon prong terrane to Stikine terrane probably before Triassic time (presuming that they were ever apart); and Stikine terrane to Cache Creek terrane in Triassic time.

The study area does not include all of these many lithotectonic elements, nor does it include all of the tectonic and magmatic elements that constitute the entire tectonic history of southeastern Alaska (fig. 3). Nevertheless, it does have some post-accretionary continental volcanic and sedimentary rocks, namely the isolated Tertiary and Quaternary volcanic fields and one very small sedimentary basin; one post-accretionary overlap assemblage, the Gravina belt rocks; and two closely related major accreted terranes, the Alexander terrane and the Alexander terrane together with the Wrangellia terrane and the boundaries between them.

The small post-accretionary isolated Tertiary and Quaternary volcanic fields and the very small coal-bearing continental sedimentary basin are dispersed over the map area and there is no evidence that they were formerly of much greater extent. The rock units involved are mapped largely as extrusive volcanic rocks (QTvo, QTvr, QTvb) and as the Kootznahoo Formation (Tk).

The post-accretionary overlap assemblage, consisting of the Gravina belt rocks (KJgc, Jgd, Jgs, Jgv) are exposed on Gravina and Revillagigedo Islands near Ketchikan and on the Cleveland Peninsula to the north. The assemblage consists of Middle Jurassic to Upper Cretaceous graywacke, argillite, and conglomerate; basaltic and andesitic breccia, tuff, flow, and volcanoclastic rocks. Sedimentary rocks are dominantly marine turbidites with minor shallow-water deposits. Coarse clastic rocks are locally derived from the stratigraphically underlying Alexander and Wrangellia terranes mainly to the west (Brew and Karl, 1988a, b, c) but may also have been derived from the Yukon prong and Stikine terranes to the east, outside the study area. The Gravina belt rocks have been interpreted by Gehrels and others (1990) to have been thrust under the Nisling terrane rocks to the east, but at least some of those faults are younger than the original juxtaposition of the Gravina belt and Nisling terrane rocks.

The two closely related major accreted terranes known as the Insular superterrane or as Superterrane II (Monger and others, 1982) make up the rest of the study area. This superterrane consists of two parts: (1) the Alexander terrane *sensu stricto*, here meaning the lower Paleozoic rocks of the terrane as originally defined by Berg and others (1978b), and (2) the Alexander terrane *sensu stricto* together with what are interpreted here to be stratigraphically overlying Wrangellia terrane rocks. Most, but not all, of the rock referred here to the Wrangellia terrane were included by Berg and others (1978b) in their Alexander terrane. Previous subdivisions of the Alexander terrane into the Annette, Craig, and Admiralty subterranes (Berg and others, 1978b; Monger and Berg, 1987) were based on local age and stratigraphic variations and do not indicate significant differences in stratigraphic, magmatic, or tectonic history.

The Insular superterrane is a composite terrane consisting of island-arc rocks between the Chugach terrane on the west and the Gravina overlap assemblage belt on the east; the Chugach terrane is not exposed at the latitude of the study area. As noted above, the superterrane includes rocks of Devonian and younger age that are here referred to the Wrangellia terrane that are in apparent stratigraphic contact with Alexander terrane *sensu stricto* rocks of Silurian and older age.

In the Wrangell Mountains, northwest of southeastern Alaska, the Wrangellia and Alexander terranes have been together since Pennsylvanian time (Gardner and others, 1988). To the south, on Vancouver Island, the Wrangellia terrane includes rocks as old as Devonian. These two facts, together with permissive evidence that the Wrangellia terrane rocks of its "type area" are facies equivalents of the rocks of the same age that are stratigraphically on the Alexander terrane *sensu stricto*, are the reasons for the interpretation used here. Although this interpretation is controversial, the sum of now-available evidence indicates that the two terranes have always been together. The Alexander terrane *sensu stricto* and Wrangellia terrane are described separately below.

The Alexander terrane *sensu stricto* as defined here is a subcontinental-size volcanic-island-arc terrane consisting of Late Proterozoic to lower Paleozoic volcanoclastic turbidites, siliceous shale, chert, limestone, and intermediate to mafic volcanic rocks, with locally developed polymictic conglomerate. Locally it includes pre-accretionary granitic plutons and batholiths. Deep-water sedimentary rocks predominate, but deep- to shallow-water volcanic rocks and shallow-water carbonate rocks represent periods of intense magmatic activity and general quiescence, respectively. As noted above, the Devonian through Lower Jurassic parts of the Alexander terrane as defined by Berg and others (1978b) are here considered to belong to the Wrangellia terrane part of the Alexander superterrane and are thus described below.

The Late Proterozoic and Lower Cambrian Wales Group in the study area consists of metamorphosed and unmetamorphosed volcanic, turbidite, and carbonate rocks intruded by small intermediate composition plutons. The overlying Ordovician through Silurian section consists of mixed volcanoclastic turbidites, carbonate rocks, and volcanic rocks intruded locally by intermediate to gabbroic and ultramafic plutons.

The Wrangellia terrane is an island-arc terrane that includes Devonian shallow-water clastic, carbonate, and sparse volcanic rocks that record local variations in the continued active volcanic-arc environment. These are succeeded upward by upper Paleozoic volcanic breccias, flows, and volcanoclastic rocks and carbonate rocks that are locally intruded by late Paleozoic granitic rocks. These are overlain by nonvolcanic Permian limestone, pelitic rocks, and chert. Regionally, these rocks are overlain by black cherty argillite and by many thousands of meters of Upper Triassic subaerial to pillowed massive tholeiitic basalt with locally abundant gabbro dikes, sills, and small plutons. These are, in turn, overlain by platformal and basinal Upper Triassic limestones that grade upward into basinal, spiculitic, argillaceous and calcareous rocks. Jurassic chert and sandstone overlie Triassic(?) rocks at one locality in northern southeastern Alaska. The terrane is overlain by the Gravina overlap assemblage and is intruded locally by extensive Late Jurassic and Early Cretaceous plutonic-arc rocks.

## MINERAL RESOURCES

The most recent summary of known deposits, identified mineral resources, and undiscovered mineral resources of the Craig and Dixon Entrance quadrangles is given by Brew and others (1991a) and Brew and Drinkwater (1991). The known-deposit and identified-resource information in those reports was derived mainly from Cobb (1972a, b; 1978a, b), Cobb and others (1968), Elliott and others (1978), Berg (1984), and U.S. Geological Survey (1989); and the basic undiscovered mineral resource information was provided by D.J. Grybeck (written commun., 1989). The results of recent geochemical studies are reported by McDanal and others (1991) and by Cathrall and others (1993a, b) and by Cathrall (1994). Philpotts and others reported (1993) on some small molybdenite occurrences near Dora Bay. Previous to all of these reports, the State of Alaska Division of Geological and Geophysical Surveys was active in the area: Roehm (1946) described some high-calcium limestone deposits in the area; Herreid (1964) studied the Niblack Anchorage area; Herreid and Rose (1966) described the Hollis and Twelvemile Creek areas; Herreid (1967) described the Dolomi area; Bufvers (1967) provided a compendium of historical mining information; Eakins (1970) reported on a geochemical experiment in the Bokan Mountain area; Herreid (1971) did geochemical work in the Hetta Inlet area; Herreid and Tribble (1973) did geochemical work in the Craig A-2 quadrangle; Herreid and others (1978) studied one area in detail, and Bundtzen (1978) briefly described the copper mining in part of the area. Horton and others (1992) evaluated the mineral resources of the Annette Island [Indian] Reserve, part of which is covered in this compilation, Karl (1992) compiled information on the known mineral deposits of Annette Island, and Godwin and Smith (1993) edited a volume containing several pertinent papers on the same topic. Most recently, Bundtzen (in Nokleberg and others, 1995) described a copper-moly porphyry system in southeasternmost Prince of Wales Island.

Briefly, the Craig and Dixon Entrance quadrangles were the sites of some of the earliest prospecting and mine development in southeastern Alaska. Mining and quarrying started in the late 1800's and has continued intermittently since then. Past production from the two quadrangles is estimated at about \$100 million (Brew and others, 1991a). At this time (March 1995), there are no mines or quarries operating, but metallic mineral exploration activity is at a sustained moderate to high level.

All or parts of five metallogenic belts are present in the map area (Brew, 1993). One belt is defined by skarn and vein deposits associated with granitic plutons of mid-Cretaceous age in westernmost Prince of Wales and adjacent islands. Another belt includes those parts of that same belt that contain evidence indicating porphyry molybdenum and (or) copper deposits. Two other belts adjoin these first belts at their southeast end and together cover all of southeasternmost Prince of Wales Island; one is defined by uranium-thorium- and rare-earth-element-bearing veins associated with Jurassic peralkalic intrusions and the other by rare-earth-element deposits associated with Ordovician and (or) Silurian alkalic intrusions. The fifth belt also is in southernmost Prince of Wales and adjacent islands; it is defined by volcanogenic massive-sulfide deposits in Paleozoic and older(?) rocks.

## DESCRIPTION OF MAP UNITS

[NOTE: These descriptions have been adapted from the citations given for each unit. In some cases the previous material has been used almost *verbatim*; however in most cases it has been shortened and altered. In every case, however, the descriptions are those of the original workers and the sources are cited.]

### UNITS EXPOSED SOUTHWEST OF CLARENCE STRAIT

- Qs**      **Surficial deposits (Holocene and (or) Pleistocene)**--Alluvium, colluvium, tidal mudflat deposits, and some possible glaciofluvial deposits. Distribution of most large areas of surficial deposits was mapped in field, but deposits have not been studied in detail; many small areas are not shown. Within Craig quadrangle, major areas have been delineated by examination of aerial photographs. [Description adapted from Eberlein and others, 1983, p. 4, unit Qag]

**Extrusive volcanic rocks (Quaternary? and Tertiary)--As mapped, divided into:**

- QTvb **Basaltic to rhyolitic breccia and tuff--**Poorly consolidated, iron-stained; weathers brown and reddish brown. Contains abundant plagioclase and minor biotite phenocrysts. Breccia consists of plagioclase porphyry fragments in crystal-lithic-lapilli tuff; breccia is associated with similar dikes that contain xenoliths of agglomerate derived from Descon Formation. Small, widely separated outcrops are probably from separate vents. Unit is exposed on northeastern Prince of Wales Island. [Description adapted from Eberlein and others, 1983, p. 16, unit QTv]
- QTvr **Rhyolite and dacite--**Flows and small plugs with locally well developed columnar jointing (especially near Arena Cove). Quartz and sanidine phenocrysts present. Obsidian in rhyolite has black glass nodules in layered greenish black and yellow glass. Associated with unit QTvo. Outcrops on southern Suemez Island. [Description adapted from Eberlein and others, 1983, p. 25-26, unit QTrd]
- QTvo **Olivine basalt and andesite--**Flows, breccia, lapilli tuff, and dikes. Layered units are flat-lying, fresh, and subaerial. Conspicuously columnar jointed porphyritic olivine basalt flows and dikes on southern Suemez Island. Flows are interbedded with massive basaltic breccia and lithic lapilli tuff. Buddington and Chapin (1929, p. 271) reported coal associated with basalt in small outcrops along streams draining into Port Refugio. Associated with unit QTvr on southern Suemez Island. At Tlevak Strait, between Dall and Prince of Wales Islands, consists of the Tlevak Basalt (Eberlein and Churkin, 1970b). Exposed on Suemez, northern Dall Island, and nearby islands. [Description adapted from Eberlein and others, 1983, p. 25, unit QTb; Brew, 1988, p. 6-7, 15, 16, 1994]
- Tk **Kootznahoo Formation (Paleogene)--**Sandstone, conglomeratic sandstone, and conglomerate. Exposed at widely spaced localities on northern Prince of Wales Island: at Coal Bay (south of Kasaan Bay) unit consists of light-gray to light-yellow-brown wacke with large fragments of carbonized logs; to northeast at head of Little Coal Bay, unit consists of conglomerate and conglomeratic sandstone with cobbles and boulders of locally derived diorite and some black argillite interbedded with wacke and several-centimeter-thick seams of lignitic coal; in Lava Creek valley (west of Thorne Bay) unit consists of conglomerate with pebbles and cobbles of locally-derived argillite gneiss, trachyte, amygdaloidal basalt, granite, schist, and chert in a wacke matrix (at this locality the unit is cut by pitchstone dikes). Unit is everywhere probably less than 30 m thick and was deposited as alluvial fans, piedmont gravels, and deltas in areas of steep relief. Name and age assignments are based on correlation of this unit with type Kootznahoo Formation on Admiralty Island (Lathram and others, 1965, p. R28) and its outliers on Kuiu, Kupreanof, Zarembo, and northern Prince of Wales Islands (Brew and others, 1984, p. 10). [Description adapted from Eberlein and others, 1983, p. 4-5, unit Ts; and Brew and others, 1984, p. 10, unit Tk]
- Intrusive rocks of Chilkat-Prince of Wales plutonic province (Early Cretaceous)--**Province named by Sonnevil (1981). K-Ar age determinations on hornblende from hornblende quartz monzodiorite with minor tonalite, granodiorite, quartz diorite, diorite, quartz monzonite, and monzodiorite unit (Kwqo) on northern Kosciusko and Prince of Wales Islands by M.A. Lanphere (written commun., 1981, 1982) give 98.7 and 100.0 Ma, respectively (Brew and others, 1984, p.10-11; Douglass and others, 1989). K-Ar age determinations on hornblende and biotite reported by Turner and others (1977) and Herreid and others (1978, p. 22) from same unit in Craig A-2 quadrangle give concordant ages averaging 102 Ma. Unit age is considered to be early Late Cretaceous. As mapped, numerous plutons on Prince of Wales Island and islands to the west are divided into:

- Kwqo** **Hornblende quartz monzodiorite with minor tonalite, granodiorite, quartz diorite, diorite, quartz monzonite, and monzodiorite**--Unit is massive to foliated, equigranular to locally porphyritic; medium-grained; color index (C.I.) 2 to 48, average (approx.) 15; locally hornblende porphyritic; contains local rounded fine-grained mafic inclusions; and includes common aplite, less common pegmatite, and several mafic dikes. Typical petrographic features are: seriate twinned and zoned plagioclase with minor alteration; K-feldspar interstitial to plagioclase and occasionally in poikilitic clots; hornblende anhedral to subhedral with some plagioclase inclusions and ubiquitous opaque inclusions; pyroxene and biotite locally present and subordinate to hornblende; pyroxene altering to hornblende and biotite to chlorite; accessories are apatite and sphene. Unit differs in general from the Late Cretaceous plutons of the Admiralty-Revillagigedo plutonic belt northeast of Clarence Strait by lack of epidote and garnet, lower color index, and by lack of local plagioclase porphyry phase. Exposed on Prince of Wales, Warren, and Coronation Islands. [Description adapted from Eberlein and others, 1983, p. 20, units Kd and MzPzg, p. 37-38, units Kgrd and Kd]
- Kwgd** **Hornblende granodiorite, hornblende diorite, and biotite-hornblende monzodiorite**--Unit is fine to medium grained, color index (C.I.) about 10-25; and includes some biotite-hornblende monzodiorite. In Kasaan Bay area includes some sodic granite (from Eberlein and others, 1983, p. 20, unit Kd); at Squaw Mountain on Dall Island: light-gray, medium- to fine-grained sphene-biotite-hornblende quartz diorite and granodiorite; at Stripe Mountain on Dall Island: homogeneous, massive to moderately foliated, medium- to coarse-grained, C.I. 5 to 20, sphene-biotite-hornblende granodiorite with U-Pb (zircon) apparent age of  $114 \pm 2$  Ma (Gehrels, 1990). Exposed on Prince of Wales and Dall Islands. [Description adapted from Eberlein and others, 1983, p. 24, unit MzPzd; Gehrels and Saleeby, 1986, p. 16-17, units Kgd and Kd; Gehrels and Saleeby, 1987, p. 134-135, diorite and granodiorite unit; Gehrels, 1990, 1991, unit Kg; Gehrels, 1992, p. 10-11, units Kgd and Kd]
- Kwgb** **Gabbro**--Unit exposed at several localities. At Thunder Mountain on northern Dall Island: deformed and altered stock- or sill-like hornblende(?) gabbro body consists of medium-grained, hypidiomorphic and subidiomorphic aggregates of calcic plagioclase and pale-green amphibole (Eberlein and others, 1983, p. 24-25, unit MzPzgb); on shore of Cordova Bay between Klakas and Kassa Inlets on southern Prince of Wales Island: a small body of altered, medium- to coarse-grained biotite-hornblende gabbro intrudes Shipwreck Point fault and rocks of Wales Group (from Gehrels and Saleeby, 1986, p. 17, unit Kgb; Gehrels, 1992, p. 11, unit Kgb); west of West Arm of Cholmondeley Sound and elsewhere: small plutons and plugs of gabbro and pyroxenite (from Eberlein and others, 1983, p. 38-39, unit Kgb)
- Intrusive rocks of Bokan Mountain plutonic province (Middle Jurassic)**--Province named in this report for Bokan Mountain on southeastern Prince of Wales Island. As mapped, divided into:
- Jbgr** **Peralkaline granite of Bokan Mountain**--Concentrically zoned pluton between Kendrick Inlet and South Arm of Moira Sound consisting of three main phases (Thompson and others, 1982; De Saint-Andre and others, 1983; Gehrels and Saleeby, 1987, p. 17-18; Gehrels, 1992, p. 11-12): an outer porphyritic aegirine-bearing albite granite, a porphyritic arfvedsonite-bearing albite granite inward, and an innermost fine-grained, but locally porphyritic, arfvedsonite- and aegirine-bearing albite granite. Veins in body have been mined for uranium, thorium, and rare-earth elements. A variety of age determinations (discussed by Armstrong, 1985) indicate a minimum age of  $151 \pm 5$  Ma
- Jbsy** **Syenite of Dora Bay**--Nepheline- and eudialyte-bearing syenite with associated pegmatites (Eberlein and others, 1983, p. 39). Hornblende K-Ar age is  $175.4 \pm 6.6$  Ma (N.B. Shew, written commun., 1990)

- Hyd(?) Group (Late Triassic?)--Consists of:**
- Thb Oligomictic breccia--**Poorly sorted, angular pebble- to boulder-size clasts of plutonic rock derived from adjacent diorite, quartz diorite, gabbro, hornblende, leucogabbro, trondhjemite, pyroxenite, and migmatite (SOdi) unit. Unit has little matrix; gabbroic and metadioritic clasts are most common; hornfels, mafic dike, mudstone, and fossiliferous limestone clasts also present and are interpreted to be derived from nearby siltstone, sandstone, polymictic conglomerate, and limestone of Clover Bay unit (Dkcb). Unit is interpreted to be a submarine talus deposit. Exposed near Clover Bay on eastern Prince of Wales Island. Originally considered to be Middle and Early(?) Devonian on basis of fossiliferous clasts, but here considered Late Triassic(?) on regional relations described by Gehrels and others (1987, p. 877). Tentative correlation with Hyd Formation is based on inferred relation of this unit to Hyd rocks on northeast side of Clarence Strait. [Description adapted from Eberlein and others, 1983, p. 29-30, unit Dbx]
- PPsy Leucosyenite of Klawock and Sukkwan Island (Early Permian and Late Pennsylvanian)--**Biotite- and hornblende-bearing syenite with C.I. 15 near Klawock has K-Ar age on biotite of  $276 \pm 8$  Ma (Churkin and Eberlein, 1975, unit P<sub>2</sub>sy). Sphene-apatite-amphibole-aegerine/augite-bearing biotite leucosyenite on Sukkwan Island has hornblende and biotite K-Ar ages of 283 and 293 Ma, respectively and is surrounded by extensive hornfels aureole (Eberlein and others, 1983, p. 39, unit Psy)
- IPk Klawak Formation (Middle and Early Pennsylvanian)--**Sandstone, siltstone, and minor limestone and chert-pebble conglomerate. Unit is about 150-300 m thick and contains small fusulinids (Douglass, 1971); abundant productid brachiopods, corals, bryozoans, conodonts (Savage and Bradley, 1985), and trace fossil *Spirophyton*. Unit named by Eberlein and Churkin (1970a, p. 54). Exposed near Klawock. [Description adapted from Eberlein and others, 1983, p. 8, unit IPk]
- IPi Ladrones Limestone (Middle and Early Pennsylvanian)--**Massive and sublithographic limestone and minor dolomite. Unit is about 300+ m thick and contains some light-gray cherty nodules. Contains oolites, endothyrid foraminifers, fusulinids, conodonts (Savage and Bradley, 1985), and shelly fossils, including trilobites (Hahn and Hahn, 1991, 1992). Unit named by Eberlein and Churkin (1970a, p. 59). Exposed in Trocadero Bay and Tlevak Strait areas. [Description adapted from Eberlein and others, 1983, p. 8, unit IPi]
- Mp Peratrovich Formation (Late and Early Mississippian)--**Limestone with dark-gray chert nodules and beds that grade downward to bedded chert. Unit is about 300+ m thick and contains abundant shelly fossils, especially rugose corals (Armstrong, 1970) and conodonts in upper part (Faulhuaber, 1977); and also trilobites (Hahn and Hahn, 1991) and microfossils (Mamet and others, 1993). Unit named by Eberlein and Churkin (1970a, p. 49); (from Eberlein and others, 1983, p. 8, unit Mp)
- Dsj St. Joseph Island Volcanics (Devonian?)--**Massive, locally amygdaloidal and pillowed, basalt flows; also basaltic breccia, tuff, agglomerate, and minor siltstone, mudstone, and sandstone; about 2,000-3,000 m thick; nonfossiliferous; cut by lamprophyre dike that has whole rock(?) K-Ar age of  $328 \pm 10$  Ma; unit named by Eberlein and Churkin (1970a, p. 27). [Description adapted from Eberlein and others, 1983, p. 16-17, unit Dsv]

- Dpr Port Refugio Formation (Late Devonian; Famennian)**--Massive to thick- and thin-bedded volcanic-detritus-rich graywacke, mudstone, siltstone, and minor polymictic conglomerate, black pyritic siltstone, calcareous siltstone, pillow basalt, tuff, fossiliferous limestone, and quartzofeldspathic arenite. Unit is several thousand meters thick; closely resembles Descon Formation and may not be separable where unfossiliferous. Northern exposures are on San Juan Bautista and Balandra Islands, southern ones include Shelikof Island, where aquagene tuff and breccia directly underlie Peratrovich Formation, and Soda Bay, where a complex section consists of amygdaloidal and scoriaceous pillow basalt, breccia, tuff, calcareous siltstone, dolomitic breccia, tuffaceous calcareous siltstone, and dark-gray to black silty limestone. Age is based on abundant brachiopods in certain limestone beds (Savage and others, 1978). Savage and Gehrels (1984) and Savage (1988) discuss general distribution of faunas. Unit named by Eberlein and Churkin (1970a, p. 43). [Description adapted from Eberlein and others, 1983, p. 8-9, 26-27, unit Dp.]  
Locally includes:
- Dprv Volcanic rocks**--Agglomerate, pillow basalt, and aquagene tuff. [Description adapted from Eberlein and others, 1983, p. 27, unit Dpv]
- Dw Wadleigh Limestone (Late to Early Devonian; Famennian to Emsian)**--Massive and thick- to medium-bedded limestone, minor argillaceous limestone and calcareous shale. Unit is about 300 m thick and is composed of abundant fragmented shelly fossils in a dark-gray lime-mudstone matrix. Locally rich in stromatoporoid *Amphipora* and corals (Oliver and others, 1975; Tchudinova and others, 1974; Savage, 1977a; Savage and Funai, 1980); massive stromatopoids locally in reefs and reef breccias; brachiopods (Soja, 1988b, c), gastropods, ostracodes, pelecypods, and crinoids locally abundant; conodonts described by Savage (1977b, 1987, 1992) and Savage and Funai (1980). Savage and Gehrels (1984) and Savage (1988) discuss general distribution of faunas. Unit named by Eberlein and Churkin (1970a, p. 37). Age assignment based on information from Savage and Gehrels (1984) and Soja (1988b, c). Exposed on west side of Prince of Wales Island near Klawock. [Description adapted from Eberlein and others, 1983, p. 9, unit Dw]
- Dc Coronados Volcanics (Middle Devonian)**--Fragmental basalt interbedded with fossiliferous limestone and amygdaloidal pillow. Basalt fragments present in matrix of aquagene tuff and breccia. Unit is about 150 m thick; limestone consists mainly of fragments of tabulate corals, massive stromatoporoids, horn corals, gastropods, crinoid columnals, and brachiopods (Tchudinova and others, 1974). Savage (1988) discusses general distribution of faunas. Unit named by Eberlein and Churkin (1970a, p. 33). Exposed in vicinity of Coronados Islands. [Description adapted from Eberlein and others, 1983, p. 17, unit Dc]
- Karheen Formation and associated rocks (Middle? and Early Devonian)**--The Karheen Formation was named by Eberlein and Churkin (1970a, p. 22) and extended geographically by Gehrels and Saleeby (1986, 1987) and Gehrels (1992) to include age-equivalent rocks of different facies south of Karheen type area. "Dkk" symbols indicate Karheen Formation rocks; the "Dk" symbols indicate associated rocks. As mapped, divided into:
- Dkr Rhyolite of Kasaan Island (Middle? and Early Devonian)**--Rhyolite dikes and sills that cut limestone of Kasaan Island unit (Dkl). Thickness not known.  
[Description adapted from Eberlein and others, 1983, p. 17, unit Dkr]
- Dkl Limestone of Kasaan Island (Middle? and Early Devonian)**--Limestone and minor rhyolite tuff associated with rhyolite of Kasaan Island (Dkr). Unit is greater than 150 m thick and thick to thin bedded. Some thin layers of welded rhyolite tuff similar to rhyolite of Kasaan Island and fragments of rhyolite present in limestone breccia that consists mainly of fossil debris. Abundant fossils, including corals, stromatoporoids, crinoids, brachiopods (Soja, 1988a, b, c), conodonts (Savage, 1981a), and ostracodes (Berdan and Copeland, 1973). Soja (1988d) discusses depositional environment in detail and Savage (1988) discusses general distribution of faunas. [Description adapted from Eberlein and others, 1983, p. 12, unit Dkl]

- Dkkt **Turbidite-facies rocks (Middle? and Early Devonian)**--Mudstone, siltstone, shale, sandstone, conglomerate, and minor limestone. Unit is greater than 2,000 m thick. Upper part of unit is dominantly interbedded andesitic flows, broken pillow breccia, and tuff; contact with younger units has not been seen and top is apparently not exposed. Basal part of unit in southern part of Craig quadrangle is conglomerate and sedimentary breccia that unconformably overlies (1) metamorphic rocks at Klakas Inlet (DSmk), (2) basaltic to andesitic volcanic rocks (SODB) and volcanoclastic graywacke, siltstone, mudstone turbidites, and minor sedimentary breccia, limestone and polymictic conglomerate (SODg) units of Descon Formation, (3) diorite, quartz diorite, gabbro, hornblende, leucogabbro, \*\*\*migmatite unit (SODi), and (4) greenstone, \*\*\*and minor marble (CZwg) unit of Wales Group. Lower contact relations are described in detail by Eberlein and others (1983, p. 27). In Dixon Entrance quadrangle, these basal strata are mapped separately as Karheen polymictic conglomerate, .\*\*\*and volcanic rocks of conglomerate facies (Dkkc) of Karheen Formation. Interbedded black pyritic slate layers contain *Monograptus pacificus*, indicating a Pragian age (Churkin and others, 1970), and graptolites and shelly faunas from limestone beds in unit support that age. Unit is considered coeval with all other units mapped here as parts of Karheen Formation and associated rocks, with Wadleigh Limestone (Dw) and perhaps Coronados Volcanics (Dc). Unit age is considered to be Early and early Middle(?) Devonian. Savage and Gehrels (1984) discuss distribution of conodonts. Exposed on southern Prince of Wales Island between Kassa Inlet and Klakas Inlet. [Description adapted from Eberlein and others, 1983, p. 27-28, unit Ds; Gehrels and Saleeby, 1986, p. 3-7, units Dsh, Dms, Ds, Dk, Dlx, Dlx; Gehrels and Saleeby, 1987, p. 124-125, p. 132: sandstone, siltstone, and limestone, and mudstone and shale units of the Karheen Formation; Gehrels, 1992, p. 2-4, units Dksh, Dkms, Dks, Dkl, Dkbx]
- Dkkr **Redbed-facies rocks (Early Devonian)**--Sandstone, shale, conglomerate, and minor well-bedded platy limestone. Unit is about 1,800 m thick. Characterized in north by red beds, calcareous cement, festoon crossbedding, ripple marks and mud cracks (Eberlein and Churkin, 1970a; Ovenshine, 1975). Clasts are mostly mafic volcanic rocks and chert, with lesser amounts of sandstone, graywacke, quartz, siltstone, limestone, and felsic to mafic granitoid rocks present as pebbles and boulders; detrital K-spar and bronze-colored biotite distinguish sandstones from those in older formations. Overlies various units conformably and unconformably, and to east may interfinger with sedimentary rocks of the Staney Creek and Tuxekan Passage area unit, as described in detail by Eberlein and others (1983, p. 10). Locally contains abundant Early Devonian brachiopods (Kirk and Amsden, 1952; Savage, 1981b), and conodonts (Savage, 1977a, 1982), but relations to other fossiliferous units suggest that deposition of unit may have begun in Late Silurian time. Exposed discontinuously from Kosciusko Island on north to southernmost Prince of Wales Island on south. [Description adapted from Eberlein and others, 1983, p. 9-11, unit Dk]
- Dkbn **Breccia of northeastern Noyes Island (Early Devonian)**--Olistostromal blocks and breccia beds of limestone rich in shelly fossils and conodonts in matrix of graptolite- and plant-bearing shale (Churkin and others, 1969), together with graywacke, sandstone, and conglomerate. Unit is more than 400 m thick and unconformable on Descon Formation. Blocks and breccia beds contain reworked shallow-water faunas and a block of Lower Silurian shale; matrix contains *Monograptus yukonensis* and other Early Devonian graptolites. [Description adapted from Eberlein and others, 1983, p. 11-12, unit Dn]

- Dksn **Sedimentary rocks of Port Nicholas area (Early Devonian)**--Sandstone overlain by massive limestone about 50 m thick, in turn overlain by graptolitic shale about 50 m thick (Churkin and others, 1970). Total thickness about 400+ m. Sandstone is calcareous, light gray, crossbedded, contains scattered chert pebbles, and is interbedded with mudstone; limestone is fossil fragmental, fetid, and contains abundant corals and stromatoporoids. Conodonts indicate an early to middle Pragian age (Savage and others, 1977). Shale contains abundant *Monograptus yukonensis* and *M. pacificus*. Unit is in part similar to *M. pacificus*-bearing shale of unit Dkkt described below. [Description adapted from Eberlein and others, 1983, p. 11, unit Dsn]
- Dkp **Plagioclase-porphyrific dacite(?), andesite, and diorite(?) (Early Devonian)**--Small plugs and associated(?) flows, breccias, and dikes. Flow layers to several hundred meters thick are interbedded with units Dkkt and Dkko (see below) and hypabyssal plugs cut same units. [Description adapted from Gehrels and Saleeby, 1986, p. 5-6, units Dpv, Dbv, p. 18, unit Dph; Gehrels and Saleeby, 1987, p. 124-125, 132-133: volcanic and hypabyssal rocks unit of Karheen Formation; Gehrels, 1992, p. 3, units Dkpv, Dkbv, p. 11-12, unit Dph]
- Dkcg **Conglomerate-facies rocks (Early Devonian)**--Polymictic conglomerate, conglomeratic sandstone, sandstone, siltstone, and volcanic rocks. Reddish-brown cobble to boulder conglomerate and pebbly sandstone; thickness ranges from a few meters in north to greater than 1 km in south. To north, moderately well rounded clasts less than 40-cm in diameter of Ordovician volcanic and plutonic rocks are most common; to south, where unit is closely associated with plagioclase-porphyrific dacite(?), andesite, and diorite(?) (Dkp) unit, clasts are larger and more angular than to north and are mostly derived from that unit. Near mouth of Klakas Inlet, conglomerate grades laterally into coarse sedimentary breccia that both overlies and contains clasts of Ordovician volcanic and plutonic rocks. Exposed on southern Prince of Wales Island between Klakas Inlet and Point Nunez. [Description adapted from Gehrels and Saleeby, 1986, p. 3-7, units Dcg, Dbx; Gehrels and Saleeby, 1987, p. 132: conglomerate and breccia unit of Karheen Formation; Gehrels, 1992, p. 4, unit Dkcg]
- Dkcb **Siltstone, sandstone, polymictic conglomerate, and limestone of Clover Bay (Early? Devonian)**--Calcareous well-bedded sandstone, siltstone with subordinate gray argillaceous limestone, banded mudstone with graded bedding. Locally cross-laminated (in sets a few centimeters to about 1 m thick) granule to pebble conglomerate. Thickness probably more than 75 m. Unit includes at least two subunits as thick as about 15 m of dark- to medium-gray, fetid argillaceous limestone with abundant fragmented shelly and coralline fossils; these fossils indicate a Devonian age, and ostracodes resembling *Beyrichia* suggest an Early Devonian age. Conglomerate clasts are mainly mafic volcanic rock, red and green chert, and minor granitic rock like that in diorite, quartz diorite, gabbro, hornblendite, leucogabbro, trondhjemite, pyroxenite, and migmatite (SOdi) unit. Exposed only at Clover Bay on eastern Prince of Wales Island. [Description adapted from Eberlein and others, 1983, p. 30-31, unit Dcb]
- DSmk **Metamorphic rocks at Klakas Inlet (Early Devonian and Silurian)**--Semischist, greenstone, and leucodiorite. Age assigned is that of metamorphism of highly altered and brecciated rocks of dominantly intermediate composition derived from volcanoclastic graywacke, \*\*\*conglomerate (SOdg) and basaltic to andesitic volcanic rocks (SOdb) units of Descon Formation of Early Silurian to Early Ordovician age and foliated hornblende granodiorite\*\*\*quartz-porphyrific granodiorite (Ogd) unit of the older part of the Cape Chacon plutonic province. Overlain by turbidite-facies rocks (Dkkt) of Karheen Formation. Exposed only at mouth of Klakas Inlet. [Description adapted from Gehrels and Saleeby, 1986, p. 12, unit DSk; and Gehrels, 1992, p. 7, unit DSk]

- DSs **Sedimentary rocks of Stoney Creek and Tuxekan Passage area (Early Devonian? and Late and Early Silurian)**--Limestone, sandstone, mudstone, and polymictic conglomerate. Unit is about 300 m thick and consists of interbedded light-gray, massive- to well-bedded, cyclically repeated limestone that resembles Heceta Limestone (Shl) and sandstone, conglomeratic sandstone, limy sandstone, and concretionary mudstone with locally well developed mudcracks and ripplemark like the redbed-facies rocks (Dkkr) in Karheen Formation and in clastic strata in Heceta Limestone (Shl). Basal limestone and shaly beds contain *Catenipora* and late Llandoveryan graptolites that are younger than highest-occurring graptolites in volcanoclastic graywacke, limestone beds in upper part of sequence contain dasycladacean algae, corals, *Amphipora*, and *Conchidium*. Stratigraphic relations discussed by Eberlein and others (1983, p. 14) and field relations suggest that this unit, the Heceta Limestone, and the Bay of Pillars Formation are, in part, facies equivalents. Savage (1988) discusses general distribution of faunas. Exposed in northern part of quadrangle on Prince of Wales Island. [Description adapted from Eberlein and others, 1983, p. 13-14, unit DSs]
- Sld **Leucodiorite at Kassa Inlet (Late Silurian)**--Medium- to coarse-grained melanite(?) -sphene-arfvedsonite (with cores of aegerine-augite) leucodiorite with C.I. <15. Age is 418±5 Ma. Exposed only at Kassa Inlet, southern Prince of Wales Island. [Description adapted from Gehrels and Saleeby, 1986, p. 18, unit Sd; Gehrels and Saleeby, 1987, p. 131-132, leucodiorite unit; Gehrels, 1992, p.12, unit Sd]
- Shl **Heceta Limestone (Late and Early Silurian)**--Unit named by Eberlein and Churkin (1970a, p. 15). Age is late Early and Late Silurian. As mapped, divided into:  
**Limestone, minor limestone breccia, sandstone, mudstone, and polymictic conglomerate**--Unit is over 3,000 m thick on western Heceta Island, thins abruptly eastward due to facies change into sedimentary rocks of Stoney Creek and Tuxekan Passage area (DSs) unit, and southward due probably to pre-Karheen erosion. Irregular masses in the Logjam Creek area east of the northern continuation of the Lake St. Nicolas fault are tentatively interpreted to be olistoliths. Limestone is massive, sublithographic, and lighter colored than other limestone units in map area; and is richly fossiliferous with corals, dasycladacean algae, brachiopods (Savage, 1989), stromatoporoids (including *Amphipora*), gastropods, pelecypods, bryozoans, trilobites, conodonts (Savage, 1985), trace fossils (Soja, 1991b), graptolites. Soja (1991a, 1990, 1993), and sphinctozoans (Rigby and others, 1994). Riding and Soja (1993) and Soja (1993b, 1994) discussed depositional environments in detail. Contact with underlying Descon Formation is generally conformable, but on west side of Heceta Island basal breccia consists of angular slabs of black graptolitic shale with monograptids of middle Early Silurian age. Exposed in northwestern Prince of Wales Island and on adjacent islands. [Description adapted from Eberlein and others, 1983, p. 12-13, unit Sh]
- Shc **Polymictic conglomerate**--Thick lenses and pods of limestone breccia, polymictic conglomerate, and sandstone in central part of type locality of Heceta Limestone on Heceta Island. Limestone breccia and conglomerate consists of reworked intraformational lithologies; polymictic conglomerate contains clasts of plutonic, volcanic, and sedimentary rock, including chert; latter is significant because chert is uncommon in older units in area. [Description adapted from Eberlein and others, 1983, p. 13, unit Shc]
- Bay of Pillars Formation (Late and Late? and Early Silurian)**--Unit named by Muffler (1967, p. C6); extended geographically into Prince of Wales Island by Ovenshine and Brew (1972) and Brew and others (1984, p. 19). Unit on northeastern Prince of Wales Island (Spg, Spc) is distinguished from that on northern and northwestern Prince of Wales Island (Spsv) by its more volcanoclastic and less calcareous composition. As mapped, divided into:

- Spsv **Sedimentary and volcanic rocks**--Graywacke, and mudstone turbidites. polymictic conglomerate, minor limestone, andesitic to basaltic volcanic flows, breccia, and tuff. Thickness probably exceeds a few thousand meters. Dominantly graywacke, mudstone, and calcareous mudstone turbidites, with subordinate conglomerate, limestone, and intermediate to mafic volcanic flows, breccia, and tuff. Sedimentary features in sandstone turbidites include massive, amalgamated beds, channelized beds, graded beds with Bouma sequences, and chaotically deformed slump deposits. Sandstones are extremely variable in composition: three dominant varieties in Petersburg quadrangle to north are calcareous graywacke, volcanoclastic graywacke, and quartzofeldspathic graywacke; proximal turbidite facies and crossbedding in conglomerates suggest shallow to moderate water depths. Karl and Giffen (1992) discuss sedimentology of this unit in Petersburg quadrangle to north. Graptolite collections made during Petersburg area study range in age from middle Llandoveryan to early Ludlovian (Claire Carter, written commun., 1980); unit is of Early and Late Silurian age. Unit differs from Descon Formation because it has significantly less volcanic debris, both in subunits and as individual clastic grains; it also is mostly younger than Descon. Exposed along El Capitan Passage on northern and northwestern Prince of Wales Island. [Description adapted from Brew and others, 1984, p. 17-18]
- Spg **Volcanoclastic graywacke and mudstone turbidites and minor limestone**--Greenish-gray, buff-weathering, volcanoclastic graywacke and argillite turbidite deposits in massive to amalgamated, graded, and rhythmic beds of Mutti and Ricchi-Lucchi A, B, C, and E facies, which suggest proximal depositional environment in moderate water depths. Thickness not known. Graptolites on some argillite bedding surfaces are all of Early Silurian age (Claire Carter, written commun., 1980); unit is considered to be of Early and Late(?) Silurian age. Exposed only in small area at northern edge of map on northeastern Prince of Wales Island. [Description adapted from Brew and others, 1984, p. 19]
- Spc **Conglomerate, agglomerate, and volcanic breccia**--Mostly volcanoclastic polymictic conglomerate and volcanic breccia of intermediate to mafic composition with feldspar- and clinopyroxene porphyry common. Thickness not known. Exposed only in small outcrop area at north edge of map. [Description adapted from Brew and others, 1984, p. 19]
- Shpc **Polymictic conglomerate between Heceta Limestone and Bay of Pillars Formation (Early? Silurian)**--Polymictic pebble and cobble conglomerate and other clastic rocks like those in polymictic conglomerate of Heceta Limestone (Spc), but present between unit Shl of Heceta Limestone and Bay of Pillars Formation (Spsv) on northern Prince of Wales Island. Thickness not known, but probably greater than several thousand meters locally. Exposed on Coronation Island and on Spanish Islands. [Description adapted from Brew and others, 1984, p. 17, unit Spc]
- Intrusive and related rocks of younger part of Cape Chacon plutonic province (Early Silurian and Late Ordovician)**--This province named in this report from locality of its southernmost exposures at Cape Chacon, Prince of Wales Island. As mapped, divided into:
- SOMk **Metamorphic rocks at Kendrick Bay**--Heterogeneous schistose biotite-hornblende-quartz-plagioclase gneiss, schist, and amphibolite. Unit consists of (1) fine- to very fine grained layered and well-foliated (muscovite-sphene-pyrite-magnetite-ilmenite-monazite(?)-) K-feldspar-biotite-hornblende-plagioclase gneiss, probably derived from sedimentary and volcanic rocks; (2) very fine grained, well-foliated, dark- and medium-gray (K-feldspar-muscovite-) biotite-hornblende-plagioclase schist of amphibolite facies; and (3) dark-green and dark-greenish-brown, coarse-grained (augite-) plagioclase-hornblende amphibolite, probably derived from mafic rocks. [Description adapted from MacKevett, 1963, p. 17-18, units Dsc, Dgn, Dam; Gehrels and Saleeby, 1986, p. 12, unit SC; Gehrels, 1992, p. 7-8, unit SOK]

- SOdi Diorite, quartz diorite, gabbro, hornblendite, leucogabbro, trondhjemite, pyroxenite, and migmatite**--Heterogeneous complex consisting of these rock types; cut by locally sheeted mafic and by felsic dike swarms. Host rock for skarn deposits on Kaasan Peninsula (Eberlein and Churkin, 1976). U-Pb (zircon) apparent age of rock from same unit on Duke Island is  $429\pm 20$  Ma (Gehrels and others, 1987, p. 872). Exposed on eastern Prince of Wales Island from Cholmondeley Sound to upper Kaasan Bay. [Description adapted from Eberlein and others, 1983, p. 20-24, unit Pzic; Gehrels and others, 1987, p. 871-872, diorite unit]
- SOsy Quartz syenite and granite**--Medium-grained, leucocratic, grayish-red to reddish-gray, massive-appearing, poorly foliated to nonfoliated (garnet-aegirine/augite-hornblende-) biotite syenite. Unit may be genetically related to unit SOgr. U-Pb (zircon) apparent age is  $438\pm 5$  Ma (Gehrels and Saleeby, 1987, p. 130). Exposed near Stone Rock Bay on southeastern Prince of Wales Island. [Description adapted from MacKevett, 1963, p. 32-33, unit Ksy; Gehrels and Saleeby, 1986, p. 19, unit SOsy; Gehrels and Saleeby, 1987, p. 130, quartz syenite unit; Gehrels, 1992, p. 12-13, unit SOsy]
- SOgr Granite**--Coarse- to medium-grained, locally well foliated and banded, porphyritic biotite granite. Unit may be genetically related to unit SOsy. U-Pb (zircon) apparent age is  $438\pm 5$  Ma (Gehrels and Saleeby, 1987, p. 130). Exposed on Kendrick Islands and south of mouth of Kendrick Bay. [Description adapted from MacKevett, 1963, p. 30-31, part of unit Kgg; Gehrels and Saleeby, 1986, p. 20, unit SOgr; Gehrels and Saleeby, 1987, p. 130, granite unit; Gehrels, 1992, p. 13, unit SOgr]
- SOqm Unfoliated and foliated quartz monzonite and granite**--Homogeneous, massive-appearing, medium-grained, light-gray hornblende granodiorite, biotite-hornblende granodiorite, leucocratic granodiorite, biotite granodiorite, biotite quartz monzonite, leucocratic quartz monzonite, and hornblende quartz monzonite. Albitized in vicinity of peralkaline granite of Bokan Mountain (Jbgr). U-Pb (zircon) apparent age is  $438\pm 4$  Ma (Gehrels and Saleeby, 1987, p. 130). [Description adapted from MacKevett, 1963, p. 24-30, unit Kqm; Gehrels and Saleeby, 1986, p. 20-21, unit SOqm; Gehrels and Saleeby, 1987, p. 130, quartz monzonite unit; Gehrels, 1992, p. 13-14, unit SOqm]
- SOpx Pyroxenite, hornblendite, and gabbro**--Composition of unit differs from locality to locality: (1) At Salt Chuck near Kasaan Bay: discordant, pipe-like stock of olivine-bearing pyroxenite and gabbro, diorite, and hornblendite; host rock for Cu, Au, Ag, and PGM (platinum-group-metal)-bearing magmatic sulfide deposits; minimum age is about  $429\pm 11$  Ma, based on K-Ar dating of biotite (Gault, 1945; Gault and others, 1992; Loney and others, 1987; Loney and Himmelberg, 1992; Himmelberg and Loney, 1995). [Description adapted from Eberlein and others, 1983, p. 18-19, unit Kpg.] (2) Near Stone Rock Bay on southeastern Prince of Wales Island: elongate body of dark-gray, medium- to coarse-grained, locally porphyritic (magnetite-biotite-hornblende-) augite pyroxenite; some parts contain magnetite in potential economic amounts; intruded by unfoliated and foliated quartz monzonite and granite (SOqm) unit. [Description adapted from MacKevett, 1963, p. 17-18, unit Kpy; Gehrels and Saleeby, 1986, p. 19, unit SOpx; Gehrels, 1992, p. 13, unit SOpx.] (3) On southern Sukkwan Island: medium- to coarse-grained hornblende gabbro, hornblendepyroxenite, and hornblendite; minimum age is about 440 to 449 Ma, based on K-Ar dating of two hornblende concentrates (M.A. Lanphere, oral commun. to G.D. Eberlein, 1981; Himmelberg and Loney, 1995). [Description adapted from Eberlein and others, 1983, p. 39-40, unit Ogb; Gehrels, 1991, unit Ogb.] (4) On western Long and western Dall Islands: medium- to coarse-grained, locally porphyritic (biotite-hornblende-) augite pyroxenite and pyroxene gabbro; minimum age is about 400 Ma, based on K-Ar dating of hornblende concentrate (M.A. Lanphere, oral commun. to R.A. Loney, 1991; Himmelberg and Loney, 1995). [Description adapted from Gehrels, 1991, unit Opx]

- SOhb**      **Hornblendite**--Altered hornblendite; probably similar to hornblende gabbro on Duke Island. Exposed on east shore of Prince of Wales Island north of Clover Bay. [Description adapted from Eberlein and others, 1983, p. 40; Gehrels and others, 1987, p. 872-873, hornblende gabbro unit]
- SObl**      **Andesitic breccia of Luck Creek (Early Silurian and Late Ordovician)**--Angular fragments of porphyritic andesite in a matrix of hornblende-diopsidic augite crystal tuff of andesitic composition. Unit is as thick as several thousand meters. Massive and structureless in outcrop, but crude large-scale layering apparent from distance west and east of Thorne River. At Stoney Cone and elsewhere 10-m-thick lenses of light-gray aphanitic limestone contain *Catenipora* sp. of Late Ordovician to Early Silurian age. K-Ar dating of hornblende from the breccia along Big Salt Lake and near Kogish Mountain gives a minimum age of about 440 Ma (M.A. Lanphere, oral commun. to G.D. Eberlein, 1981; Eberlein and others, 1983, p.18). May be in part extrusive equivalent of diorite, quartz diorite, gabbro, hornblendite, leucogabbro, trondhjemite, pyroxenite, and migmatite (SOdi) unit, and more proximal equivalent of the Descon Formation (SOdg). However, close association and lithologic similarity of this unit with Descon volcanic and other rocks make the two units difficult to distinguish. As presently mapped, exposed in large areas in central Prince of Wales Island. [Description adapted from Eberlein and others, 1983, p. 17-18]
- SOdg**      **Descon Formation (Early Silurian to Early Ordovician)**--Unit named by Eberlein and Churkin (1970a, p. 5) and geographically extended by Herreid and others (1978, p. 10) into southern Prince of Wales Island. As mapped, divided into:
- SOdg**      **Volcaniclastic graywacke, siltstone, mudstone turbidites, and minor sedimentary breccia, limestone, and polymictic conglomerate**--Mainly graywacke and mudstone with some interbedded massive basalt lava and pyroclastic flows that cannot be mapped separately at this scale. Unit is at least 3,000 m thick, and base is not exposed. Graywacke consists of dominantly basaltic detritus in a noncalcareous chlorite-rich matrix; graded bedding common; interbedded mudstone is rhythmically bedded and has graded bedding and penecontemporaneous slump features. Conglomerate and sedimentary breccia range from crudely layered augite-phenocryst-rich andesite breccia like that of andesitic breccia of Luck Creek (SObl) to polymictic types with clasts of chert, felsic volcanic rock, graywacke, and gabbro. Minor black graptolite-bearing chert and siliceous shale present. Orange-weathering quartzofeldspathic wacke present in vicinities of Heceta and Lulu Islands and near Salt Chuck Bay on western Prince of Wales Island. To south and west, unit is less indurated, less metamorphosed, and more calcareous. Abundant and well-preserved graptolites establish presence of virtually all of the Ordovician and Lower Silurian graptolite zones (Churkin and others, 1970; Carter and others, 1980; Eberlein and others, 1983, table 1, localities G1, 2, 4-6, 8-10, 12-23, 25-48). Locally penetratively deformed and metamorphosed to hornfels, slate, phyllite, semischist, schist, and greenschist. Close association and lithologic similarity of this unit with andesitic breccia of Luck Creek (SObl) make the two units difficult to distinguish. Exposed throughout Prince of Wales and adjacent islands north of about latitude 55° N. (Craig-Dixon Entrance quadrangle boundary). [Description adapted from MacKevett, 1963, p. 8-10, unit Dsl; Eberlein and others, 1983, p. 15-16, unit SOd, p. 31-33, unit SOs; Gehrels and Saleeby, 1986, p. 7-9, units SOa, SOms, SOgw, SOcs; Gehrels and Saleeby, 1987, p. 127-128, argillite, and mudstone and siltstone units of Descon Formation; Gehrels, 1991, units SOda, SOdm; Gehrels, 1992, p. 4-6, units SOda, SOdms, SOdgw, SOdl]
- SOdc**      **Volcaniclastic conglomerate**--Mapped by J.A. Baichtal (written comm., 1995) inland to the east of Tuxekan Passage and south of Karheen Passage. Mapped by Eberlein and others (1983) as parts of units DSs and SObl.
- SOdr**      **Dacitic to rhyolitic volcanic rocks**--Silicic (dacite to rhyolite) breccia, tuff-breccia, tuff, and extrusive domes. Exposed in southeasternmost part of Prince of Wales Island, and in one locality at Augustine Bay on Dall Island. [Description adapted from Gehrels and Saleeby, 1986, p. 10-11, unit SOsv; Gehrels and Saleeby, 1987, p.127, rhyolitic volcanic rocks unit; Gehrels, 1991, unit SOdr; Gehrels, 1992, p. 6-7, unit SOdsv]

- SOda Andesitic to dacitic volcanic rocks**--Intermediate (andesite to dacite) flows and breccia. Exposed south of Moira Sound and near Klakas Inlet on southern Prince of Wales Island, and in Barrier Islands. [Description adapted from Gehrels and Saleeby, 1986, p. 10, part of unit SOiv; Gehrels and Saleeby, 1987, p.127, included these rocks in their rhyolitic volcanic rocks unit of Descon Formation; Gehrels, 1992, p. 6, part of unit SOdiv]
- SOdb Basaltic to andesitic volcanic rocks**--Augite-bearing basaltic to andesitic pillow flows, pillow breccia, agglomerate, breccia, and tuff-breccia; some massive lava flow deposits. Exposed in vicinity of El Capitan and Esquibel Islands and at Salt Lake Bay on northern Prince of Wales Island, on Sukkwan Island, and in central and southeasternmost parts of Prince of Wales Island. [Description adapted from MacKevett, 1963, p. 6-8, unit Dv; Eberlein and others, 1983, p. 15, 16, unit SOdv, p. 33-34, unit SOv; Gehrels and Saleeby, 1986, p. 10, units SObv and SOms, part of unit SOiv; Gehrels and Saleeby, 1987, p.127, basaltic to andesitic volcanic rocks unit of Descon Formation; Gehrels, 1991, unit SOdb; Gehrels, 1992, p. 5-6, units SOdbv and SOdms, part of unit SOdiv]
- Intrusive and related rocks of older part of Cape Chacon plutonic province (Late and Middle Ordovician)**--This province named in this report from locality of its southernmost exposures at Cape Chacon, Prince of Wales Island. As mapped, divided into:
- Oqd Hornblende quartz diorite, diorite, and quartz monzonite (Late Ordovician)**--Massive, fine- to medium-grained, C.I. about 20, biotite-hornblende quartz diorite and minor diorite. U-Pb (zircon) apparent age is  $445\pm 5$  Ma (Gehrels and Saleeby, 1987, p. 130). K-Ar dating of hornblende there gave apparent age of  $439\pm 21$  Ma (Lanphere and others, 1964, p. 705, recalculated by Gehrels and Saleeby, 1986). Exposed at and near Kendrick Bay. [Description adapted from MacKevett, 1963, p. 22-28, unit Kqm; Gehrels and Saleeby, 1986, p. 21-22, unit Oqd; Gehrels and Saleeby, 1987, p. 128-130, part of quartz diorite unit; Gehrels, 1992, p. 14-15, unit Oqd]
- Odi Foliated and layered hornblende diorite and quartz diorite, heterogeneous diorite and gabbro, and minor foliated granodiorite (Late and Middle? Ordovician)**--Massive and homogeneous bodies and heterogeneous masses of medium-grained, C.I. 30-50, foliated diorite and hornblende gabbro. U-Pb (zircon) apparent age is  $446\pm 5$  Ma (Gehrels and Saleeby, 1987, p. 130). Exposed at and near Tah Bay. [Description adapted from Gehrels and Saleeby, 1986, p. 24-25, 22-23, units Od, Ofd, Ofgd; Gehrels and Saleeby, 1987, p.128-130, diorite unit and part of quartz diorite unit; Gehrels, 1992, p. 15-16, units Od, Ofd, Ofgd]
- Ogd Foliated hornblende granodiorite, chlorite leucogranodiorite, and quartz-porphyritic granodiorite (Middle Ordovician)**--Homogeneous, massive, medium- to coarse-grained, C.I. 5-20, hornblende granodiorite. U-Pb (zircon) apparent ages from this unit are  $462\pm 15$  Ma,  $468\pm 15$  Ma, and  $472\pm 5$  Ma (Gehrels and Saleeby, 1987, p. 128). Exposed on southern and southeastern Prince of Wales Island. [Description adapted from Gehrels and Saleeby, 1986, p. 24-25, 23-24, units Oqgd and Ogd; Gehrels and Saleeby, 1987, p. 128-129, granodiorite unit; Gehrels, 1992, p. 16-17, unit Ogd]
- Omr Metamorphic rocks at Ruth Bay (Middle Ordovician)**--Foliated and layered metaplutonic and minor metavolcanic and metasedimentary rocks: about 60 percent foliated and slightly layered hornblende gabbro; about 35 percent well-foliated, C.I. less than 15, hornblende-biotite granodiorite; and about 5 percent hornblende-bearing metavolcanic and metasedimentary rocks. U-Pb (zircon) apparent age of a granodiorite sill is  $465\pm 7$  Ma (Gehrels and Saleeby, 1987, p. 130). Exposed between mouths of Kassa and Klakas Inlets. [Description adapted from Gehrels and Saleeby, 1986, p. 12-13, unit Or; Gehrels and Saleeby, 1987, p. 130-131, Ordovician metaplutonic rocks; Gehrels, 1992, p. 8-9, unit Or]

- Qmpw Metamorphic rocks on eastern Prince of Wales Island (Late and Middle Cambrian)**--Foliated, slightly layered, metamorphosed diorite, microdiorite, quartz diorite, gabbro, microgabbro, and hornblendite. Correlates with unit on southern tip of Gravina Island, where a U-Pb (zircon) apparent age of a sample from Bronough Islands (off south tip of Gravina Island) is in range 540 to 510 Ma (Gehrels and others, 1987, p. 868-869). Exposed to west and east of entrance to Cholmondeley Sound on southeastern Prince of Wales Island. [Description adapted from Gehrels and others, 1987, p. 867, 869, Cambrian metaplutonic rocks unit]
- Intrusive rocks of southern Dall Island plutonic province (Early Cambrian)**-- This province named in this report from its only exposures on southern Dall Island. As mapped, divided into:
- Qqd Foliated, lineated, and gneissic biotite-hornblende quartz diorite, granodiorite, and minor diorite**--Massive, medium-grained, C.I. 20-40, hornblende-biotite quartz diorite and granodiorite with no fabric grades to strongly foliated and locally layered quartzofeldspathic schist. U-Pb (zircon) apparent age of a gneissic granodiorite that intrudes rocks of Wales Group at Cape Muzon is  $555 \pm 4$  Ma. [Description adapted from Gehrels, 1991, unit Qq]
- Qdi Foliated and layered biotite-hornblende diorite and quartz diorite**-- Foliated, compositionally-layered and grain-size-layered biotite-hornblende diorite and lesser quartz diorite. [Description adapted from Gehrels, 1991, unit Qd]
- Qgb Hornblende gabbro, microgabbro, and quartz diorite**--Gabbro is foliated, lineated, and metamorphosed to chlorite and locally biotite schist; grades into heterogeneous quartz diorite, granodiorite, and gabbro containing as much as 50 percent screens and pendants of metavolcanic rocks. [Description adapted from Gehrels, 1991, units Qgb and Qqb]
- Wales Group (Early Cambrian? and Late Proterozoic)**--Unit named by Brooks (1902) as Wales Series and redefined by Buddington and Chapin (1929, p. 45-49) as Wales Group. As mapped, divided into:
- Qzwgb Metagabbro south of Kassa Inlet**--Heterogeneous deformed and metamorphosed hornblende-clinopyroxene gabbro. Intrusive into Wales Group, but included here as part of Wales Group because it has been metamorphosed together with surrounding rocks whereas other intrusive rocks in the area (those of Cape Chacon plutonic province) were not. [Description adapted from Gehrels and Saleeby, 1986, p. 25, unit pmOgb; Gehrels, 1992, p. 17-18, unit pOgb]
- Qzwg Greenstone, greenschist, black phyllite, quartz-sericite schist, metakeratophyre, and minor marble**--Folded and otherwise-deformed, light to dark-green, greenschist- and local amphibolite-facies schist and gneiss derived from basaltic to andesitic volcanic rocks, volcanoclastic rocks, and some sedimentary rocks; also some schist is derived from meter-thick layered silicic volcanic rocks. Most common rock type is fine-grained (actinolite-)epidote-albite-chlorite schist. Highest grade rocks are garnet-biotite-hornblende-plagioclase gneiss near Kassa Inlet. Metamorphosed silicic volcanic rocks are quartz-porphyrific quartz-albite schist (that has been called metakeratophyre) that have locally preserved protolithic volcanic features. Unit thickness is at least several thousand meters. Age assignment used here is based on interpretation of intrusive relation described above for foliated, lineated, and gneissic biotite-hornblende quartz diorite, granodiorite, and minor diorite (Qqd) unit into this unit on southern Dall Island; that relation establishes a pre-Early Cambrian age for at least some protoliths of Wales Group (Gehrels, 1991). D.J. Grybeck reported (oral commun., April 22, 1993) that G.E. Gehrels has obtained a U-Pb age of about 595 Ma on a zircon sample from this unit near Niblack Anchorage on southeastern Prince of Wales Island; this determination is interpreted here to indicate that unit is of Late Proterozoic and Early Cambrian(?) age. Exposed on southern Prince of Wales and southern Dall Islands. [Description adapted from Eberlein and others, 1983, p. 23-24, unit PzpCs, p. 34-37, unit PzpOw; Gehrels and Saleeby, 1986, p. 14-15, unit pmOw; Gehrels and Saleeby, 1987, p. 126-127, informally named Wales metamorphic suite; Gehrels, 1991, units pOwb, pOwd, pOwr, pOws, pOwp, pOwu; Gehrels, 1992, p. 9-10, unit pOw]

**Marble and minor calcisilicate rocks**--Medium- to fine-grained, medium- to light-gray and tan marble. Present both as centimeter- to meter-thick layers in greenstone, greenschist, black phyllite, quartz-sericite schist, metakeratophyre, and minor marble (CZwg) unit and as massive to very thick (200 m) masses in localities shown on map at head of Hetta Inlet on Prince of Wales Island and on Long and Dall Islands. Massive parts have 5-cm- to 2-m-thick compositional layering. At Hetta Inlet unit is host to skarn deposits associated with a pluton of hornblende quartz monzodiorite with minor tonalite, granodiorite, quartz diorite, diorite, quartz monzonite, and monzodiorite (Kwqo) unit. [Description adapted from Eberlein and others, 1983, p. 37; Gehrels, 1991, unit pOwm]

#### UNITS EXPOSED NORTHEAST OF CLARENCE STRAIT

- Qs Surficial deposits (Holocene and (or) Pleistocene)**--Alluvium, colluvium, tidal mudflat deposits, and some glaciofluvial deposits. Distribution of most large areas of surficial deposits was mapped in field, but deposits have not been studied in detail. Many small areas of surficial deposits are not shown
- QTV Extrusive volcanic rocks (Quaternary? and Tertiary)**--Basalt and andesite exposed on Eagle and Muffin Islands at juncture of Clarence Strait and Ernest Sound. [Description adapted from Eberlein and others, 1983, p. 5-6, unit Tba; Brew, 1988, p. 6-7, 15, 16, unit QTb, 1994]
- Intrusive and related rocks of Kuiu-Etolin volcanic-plutonic province (Miocene and (or) Oligocene)**--Province named by Brew and others (1979) for area of its greatest exposure on Kuiu and Etolin Islands. Brew and Morrell (1983) and Brew (1988) provide general background information for these rocks. As mapped, divided into:
- Tsh Hornfelsed sedimentary rocks**--Composition of unit differs from locality to locality: (1) On southern Etolin, Brownson, and Deer Islands its protoliths are turbidites of Seymour Canal Formation of original Late Jurassic and Early Cretaceous age that are hornfelsed to albite-epidote hornfels facies metamorphic rocks and generally preserve both original structures and textures as well as metamorphic effects of Cretaceous metamorphic events. [Description adapted from Brew and others, 1984, p. 9, unit Tsh.] (2) On southwestern Revillagigedo Island metasedimentary rocks in vicinity of leucogabbro (Tlgb) unit are hornfelsed to spotted schist and hornfels. [Description adapted from Berg and others, 1988, p. 2, 17-18, hornfelsed unit MzPzms]
- Tvh Hornfelsed volcanic rocks**--Hornfelsed metavolcanic rocks in vicinity of leucogabbro (Tlgb) unit on southwestern Revillagigedo Island. [Description adapted from Berg and others, 1988, p. 2, 17-19, hornfelsed unit MzPzmv]
- Tme Migmatitic granitic rocks of southern Etolin Island**--Hornblende-biotite-pyroxene quartz monzodiorite, quartz monzonite, granodiorite, quartz diorite, and diorite paleosomes invaded by neosomes of these same compositions as well as of granite, alkali granite, and quartz syenite. Massive, extremely heterogeneous, and generally nonfoliated; hypidiomorphic to allotriomorphic; equigranular to seriate to porphyritic; generally fine- to medium-grained; C.I. 10 to 50 (paleosomes), 03 to 25 (neosomes). [Description adapted from Brew and others, 1984, p. 7, unit Tmme]
- Tge Granitic rocks of southern Etolin Island**--Hornblende-biotite granite, alkali granite, quartz syenite, and alkali quartz syenite. Massive, nonfoliated; allotriomorphic to hypidiomorphic; equigranular to seriate; medium- to coarse-grained; C.I. 01 to 07. Weathers to a distinctive pale orange to white; miarolitic cavities common, often rusty weathering; generally quite homogeneous at outcrop scale. [Description adapted from Brew and others, 1984, p. 7, unit Tmme]
- Tlgb Leucogabbro near Ketchikan**--Olivine-bearing (biotite-)two-pyroxene leucogabbro; forms crudely zoned stock at Deer Mountain and a small body on northeast shore of Gravina Island. [Description adapted from Koch and Elliott, 1984; Berg and others, 1988, p. 10, 15, unit Tgb]

**Tk Kootznahoo Formation (Paleogene)**--Conglomerate, sandstone, siltstone, shale, and minor lignite that are in general similar to rocks in Kootznahoo Formation southwest of Clarence Strait. Unit is everywhere probably less than 30 m thick and was deposited as alluvial fans, piedmont gravels, and deltas in areas of steep relief. Name and age assignments are based on correlation of these rocks with type Kootznahoo Formation on Admiralty Island and its outliers on Kuiu, Kupreanof, Zarembo, and northern Prince of Wales Islands. Exposed on northeastern Eagle Island, off south tip of Etolin Island, and south of Union Point, northeast headland of Union Bay. [Description adapted from Eberlein and others, 1983, p. 4-5, unit Ts; Brew and others, 1984, p. 10, unit Tk]

**Intrusive and related rocks of Admiralty-Revillagigedo plutonic province (Late Cretaceous)**--Province named by Brew and Morrell (1983) for northern and southern ends of province in southeastern Alaska. Table 1 gives age information; the first five-listed K-Ar ages in Ketchikan quadrangle are interpreted to be either cooling ages or to have been reset in by later magmatic events to northeast (Smith and others, 1979). Unit Ktif does not occur in this map area, but does crop out not far to the north (Brew and others, 1984, p. 21). Similarly dated rocks also present in lithically correlative units to northeast in the Bradfield Canal quadrangle (R.L. Elliott and R.D. Koch, oral commun., 1982). Rocks in these units in Petersburg quadrangle to north are described in detail by Burrell (1984a, b, c). As mapped, divided into:

Table 1. K-Ar and other age determinations of rocks from the Admiralty-Revillagigedo plutonic province. [Determinations from nearby in Petersburg quadrangle are by M. A. Lanphere (written commun., 1981, 1982, cited in Brew and others (1984)) and from this part of the Ketchikan quadrangle are those cited by Berg and others (1988). Map unit symbols are those on cited maps.]

Map unit	General location	K-Ar (Ma)			Pb-U (Ma) (Zircon)
		Biotite	Hornblende	Muscovite	
Petersburg quadrangle					
Ktif	Wrangell Is.	83.2	91.6	- - -	- - -
Ktif	Mitkof Is.	- - -	89.1	- - -	- - -
Ktef	Zarembo Is.	90.4	93.0	- - -	- - -
Ketchikan quadrangle					
Ktef	Bell Is.	74.2	80.2	- - -	- - -
Ktef	Revillagigedo Is.	70.9	85.4	- - -	- - -
Ktef	Cleveland Peninsula	- - -	- - -	89, 91	- - -
Ktef	Cleveland Peninsula	70.4	79.5	- - -	- - -
Ktef	Cleveland Peninsula	74.2	85.9	- - -	- - -
Ktef	Black Is.	74.4	81.8	- - -	- - -
Ktoc	Revillagigedo Is.	96.5	94.5	- - -	- - -
Ktqd	Spire Is.	- - -	- - -	- - -	80.1

**Krtn Nonfoliated plagioclase-porphyritic (hornblende-)(epidote-)garnet-biotite tonalite, quartz diorite, and minor granodiorite**--Nonfoliated crowded-plagioclase-phenocryst rock; inequigranular to porphyritic; very fine to medium-grained; C.I. 14 to 29; medium gray fresh; weathers light gray. Generally forms small elongate bodies less than 3 km<sup>2</sup> in area. Minerals include reddish-brown garnet, clinozoisite (or rarely epidote) and local muscovite; biotite and quartz commonly interstitial to closely spaced plagioclase laths. Some of epidote is magmatic (Zen and Hammarstrom, 1984). Unit is similar to the porphyritic, foliated biotite tonalite, quartz diorite, and granodiorite (Krtf) unit mineralogically, but differs texturally by its finer grain size and lack of large phenocrysts. Exposed on Brownson Island, Cleveland Peninsula, and Revillagigedo Island. [Description adapted from Brew and others, 1984, p. 21-22, unit Ktoc; Berg and others, 1988, p. 16, part of unit Kpg]

- Krtp Porphyritic, foliated biotite tonalite, quartz diorite, and granodiorite--** Prominently porphyritic, foliated, medium- to coarse-grained, C.I. 11 to 35, local cataclastic texture. Minerals include zoned, complexly twinned plagioclase, quartz, interstitial K-feldspar, partly chloritized biotite, epidote, minor local hornblende; and garnet, sphene, apatite, and allanite as accessories; in this map area lacks K-feldspar and shows moderate to extreme alteration of plagioclase, biotite, and garnet. Locally cut by pegmatite and basalt dikes; local inclusions of country rock. Exposed on southern Etolin Island and on Cleveland Peninsula. [Description adapted from Brew and others, 1984, p. 22, unit Ktgp; Berg and others, 1988, p. 16, part of unit Kpg]
- Krtf Foliated to massive hornblende-biotite tonalite and granodiorite, quartz monzonite, and quartz diorite--**Foliated to massive equigranular, average grain size is medium, but is fine grained near some margins, C.I. 17 to 50, light to medium gray where fresh, weathers brownish to dark gray. Minerals include zoned, complexly twinned plagioclase with minor alteration to sericite, mafic minerals usually biotite greater than hornblende, subhedral epidote, and local garnet and pyroxene; some of epidote is magmatic (Zen and Hammarstrom, 1984); accessory minerals are sphene, apatite, opaque mineral, and allanite. Foliation varies both in direction and development; locally semischistose and cataclastic. Contains aplite dikes, pegmatite dikes and veins, and rounded very fine grained hornblende diorite inclusions near pluton margins. Exposed on Cleveland Peninsula, Bell Island, and elsewhere in northeastern corner of map area. [Description adapted from Brew and others, 1984, p. 36, unit Ktef; Berg and others, 1988, p. 16, unit Kq]
- Krqd Quartz diorite--**Strongly sheared and altered brownish-gray, medium-grained, quartz diorite. Minerals include albite, quartz, muscovite, chlorite, epidote-clinozoisite, calcite, apatite, sphene, and pyrite. Exposed only on Spire Island, to northeast of Annette Island. [Description adapted from Berg and others, 1988, p. 10, unit Kqd]
- Krmg Migmatite--**Mainly agmatite and irregular banded gneiss, in zones between foliated to massive hornblende-biotite tonalite and granodiorite, quartz monzonite, and quartz diorite (Krtf), porphyritic, foliated biotite tonalite, quartz diorite, and granodiorite (Krtp), and hornfelsed sedimentary rocks (Tsh) units. Exposed only on east side of Ernest Sound. [Description adapted from Brew and others, 1984, p. 37, unit Kmgf]
- Intrusive rocks of Klukwan-Duke plutonic province (Early Cretaceous)--** Province named by Brew and Morrell (1983) for northern and southern ends in southeastern Alaska. Himmelberg and Loney (1995) summarized petrology of rocks in this belt. A variety of K-Ar and other dates (Lanphere and Eberlein, 1986; Saleeby, 1991; Meen and others, 1991a, b) indicate that age of ultramafic rocks is about 110-125 Ma (late Early Cretaceous). As mapped, divided into:
- Kku Ultramafic rocks, mainly dunite, peridotite, clinopyroxenite, and hornblendite--**(Olivine-)(chromite-)peridotite, (chromite-)dunite, (magnetite-)clinopyroxenite, and (magnetite-)hornblendite. Composition apparently differs from locality to locality: (1) On Annette Island: occurs in concentrically zoned complex with dunite and pyroxenite core and hornblende gabbro margin; (2) On Percy, and Revillagigedo Islands: present, but no detailed studies available for those localities. (3) On Duke Island not far to east of map area unit: Present in concentrically zoned complex with dunite and peridotite core and clinopyroxenite and hornblendite margin. [General description adapted from Berg and others, 1988, p. 11, unit Kum; description for Annette Island from Karl, 1992, units Kdu, Kpx, and Khg; comprehensive description of Duke Island body in Irvine, 1974]
- Kkdu Dunite, pyroxenite, and peridotite at Union Bay--**(Magnetite-)hornblende pyroxenite, (magnetite-)pyroxenite, (magnetite-)olivine pyroxenite, peridotite, and dunite. Rock types present in concentrically zoned pipe and lopolith with dunite in central part and pyroxenite and hornblende pyroxenite at margin. [Description adapted from Eberlein and others, 1983, p. 7, unit Kum; comprehensive description in Ruckmick and Noble, 1959]

- Kkgu Gabbro and diorite at Union Bay**--Moderately foliated to massive, medium-grained, (magnetite-) biotite-)gabbro and hornblende. Forms partial margin and carapace of younger dunite, pyroxenite, and peridotite (Kkdu) unit. [Description adapted from Eberlein and others, 1983, p. 7, unit Kgb; comprehensive description in Ruckmick and Noble, 1959]
- Sedimentary, volcanic, and related rocks of Gravina belt**--Gravina belt named by Berg and others (1972) for Gravina Island, which is in this map area. Brew and Karl (1988a, b, c) summarized stratigraphic relations; Ford and Brew (1978) reviewed volcanic rock compositions. Composite units of Berg and others (1988) reinterpreted by D.A. Brew and A.B. Ford (unpublished data, 1988) to include separate Upper Jurassic and Lower Cretaceous and Middle and Upper Jurassic units. As mapped, divided into:
- KJgg Gravina Island Formation (Early Cretaceous and Late Jurassic)**--Sedimentary and minor volcanic rocks. Basal, locally derived, polymictic conglomerate and grit grade upward and laterally into slaty but otherwise unmetamorphosed, dark-gray, thin-bedded graywacke and minor limestone. Graywacke locally contains *Buchia* of Late Jurassic age. Unit named by Berg (1973, p. 27-28). Exposed on southwestern Gravina Island. [Description adapted from Berg and others, 1988, p. 11-12, part of unit KJgs]
- Jgs Sedimentary and minor volcanic rocks (Late and Middle Jurassic)**--Regionally metamorphosed and deformed greenschist-facies metaflysch and meta-andesitic tuff. Metaflysch consists of phyllitic graywacke, argillite, polymictic conglomerate, grit, and minor silty limestone. Unit locally contains poorly preserved pectinid clams and belemnoids of possible Middle or Late Jurassic age (Berg, 1973, p. 27-28). Unit named by Berg (1973, p. 28). Exposed on northeastern Gravina and northern Annette Islands. [Description adapted from Berg and others, 1988, p. 11-12, part of unit KJgs]
- Jgv Andesitic to basaltic volcanic and volcanoclastic rocks and minor sedimentary rocks (Late and Middle Jurassic)**--Locally massive, but generally foliated and schistose metabasaltic tuff and agglomerate and minor metaandesitic tuff and agglomerate; flows are rare; also contains minor grayish-green and black phyllite and phyllitic siltstone that intertongue with metavolcanic rocks. Exposed on northeastern Gravina and northern Annette Islands. [Description adapted from Berg and others, 1988, p. 11-12, part of unit KJgv]
- Jgd Altered diorite and quartz diorite (Late and Middle Jurassic)**--Massive to locally moderately foliated, hydrothermally altered, actinolite-epidote-chlorite-sericite-sphene-amphibole(?) albite-quartz rock; present in elongate, zoned (from hypidiomorphic core to porphyritic margin) stock exposed on northern Annette Island. Relict minerals suggest that original composition was mostly pyroxene(?) hornblende diorite. Field relations suggest that it grades into rocks of adjacent andesitic to basaltic volcanic and volcanoclastic rocks and minor sedimentary rocks (Jgv) unit. [Description adapted from Berg and others, 1988, p. 11, unit KJgd]
- Hyd Group (Late Triassic)**--Unit named by Loney (1964, p. 43) on Admiralty Island and extended by Gehrels and others (1987, p. 877) into the area. As mapped, divided into:
- Tbc Chapin Peak Formation**--Basaltic volcanic and minor sedimentary rocks. Mainly basaltic pillow flows, calcareous agglomerate and volcanic breccia, and aquagene tuff that intertongue with minor conglomerate, limestone, and calcareous detrital sedimentary rocks near base. Unit is 150- to 500-m thick. Limestone beds and limestone clasts in conglomerate contain a variety of well-preserved Late Triassic (late Norian) fossils (Berg, 1973, p. 24). Unit named by Berg (1973, p. 23-26). Exposed on western Gravina and northern Annette Islands. [Description adapted from Berg and others, 1988, p. 12, unit Tbc]

- T<sub>hsv</sub>** **Sedimentary and mafic volcanic rocks**--Unit consists of Nehenta Formation, a unit named by Berg (1973, p. 19-23) on Gravina Island, and correlative rocks on northwestern Annette Island. Unit is about 500-m thick. As described includes three members: (1) carbonaceous limestone and siltstone member that grades into limestone, limestone breccia, and calcareous conglomerate, grit, and sandstone; (2) coarse-grained, trondjemite-clast-rich grit and conglomerate member; and (3) basaltic pillow flows, agglomerate, and aquagene tuff member. More metamorphosed and deformed on Annette Island than on Gravina. Age assignment is based on *Heterastridium* (late Norian) from upper part, and *Halobia* (of early and middle Norian age) from lower part of carbonaceous limestone and siltstone member, and on conodonts of latest Carnian and early Norian age from near base of unit on Annette Island (Savage and Gehrels, 1987). [Description adapted from Berg and others, 1988, p. 12-13, unit T<sub>sv</sub>]
- T<sub>hl</sub>** **Limestone and dolomite**--Massive to moderately thick bedded, dark-bluish-gray, very fine grained recrystallized limestone or marble. Unit is about 70 m thick. Age assignment is based on conodonts collected from unit to east of this map area (Berg, 1982, p. 5-7). Exposed on southern Gravina and northwestern Annette Islands. [Description adapted from Berg and others, 1988, p. 13, unit T<sub>l</sub>]
- T<sub>hv</sub>** **Felsic volcanic rocks**--Unit consists of Puppets Formation, a unit named by Berg (1973, p. 10-14) on Gravina Island, and correlative rocks on northwestern Annette Island. Unit is about 350 m thick. As described includes two main intertonguing members: (1) massive-appearing, thinly layered recrystallized rhyolite member, and (2) recrystallized rhyolitic to latitic tuff, and minor bedded tuffaceous limestone and calcareous tuff member. Second member occurs as discontinuous layers, generally near base of unit, that locally contain trondjemite clasts. Age assignment based on stratigraphic relation to overlying fossiliferous unit and a U-Pb apparent age of 225±3 Ma on zircon from a sample on Gravina Island (Gehrels and others, 1987, p. 877). [Description adapted from Berg and others, 1988, p. 13, unit T<sub>v</sub>]
- Metamorphosed sedimentary, volcanic, and intrusive rocks (Mesozoic and (or) Paleozoic)**--Relations of these unfossiliferous rocks to nearby unmetamorphosed paleontologically or otherwise-dated units are discussed by Berg and others (1988, p. 17-18). On this map, units are depicted more or less as shown by Berg and others (1988), however, Rubin and Saleeby (1991) presented a significantly different alternative interpretation. As mapped, divided into:
- MzP<sub>2</sub>ms** **Metasedimentary rocks**--Metamorphosed pelitic and semipelitic flysch with minor interlayered basaltic and (or) andesitic volcanic or volcanoclastic rocks. Dominantly dark-gray phyllite and fine-grained semischist with minor green phyllite and semischist to southwest on Etoin and Revillagigedo Islands and Cleveland Peninsula, grades into schist and gneiss to north and east. Typical metamorphic minerals to southwest are quartz, feldspar, biotite, garnet, muscovite, hornblende or actinolite, calcite, and chlorite; typical metamorphic minerals to northeast are quartz, plagioclase, biotite, garnet, hornblende, and muscovite. Sillimanite and other minerals occur in an aureole adjacent to the large pluton (Krtf) in the northeastern part of the map area. Protoliths of this unit probably include Gravina belt rocks, other Mesozoic rocks, and nearby Paleozoic rocks. Lower grade metamorphism to the southwest is post-mid-Cretaceous and pre-95 Ma in age; that to northeast is about 95 Ma (Brew and others, 1989a, 1992b). [Description adapted from Berg and others, 1988, p. 18, unit MzP<sub>2</sub>ms]

- MzPzmv**      **Metavolcanic rocks**--Metamorphosed marine andesitic and basaltic lava flows, agglomerate, and tuff, together with minor pelitic and semipelitic flysch. Exposures to southwest are massive-appearing layers and lenses of coarse, blocky, dark-green breccia with conspicuous relict augite phenocrysts and light-green laminated and lineated semischist and phyllite. Common greenschist-facies metamorphic minerals there are albite, epidote, calcite, quartz, chlorite, actinolite, and sphene. Exposures to northeast are dark-green, greenish-gray, and gray phyllite, semischist, and schist, together with minor marble. Common greenschist-facies metamorphic minerals there are albite, epidote, quartz, chlorite, blue-green hornblende and actinolite, and sphene. Exposures farthest north in map area are dark-green and dark-gray amphibolitic and quartzofeldspathic schist and gneiss with minor marble and pelitic schist. Common amphibolite-facies metamorphic minerals there are hornblende, biotite, quartz, plagioclase, and sphene. Protoliths of this unit probably include Gravina belt rocks, other Mesozoic rocks, and nearby Paleozoic rocks. Lower grade metamorphism to southwest is post-mid-Cretaceous and pre-95 Ma in age; that to northeast is about 95 Ma (Brew and others, 1989a, 1992b). Exposed on Cleveland Peninsula and Onslow and Revillagigedo Islands. [Description adapted from Berg and others, 1988, p. 18-19, unit MzPzmv]
- MzPzmc**      **Metapolyimictic conglomerate**--Spheroidal to ellipsoidal clasts to 25 cm maximum dimension of fine- to coarse-grained leucocratic (trondjemitic?) plutonic rocks, phyllite, quartzite, marble, pelitic schist, and semischist in matrix of fine-grained schist consisting of quartz, plagioclase, biotite, epidote-clinozoisite, garnet, calcite, muscovite, and pyrite. Generally in lenses less than 1-m thick. Exposed at Helm Bay, on Back Island, and at one locality north of Ketchikan on Revillagigedo Island. [Description adapted from Berg and others, 1988, p. 19-20, unit MzPzmc]
- MzPzmi**      **Metaintrusive rocks**--Regionally metamorphosed, recrystallized, or altered diorite, quartz diorite, and gabbro. Ages of protoliths and of metamorphism(s) are uncertain. Exposed on Cleveland Peninsula and Revillagigedo Island. [Description adapted from Berg and others, 1988, p. 20, unit MzPzmi]
- Pp**      **Pybus(?) Formation (Early Permian)**--Light-brown weathering, massive to laminated, light- dark-, and bluish-gray marble, together with minor phyllite, semischist, metapolyimictic conglomerate or sedimentary breccia, and quartzite-or-felsic-volcanic-rock fragmental schist and semischist. Age assignment is based on conodonts and brachiopods of late Early Permian (Leonardian) age. Unit named Pybus Dolomite by Loney (1964, p. 36) and redefined as Pybus Formation by Muffler (1967, p. C25). Unit questionably extended geographically into this area of this report to include age- and lithologically-equivalent rocks. [Description adapted from Berg and others, 1988, p. 20-21, unit Pm]
- Dka**      **Karheen Formation and associated rocks (Middle? and Early Devonian)**-- Karheen Formation was named by Eberlein and Churkin (1970a, p. 22) on Prince of Wales Island and was extended geographically by Gehrels and others (1987, p. 876-877) to age-equivalent rocks in vicinity of Hotspur Island. "Dkk" symbol indicates Karheen Formation rocks; "Dk" symbol indicates the associated rocks. As mapped, divided into:
- Dka**      **Andesitic flows and breccia (Middle? Devonian)**--Slightly deformed and metamorphosed andesitic pillow flows, tuff, tuff breccia, massive flows, hypabyssal intrusive rocks, and minor clastic beds. Unit is at least 800 m thick. Exposed on Hotspur Island. [Description adapted from Berg and others, 1988, p. 13-14, part of unit Dsv; Gehrels and others, 1987, p. 875-877, upper part of Early Devonian Karheen Formation and associated volcanic rocks unit]

- Dkkm **Mudstone-and-siltstone-facies rocks (Middle? and Early Devonian)--**  
Slightly deformed and metamorphosed interbedded gray mudstone and siltstone, crossbedded and channeled brownish-gray sandstone, gray tuffaceous mudstone, massive to thinly bedded limestone and marble, and olistostromal rocks consisting of meter-size olistoliths of intraformational limestone, siltstone, and sandstone in a matrix of chaotically deformed mudstone, siltstone, and sandstone. Unit is about 1,000 m thick. Megafossils (Berg, 1972) and conodonts (Savage and Gehrels, 1987) indicate an Early Devonian and early Middle(?) Devonian age. Exposed on Hotspur Island. [Description adapted from Berg and others, 1988, p. 13-14, part of unit Dsv; Gehrels and others, 1987, p.875-877, Karheen Formation]
- Stj **Trondhjemite, diorite, and granite of Annette Island (Late Silurian)--**Medium- to coarse-grained, light-gray, generally leucocratic (chlorite-)(biotite-)(hornblende) trondhjemite, quartz diorite, diorite, granite, quartz monzonite, and granodiorite. Pb-U (zircon) ages of 409±30, 410±5, and 415±5 Ma are reported by Gehrels and others (1987, p. 873) from this unit in this general area. Exposed on Annette Island and on southern and western Gravina Islands. [Description adapted from Berg and others, 1988, p. 14, unit St; Gehrels and others, 1987, p. 873, trondhjemite to granite unit]
- Intrusive rocks of younger part of Cape Chacon plutonic province (Early Silurian and Late Ordovician)--**Consists of:
- SOdi **Diorite, quartz diorite, gabbro, hornblendite, leucogabbro, trondhjemite, pyroxenite, and migmatite--**Locally heterogeneous, locally homogeneous; variably deformed and metamorphosed; locally foliated and layered. Biotite, hornblende, chlorite, quartz, plagioclase, and K-feldspar are dominant primary minerals; secondary minerals are epidote, white mica and calcite. Pb-U (zircon) ages of 429±20 and 426±15 Ma are reported by Gehrels and others (1987, p. 872) from this unit in this general area. Exposed on Annette and Gravina Islands. [Description adapted from Berg and others, 1988, p. 14-15, units SOu and SOui; Gehrels and others, 1987, p. 871-872, "Ordovician-Early Silurian" diorite unit]
- Metamorphosed rocks on Cleveland Peninsula (Silurian and (or) Ordovician)--**Contacts for units shown on map are based on Berg and others (1976) and on unpublished mapping and interpretation by D.A. Brew, A.B. Ford, and S.M. Karl (1987). Age assignment is based on zircon Pb-U age of cross-cutting dike (Rubin and Saleeby, 1987). As mapped, divided into:
- SOms **Metasedimentary rocks--**Metamorphosed mudstone, siltstone, sandstone, grit, conglomerate, and minor limestone. Generally phyllite to southwest and schist to northeast; hornfelsed adjacent to plutons. May include some rocks that should be assigned to unit MzPzms. [Description adapted from Eberlein and others, 1983, p. 5-6, part of metasedimentary rocks (MzPzms) unit; Berg and others, 1988, p. 14, part of unit SOu; p. 18, and part of metasedimentary rocks (MzPzms) unit; Rubin and Saleeby, 1991, fig. 2, includes parts of their "Ordovician-Silurian tonalite, diorite, & gabbro" unit (which is like unit SOdi of this map) and of their "Ordovician-Silurian basaltic andesite, tuff, breccia, pillowed flows and hypabyssal rocks" unit]
- SOmv **Metavolcanic rocks--**Metamorphosed andesitic and basaltic metatuff and agglomerate with minor pelitic rocks and limestone. Relict euhedral augite(?) phenocrysts and coarse- to fine-grained fragmental textures preserved locally. Now green phyllite to semischist with greenschist-facies minerals albite, epidote, chlorite, actinolite, quartz, muscovite, biotite, and calcite. [Description adapted from Eberlein and others, 1983, p. 5-6, part of andesitic and basaltic metatuff and agglomerate (MzPzvm) unit; Rubin and Saleeby, 1991, fig. 2, includes parts of their "Ordovician-Silurian tonalite, diorite, & gabbro" unit (which is like unit SOdi of this map) and of their "Ordovician-Silurian basaltic andesite, tuff, breccia, pillowed flows & hypabyssal rocks" unit]
- SOmc **Metacarbonate rocks--**Metamorphosed limestone; now fine-grained gray marble; exposed near Caamano Point. [Description from Brew and others, unpublished mapping; Berg and others, 1988, p. 14, subunit "m" of unit MzPzu]

**Descon Formation (Early Silurian to Early Ordovician)**--Unit named by Eberlein and Churkin (1970a, p. 5) on Prince of Wales Island and geographically extended by Gehrels and others (1987, p. 869-871) across Clarence Strait to east. As mapped, divided into:

- S0dr      **Dacitic to rhyolitic volcanic rocks**--Moderately deformed and metamorphosed dacitic to rhyolitic tuff and tuff-breccia with well preserved pyroclastic fragments and tuffaceous lamination and layering; some interbedded basaltic to andesitic pillow flows and tuff-breccia. Generally similar to dacitic to rhyolitic volcanic rocks unit described southwest of Clarence Strait. Exposed on western Annette and southern Gravina Islands. [Description adapted from Berg and others, 1988, p. 14, part of unit SOu; Gehrels and others, 1987, p. 869-871, Dacitic to rhyolitic volcanic rocks of Descon Formation]
- S0db      **Basaltic to andesitic volcanic rocks**--Basaltic to andesitic pillow flows, tuff, and tuff breccia with interbedded black argillite and mudstone; exposed on western and northern Annette Island. [Description adapted from Gehrels and others, 1987, p. 869-871, Basaltic to andesitic volcanic rocks of Descon Formation]
- Omgi      **Metamorphic rocks on southern Gravina Island (Late and Middle Cambrian)**--Foliated, slightly layered, metamorphosed diorite, microdiorite, quartz diorite, gabbro, microgabbro, and hornblendite. U-Pb (zircon) apparent age of a sample from Bronough Islands (off south tip of Gravina Island) is in range 540 to 510 Ma (Gehrels and others, 1987, p. 868-869). Correlates with unit to west and east of entrance to Cholmondeley Sound on southeastern Prince of Wales Island. Exposed on southern tip of Gravina Island. [Description adapted from Gehrels and others, 1987, p. 867, 869, Cambrian metaplutonic rocks unit]

## REFERENCES CITED

- Armstrong, R.L., 1985, Rb-Sr dating of the Bokan Mountain granite complex and its country rocks: Canadian Journal of Earth Sciences, v. 22, p. 1233-1236.
- Armstrong, A.K., 1970, Mississippian rugose corals, Peratrovich Formation, west coast, Prince of Wales Island, southeastern Alaska: U.S. Geological Survey Professional Paper 534, 44 p.
- Barker, J.C., and Mardock, C., 1988, Lithophile metal, REE-Y-Nb deposits on southern Prince of Wales Island, Alaska: The Minerals, Metals, and Materials Society, Process Mineralogy VIII, p. 139-157.
- Barker, J. C., and Mardock, C., 1990, Rare-earth element- and yttrium-bearing pegmatite dikes near Dora Bay, southern Prince of Wales Island: U.S. Bureau of Mines Open-File Report 19-97, 41 p.
- Barnes, D.F., Brew, D.A., and Morin, R.L., 1989, Bouguer gravity map of the Petersburg quadrangle and parts of the Port Alexander, Sitka, and Sumdum quadrangles, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1970-A, 21 p., scale 1:250,000.
- Barnes, D.F., Erwin, M.J., Holden, K.D., and Morin, R.L., 1975, USGS Alaskan gravity data maps of the Port Alexander, Sitka, Juneau, Mt. Fairweather, and Skagway 1:250,000 quadrangles, Alaska: U.S. Geological Survey Open-File Report 75-6, 5 maps, 58 p.
- Barnes, D.F., Olson, R.C., Holden, K.D., Morin, R.L., and Erwin, M.J., 1972a, Tabulated gravity data from southeastern Alaska obtained during the 1968 field season: U.S. Geological Survey Open-File Report, 82 p.
- Barnes, D.F., Popenoe, Peter, Olson, R.C., MacKenzie, M.V., and Morin, R.L., 1972b, Tabulated gravity data from southeastern Alaska obtained during the 1969 field season, 91 p.
- Beikman, H.M., 1974, Preliminary geologic map of the southeast quadrant of Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-612, 1 sheet, scale 1:1,000,000.
- Beikman, H.M., 1975, Preliminary geologic map of southeastern Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-673, 2 sheets, scale 1:1,000,000.
- Berdan, J.M., and Copeland, J.M., 1973, Ostracodes from Lower Devonian formations in Alaska and Yukon Territory: U.S. Geological Survey Professional Paper 825, 47 p.
- Berg, H.C., 1972, Geologic map of Annette Island, Alaska: U.S. Geological Survey Miscellaneous Investigations Map I-684, scale 1:63,360, 8 p.
- \_\_\_\_\_ 1973, Geology of Gravina Island, Alaska: U.S. Geological Survey Bulletin 1373, 41 p.
- \_\_\_\_\_ 1978, Map and tables describing areas of metalliferous mineral resource potential in the Ketchikan and Prince Rupert quadrangles, Alaska: U.S. Geological Survey Open-File Report 78-73-M, 52 p., 1 sheet, scale 1:250,000.
- \_\_\_\_\_ 1980, The Alaska Mineral Resource Assessment Program: Guide to information about the geology and mineral resources of the Ketchikan and Prince Rupert quadrangles, southeastern Alaska: U.S. Geological Survey Open-File Report 80-794, 43 p.
- \_\_\_\_\_ 1982, The Alaska Mineral Resource Assessment Program: Guide to information about the geology and mineral resources of the Ketchikan and Prince Rupert quadrangles, southeastern Alaska: U.S. Geological Survey Circular 855, 24 p.
- \_\_\_\_\_ 1984, Regional geologic summary, metallogenesis, and mineral resources of southeastern Alaska: U.S. Geological Survey Open-File Report 84-572, 298 p.

- Berg, H.C., Decker, J.E., and Abramson, B.S., 1981, Metallic mineral deposits of southeastern Alaska: U.S. Geological Survey Open-File Report 81-122, 136 p., 1 map, scale 1:1,000,000.
- Berg, H.C., Elliott, R.L., and Koch, R.D., 1978a, Table describing metalliferous and selected nonmetalliferous mineral deposits in the Ketchikan and Prince Rupert quadrangles, Alaska: U.S. Geological Survey Open-File Report 78-73B, 1 sheet, scale 1:250,000, 17 p. pamphlet.
- Berg, H.C., Elliott, R.L., Koch, R.D., Carten, R.B., and Wahl, F.A., 1976, Preliminary geologic map of the Craig D-1 and parts of the Craig C-1 and D-2 quadrangles, Alaska: U.S. Geological Survey Open-File Report 76-430, 1 sheet, scale 1:63,360.
- Berg, H.C., Elliott, R.L., and Koch, R.D., 1988, Geologic map of the Ketchikan and Prince Rupert quadrangles, southeastern Alaska: U.S. Geological Survey Map I-1807, 2 sheets, 27 p.
- Berg, H.C., Jones, D.L., and Richter, D.H., 1972, Gravina-Nutzotin belt--tectonic significance of an upper Mesozoic sedimentary and volcanic sequence in southern and southeastern Alaska, *in* Geological Survey Research 1972: U.S. Geological Survey Professional Paper 800-D, p. D1-D24.
- Berg, H.C., Jones, D.L., and Coney, P.J., 1978b, Map showing pre-Cenozoic tectono-stratigraphic terranes of southeastern Alaska and adjacent areas: U.S. Geological Survey Open-File Report 78-1085, scale 1:1,000,000.
- Berry, W.B.N., and Boucot, A.J., eds., 1970, Correlation of the North American Silurian rocks: Geological Society of America Special Paper 102, 289 p.
- Brew, D.A., 1988, Latest Mesozoic and Cenozoic igneous rocks of southeastern Alaska--a synopsis: U.S. Geological Survey Open-File Report 88-405, 53 p.
- \_\_\_\_\_, 1993, Regional geologic setting of mineral resources in southeastern Alaska, *in* Godwin, L.H., and Smith, B. D., eds., Economic mineral resources of the Annette Islands Reserve, Alaska: U.S. Dept. of the Interior, Bureau of Indian Affairs, Division of Energy and Mineral Resources Publication, p. 13-20.
- \_\_\_\_\_, 1994, Latest Mesozoic and Cenozoic magmatism in southeastern Alaska, *in* Platker, G., and Berg, H.C., eds., The geology of Alaska: Boulder, Colorado, Geological Society of America, The Geology of North America, v. G-1, Chap. 19, p. 621-656.
- Brew, D.A., Berg, H.C., Morrell, R.P., Sonnevill, R.S., and Hunt, S.J., 1979, The mid-Tertiary Kuiu-Etolin volcanic-plutonic belt, southeastern Alaska, *in* Johnson, K.M., and Williams, J.R., eds., The United States Geological Survey in Alaska: Accomplishments during 1978: U.S. Geological Survey Circular 804-B, p. B129-B130.
- Brew, D.A., Drew, L.J., and Ludington, S.D., 1992a, The study of the undiscovered mineral resources of the Tongass National Forest and adjacent lands, southeastern Alaska: Nonrenewable Resources, v. 1, no. 4, p. 303-321.
- Brew, D.A., Drew, L.J., Schmidt, L.M., Root, D.H., and Huber, D.F., 1991a, Undiscovered locatable mineral resources of the Tongass National Forest and adjacent areas, southeastern Alaska: U.S. Geological Survey Open-File Report 91-10, 370 p., 16 sheets, scales 1:250,000 and 1:500,000.
- Brew, D.A., and Drinkwater, J.L., 1991, Tongass Timber Reform Act Wilderness Areas supplement to U.S. Geological Survey Open-File Report 91-10 (Undiscovered locatable mineral resources of the Tongass National Forest and adjacent lands, southeastern Alaska): U.S. Geological Survey Open-File Report 91-343: 56 p.
- Brew, D.A., and Ford, A.B., 1978, Megalineament in southeastern Alaska marks southwest edge of Coast Range batholithic complex: Canadian Journal Earth Science, v. 15, no. 11, p. 1763-1772.

- \_\_\_\_\_ 1983, Comment on Monger, J.W.H., Price, R. A., and Tempelman-Kluit, D.J., 1982, Tectonic accretion and the origin of the two major metamorphic and plutonic belts in the Canadian Cordillera: *Geology*, v. 11, p. 427-429.
- \_\_\_\_\_ 1984, Tectonostratigraphic terranes in the Coast plutonic-metamorphic complex, southeastern Alaska, *in* Bartsch-Winkler, S., and Reed, K.M., eds., *The United States Geological Survey in Alaska: Miscellaneous geologic research 1982: U.S. Geological Survey Circular 939*, p. 90-93.
- Brew, D.A., Ford, A.B., and Himmelberg, G.R., 1989a, Evolution of the western part of the Coast plutonic-metamorphic complex, southeastern Alaska, U.S.A.--A synopsis, *in* Daly, S.R., ed., *Evolution of metamorphic belts: Geological Society of London Special Paper 43*, p. 447-452.
- Brew, D.A., Grybeck, D.J., Cathrall, J.B., Karl, S.M., Koch, R.D., Barnes, D.F., Newberry, R.J., Griscom, A., and Berg, H.C., 1989b, Mineral-resource map of the Petersburg and parts of the Port Alexander and Sumdum 1:250,000 quadrangles, southeastern Alaska: U.S. Geological Survey MF-1970-B, scale 1:250,000, 1 sheet, 47 p.
- Brew, D.A., Himmelberg, G.R., Loney, R.A., and Ford, A.B., 1992b, Styles of metamorphism in the southeastern Alaska part of the Cordilleran orogen: *Journal of Metamorphic Geology*, v. 10, p. 465-482.
- Brew, D.A., and Karl, S.M., 1988a, A reexamination of the contacts and other features of the Gravina belt, southeastern Alaska, *in* Galloway, J.P., and Hamilton, T.D., eds., *Geologic studies in Alaska by the U.S. Geological Survey during 1987: U.S. Geological Survey Circular 1016*, p. 143-146.
- \_\_\_\_\_ 1988b, A reexamination of the contacts and other features of the Gravina belt, southeastern Alaska, Supplemental data: U.S. Geological Survey Open-File Report 88-652, 8 p.
- \_\_\_\_\_ 1988c, A reexamination of the contacts and other features of the Gravina Belt, southeastern Alaska (abs.): *Geological Society of America, Abstracts with Programs*, v. 20, no. 7, p. 111.
- Brew, D.A., Karl, S.M., Barnes, D.F., Jachens, R.C., Ford, A.B., and Horner, R., 1991b, A northern Cordilleran ocean-continent transect: Sitka Sound to Atlin Lake, British Columbia: *Canadian Journal of Earth Sciences*, v. 28, no. 6, p. 840-853.
- Brew, D.A., and Morrell, R.P., 1983, Intrusive rocks and plutonic belts in southeastern Alaska. *in* Roddick, J.A., ed., *Circum-Pacific plutonic terranes: Geological Society of America Memoir 159*, p. 171-193.
- Brew, D.A., Ovenshine, A.T., Karl, S.M., and Hunt, S.J., 1984, Preliminary reconnaissance geologic map of the Petersburg and parts of the Port Alexander and Sumdum 1:250,000 quadrangles, southeastern Alaska: U.S. Geological Survey Open-File Report 84-405, 2 sheets, 43 p. pamphlet.
- Brooks, A.H., 1902, Preliminary report on the geology of the Ketchikan mining district, Alaska, with an introductory sketch of the geology of southeastern Alaska: U.S. Geological Survey Professional Paper 1, 120 p.
- Buddington, A.F., and Chapin, T., 1929, Geology and mineral deposits of southeastern Alaska: U.S. Geological Survey Bulletin 800, 398 p.
- Bufvers, J., 1967, History of mines and prospects, Ketchikan district, prior to 1952: Alaska Division of Geological and Geophysical Surveys Special Report SR 1, 32 p.
- Bundtzen, T.K., 1978, The Prince of Wales copper mining industry: *Alaska Mines and Geology Bulletin*, v. 27, no. 2, p. 11-13.

- Burchard, E.F., 1913, Marble resources of the Ketchikan and Wrangell districts: U.S. Geological Survey Bulletin 542, p. 52-77.
- Burchard, E.F., 1920, Marble resources of southeastern Alaska: U.S. Geological Survey Bulletin 682, 118 p.
- Burrell, P.D., 1984a, Cretaceous or younger plutonic rocks, Mitkof and Kupreanof Islands, southeast Alaska, *in* Coonrad, W.C., and Elliott, R.L., eds., The United States Geological Survey in Alaska: Accomplishments during 1981: U.S. Geological Survey Circular 868, p. 124-126.
- \_\_\_\_\_ 1984b, Late Cretaceous plutonic rocks, Petersburg quadrangle, southeast Alaska, *in* Bartsch-Winkler, S., and Reed, K.M., eds., The United States Geological Survey in Alaska: Accomplishments during 1982: U.S. Geological Survey Circular 939 p. 93-96.
- \_\_\_\_\_ 1984c, Map and tables describing the Admiralty-Revillagigedo intrusive belt plutons in the Petersburg 1:250,000 quadrangle, southeastern Alaska: U.S. Geological Survey Open-File Report 84-171, 6 p., 1 sheet, scale 1:250,000.
- Carter, C., Trexler, J.H., Jr., and Churkin, M., Jr., 1980, Dating of graptolite zones by sedimentation rates: implications for rates of evolution: *Lethaia*, v. 13, p. 279-287.
- Cathrall, J.B., 1994, Geochemical survey of the Craig study Area--Craig, Dixon Entrance, and western edges of the Ketchikan and Prince Rupert quadrangles, southeast Alaska: U.S. Geological Bulletin 2082, 52 p., 7 tables, 1 sheet scale 1:250,000.
- Cathrall, J.B., Arbogast, B.F., VanTrump, G., Jr., and McDanal, S.K., 1993a, Geochemical maps showing the distribution of selected anomalous elements in stream sediments from the Craig, Dixon Entrance, and western edges of the Ketchikan and Prince Rupert quadrangles, southeast Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-2217-A, 2 sheets, scale 1:250,000.
- Cathrall, J.B., McDanal, S.K., VanTrump, G., Jr., Arbogast, B.F., and Grybeck, D., 1993b, Geochemical maps showing the distribution of selected anomalous elements in nonmagnetic heavy-mineral concentrates from the Craig study area, southeast Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-2217-B, 2 sheets, scale 1:250,000.
- Churkin, M., Jr., Carter, C., and Eberlein, G.D., 1970, Graptolite succession across the Ordovician-Silurian boundary in southeastern Alaska: *Quarterly Journal of the Geological Society of London*, v. 126, p. 319-330.
- Churkin, M., Jr., and Eberlein, G.D., 1975, Geologic map of the Craig C-4 quadrangle, Alaska: U.S. Geological Survey Geologic Quadrangle Map GQ-1169, scale 1:63,360.
- Churkin, M., Jr., Eberlein, G.D., Hueber, F.M., and Mamay, S.H., 1969, Lower Devonian land plants from graptolitic shale in south-eastern Alaska: *Paleontology*, v. 12, p. 559-573.
- Churkin, M., Jr., Jager, H., and Eberlein, G.D., 1970, Lower Devonian graptolites from southeastern Alaska: *Lethaia*, v. 3, no. 2, p. 183-202.
- Clark, A.L., Berg, H.C., Grybeck, D., and Ovenshine, A.T., 1971, Reconnaissance geology and geochemistry of Forrester Island National Wildlife Refuge, Alaska: U.S. Geological Survey Open-File Report 71-67, 9 p., 1 sheet, scale 1:63,360.
- Clark, A.L., and Greenwood, W.R., 1972, Geochemistry and distribution of platinum-group metals in mafic to ultramafic complexes of southern and southeastern Alaska, *in* Geological Survey Research: U.S. Geological Survey Professional Paper 800-C, p. C157-C160.

- Clautice, K.H., Gilbert, W.G., Wiltse, M.A., Werdon, M.B., Hozma, T.X., 1994, Geology of the Helm Bay area, portions of the Craig C-1 and Ketchikan C-6 quadrangles, southeastern Alaska: Alaska Division of Geological and Geophysical Surveys Public-Data File 94-41, one sheet, scale 1:63,360.
- Cobb, E.H., 1972a, Metallic mineral resources map of the Craig quadrangle, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-433, 1 sheet, scale 1:250,000.
- \_\_\_\_\_ 1972b, Metallic mineral resources map of the Dixon Entrance quadrangle, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-434, 1 sheet, scale 1:250,000.
- \_\_\_\_\_ 1972c, Metallic mineral resources map of the Ketchikan quadrangle, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-420, 1 sheet, scale 1:250,000.
- \_\_\_\_\_ 1972d, Metallic mineral resources map of the Prince Rupert quadrangle, Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-437, 1 sheet, scale 1:250,000.
- \_\_\_\_\_ 1978a, Summary of references to mineral occurrences (other than mineral fuels and construction materials) in the Craig quadrangle, Alaska: U.S. Geological Survey Open-File Report 78-869, 262 p.
- \_\_\_\_\_ 1978b, Summary of references to mineral occurrences (other than mineral fuels and construction materials) in the Dixon Entrance quadrangle, Alaska: U.S. Geological Survey Open-File Report 78-863, 34 p.
- Cobb, E.H., and Elliott, R.L., 1980, Summaries of data and lists of references to metallic and selected nonmetallic mineral deposits in the Ketchikan and Prince Rupert quadrangles, Alaska: U.S. Geological Survey Open-File Report 80-1053, 157 p.
- Cobb, E.H., Wanek, A.A., Grantz, A., and Carter, C., 1968, Summary report on the geology and mineral resources of the Bering Sea, Bogoslof, Simeonof, Semidi, Tuxedni, St. Lazaria, Hazy Islands, and Forrester Island National Wildlife Refuges, Alaska: U.S. Geological Survey Bulletin 1260-K, p. K1-K28.
- Coldwell, J.R., 1990, An economic analysis—Tongass land management plan—mineral resource inventory—inferred (identified) reserves: Unpublished U.S. Bureau of Mines Report, 154 p. (This report is similar to that referred to as Coldwell (1989) in U.S. Forest Service (1990).)
- Condon, W.H., 1961, Geology of the Craig quadrangle, Alaska: U.S. Geological Survey Bulletin 1108-B, 43 p.
- Decker, John, 1979, Preliminary aeromagnetic map of southeastern Alaska: U.S. Geological Survey Open-File Report 79-1694, 1 sheet, scale 1:1,000,000.
- Decker, John, Mullen, M.W., and Schwab, C.E., 1981, Aeromagnetic profile map of southeastern Alaska: U.S. Geological Survey Open-File Report 82-505, scale 1:1,000,000.
- De Saint Andre, B., Lancelot, J.R., and Collot, B., 1983, U-Pb chronology of the Bokan Mountain peralkaline granite, southeastern Alaska: Canadian Journal of Earth Sciences, v.20, p. 236-245.
- Detra, D.E., Motooka, J.M., and Cathrall, J.B., 1992, Supplemental analytical results and sample locality map of stream-sediment and heavy-mineral-concentrate samples from the Craig study area: Dixon Entrance, Ketchikan, and Prince Rupert quadrangles, Alaska: U.S. Geological Survey Open-File Report 92-552-A, 16 p., scale 1:250,000, 1 sheet, tables 1-4 in digital format or 5.25" floppy disk in pocket.
- Douglass, R.C., 1971, Pennsylvanian fusulinids from southeastern Alaska: U.S. Geological Survey Professional Paper 706, 20 p.

- Douglass, S.L., Webster, J.H., Burrell, P.D., Lanphere, M.L., and Brew, D.A., 1989, Major element chemistry, radiometric values, and locations of samples from the Petersburg and parts of the Port Alexander and Sumdum quadrangles, southeastern Alaska: U.S. Geological Survey Open-File Report 89-527, 66 p., scale 1:250,000.
- Drinkwater, J.L., and Calzia, J.P., 1994, Geochemical reconnaissance of alkalic plutons on Prince of Wales Island, southeastern Alaska, *in* Till, A.B., and Moore, T.E., eds., *Geologic studies in Alaska by the U.S. Geological Survey, 1993*: U.S. Geological Survey Bulletin 2107, p.173-184.
- Eakins, G.R., 1970, An experiment in geobotanical prospecting for uranium, Bokan Mountain area, southeastern Alaska: Alaska Division of Geological and Geophysical Surveys Report 41, 52 p.
- Eberlein, G.D., and Churkin, M., Jr., 1970a, Paleozoic stratigraphy of the northwest coastal area of Prince of Wales Island, southeastern Alaska: U.S. Geological Survey Bulletin 1284, 67 p.
- \_\_\_\_\_ 1970b, Tlevak basalt, west coast of Prince of Wales Island, *in* Changes in stratigraphic nomenclature by the U.S. Geological Survey, 1968: U.S. Geological Survey Bulletin 1294-A, p. A25-A28, 67 p.
- \_\_\_\_\_ 1976, Early Paleozoic volcanic center and associated iron-copper deposits, *in* Geological Survey Research 1976: U.S. Geological Survey Professional Paper 1000, p. 89.
- Eberlein, G.D., Churkin, M., Jr., Carter, C., Berg, H.C., and Ovenshine, A.T., 1983, Geology of the Craig quadrangle, Alaska: U.S. Geological Survey Open-File Report 83-91, 4 sheets, 50 p.
- Elliott, R.L., Berg, H.C., and Karl, S.M., 1978, Map and table describing metalliferous and selected non-metalliferous mineral deposits, Ketchikan and Prince Rupert quadrangles, Alaska: U.S. Geological Survey Open-File Report 78-73-B, 18 p., 1 sheet, scale 1:250,000.
- Faulhuaber, J.J., 1977, Late Mississippian (Late Osage through Chesterian) conodonts from the Peratrovich Formation, southeastern Alaska: Unpublished M.S thesis, University of Oregon, 126 p.
- Ford, A.B., and Brew, D.A., 1988, The Douglas Island Volcanics; basaltic-rift--not andesitic-arc--volcanism of the "Gravina-Nutzotin belt," northern southeastern Alaska (abs.): Geological Society of America Abstracts with Programs, v. 20, no. 7, p. 111.
- Gardner, M.C., Bergman, S.C., Cushing, G.W., MacKevett, E.M., Jr., Plafker, G., Campbell, F.B., Dodds, C.J., McClelland, W.C., and Mueller, P.A., 1988, Pennsylvanian pluton stitching of Wrangellia and the Alexander terrane, Wrangell Mountains, Alaska: *Geology*, v. 16, p. 967-971.
- Gault, H.R., 1945, The Salt Chuck copper-palladium mine, Prince of Wales Island, southeastern Alaska: U.S. Geological Survey Open-File Report 46-19, 16 p..
- Gault, H.R., Wahrhaftig, C., and Loney, R.A., 1992, The Salt Chuck copper-palladium mine, Prince of Wales Island, southeastern Alaska: U.S. Geological Survey Open-File Report 92-293, 10 p.
- Gehrels, G.E., 1990, Late Proterozoic-Cambrian metamorphic basement of the Alexander terrane on Long and Dall Islands, southeast Alaska: *Geological Society of America Bulletin*, v. 102, p. 760-767.
- \_\_\_\_\_ 1991, Geologic map of Long Island and southern and central Dall Island, southeastern Alaska: U.S. Geological Survey Map MF-2146, 1 sheet.
- \_\_\_\_\_ 1992, Geologic map of the southern Prince of Wales Island, southeastern Alaska: U.S. Geological Survey Miscellaneous Investigations Series Map I-2169, 1 sheet, scale 1:63,360, 23 p. [Supersedes Gehrels and Saleeby (1986), USGS OFR 86-275.]

- Gehrels, G.E., Berg, H.C., and Saleeby, J.B., 1983, Ordovician-Silurian volcanogenic marine sulfide deposits on the southern Prince of Wales Island and the barrier islands, southeastern Alaska: U.S. Geological Survey Open-File Report 83-318, 10 p., 1 sheet.
- Gehrels, G.E., McClelland, W.C., Samson, S.D., Patchett, P.J., and Jackson, J.L., 1990, Ancient continental margin assemblage in the northern Coast Mountains, southeast Alaska and northwest Canada: *Geology*, v. 18, p. 208-211.
- Gehrels, G.E., and Saleeby, J.B., 1986, Geologic map of southern Prince of Wales Island, southeastern Alaska: U.S. Geological Survey Open-File Report 86-275, 1 sheet, 33 p. [Superseded by Gehrels (1992), USGS Map I-2169.]
- \_\_\_\_\_ 1987, Geology of southern Prince of Wales Island, southeastern Alaska: *Geological Society of America Bulletin*, v. 98, p. 123-137.
- Gehrels, G.E., Saleeby, J.B., and Berg, H.C., 1987, Geology of Annette, Gravina, and Duke Islands, southeastern Alaska: *Canadian Journal of Earth Sciences*, v. 24, p. 866-881.
- Godwin, L.H., and Smith, B. D., eds., Economic mineral resources of the Annette Islands Reserve, Alaska: U.S. Dept. of the Interior, Bureau of Indian Affairs, Division of Energy and Mineral Resources Publication, 136 p.
- Grybeck, D., Berg, H.C., and Karl, S.M., 1984, Map and description of the mineral deposits in the Petersburg and eastern Port Alexander quadrangles, southeastern Alaska: U.S. Geological Survey Open-File Report 84-0837, 87 p., 1 sheet, scale 1:250,000.
- Hahn, G., and Hahn, R., 1991, Trilobiten aus dem Karbon von SE-Alaska, Teil 1: *Geologica et Palaeontologica*, v. 25, p. 147-191.
- Hahn, G., and Hahn, R., 1992, Trilobiten aus dem Karbon von SE-Alaska, Teil 2: *Geologica et Palaeontologica*, v. 26, p. 99-133.
- Hansen, W.R., 1991, Suggestions to authors of the reports of the United States Geological Survey, Seventh edition: U.S. Government Printing Office, 289 p.
- Heaton, T.H., 1995, Middle Wisconsin bear and rodent remains discovered on Prince of Wales Island, Alaska: *Current Research in the Pleistocene*, v. 12, p. 92-95.
- Heaton, T.H., and Grady, F., 1992a, Preliminary report on the fossil bears of El Capitan Cave, Prince of Wales Island, Alaska: *Current Research in the Pleistocene*, v. 9, p. 97-99.
- Heaton, T.H., and Grady, F., 1992b, Two species of bear found in Late Pleistocene/Early Holocene den in El Capitan Cave, Prince of Wales Island, southern Alaska coast (abs.): *Journal of Vertebrate Paleontology*, v. 12, p. 32A.
- Heaton, T.H., and Grady, F., 1993, Fossil grizzly bears (*Ursus arctos*) from Prince of Wales Island, Alaska, offer new insights into animal dispersal, interspecific competition, and age of deglaciation: *Current Research in the Pleistocene*, v. 10, p. 98-100.
- Herreid, G., 1964, Geology of the Niblack Anchorage area, southeastern Alaska: Alaska Division of Geology and Geophysical Surveys Geological Report 5, 10 p.
- \_\_\_\_\_ 1967, Geology and mineral deposits of the Dolomi area, Prince of Wales Island, Alaska: Alaska Division of Geology and Geophysical Surveys Geological Report 27, 29 p., 2 sheets, scale 1:24,000.
- \_\_\_\_\_ 1971, Analyses of rock and stream-sediment samples, Hetta Inlet area, Prince of Wales Island, Alaska: Alaska Division of Geology and Geophysical Surveys Geochemical Report 24, 1 sheet, scale 1:24,000.

- Herreid, G., Bundtzen, T.K., and Turner, D.L., 1978, Geology and geochemistry of the Craig A-2 quadrangle and vicinity, Prince of Wales Island, Craig quadrangle, Alaska: Alaska Division of Geological and Geophysical Surveys Geochemical Report 48, 49 p.
- Herreid, G., and Rose, A.W., 1966, Geology and geochemistry of the Hollis and Twelvemile Creek areas, Prince of Wales Island, southeastern Alaska: Alaska Division of Geological and Geophysical Surveys Geologic Report 17, 35 p.
- Herreid, G., and Tribble, T.C., 1973, Analyses of rock and stream-sediment samples, Craig A-2 and vicinity, Prince of Wales Island, southeastern Alaska: Alaska Division of Geology and Geophysical Surveys Geochemical Report 27, 1 sheet, scale 1:63,360.
- Himmelberg, G.R., and Loney, R.A., 1995, Characteristics and petrogenesis of Alaskan-type ultramafic intrusions, southeastern Alaska: U.S. Geological Survey Professional Paper 1564, 47 p..
- Horton, R., Karl, S., Griscom, A., Taylor, C., Bond, B., and Senterfit, M, 1992, Mineral resource assessment of the Annette Islands Reserve, *in* Manydeeds, S.A., ed., 1992 Mineral Frontiers on Indian Lands: U.S. Department of the Interior, Bureau of Indian Affairs, Division of Energy and Mineral Resources, Publication G-92-2, p. 11-19.
- Irvine, T.N., 1974, Petrology of the Duke Island ultramafic complex, southeastern Alaska: Geological Society of America Memoir 138, 240 p.
- Jones, D.L., Irwin, W.P., and Ovenshine, A.T., 1972, Southeastern Alaska-A displaced continental fragment?, *in* Geological Survey Research 1972: U.S. Geological Survey Professional Paper 800-B, p. B211-B217.
- Karl, S.M., 1992, Map and table of mineral deposits on Annette Island, Alaska: U.S. Geological Survey Open-File Report 92-690, 57 p. pamphlet, 1 sheet, scale 1:63,360.
- Karl, S.M., and Giffen, C.F., 1992, Sedimentology of the Bay of Pillars and Point Augusta Formations, Alexander Archipelago, Alaska, *in* Bradley, D.C., and Dusel-Bacon, C., eds., Geologic studies in Alaska by the U.S. Geological Survey, 1991: U.S. Geological Survey Bulletin 2041, p. 171-185.
- Karl, S.M., and Koch, R.D., 1990, Maps and preliminary interpretation of anomalous rock geochemical data from the Petersburg quadrangle, and parts of the Port Alexander, Sitka, and Sumdum quadrangles, southeastern Alaska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1970-C, 40 p., 7 sheets, scale 1:250,000.
- Kennedy, G.C., 1953, Geology and mineral deposits of Jumbo basin, southeastern Alaska: U.S. Geological Survey Professional Paper 251, 46 p.
- Kirk, E., and Amsden, T.W., 1952, Upper Silurian brachiopods from southeastern Alaska: U.S. Geological Survey Professional Paper 233-C, p. 53-66.
- Koch, R.D., and Elliott, R.L., 1984, Late Oligocene gabbro near Ketchikan, southeastern Alaska, *in* Coonrad, W.L., and Elliott, R.L., eds., The United States Geological Survey in Alaska: Accomplishments during 1981: U.S. Geological Survey Circular 868, p. 126-128.
- Lanphere, M.A., and Eberlein, G.D., 1966, Potassium-argon ages of magnetite-bearing ultramafic complexes in southeastern Alaska (abs.): Geological Society of America Special Paper 87, p. 94.
- Lanphere, M.A., MacKevett, E.M., Jr., and Stern, T.W., 1964, Potassium-argon and lead-alpha ages of plutonic rocks, Bokan Mountain area, Alaska: Science, v. 145, p. 705-707.
- Lathram, E.H., Pomeroy, J.S., Berg, H.C., and Loney, R.A., 1965, Reconnaissance geology of Admiralty Island, Alaska: U.S. Geological Survey Bulletin 1181-R, p. B1-R48, 2 pls., scale 1:250,000.

- Loney, R.A., 1964, Stratigraphy and petrography of the Pybus-Gambier area, Admiralty Island, Alaska: U.S. Geological Survey Bulletin 1178, 103 p.
- Loney, R.A., and Himmelberg, G.R., 1992, Petrogenesis of the Pd-rich intrusion at Salt Chuck, Prince of Wales Island: An Early Paleozoic Alaskan-type ultramafic body: *Canadian Mineralogist*, v. 30, p. 1005-1022.
- Loney, R.A., Himmelberg, G.R., and Shew, N., 1987, Salt Chuck palladium-bearing ultramafic body, Prince of Wales Island, *in* Hamilton, T.D., and Galloway, J.P., eds., *Geologic studies in Alaska by the U.S. Geological Survey during 1986*: U.S. Geological Survey Circular 998, p. 126-127.
- Maas, K.M., Still, J.C., Clough, A.H., and Oliver, L.K., 1991, Mineral investigations in the Ketchikan mining district, Alaska, 1990: Southern Prince of Wales Island and vicinity--Preliminary sample location maps and descriptions: U.S. Bureau of Mines Open-File Report 33-91, 139 p.
- MacKevett, E.M., Jr., 1963, Geology and ore deposits of the Bokan Mountain uranium-thorium area, southeastern Alaska: U.S. Geological Survey Bulletin 1154, 125 p.
- Mamet, B.L., Pinard, S., and Armstrong, A.K., 1993, Micropaleontological zonation (Foraminifers, algae) and stratigraphy, Carboniferous Peratrovich Formation, southeastern Alaska: U.S. Geological Survey Bulletin 2031, 64 p.
- McDanaI, S.K., Arbogast, B.F., and Cathrall, J.B., 1991, Analytical results and sample locality map of stream-sediment, heavy-mineral-concentrate, pebble, and rock samples from the Craig study area; Craig, Dixon Entrance, Ketchikan, and Prince Rupert quadrangles, Alaska: U.S. Geological Survey Open-File Report 91-36A, 1 sheet, scale 1:250,000, 122 p.
- Meen, J.K., Ross, D.K., Elthon, D., 1991a, Gross isotopic heterogeneity in layered ultramafic cumulates (abs.): EOS, American Geophysical Union Transactions, v. 72, no. 44, p. 521.
- Meen, J.K., Snee, L.W., Ross, D.K., Elthon, D., 1991b, Age and geologic relations of the Hall Cove Complex, Duke Island, southeastern Alaska (abs.): Geological Society of America Abstracts with Programs, v. 23, no. 5, p. A389.
- Monger, J.W.H., and Berg, H.C., 1987, Lithotectonic terrane map of western Canada and southeastern Alaska: U.S. Geological Survey Map MF-1874-B, one sheet, scale 1:2,500,000, 12 p.
- Monger, J.W.H., Price, R.A., and Tempelman-Kluit, D.J., 1982, Tectonic accretion and the origin of the two major metamorphic and plutonic belts in the Canadian Cordillera: *Geology*, v. 10, p. 70-75.
- Monger, J.W.H., and Ross, C.A., 1971, Distribution of fusulinaceans in the western Canadian Cordillera: *Canadian Journal of Earth Sciences*, v. 8, p. 259-278.
- Motyka, R.J., and Moorman, M.A., 1987, Geothermal resources of southeastern Alaska: Alaska Division of Geological and Geophysical Surveys Professional Paper 93, scale 1:1,000,000.
- Muffler, L.J.P., 1967, Stratigraphy of the Keku Islets and neighboring parts of Kuiu and Kupreanof Islands, southeastern Alaska: U.S. Geological Survey Bulletin 1241-C, p. C1-C52.
- Nokleberg, W.J., Bundtzen, T.K., Berg, H.C., Brew, D.A., Grybeck, D., Robinson, M.S., Smith, T.E., and Yeend, W., 1987, Significant metalliferous lode deposits and placer districts of Alaska: U.S. Geological Survey Bulletin 1786, 104 p., 2 pls, scale 1:5,000,000.
- Nokleberg, W.J., Bundtzen, T.K., Berg, H.C., Brew, D.A., Grybeck, D., Robinson, M.S., Smith, T.E., and Yeend, W., and 33 contributors, 1994a, Metallogeny and major mineral deposits of Alaska, *in* Plafker, G., and Berg, H.C., eds., *The geology of Alaska: Boulder, Colorado, Geological Society of America, The Geology of North America*, v. G-1, Chap. 29, p. 855-903.

- Nokleberg, W.J., Bundtzen, T.K., Berg, H.C., Brew, D.A., Grybeck, D.J., Robinson, M.S., Smith, T.E., Yeend, W., 1994b, Map showing locations of metalliferous lode deposits and placer districts of Alaska, *in* Plafker, G., and Berg, H.C., eds., *The geology of Alaska: Boulder, Colorado, Geological Society of America, The Geology of North America, v. G-1 (GNA-G1), Plate 11, 1:2,500,000 scale.*
- Nokleberg, W.J., Bundtzen, T.K., Brew, D.A., and Plafker, G., 1995, Metallogeneses and tectonics of porphyry Cu and Mo (Au, Ag) and granitoid-hosted Au deposits of Alaska, *in* Schroeter, T.G., ed., *Porphyry deposits of the northwestern Cordillera-A sequel to CIM Special Volume 15, Canadian Institute of Mining, Metallurgy, and Petroleum Special Volume 46, p. 103-141.*
- Nokleberg, W.J., Moll-Stalcup, E.J., Miller, T.P., Brew, D.A., Grantz, A., Reed, J.C., Plafker, G., Moore, T.E., Silva, S.R., and Patton, W.W., Jr., 1994c, Tectonostratigraphic terrane and overlap assemblage map of Alaska: U.S. Geological Survey Open-File Report 94-194, 1 sheet, scale 1:2,500,000, 53p.
- Nokleberg, W.J., Moll-Stalcup, E.J., Grantz, A., Plafker, G., Miller, T.P., Brew, D.A., and Moore, T.E., in press, Tectonic-magmatic map of Alaska: U.S. Geological Survey Miscellaneous Investigations Series Map, scale 1:2,500,000.
- North American Commission on Stratigraphic Nomenclature, 1983, North American stratigraphic code: American Association of Petroleum Geologists Bulletin, v. 67, no. 5, p. 841-875.
- Oliver, W.A., Jr., Merriam, C.W., and Churkin, M., Jr., 1975, Ordovician, Silurian, and Devonian corals of Alaska: U.S. Geological Survey Professional Paper 823-B, p. B12-B57.
- Ovenshine, A.T., 1975, Tidal origin of part of the Karheen Formation (Lower Devonian), southeastern Alaska, *in* Ginsburg, R.N., ed., *Tidal deposits: a case book of recent examples of fossil counterparts: Springer-Verlag, p. 141-148.*
- Ovenshine, A.T., and Brew, D.A., 1972, Separation and history of the Chatham Strait fault, southeast Alaska, North America: 24th International Geological Congress Section 3, p. 245-254.
- Ovenshine, A.T., and Webster, G.D., 1970, Age and stratigraphy of the Heceta Limestone in northern Sea Otter Sound, southeastern Alaska, *in* Geological Survey Research 1970: U.S. Geological Survey Professional Paper 700-C, p. C170-C174.
- Philpotts, J., Taylor, C., Evans, J., and Emsbo, P., 1993, Newly discovered molybdenite occurrences at Dora Bay, Prince of Wales Island, southeast Alaska, and preliminary scanning electron microscope studies, *in* Dusel-Bacon, C., and Till, eds., *Geologic studies in Alaska by the U.S. Geological Survey, 1992: U.S. Geological Survey Bulletin 2068, p.187-196.*
- Redman, E., 1982, The Keete Inlet thrust fault, Prince of Wales Island: Alaska Division of Geological and Geophysical Surveys Geologic Report 73, p. 17-18.
- Riding, R., and Soja, C.M., 1993, Silurian calcareous algae, cyanobacteria, and microproblematica from the Alexander terrane, Alaska: *Journal of Paleontology, v. 67, no. 5, p. 710-728.*
- Rigby, J.K., Nitecki, M.H., Soja, C.M., and Blodgett, R.B., 1994, Silurian aphrosalpingid sphinctozoans from Alaska and Russia: *Acta Palaeontologica Polonica, v. 39, no. 4, p. 341-391.*
- Robinson, G.D., and Twenhofel, W.S., 1953, Some lead-zinc and zinc-copper deposits of the Ketchikan and Wales districts, Alaska: U.S. Geological Survey Bulletin 998-C, p. 59-84.
- Roehm, J.C., 1946, Some high calcium limestone deposits in southeastern Alaska: Alaska Division of Geological and Geophysical Surveys Pamphlet no. 6, 85 p.

- Rosman, D.L., Henderson, J.R., and Walton, M.S., Jr., 1956, Reconnaissance total intensity aeromagnetic map of the southern part of Prince of Wales Island, Alaska: U.S. Geological Survey Geophysical Investigations Map GP-135, scale 1:126,720.
- Rubin, C.M., and Saleeby, J.B., 1987, The inner boundary of the Alexander terrane, southern SE Alaska-A newly discovered thrust belt (abs.): Geological Society of America Abstracts with Programs, v. 19, no. 6, p. 445.
- \_\_\_\_\_ 1991, Tectonic framework of the upper Paleozoic and lower Mesozoic Alava sequence: a revised view of the polygenetic Taku terrane in southern southeast Alaska: Canadian Journal of Earth Sciences, v. 28, p. 881-893.
- \_\_\_\_\_ 1992, Tectonic history of the eastern edge of the Alexander terrane, southeast Alaska: Tectonics, v. 11, no. 3, p. 586-602.
- Ruckmick, J.C., and Noble, J.A., 1959, Origin of the ultramafic complex at Union Bay, southeastern Sainsbury, C.L., 1961, Geology of part of the Craig C-2 quadrangle and adjoining areas, Prince of Wales Island, southeastern Alaska: U.S. Geological Survey Bulletin 1058-H, p. 299-362.
- Saleeby, J.B., 1991, Age and tectonic setting of the Duke Island ultramafic intrusion, southeast Alaska (abs.): EOS, American Geophysical Union Transactions, v. 72, no. 44, p. 521.
- Saleeby, J.B., 1992, Age and tectonic setting of the Duke Island ultramafic intrusion, southeast Alaska: Canadian Journal of Earth Sciences, v. 29, p. 506-522.
- Savage, N.M., 1977a, Lower Devonian conodonts from the Karheen Formation, southeastern Alaska: Canadian Journal of Earth Sciences, v. 15, p. 278-284.
- \_\_\_\_\_ 1977b, Middle Devonian (Eifelian) conodonts of the genus *Polygnathus* from the Wadleigh Limestone, southeastern Alaska: Canadian Journal of Earth Sciences, v. 15, p. 1343-1355.
- \_\_\_\_\_ 1981a, Lower Devonian conodonts from Kasaan Island, southeastern Alaska: Journal of Paleontology, v. 55, p. 848-852.
- \_\_\_\_\_ 1981b, A reassessment of the age of some Paleozoic brachiopods from southeastern Alaska: Journal of Paleontology, v. 55, p. 353-369.
- \_\_\_\_\_ 1982, Lower Devonian (Lochkovian) conodonts from Lulu Island, southeastern Alaska: Journal of Paleontology, v. 56, p. 983-988.
- \_\_\_\_\_ 1985, Silurian (Llandovery-Wenlock) conodonts from the base of the Heceta Limestone, southeastern Alaska: Canadian Journal of Earth Sciences, v. 22, p. 711-727.
- \_\_\_\_\_ 1987, New polygnathid conodonts from the Frasnian (Upper Devonian) of southeastern Alaska: Canadian Journal of Earth Sciences, v. 24, p. 2323-2328.
- \_\_\_\_\_ 1988, Devonian faunas and major depositional events in the southern Alaska terrane, southeastern Alaska: in McMillan, N.J., Embry, A.T., and Glass, D.J., eds., Devonian of the World, v. III: Paleontology, Paleoecology, and Biostratigraphy: Canadian Society of Petroleum Geologists, Memoir 14, p. 257-264.
- \_\_\_\_\_ 1989, The occurrence of the brachiopods *Nanukidium* and *Atrypoides* in the Late Silurian of southeastern Alaska, Alexander terrane: Journal of Paleontology, v. 63(4), p. 530-533.
- \_\_\_\_\_ 1992, Late Devonian (Frasnian and Fammenian) conodonts from the Wadleigh Limestone, southeastern Alaska: Journal of Paleontology, v. 66(2), p. 277-292.
- Savage, N.M., and Bradley, S.J., 1985, Early to Middle Pennsylvanian conodonts from the Klawak Formation and the Ladrones Limestone, southeastern Alaska: Journal of Paleontology, v. 59, p. 1451-1475.

- Savage, N.M., Churkin, M., Jr., and Eberlein, G.D., 1977, Lower Devonian conodonts from Port St. Nicholas, southeastern Alaska: *Canadian Journal of Earth Sciences*, v. 14, p. 2928-2936.
- Savage, N.M., Eberlein, G.D., and Churkin, M., Jr., 1978, Upper Devonian brachiopods from the Port Refugio Formation, Suemez Island, southeastern Alaska: *Journal of Paleontology*, v. 52, p. 370-393.
- Savage, N.M., and Funai, C., 1980, Devonian conodonts of probable early Frasnian age from the Coronados Islands of southeastern Alaska: *Journal of Paleontology*, v. 54, p. 806-813.
- Savage, N.M., and Gehrels, G.E., 1984, Early Devonian conodonts from Prince of Wales Island, southeastern Alaska: *Canadian Journal of Earth Sciences*, v. 21, p. 1415-1425.
- \_\_\_\_\_ 1987, Early Devonian and Late Triassic conodonts from Annette and Hotspur Islands, southeastern Alaska (abs.): *Geological Society of America, Abstracts with Programs*, v. 19, no. 6, p. 446.
- Smith, J.G., Stern, T.W., and Arth, J.G., 1979, Isotopic ages indicate multiple episodes of plutonism and metamorphism in the Coast Mountains near Ketchikan, Alaska (abs.): *Geological Society of America Abstracts with Programs*, v. 11, no. 7, p. 519.
- Soja, C.M., 1988a, Lower Devonian (Emsian) benthic communities from Kasaan Island, southeastern Alaska, in McMillan, N.J., Embry, A.T., and Glass, D.J., eds., *Devonian of the World, v. III: Paleontology, Paleoecology, and Biostratigraphy*: Canadian Society of Petroleum Geologists, *Memoir* 14, p. 265-279.
- \_\_\_\_\_ 1988b, Lower Devonian (Emsian) brachiopods from southeastern Alaska, U.S.A.: *Paleontographica, Series A*, v. 201, p. 129-193.
- \_\_\_\_\_ 1988c, Early Devonian benthic communities of the Alexander terrane, southeastern Alaska: *Lethia*, v. 21, p. 319-338.
- \_\_\_\_\_ 1988d, Lower Devonian platform carbonates from Kasaan Island, southeastern Alaska, Alexander terrane: *Canadian Journal of Earth Sciences*, v. 25, p. 639-656.
- \_\_\_\_\_ 1990, Island arc carbonates from the Silurian Heceta Formation of southeastern Alaska (Alexander terrane): *Journal of Sedimentary Petrology*, v. 60, p. 235-249.
- \_\_\_\_\_ 1991a, Origin of Silurian reefs in the Alexander terrane of southeastern Alaska: *Palaios*, v. 6, p. 111-125.
- \_\_\_\_\_ 1991b, Silurian trace fossils in carbonate turbidites from the Alexander arc of southeastern Alaska: *Ichnos*, v. 1, p. 173-181.
- \_\_\_\_\_ 1993a, Silurian microbial associations from the Alexander terrane, Alaska: *Journal of Paleontology*, v. 67, no. 5, p. 728-738.
- \_\_\_\_\_ 1993b, Carbonate platform evolution in a Silurian oceanic island: A case study from Alaska's Alexander terrane: *Journal of Sedimentary Petrology*, v. 63, no. 6, p. 1078-1088.
- \_\_\_\_\_ 1994, Significance of Silurian stromatolite-sphinctozoan reefs: *Geology*, v. 22, p. 355-358.
- Sonnevil, R.A., 1981, The Chilkat-Prince of Wales plutonic province, southeastern Alaska, in Albert, N.R.D., and Hudson, T., eds., *The United States Geological Survey in Alaska: Accomplishments during 1979*: U.S. Geological Survey Circular 823-B, p. B112-B115.
- Sullivan, A.B., 1991, Mid-crustal deformation and intrusion in southeastern Alaska and southeastern Pennsylvania: Bryn Mawr College, Bryn Mawr, PA, A.B. Thesis (Geology), 92 p.

- Taylor, H.P., Jr., 1967, The zoned ultramafic complexes of southeastern Alaska, *in* Wylie, P.J., ed., *Ultramafic and related rocks*, John Wiley and Sons, New York, p. 97-121.
- Tchudinova, I.I., Churkin, M., Jr., and Eberlein, G.D., 1974, Devonian syringoporoid corals from southeastern Alaska: *Journal of Paleontology*, v. 48, p. 124-134.
- Thompson, T.B., Pierson, J.R., and Lyttle, T., 1982, Geology and petrogenesis of the Bokan Mountain Granite complex, southeastern Alaska: *Geological Society of America Bulletin*, v. 93, p. 898-908.
- Turner, D.L., Herreid, G., and Bundtzen, T.K., 1977, Geochronology of southern Prince of Wales Island: Alaska Division of Geological and Geophysical Surveys Geologic Report 55, p. 11-16.
- U.S. Forest Service, 1987, Prince of Wales Island road guide, Tongass National Forest: U.S. Department of Agriculture, Forest Service, Map R10-RG-10, 1 sheet, scale 1:250,000.
- \_\_\_\_\_ 1990, Draft environmental impact statement for Tongass Forest Plan revision: U.S. Forest Service, Alaska Region, Juneau, Alaska, June 25, 1990, 6 pl. scale 1:500,000, many p.
- \_\_\_\_\_ 1991, Tongass Land Management Plan revision--Supplement to the Draft Environmental Impact Statement: U.S. Forest Service, Alaska Region, Juneau, Alaska, August, 1991, 5 v., 9 pl., scale 1:500,000, many p.
- U.S. Geological Survey, 1977, Aeromagnetic map of Ketchikan, Prince Rupert, and northeastern Craig quadrangles, Alaska: U.S. Geological Survey Open-File Report 77-359, 1 sheet, scale 1:250,000.
- \_\_\_\_\_ 1979, Aeromagnetic maps of parts of the Ketchikan, Prince Rupert, and Craig quadrangles, southeastern Alaska: U.S. Geological Survey Open-File Report 79-937, 19 sheets, scale 1:63,360.
- \_\_\_\_\_ 1984, Aeromagnetic map of the Craig area, Alaska: U.S. Geological Survey Open-File Report 84-666, 3 sheets, scale 1:250,000.
- \_\_\_\_\_ 1989, Mineral Resources Data System (MRDS), Branch of Resource Analysis, Mail Stop 920, Reston, VA 22092.
- Wahrhaftig, C., 1965, Physiographic divisions of Alaska: U.S. Geological Survey Professional Paper 482, 52 p.
- Warner, J.D., and Barker, J.C., 1989, Columbium- and rare earth-element-bearing deposits at Bokan Mountain, southeast Alaska: U.S. Bureau of Mines Open-File Report 33-89, 196 p.
- Warner, L.A., Goddard, E.N., and others, 1961, Iron and copper deposits of Kasaan Peninsula, Prince of Wales Island, southeastern Alaska: U.S. Geological Survey Bulletin 1090, 136 p.
- Wright, C.W., 1915, Geology and ore deposits of Copper Mountain and Kasaan Peninsula, Alaska: U.S. Geological Survey Professional Paper 87, 110 p.
- Wright, F.E., and Wright, C.W., 1908, The Ketchikan and Wrangell mining districts, Alaska: U.S. Geological Survey Bulletin 347, 210 p.
- Zen, E., and Hammarstrom, J.M., 1984, Magmatic epidote and its petrologic significance: *Geology*, v. 12, p. 515-518.

## INDEX TO MAP UNITS, BY NAME

[An index to map units, by symbol, is part of the "Brief Description of Map Units" section on the map sheet]

<u>UNIT NAME</u>	<u>SYMBOL</u>	<u>PAGE(S)</u>
Altered diorite and quartz diorite (SEDIMENTARY, VOLCANIC, AND RELATED ROCKS OF GRAVINA BELT)	Jgd	2 6
ANDESITIC BRECCIA OF LUCK CREEK	SObl	2 0
Andesitic flows and breccia (KARHEEN FORMATION AND ASSOCIATED ROCKS)	Dka	2 8
Andesitic to basaltic volcanic and volcanoclastic rocks and minor sedimentary rocks (SEDIMENTARY, VOLCANIC, AND RELATED ROCKS OF GRAVINA BELT)	Jgv	2 6
Andesitic to dacitic volcanic rocks (DESCON FORMATION)	SOda	2 1
Basaltic to andesitic volcanic rocks (DESCON FORMATION)	SOdb	21, 30
Basaltic to rhyolitic breccia and tuff (EXTRUSIVE VOLCANIC ROCKS)	QTVb	1 1
Breccia of northeastern Noyes Island (KARHEEN FORMATION, ETC.)	Dkbn	1 5
Chapin Peak Formation (HYD GROUP)	Fhc	26
Conglomerate, agglomerate, and volcanic breccia (BAY OF PILLARS FORMATION)	SpC	1 8
Conglomerate-facies rocks, (KARHEEN FORMATION, ETC.)	Dkcg	1 6
CORONADOS VOLCANICS	Dc	1 4
Dacitic to rhyolitic volcanic rocks (DESCON FORMATION)	SOdr	20, 30
Diorite, quartz diorite, gabbro, hornblendite, leucogabbro, trondhjemite, pyroxenite, and migmatite (INTRUSIVE AND RELATED ROCKS OF YOUNGER PART OF CAPE CHACON PLUTONIC PROVINCE)	SOdi	19, 29
Dunite, pyroxenite, and peridotite at Union Bay (INTRUSIVE ROCKS OF KLUKWAN-DUKE PLUTONIC PROVINCE)	Kkdu	2 5

EXTRUSIVE VOLCANIC ROCKS (Northeast of Clarence Strait)	QTV	23
Felsic volcanic rocks (HYD GROUP)	Fhv	27
Foliated and layered biotite-hornblende diorite and quartz diorite (INTRUSIVE ROCKS OF SOUTHERN DALL ISLAND PLUTONIC PROVINCE)	Cdi	22
Foliated and layered hornblende diorite and quartz diorite and heterogeneous diorite and gabbro (INTRUSIVE AND RELATED ROCKS OF OLDER PART OF CAPE CHACON PLUTONIC PROVINCE)	Odi	21
Foliated hornblende granodiorite, chlorite leucogranodiorite, and quartz-porphyritic granodiorite (INTRUSIVE AND RELATED ROCKS OF OLDER PART OF CAPE CHACON PLUTONIC PROVINCE)	Ogd	21
Foliated, lineated, and gneissic biotite-hornblende quartz diorite, granodiorite, and minor diorite (INTRUSIVE ROCKS OF SOUTHERN DALL ISLAND PLUTONIC PROVINCE)	Cqd	22
Foliated to massive hornblende-biotite tonalite and granodiorite, quartz monzonite, and quartz diorite (INTRUSIVE ROCKS OF ADMIRALTY-REVILLAGIGEDO PLUTONIC PROVINCE)	Krtf	25
Gabbro (INTRUSIVE ROCKS OF CHILKAT-PRINCE OF WALES PLUTONIC PROVINCE)	Kwgb	12
Gabbro and diorite at Union Bay (INTRUSIVE ROCKS OF KLUKWAN-DUKE PLUTONIC PROVINCE)	Kkgu	26
Granite (INTRUSIVE AND RELATED ROCKS OF YOUNGER PART OF THE CAPE CHACON PLUTONIC PROVINCE)	SOgr	19
Granitic Rocks of Southern Etolin Island (INTRUSIVE GRANITIC AND RELATED ROCKS OF KUIU-ETOLIN VOLCANIC-PLUTONIC PROVINCE)	Tge	23
Gravina Island Formation (SEDIMENTARY, VOLCANIC, AND RELATED ROCKS OF GRAVINA BELT)	KJgg	26
Greenstone, greenschist, black phyllite, quartz-sericite schist, metakeratophyre, and minor marble (WALES GROUP)	CZwg	22

Hornblende gabbro, microgabbro, and quartz diorite (INTRUSIVE ROCKS OF SOUTHERN DALL ISLAND PLUTONIC PROVINCE)	Ɔgb	22
Hornblendite (INTRUSIVE AND RELATED ROCKS OF YOUNGER PART OF CAPE CHACON PLUTONIC PROVINCE)	SOhb	20
Hornblende granodiorite and diorite and biotite- hornblende monzodiorite (INTRUSIVE ROCKS OF CHILKAT-PRINCE OF WALES PLUTONIC PROVINCE)	Kwgd	12
Hornblende quartz diorite, diorite, and quartz monzonite (INTRUSIVE AND RELATED ROCKS OF OLDER PART OF CAPE CHACON PLUTONIC PROVINCE)	Oqd	21
Hornblende quartz monzodiorite with minor tonalite, granodiorite, quartz diorite, diorite, quartz monzonite, and monzodiorite (INTRUSIVE ROCKS OF CHILKAT-PRINCE OF WALES PLUTONIC PROVINCE)	Kwqo	12
Hornfelsed sedimentary rocks (INTRUSIVE AND RELATED ROCKS OF KUIU- ETOLIN VOLCANIC-PLUTONIC PROVINCE)	Tsh	23
Hornfelsed volcanic rocks (INTRUSIVE GRANITIC AND RELATED ROCKS OF KUIU-ETOLIN VOLCANIC-PLUTONIC PROVINCE)	Tvh	23
KLAWAK FORMATION	ƆPk	13
KOOTZNAHOO FORMATION	Tk	11, 24
LADRONES LIMESTONE	ƆPl	13
LEUCODIORITE AT KASSA INLET	Sld	17
Leucogabbro near Ketchikan (INTRUSIVE AND RELATED ROCKS OF KUIU-ETOLIN VOLCANIC-PLUTONIC PROVINCE)	Tlgb	23
LEUCOSYENITE OF KLAWOCK AND SUKKWAN	ƆPsy	13
Limestone and dolomite (HYD GROUP)	Tɦl	27
Limestone, minor limestone breccia, sandstone, mudstone, and polymictic conglomerate (HECETA LIMESTONE)	Shl	17
Limestone of Kasaan Island (KARHEEN FORMATION, ETC.)	Dkl	14
Marble and minor calcisilicate rocks in Wales Group (WALES GROUP)	ƆZwm	23

Metacarbonate rocks (METAMORPHOSED ROCKS ON CLEVELAND PENINSULA)	SOmc	29
Metagabbro in Wales Group south of Kassa Inlet (WALES GROUP)	ØZwgb	22
Metaintrusive rocks (METAMORPHOSED SEDIMENTARY, VOLCANIC, AND INTRUSIVE ROCKS)	MzPzmi	28
Metamorphic rocks at Kendrick Bay (INTRUSIVE AND RELATED ROCKS OF THE YOUNGER PART OF THE CAPE CHACON PLUTONIC PROVINCE)	SOmk	18
METAMORPHIC ROCKS AT KLAKAS INLET	DSmk	16
Metamorphic rocks at Ruth Bay (INTRUSIVE AND RELATED ROCKS OF OLDER PART OF CAPE CHACON PLUTONIC PROVINCE)	Omr	21
METAMORPHIC ROCKS ON EASTERN PRINCE OF WALES ISLAND	Ømpw	22
METAMORPHIC ROCKS ON SOUTHERN GRAVINA ISLAND	Ømgi	30
Metapolyimictic conglomerate (METAMORPHOSED SEDIMENTARY, VOLCANIC, AND INTRUSIVE ROCKS)	MzPzmc	28
Metasedimentary rocks (METAMORPHOSED SEDIMENTARY, VOLCANIC, AND INTRUSIVE ROCKS)	MzPzms	27
Metasedimentary rocks (METAMORPHOSED ROCKS ON CLEVELAND PENINSULA)	SOms	29
Metavolcanic rocks (METAMORPHOSED SEDIMENTARY, VOLCANIC, AND INTRUSIVE ROCKS)	MzPzmv	28
Metavolcanic rocks (METAMORPHOSED ROCKS ON CLEVELAND PENINSULA)	SOmv	29
Migmatite (INTRUSIVE AND RELATED ROCKS OF ADMIRALTY-REVILLAGIGEDO PLUTONIC PROVINCE)	Krmg	25
Migmatitic Granitic Rocks of Southern Etolin Island (INTRUSIVE AND RELATED ROCKS OF KUIU-ETOLIN VOLCANIC-PLUTONIC PROVINCE)	Tme	23
Mudstone-and-siltstone-facies rocks (KARHEEN FORMATION, ETC.)	Dkkm	29

Nonfoliated plagioclase-porphyritic (hornblende-)(epidote-) garnet-biotite tonalite, quartz diorite, and minor granodiorite (INTRUSIVE ROCKS OF ADMIRALTY-REVILLAGIGEDO PLUTONIC PROVINCE)	Krtn	24
Oligomictic breccia (HYD(?) GROUP)	Fhb	13
Olivine basalt and andesite (EXTRUSIVE VOLCANIC ROCKS)	QTvo	11
Peralkaline granite of Bokan Mountain (INTRUSIVE ROCKS OF BOKAN MOUNTAIN PLUTONIC PROVINCE)	Jbgr	12
PERATROVICH FORMATION	Mp	13
Plagioclase-porphyritic dacite (?), andesite, and diorite(?) (KARHEEN FORMATION, ETC.)	Dkp	16
POLYMICTIC CONGLOMERATE BETWEEN HECETA LIMESTONE AND BAY OF PILLARS FORMATION	Shpc	18
Polymictic conglomerate in Heceta Limestone (HECETA LIMESTONE)	Shc	17
Porphyritic foliated biotite tonalite, quartz diorite, and granodiorite (INTRUSIVE ROCKS OF ADMIRALTY- REVILLAGIGEDO PLUTONIC PROVINCE)	Krtp	25
PORT REFUGIO FORMATION	Dpr	14
PYBUS(?) FORMATION	Pp	28
Pyroxenite, hornblendite, and gabbro (INTRUSIVE AND RELATED ROCKS OF YOUNGER PART OF CAPE CHACON PLUTONIC PROVINCE)	SOpx	19
Quartz diorite (INTRUSIVE ROCKS OF ADMIRALTY- REVILLAGIGEDO PLUTONIC BELT)	Krqd	25
Quartz syenite and granite (INTRUSIVE AND RELATED ROCKS OF YOUNGER PART OF CAPE CHACON PLUTONIC PROVINCE)	SOsy	19
Redbed-facies rocks (KARHEEN FORMATION, ETC.)	Dkkr	15
Rhyolite and dacite (EXTRUSIVE VOLCANIC ROCKS)	QTvr	11
Rhyolite of Kasaan Island (KARHEEN FORMATION, ETC.)	Dkr	14
ST. JOSEPH ISLAND VOLCANICS	Dsj	13

Sedimentary and mafic volcanic rocks (HYD GROUP)	Thsv	27
Sedimentary and minor volcanic rocks (SEDIMENTARY, VOLCANIC, AND RELATED ROCKS OF GRAVINA BELT)	Jgs	26
Sedimentary and volcanic rocks (BAY OF PILLARS FORMATION)	Spsv	18
Sedimentary rocks of the Port Nicholas area (KARHEEN FORMATION, ETC.)	Dksn	16
SEDIMENTARY ROCKS OF STANEY CREEK AND TUXEKAN PASSAGE AREA	DSs	17
Siltstone, sandstone, polymictic conglomerate, and limestone of Clover Bay (KARHEEN FORMATION, ETC.)	Dkcb	16
SURFICIAL DEPOSITS	Qs	10, 23
Syenite of Dora Bay (INTRUSIVE ROCKS OF BOKAN MOUNTAIN PLUTONIC PROVINCE)	Jbsy	12
TRONDHJEMITE, DIORITE, AND GRANITE OF ANNETTE ISLAND	Stj	29
Turbidite-facies rocks (KARHEEN FORMATION, ETC.)	Dkkt	15
Ultramafic rocks, mainly dunite, peridotite, clinopyroxenite, and hornblendite (INTRUSIVE ROCKS OF KLUKWAN-DUKE PLUTONIC PROVINCE)	Kku	25
Unfoliated and foliated quartz monzonite and granite (INTRUSIVE AND RELATED ROCKS OF YOUNGER PART OF CAPE CHACON PLUTONIC PROVINCE)	SOqm	19
UNITS EXPOSED NORTHEAST OF CLARENCE STRAIT		23
UNITS EXPOSED SOUTHWEST OF CLARENCE STRAIT		10
Volcanic rocks of Port Refugio Formation (PORT REFUGIO FORMATION)	Dprv	14
Volcaniclastic conglomerate (DESCON FORMATION)	SOdc	20
Volcaniclastic graywacke and mudstone turbidites and minor limestone (BAY OF PILLARS FORMATION)	Spg	18

Volcaniclastic graywacke, siltstone, and mudstone turbidites, minor sedimentary breccia, limestone, and polymictic conglomerate (DESCON FORMATION)	SOdg	20
WADLEIGH LIMESTONE	Dw	14





Figure 1. Index map showing location of study area and physiographic provinces and sections (from Wahrhaftig, 1965).



**EXPLANATION**

- |   |   |
|---|---|
| 1. George Moerlein (written commun. to G.D. Eberlein, 1981) | 8. Horton and others (1992) and Karl (1992)                   |
| 2. Clark and others (1971)                                  | 9. MacKevett (1963)   |
| 3. Eberlein and others (1983)                               | 10. Gehrels and Saleeby (1986) and Gehrels (1992)             |
| 4. Berg and others (1988)                                   | 11. Gehrels (1991)  |
| 5. Redman (1982)  | 12. Berg and others (1976)                                    |
| 6. Gehrels and others (1987)                                | 13. J.F. Baichtal (written comm., 1995, selected units shown) |
| 7. Sullivan (1991)  |   |

Figure 2. Sources of geologic information, 1983-1995.




Figure 3. Lithotectonic terrane and major fault map of southeastern Alaska and adjacent regions. Dashed line is boundary of the Craig, Dixon Entrance, and parts of the Ketchikan and Prince Rupert quadrangles map area (after Brew and others, 1991a, b, 1992a). Major faults are labeled as follows: BR, Border Ranges; CHS, Chatham Strait; CLS, Clarence Strait; CRML, Coast Range megalineament; FQ, Fairweather-Queen Charlotte; NA, Nahlin; NS, Neva Strait; THL, Tally Ho-Llewellyn; and TR, Transitional.

