



MAP A. DISTRIBUTION OF GOLD IN STREAM-SEDIMENT SAMPLES



MAP B. DISTRIBUTION OF ANTIMONY IN STREAM-SEDIMENT SAMPLES



MAP C. DISTRIBUTION OF ARSENIC IN STREAM-SEDIMENT SAMPLES

EXPLANATION

- Nonanomalous value
- Anomalous value




Figure 1. Histogram showing gold concentrations in 161 stream-sediment samples collected from the Delta quadrangle, Utah. N, not detected at 0.1 ppm; L, detected, but less than 0.1 ppm.

STUDIES RELATED TO CUSMAP

This map is part of a folio of maps of the Delta 1° x 2° quadrangle, Utah, prepared under the Comprehensive United States Mineral Assessment Program (CUSMAP). Other publications in this folio include an aeromagnetic-anomaly map and related geophysical maps (Kocks, 1991), and a complete Bouguer-gravity-anomaly map and related geophysical maps (Bankey and Cook, 1989).

INTRODUCTION

The Delta quadrangle is located in the eastern part of the Basin and Range physiographic province, near the boundary of the Basin and Range province with the Rocky Mountain and Colorado Plateau provinces. Much of the Timne-Deery Creek igneous and mineral belt is within the northern part of the quadrangle.

GEOLOGIC SETTING

Bedrock in the quadrangle consists predominantly of a sequence of late Precambrian and Paleozoic strata that was thrust eastward during the Cretaceous Sevier orogeny. These rocks were thrust onto an attachment of Mesozoic sedimentary rocks in the southeastern part of the quadrangle and onto Precambrian and Paleozoic rocks in other parts of the quadrangle. Much of the northern part of the quadrangle is composed of Oligocene and younger volcanic rocks and related intrusions; a single Jurassic-age stock is present in the southwestern part of the quadrangle. Late Cretaceous extensional tectonics produced non-trending fault blocks; the highlands of these blocks were deeply eroded and the resulting debris was deposited in adjacent basins. Most of the mineral deposits in the Timne-Deery Creek igneous and mineral belt were formed during the middle and late Cretaceous.

GEOCHEMICAL STUDIES

Data obtained from this regional geochemical study of stream-sediment samples can be used to define broad geochemical patterns and trends that can be combined with geologic and geophysical data to assess the mineral resource potential of this quadrangle. These maps show the regional distributions of gold (map A), antimony (map B), arsenic (map C), bismuth (map D), cadmium (map E), and zinc (map F).

Collection of Samples

A total of 161 stream-sediment samples were used to generate the data presented in this report. These samples were collected by the U.S. Geological Survey (USGS) during fieldwork in wilderness study areas. Reports covering other elements analyzed in stream-sediment samples are given in Zimbelman (1993a) and a summary of anomalous abundances in heavy-mineral-concentrate samples is presented in Zimbelman (1993b); both of these reports include data from samples collected during the National Uranium Resource Evaluation (NURE) program. However, the NURE samples did not have sufficient material to analyze for gold, antimony, arsenic, bismuth, cadmium, or zinc using the methods described in this report, although they were analyzed for these elements by emission spectrographic methods using lower limits of determination.

The samples consisted of active alluvium collected primarily from first-order (unbranched) and second-order (below the junction of two first-order streams) streams as shown on USGS topographic maps (scale = 1:24,000). Each sample was collected from several localities within an area that may extend as much as 17 m from the site plotted on the map.

ANALYTICAL PROCEDURES

The stream-sediment samples were analyzed by various chemical methods, including an atomic absorption spectroscopy method (Thompson and others, 1968) and an inductively coupled plasma-atomic emission spectrometric method (Cook and others, 1987).

These samples, along with 597 additional NURE samples, were also analyzed by a semiquantitative, direct-current arc emission spectrographic method. Results of analyses for barium, beryllium, copper, lead, molybdenum, silver, and tin are given in Zimbelman (1993a). In general, the samples collected during the NURE program did not have enough material to allow complete analysis by methods other than emission spectrography, and additional chemical analyses were not performed on these samples.

GEOCHEMICAL MAPS

Point-plot maps for the data were prepared using a computer program in the USGS's STATPAC system (VanTrump and Misch, 1977). The point-plot maps show the locations and magnitudes of geochemical anomalies. Maps showing the distributions of analytical results for barium, beryllium, copper, lead, molybdenum, silver, and tin in stream-sediment samples are given in Zimbelman (1993a). Maps showing the distributions of analytical results for barium, beryllium, cadmium, copper, lead, molybdenum, silver, tin, tungsten, and zinc in heavy-mineral-concentrate samples are presented in Zimbelman (1993b).

GEOCHEMICAL IMPLICATIONS OF THE DATA

Reconnaissance geochemical surveys are valuable tools in mineral exploration, but they should be used in conjunction with data from other earth science disciplines. In particular, outlining exploration targets generally involves considerable additional, more detailed investigations.

REFERENCES CITED

- Bankey, Viki, and Cook, K.L., 1989, Complete Bouguer gravity map and related geophysical maps of the Delta 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-2081-A, scale 1:250,000 and 1:500,000.
- Cook, J.C., Bankey, V.K., Jackson, L.L., and Lauer, F.E., 1987, Analytical for the analysis of stream sediments and rocks from Wilderness Study Areas: U.S. Geological Survey Open-File Report 87-84, 35 p.
- Kocks, R.P., 1991, Aeromagnetic anomaly map and related geophysical maps of the Delta 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-2081-B, scale 1:250,000.
- Thompson, C.E., Nakagawa, H.M., and Van Sickle, G.H., 1968, Rapid analysis for 16 elements in U.S. Geological Survey Professional Paper 609-B, p. 19-20.
- VanTrump, George, and Misch, A.T., 1977, The U.S. Geological Survey KASS-STATPAC system for management and statistical reduction of geochemical data: Computers and Geosciences, v. 3, p. 475-488.
- Zimbelman, D.R., 1993a, Maps showing distribution of barium, beryllium, copper, lead, molybdenum, silver, and tin in stream-sediment samples, Delta 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-2081-C, scale 1:250,000.
- Zimbelman, D.R., 1993b, Maps showing distribution of antimony, arsenic, barium, beryllium, bismuth, cadmium, copper, lead, molybdenum, silver, tin, tungsten, and zinc in heavy-mineral-concentrate samples, Delta 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-2081-E, scale 1:250,000.

EXPLANATION

- Nonanomalous value
- Anomalous value




Figure 1. Histogram showing antimony concentrations in 161 stream-sediment samples collected from the Delta quadrangle, Utah. N, not detected at 2 ppm; L, detected, but less than 2 ppm.

- DESCRIPTION OF MAP UNITS**
- Qv Quaternary volcanic deposits (Pleistocene and Holocene): Fluvial and pyroclastic deposits of basalt and basaltic andesite.
- Qts Alluvium and valley fill deposits (Holocene, Pleistocene, and Pliocene): Colluvium and alluvium in upland areas and floodplains; lacustrine units, well-sorted, and well-sorted sand in valley and floodplains.
- Ttp Tertiary volcanic deposits (Pliocene through Pliocene): Fluvial, igneous, pyroclastic, and sedimentary units; basaltic to alkali rhyolite; generally limited to well-defined negative centers in northern part of quadrangle.
- Ts Tertiary sedimentary deposits (Pliocene through Pliocene): Fluvial, igneous, pyroclastic, and sedimentary units; basaltic to alkali rhyolite; generally limited to well-defined negative centers in northern part of quadrangle.
- Tk North Basin Formation (Tertiary and Quaternary): Red and green post-tectonic shales, sandstones, conglomerates, and limestones. Considerable mineral resources in upper part and dominant resource in the lower part.
- Ku Cretaceous strata, undivided (but separate conglomerates, sandstones, and shales): Characterized by large, semi-circular clasts of quartzite and various rocks. Chiefly consists of Canyon Bench, Fish River, and Indian (Conglomerates); locally includes some Paleozoic strata.
- Jl Jurassic plutons: Felsic to white, coarse crystalline, quartz monzonitic porphyry stock cutting Canyon Bench in House Range. Age is estimated to be 165 Ma.
- Jbu Jurassic and Triassic strata, undivided: Includes the House Range in western part of quadrangle and Mesquite, Chinle, and Navajo Formations in eastern part of quadrangle.
- Pu Paleozoic strata, undivided (Pliocene through Cambrian): Includes strata deposited during each period of the Paleozoic Era. These strata consist of about 60 percent carbonate rock, 30 percent quartzite and shale, and 10 percent igneous and are subdivided into as many as 20 or 30 formational units in any one area.
- Zy Precambrian 2 and 3 strata, undivided: Copper and molybdenum-bearing, igneous and metamorphic rocks; some local beds of algal limestone (all subdivided into 5 or more formational units or sequences).

EXPLANATION

- Nonanomalous value
- Moderately anomalous value
- Highly anomalous value




Figure 1. Histogram showing arsenic concentrations in 161 stream-sediment samples collected from the Delta quadrangle, Utah. L, detected, but less than 5 ppm.

- Contact—dashed where approximately located
- High-angle fault—bar and ball on downthrown side; arrow indicates direction of relative horizontal displacement
- Thrust fault—contact on upper plate; dotted where concealed; groups of three dots where inferred and concealed; short concealed positions approximately located
- Anticline—showing trace of axial plane; dotted where concealed
- Syncline—showing trace of axial plane; dotted where concealed

MAPS SHOWING DISTRIBUTION OF GOLD, ANTIMONY, ARSENIC, BISMUTH, CADMIUM, AND ZINC IN STREAM-SEDIMENT SAMPLES, DELTA 1° x 2° QUADRANGLE, UTAH

By
David R. Zimbelman
1993