

INTRODUCTION

This map of the Richfield 1° x 2° quadrangle shows the regional distribution of thorium in the less-than-0.180-mm (minus-80-mesh) fraction of stream-sediment samples. It is part of a folio of maps of the Richfield 1° x 2° quadrangle, Utah, prepared under the Continuous United States Mineral Assessment Program. Other published geochemical maps in this folio are listed in the references (this publication).

The Richfield quadrangle is located in west-central Utah and includes the eastern part of the Pioche-Marysvale igneous and mineral belt, which extends from the vicinity of Pioche in southeastern Nevada, east-northeastward for 155 miles into central Utah. The western two-thirds of the Richfield quadrangle is part of the Basin and Range province, whereas the eastern third is part of the High Plateaus of Utah, a subprovince of the Colorado Plateau.

Bedrock in the northern part of the Richfield quadrangle consists predominantly of Late Proterozoic and Paleozoic sedimentary strata that were thrust eastward during the Sevier orogeny in Cretaceous time onto an autochthon of Mesozoic sedimentary rocks located in the eastern part of the quadrangle. The southern part of the quadrangle is largely underlain by Tertiary and younger volcanic rocks and related intrusions. Extensive tectonism in late Cenozoic time broke the bedrock terrain into a series of north-trending fault blocks; the uplifted mountain areas were eroded to various degrees and the resulting debris was deposited in adjacent basins. Most mineral deposits in the Pioche-Marysvale mineral belt were formed as a result of igneous activity in middle and late Cenozoic time. A more complete description of the geology and a mineral-resource appraisal of the Richfield quadrangle appears in Steven and Morris (1984 and 1987).

The regional sampling program was designed to define broad geochemical patterns and trends that can be utilized along with geological and geophysical data to assess the mineral-resource potential for this quadrangle. Reconnaissance geochemical surveys are valuable tools in mineral exploration, especially when used in conjunction with data obtained from other earth science disciplines. Identifying specific exploration targets, however, generally involves additional, more detailed investigations.

SAMPLE COLLECTION AND PREPARATION

Stream-sediment samples were collected at 1,462 sites throughout the Richfield quadrangle. The sample sites are located along small, normally unbranched or first-order stream drainages that range from 1 to 2 miles in length and whose stream courses are 2 to 12 feet wide. Sample density within the bedrock areas was one sample per 3 square miles. Intermountain basins containing sediments were not sampled. Each sample is a composite of material collected at four or five sites (usually within 30 feet of each other) across and along the active channel. About 1 to 2 pounds of bulk sediments were collected at each site. The geochemical sampling was conducted by G.K. Lee, W.A. Miller, J.B. McHugh, R.E. Tucker, J.D. Tucker, and J.F. Gussadpoull.

The less-than-0.180-mm fraction of stream sediments was prepared by drying the bulk sediment and sieving it to less than 0.180 mm. This fraction was then pulverized in a vertical ceramic-plate mill to a powder (less than 0.105 mm) and analyzed.

ANALYTICAL PROCEDURES

For this study, thorium was determined by delayed neutron counting, which is a method of nuclear-activation analysis used to measure thorium in a matrix without chemical processing (conducted under the direction of H.T. Millard, Jr.). A complete listing of analytical results appears in McHugh and others (1989). Ten-gram sample aliquots were used, and the results exhibited a detection limit of about 1 ppm. Analytical precision of (+ or -) 10 percent may be achieved for the determination of thorium at concentrations greater than about 10 ppm. A more detailed description of this analytical procedure appears in McKown and Millard (1987).

GENERATION OF MAPS

A computer-generated point-plot map for thorium in the less-than-0.180-mm fraction of stream sediments was prepared using the computerized mapping program within the U.S. Geological Survey's STATMAP system (VanTrump and Miesch, 1977). Thorium concentrations ranged from less than 2.4 to 317 ppm. Approximately 50 percent of the samples were within the range of background values; the remaining 50 percent are divided into six classifications that range from highly anomalous to very weakly anomalous. Each classification is represented by a symbol or size of symbol on the histogram for thorium (fig. 1). The least anomalous classification only indicates elevated values above background and does not indicate mineralization. The most anomalous classification represents one percent of the total population followed by less anomalous classifications at approximately 2.5, 5, 10, 25, and 50 percent of the total population. Thus, each succeeding classification contains approximately twice the population of the preceding less anomalous classification.

REFERENCES

McHugh, J.B., Miller, W.R., Vaughn, R.B., McKown, D.M., Tucker, R.E., and Bronfield, C.S., 1989, Analytical results for uranium and thorium in 1,462 stream-sediment samples, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Open-File Report 89-366, 18 p.

McKown, D.M., and Millard, Jr., H.T., 1987, Determination of uranium and thorium by delayed neutron counting, in Badocker, F.A., ed., Methods for geochemical analysis: U.S. Geological Survey Bulletin 1770, p. 11-112.

Miller, W.R., and McHugh, J.B., 1990, Map showing the distribution of uranium in stream-sediment samples, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-2138-J, scale 1:250,000.

Miller, W.R., Motooka, J.M., and McHugh, J.B., 1980, Distribution of cobalt in heavy-mineral concentrates, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-1246-A, scale 1:500,000.

1985, Maps showing distribution of arsenic in heavy-mineral concentrates, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-1246-B, scale 1:500,000.

1985, Maps showing distribution of barium in heavy-mineral concentrates, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-1246-C, scale 1:500,000.

1985, Maps showing distribution of beryllium in heavy-mineral concentrates and stream sediments, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-1246-D, scale 1:500,000.

1985, Maps showing distribution of bismuth in heavy-mineral concentrates, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-1246-E, scale 1:500,000.

1985, Maps showing distribution of copper in heavy-mineral concentrates, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-1246-F, scale 1:500,000.

1985, Maps showing distribution of lead in heavy-mineral concentrates, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-1246-G, scale 1:500,000.

1985, Maps showing distribution of thorium in heavy-mineral concentrates, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-1246-H, scale 1:500,000.

1985, Maps showing distribution of tin in heavy-mineral concentrates, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-1246-I, scale 1:500,000.

1985, Maps showing distribution of tungsten in heavy-mineral concentrates, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-1246-J, scale 1:500,000.

1985, Maps showing distribution of zinc in heavy-mineral concentrates, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-1246-K, scale 1:500,000.

1990, Map showing the distributions of cadmium and antimony in the nonmagnetic fraction of heavy-mineral concentrates, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-2137-A, scale 1:250,000.

1990, Map showing the distribution of silver in the nonmagnetic fraction of heavy-mineral concentrates, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-2137-B, scale 1:250,000.

1990, Map showing the distribution of barium in stream-sediment samples, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-2138-A, scale 1:250,000.

1990, Map showing the distributions of bismuth and cadmium in stream-sediment samples, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-2138-B, scale 1:250,000.

1990, Map showing the distribution of copper in stream-sediment samples, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-2138-C, scale 1:250,000.

1990, Map showing the distribution of molybdenum in stream-sediment samples, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-2138-D, scale 1:250,000.

1990, Map showing the distribution of lead in stream-sediment samples, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-2138-E, scale 1:250,000.

1990, Map showing the distribution of silver in stream-sediment samples, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-2138-F, scale 1:250,000.


1990, Map showing the distribution of tin in stream-sediment samples, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-2138-G, scale 1:250,000.

1990, Map showing the distribution of zinc in stream-sediment samples, Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-2138-H, scale 1:250,000.

Steven, T.A., and Morris, H.T., 1984, Mineral resource potential of the west-central Utah: U.S. Geological Survey Circular 916, 24 p.


Steven, T.A., Rowley, P.D., Hintze, L.F., Best, M.G., Nelson, M.G., and Cunningham, C.G., compilers, 1978, Preliminary geologic map of the Richfield 1° x 2° quadrangle, Utah: U.S. Geological Survey Open-File Report 83-602, 1 sheet, scale 1:250,000.

VanTrump, G., and Miesch, A.T., 1977, The U.S. Geological Survey RASS-STATMAP system for management and statistical reduction of geochemical data: Computers and Geosciences, v. 3, p. 475-488.



- EXPLANATION**
- SAMPLE LOCALITIES FOR THORIUM
 - Highly anomalous value
 - ◻ Moderately high anomalous value
 - ◻ Moderately anomalous value
 - ◻ Moderately weak anomalous value
 - ◻ Weakly anomalous value
 - ◻ Very weakly anomalous value
 - ◻ Nonanomalous value

- LIST OF MAP UNITS**
- QTa Surficial deposits, undivided (Quaternary and Tertiary)
 - QTV Volcanic rocks, undivided (Quaternary and Tertiary)
 - Ti Intrusive igneous rocks, undivided (Tertiary)
 - Tzs Sedimentary rocks, undivided (Tertiary to Late Proterozoic)
 - Contact



MAP SHOWING DISTRIBUTION OF THORIUM IN STREAM-SEDIMENT SAMPLES, RICHFIELD 1° X 2° QUADRANGLE, UTAH

By
William R. Miller, Jerry M. Motooka, and John B. McHugh
1990

Figure 1.—Histogram showing thorium concentrations in stream-sediment samples from the Richfield 1° x 2° quadrangle, Utah. Number of samples, 1,462; L, not detected at 2.4 ppm; G, greater than 317 ppm.

Any use of trade names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Geological Survey.

INTERIOR—GEOLOGICAL SURVEY, RESTON, VIRGINIA—1990

For sale by U.S. Geological Survey Map Distribution, Box 24286, Federal Center, Denver, CO 80225