

Table 1. Characteristics of young faults in the Lovelock 1° by 2° quadrangle  
[See figure 1 for area boundaries]

Orientation	Northwest		Central		East	
	Length (km)	Percent	Length (km)	Percent	Length (km)	Percent
90.0 to 80°	---	---	7.0	1.4	---	---
79.9 to 70°	---	---	2.0	0.4	---	---
69.9 to 60°	---	---	---	---	---	---
59.9 to 50°	5.4	1.3	7.4	1.5	21.6	16.9
49.9 to 40°	---	---	16.4	3.2	9.4	7.3
39.9 to 30°	6.6	1.6	42.6	8.4	---	---
29.9 to 20°	25.4	6.0	126.0	24.7	26.4	20.6
19.9 to 10°	42.0	9.9	119.0	23.3	11.8	9.2
09.9 to 0°	72.4	17.1	94.8	18.6	22.4	17.5
359.9 to 350°	103.2	24.4	29.0	5.7	9.6	7.5
349.9 to 340°	80.2	19.0	24.8	4.9	12.0	9.4
339.9 to 330°	23.8	5.6	22.8	4.5	10.8	8.4
329.9 to 320°	17.6	4.2	3.4	0.7	---	---
319.9 to 310°	2.0	0.5	---	---	---	---
309.9 to 300°	32.6	7.7	2.0	0.4	---	---
299.9 to 290°	3.6	0.9	---	---	4.0	3.1
289.9 to 280°	5.2	1.2	---	---	---	---
279.9 to 270°	2.2	0.5	12.6	2.5	---	---
Total (km)	422.2	100.0	509.8	100.0	128.0	100.0
Area (km <sup>2</sup> )	4880	---	10630	---	3440	---
Density (km/km <sup>2</sup> )	0.087	---	0.048	---	0.037	---
Number	82	---	107	---	23	---
Density (no./km <sup>2</sup> )	0.017	---	0.010	---	0.007	---
Mean length (km)	5.1	---	4.8	---	5.6	---

Table 2. General photogeologic and geomorphic criteria used to estimate general ages of piedmont surfaces

Map unit	Age	Depth of dissection	Drainage net morphology	Interflow morphology	Geomorphic relations	Typical geomorphic environments	General field criteria
Unit Q3	Holocene (0 to 10 ka)	Shallow to none; generally < 5 m	Predominantly radial from fan apex; channels typically poorly to moderately well defined	Typically poorly defined; bar and swale micro-topography common on most surfaces	Surfaces cut pluvial shorelines and (or) late Pleistocene glacial moraines	Distal piedmont surfaces; channels and terraces in proximal areas	Unweathered to slightly weathered; very weak to weak soil development
Unit Q2	Late Pleistocene (10 to 130 ka)	Shallow to moderate; typically 2-6 m	Predominantly distributary; however, some channels head on piedmont; well-defined channels	Typically well defined; surfaces broad and flat with abrupt margins	Surfaces overlain by pluvial shorelines and (or) latest Pleistocene glacial moraines	Proximal to distal piedmont surfaces; some inset terraces	Weakly to moderately well developed soils; interlocking stone pavements
Unit Q1	Early to middle Pleistocene (0.15 to 1.5 Ma)	Moderate to deep; commonly > 10 m	Predominantly subparallel; well-defined channels	Well defined; older interflow surface (ballenas) commonly narrow and irregular	Surfaces overlain by pluvial shorelines and (or) latest Pleistocene glacial moraines	Generally confined to intermediate and proximal piedmont areas	Moderately to very well developed soils; interlocking to highly degraded stone pavements




Figure 1. Rose diagrams summarizing orientation of young faulting. Number indicates mean fault trend. See table 1 for data.

## INTRODUCTION

The State of Nevada occupies most of the central and western parts of the Great Basin, the largest tectonically active region within the Basin and Range province of North America. The topography of this region is typified by generally north-northwest- to northeast-trending, subparallel mountain ranges separated by alluvial basins of similar plan form and orientation. This classic Basin and Range physiography is the product of at least two phases of middle- to late Cenozoic extensional faulting (Zoback and others, 1981; Eaton, 1982; Stewart, 1983), and most of the basins and ranges of the region are at least partly bounded by late Cenozoic faults. The earlier phase of extension was marked by widespread shallow detachment faulting that began at approximately 35 Ma and locally continued into the time of the later phase (Eaton, 1982; Stewart, 1983). The later phase, which was dominated by high-angle, more deeply penetrating block faulting, may have begun locally at about 17 Ma and continues episodically to the present time (Christiansen and McKee, 1978; Eaton, 1982). This map, one of a series of 1° by 2° quadrangle maps showing young faults in Nevada, provides a generalized picture of the late Tertiary and Quaternary faulting that is associated with the later extensional phase. These young faults are a primary determinant of the present configuration of ranges and basins within the quadrangle.

## MAPPING PROCEDURE

Young faults are herein defined as those faults that have undergone latest Tertiary and (or) Quaternary movement. These faults are commonly marked by a variety of diagnostic structural landforms and other surficial phenomena that can be readily identified and mapped on aerial photographs. These features include (1) scarps on latest Tertiary and (or) Quaternary surficial deposits, volcanic strata, or geomorphic surfaces (either erosional or depositional); (2) prominent alignments of linear drainageways, ridges and swales, active springs and (or) spring deposits, and linear discontinuities of structure, rock type, and vegetation; and (3) abrupt, steeply sloping range fronts with basal scarps, faceted spurs, winged valleys, and elongate drainage basins with narrow valley floors (Thornbury, 1969; Bull, 1977; Bull and McFadden, 1977; Wallace, 1977, 1978).

National High Altitude Program (NHAP), 1:58,000-nominal-scale, color-infrared photographs were used for photogeologic interpretation. This mapping was transferred directly to 12" by 1" topographic quadrangle maps that were enlarged to the scale of the photographs. These maps were reduced and compiled at 1:250,000 scale. This compilation was then compared with previous mapping of young faults within the quadrangle (Wallace, 1979), and significant differences between maps were resolved. Following comparison and resolution with this previous mapping, the final 1:250,000-scale compilation was digitized using a GTCO digitizing board connected to a Macintosh II minicomputer. The resulting vector file was converted to raster format (cell size = 200 m by 200 m) and analyzed to determine the approximate length and average orientation of each fault segment. These data are summarized in table 1 and figure 1.

General ages of surficial deposits and erosion surfaces cut by young faults were estimated using a variety of photogeologic and geomorphic criteria (table 2). These age estimates provide a general indication of the approximate timing of young faulting throughout the quadrangle. However, it should be emphasized that these data do not necessarily reflect the age of most recent surface rupture along any particular fault segment. Rather, they provide only very general (and commonly somewhat biased) age constraints on this surface faulting. Age estimates based on photogeologic analysis of surficial deposits and erosion surfaces are at best, both tentative and imprecise. Moreover, the distribution of these deposits and surfaces is inherently biased by geomorphic process and environment. For example, in those areas of the Great Basin where range uplift rates are low to moderate, remnants of older geomorphic surfaces tend to be concentrated in proximal piedmont areas, whereas younger surficial deposits tend to accumulate on distal piedmonts and basin flats. Consequently, young faults located in intrabasin areas are more likely to offset younger surface deposits than are faults located along range fronts or in proximal piedmont areas. Therefore, inferences based on these data regarding the temporal distribution of young fault activity should be used with caution.

In addition to limitations imposed by map and photo scales, one other factor also significantly constrains the resolution of the present map. The photography used in this analysis, which was acquired under high sun-angle conditions, is not well suited for the discrimination and mapping of subtle topographic features. Consequently, reexamination of any of the fault systems shown on this map using larger scale and (or) lower sun-angle aerial photography would very likely reveal a substantial number of additional young fault segments.

## PATTERNS OF LATEST TERTIARY AND QUATERNARY FAULTING

Several factors significantly influence the preservation of fault-related landforms and, therefore, the apparent distribution of young faults as indicated by the distribution of these landforms can be significantly biased. These factors include (1) composition, induration, and structural integrity of the rock or sediment type(s) underlying fault scarps; (2) local geomorphic environment of the scarp or other fault-related landform; (3) regional climatic conditions and paleoclimatic variations; and (4) magnitude and recurrence of fault movement (Wallace, 1977; Bucknam and Anderson, 1979; Nash, 1980, 1984; Hanks and others, 1984; Mayer, 1984; Pierce and Colman, 1986; Machette, 1986, 1988, 1989). Therefore, the distribution of young faults shown on this map provides, at best, only an approximate and somewhat biased picture of late Tertiary and Quaternary faulting within the quadrangle. Specifically, faults having a long history of recurrent movement, juxtaposing bedrock and alluvium, or cutting upper Cenozoic lava flows and (or) welded ash-flow tuffs tend to be overrepresented, whereas faults of pre-late Pleistocene age cutting unconsolidated surficial deposits and having either short histories of recurrent movement or long recurrence intervals tend to be underrepresented. Scarps developed on volcanic rocks may be preserved for periods of as much as 10 m.y. By comparison, scarps on the fluvially active parts of piedmont surfaces would likely be completely destroyed within a few thousand years at most, and even on inactive piedmont surfaces, fault scarps on unconsolidated alluvial fill are significantly rounded within 10,000 years (Wallace, 1977), and low scarps would be sufficiently degraded to be unrecognizable on standard aerial photography within a few hundred years (Wallace, 1977; Nash, 1980; Hanks and others, 1984; Machette, 1986).

Young faulting within the Lovelock quadrangle is variable (table 1 and fig. 1). Young fault densities west and northwest of the Black Rock and Smoke Creek Deserts average approximately 0.09 km/km<sup>2</sup>, and fault trace orientations average approximately N10°W. The area is largely underlain by volcanic rocks of late Miocene age. In contrast, young faults are almost completely absent from a 20- to 25-km-wide, north-northeast-trending zone of extensive pediments that extends westward from and roughly parallels the valley of the Humboldt River in the east-central part of the quadrangle. In the east-central part of the quadrangle, however, major range-bounding fault zones are almost uniformly distributed; young fault densities average approximately 0.05 km/km<sup>2</sup> and fault trace orientations average about N15°E (table 1 and fig. 1). All of these fault zones have been active during Quaternary (and, quite possibly, late Pleistocene) time, and at least five of them display abundant evidence of late Holocene (10-30 ka) and (or) Holocene (0-10 ka) movement (for example, the west flanks of the Fox and Lake Ranges, southwest flank of the Granite Range, east flank of the Shawave Mountains, and west flank of the Humboldt Range). Late Pleistocene and (or) Holocene movement also has occurred within two north-northeast-trending systems of distributed intrabasin faulting. These intrabasin systems are located near the center of the quadrangle (east of the Selenite Range and northeast of the Trinity Range).

## REFERENCES CITED

- Bucknam, R.C., and Anderson, E., 1979, Estimation of fault-scar ages from a scarp-height-slope-angle relationship, *Geology*, v. 7, p. 11-14.
- Bull, W.B., 1977, Tectonic geomorphology of the Mojave Desert, California: U.S. Geological Survey, Office of Earthquakes, Volcanoes, and Engineering, Contract Report 14-08-001-G-394, Menlo Park, Calif., 183 p.
- Bull, W.B., and McFadden, L.D., 1977, Tectonic geomorphology north and south of the Garlock fault, California, in Doehring, D.C., ed., *Geomorphology in arid regions: Proceedings 8th Annual Geomorphology Symposium*, State University of New York at Binghamton, p. 115-137.
- Christiansen, R.L., and McKee, E.H., 1978, Late Cenozoic volcanic and tectonic evolution of the Great Basin and Columbia intermontane basin: *Geological Society of America Memoir* 152, p. 283-311.
- Eaton, G.P., 1982, The Basin and Range Province: Origin and tectonic significance: *Annual Reviews Earth and Planetary Science*, v. 10, p. 409-440.
- Hanks, T.C., Bucknam, R.C., Lajoie, K.R., and Wallace, R.E., 1984, Modification of wave-cut and faulting-controlled landforms: *Journal of Geophysical Research*, 89 (B7), p. 5771-5790.
- Machette, M. N., 1986, History of Quaternary offset and paleoseismically along the La Jencia fault, central Rio Grande rift, New Mexico: *Seismology Society of America Bulletin*, v. 76, p. 259-272.
- 1988, Quaternary movement along the La Jencia fault, central New Mexico: U. S. Geological Survey Professional Paper 1440, 81 p.
- 1989, Slope-morphometric dating, in Forman, S.L., ed., *Dating methods applicable to Quaternary geologic studies in the western United States: Utah Geological and Mineral Survey Miscellaneous Publication* 89-7, p. 30-42.
- Mayer, L., 1984, Dating Quaternary fault scarps formed in alluvium using morphological parameters: *Quaternary Research*, v. 22, p. 303-313.
- Nash, D.B., 1980, Morphological dating of degraded normal fault scarps: *Journal of Geology*, v. 88, p. 353-360.
- 1984, Morphologic dating of fluvial terrace scarps and fault scarps near West Yellowstone, Montana: *Geological Society of America Bulletin*, v. 95, p. 1413-1424.
- Pierce, K.L., and Colman, S.M., 1986, Effect of height and orientation (microclimate) on geomorphic degradation rates and processes, late-glacial terrace scarps in central Idaho: *Geological Society of America Bulletin*, v. 97, p. 869-885.
- Stewart, J.H., 1983, Cenozoic structure and tectonics of the northern Basin and Range Province, California, Nevada and Utah, in *The role of heat in the development of energy and mineral resources in the northern Basin and Range Province: Geothermal Resources Council, Special Report* 13, p. 25-40.
- Thornbury, W.D., 1969, *Principles of Geomorphology*, John Wiley & Sons, Inc., New York, 594 p.
- Wallace, R.E., 1977, Profiles and ages of young fault scarps, north-central Nevada: *Geological Society of America Bulletin*, v. 88, p. 1267-1281.
- 1978, Geometry and rates of change of fault-generated range fronts, north-central Nevada: *U.S. Geological Survey Journal of Research*, v. 6, p. 637-650.
- 1979, Map of young fault scarps related to earthquakes in north central Nevada: U.S. Geological Survey Open File Report 79-1554, scale 1:125,000.
- Zoback, M.L., Anderson, R.E., and Thompson, G.A., 1981, Cenozoic evolution of the state of stress and style of tectonism of the Basin and Range Province of the western United States, in Vine, F.J., and Smith, A.D., eds., *Extensional tectonics associated with convergent plate boundaries*: The Royal Society, London, p. 189-216.

Fault scarps on Quaternary surficial deposits or Quaternary erosional surfaces—Scarps on Quaternary deposits or Quaternary erosional surfaces other than those displacements associated with known historic earthquakes. Hachures indicate downslope direction of scarp. Symbol indicates approximate age of offset surficial deposit or erosion surface:

- Q3 Holocene (0-10 ka)
- Q2-3 Latest Pleistocene and (or) Holocene (0-30 ka)
- Q2 Late Pleistocene (10-130 ka)
- Q1-2 Early to middle and (or) late Pleistocene (0.01-1.5 Ma)
- Q1 Early to middle Pleistocene (0.13-1.5 Ma)

Photogeologic and geomorphic criteria used to estimate ages are summarized in table 2

Fault-related lineaments on Quaternary surficial deposits or on Quaternary erosion surfaces—Alignments, in Quaternary surficial deposits or across Quaternary erosional surfaces, of one or more of the following features: linear reaches of stream channels, linear stream valleys, shallow linear swales, springs, vegetation discontinuities. Commonly associated with fault scarps on Quaternary deposits and (or) erosional surfaces. Age designations as above

Major range-front faults—Faults bounding tectonically active fronts of major mountain ranges. These range fronts are characterized by: fault juxtaposition of Quaternary alluvium against bedrock; fault scarps and lineaments on surficial deposits along or immediately adjacent to range front, a general absence of pediments, abrupt piedmont-hillslope transitions, steep bedrock slopes, faceted spurs, winged valleys, and subparallel systems of high-gradient, narrow, step-sided canyons orthogonal to range front. Only mapped in areas along range front where fault scarps and (or) lineaments on Quaternary surficial deposits or on Quaternary erosion surfaces are absent

Faults juxtaposing Quaternary alluvium against bedrock (other than major range-front faults)—Morphologically similar to major range-front faults except that associated fault systems are extensive and fault scarps are substantially lower, shorter, and less continuous. Solid lines indicate locations where fault scarps are abrupt and well-defined; long dashes indicate locations where scarps are less well defined

Faults forming scarps and (or) prominent topographic lineaments on Tertiary volcanic or sedimentary rocks—These scarps have morphologies ranging from partly rounded to moderately dissected. Topographic lineaments are composed of alignments of one or more of the following landforms: abrupt scarps, linear hillside ridges, benches and trenches, linear reaches of stream channels and small stream valleys, ridge-crest saddles and cols, linear depressions and small closed basins. Many of these faults form closely spaced groups in areas underlain by Tertiary ash-flow tuffs and lava flows

## RECONNAISSANCE PHOTOGEOLOGIC MAP OF YOUNG FAULTS IN THE LOVELOCK 1° BY 2° QUADRANGLE, NEVADA AND CALIFORNIA

By  
John C. Dohrenwend, Mary Anne McKittrick, and Barry C. Moring  
1991

M(200)

MF

no. 2178

C.I.