

Base from U.S. Geological Survey, 1956 Transverse Mercator projection
MAPPED IN 1988-90 BY J.C. DOHRENWEND AND B.A. SCHELL
EDITED BY C.L. OSTERGREN, PREPARED BY LORI MOORE
MANUSCRIPT APPROVED FOR PUBLICATION JUNE 28, 1991

Table 1. Characteristics of young faults in the Lund 1° by 2° quadrangle (See figure 1 for area boundaries)

Orientation	West		East	
	Length (km)	Percent	Length (km)	Percent
90.0 to 80°	6.0	0.7	4.6	1.1
79.9 to 70°	12.0	1.3
69.9 to 60°	24.2	2.7
59.9 to 50°	57.8	6.5	10.4	2.5
49.9 to 40°	62.8	7.0	11.4	2.7
39.9 to 30°	50.6	5.7	2.0	0.5
29.9 to 20°	101.6	11.4	24.2	5.7
19.9 to 10°	169.0	18.9	32.2	7.6
09.9 to 0.0°	181.8	20.3	110.6	26.1
359.9 to 350°	139.8	15.6	111.2	26.2
349.9 to 340°	49.2	5.5	42.6	10.1
339.9 to 330°	30.6	3.4	47.0	11.1
329.9 to 320°	2.6	0.3	15.8	3.7
319.9 to 310°	8.6	2.0
309.9 to 300°
299.9 to 290°	2.6	0.3
289.9 to 280°	3.2	0.4	3.2	0.8
279.9 to 270°
Total (km)	892.8	100.0	423.8	100.0
Area (km ²)	10825	...	8565	...
Density (km/km ²)	0.083	...	0.049	...
Number	186	...	88	...
Density (no./km ²)	0.017	...	0.010	...
Mean length (km)	4.8	...	4.8	...

Table 2. General photogeologic and geomorphic criteria used to estimate general ages of piedmont surfaces

Map unit	Age	Depth of dissection	Drainage net morphology	Interfluvial morphology	Geomorphic relations	Typical geomorphic environments	General field criteria
Unit Q3	Holocene (0 to 10 ka)	Shallow to none; generally <3 m	Predominantly radial from fan apex; channels typically poorly to moderately well defined	Typically poorly defined; bar and swale micro-topography common on most surfaces	Surfaces cut by pluvial shorelines and (or) late Pleistocene glacial moraines	Distal piedmont surfaces; proximal areas; (proximal surfaces along some highly active range fronts)	Unweathered to slightly weathered; very weak to weak soil development
Unit Q2	Late Pleistocene (10 to 130 ka)	Shallow to moderate; typically 2-6 m	Predominantly distributary; however, some channels head on piedmont; well-defined channels	Typically well defined; surfaces broad and flat with abrupt margins	Surfaces overlain by pluvial shorelines and (or) latest Pleistocene glacial moraines	Proximal to distal piedmont surfaces; some inset terraces	Weakly to moderately well developed soils; interlocking stone pavements
Unit Q1	Early to middle Pleistocene (0.13 to 1.3 Ma)	Moderate to deep; commonly >10 m	Predominantly subparallel; well-defined channels	Well defined; older interfluvial surfaces (bollinas) commonly narrow and irregular	Surfaces overlain by pluvial shorelines and (or) latest Pleistocene glacial moraines	Generally confined to intermediate and proximal piedmont areas	Moderately to very well developed soils; interlocking to highly degraded stone pavements




Figure 1. Rose diagrams summarizing orientation of young faulting. Number indicates mean fault trend. See table 1 for data.

INTRODUCTION

The State of Nevada occupies most of the central and western parts of the Great Basin, the largest tectonically active region within the Basin and Range geomorphic province of North America. The topography of this region is typified by generally north-northwest-trending, subparallel mountain ranges separated by alluvial basins of similar plan form and orientation. This classic Basin and Range physiography is the product of at least two phases of middle to late Cenozoic extensional faulting (Zoback and others, 1981; Eaton, 1982; Stewart, 1983), and most of the basins and ranges of the region are at least partly bounded by late Cenozoic faults. The earlier phase of extension was marked by widespread shallow detachment faulting that began at approximately 35 Ma and locally continued into the time of late Cenozoic extensional faulting (Zoback and others, 1981; Eaton, 1982; Stewart, 1983). The later phase, which was dominated by high-angle, more deeply penetrating block faulting, may have begun locally at about 17 Ma and continues episodically to the present time (Christiansen and McKee, 1978; Eaton, 1982). This map, one of a series of 1° by 2° quadrangle maps showing young faults in Nevada, provides a generalized picture of the late Tertiary and Quaternary faulting that is associated with the later extensional phase. These young faults are a primary determinant of the present configuration of ranges and basins within the quadrangle.

MAPPING PROCEDURE

Young faults are herein defined as those faults that have undergone latest Tertiary and (or) Quaternary movement. These faults are commonly marked by a variety of diagnostic structural landforms and other surficial phenomena that can be readily identified and mapped on aerial photographs. These features include (1) scarps on latest Tertiary and Quaternary surficial deposits, volcanic tuffs, or geomorphic surfaces (either erosional or depositional); (2) prominent alignments of linear drainageways, ridges and swales, active springs and (or) spring deposits, and linear discontinuities of structure, rock type, and vegetation; and (3) abrupt, steeply sloping range fronts with basal scarps, faceted spurs, winglike valleys, and elongate drainage basins with narrow valley floors (Thornbury, 1969; Bull, 1977; Bull and McFadden, 1977; Wallace, 1977, 1978).

National High Altitude Program (NHAP), 1:58,000-nominal-scale, color-infrared photography was used for photogeologic interpretation. This mapping was transferred directly to 1/2° by 1° topographic quadrangle maps that were enlarged to the scale of the photographs. These maps were reduced and compiled at 1:250,000 scale. This compilation was then compared with previous mapping of young faults within the quadrangle (Ertco Western, Inc., 1981) and significant differences between maps were resolved. Because the previous mapping was based on photogeologic analysis of 1:24,000-scale color aerial photography and extensive field verification, more young faults were identified by this mapping than by the present reconnaissance study. Many of these additional faults are located on distal piedmonts and basin flats and most are included on our map to show their approximate location and general average orientation of each fault segment. These data are summarized in table 1 and figure 1.

General ages of surficial deposits and erosion surfaces cut by young faults were estimated using a variety of photogeologic and geomorphic criteria (table 2). These age estimates provide a general indication of the approximate timing of young faulting throughout the quadrangle. However, it should be emphasized that these data do not necessarily reflect the age of most recent surface rupture along any particular fault segment. Rather, they provide only very general (and commonly somewhat biased) age constraints on this surface rupture. Age estimates based on photogeologic analysis of surficial deposits and erosion surfaces are, at best, both tentative and imprecise. Moreover, the distribution of these deposits and surfaces is inherently biased by geomorphic process and environment. For example, in those areas of the Great Basin where uplift rates are low to moderate, remnants of older geomorphic surfaces tend to be concentrated in proximal piedmont areas, whereas younger surficial deposits tend to accumulate on distal piedmonts and basin flats. Consequently, young faults located in intrabasin areas are more likely to offset younger surface deposits than are faults located along range fronts or in proximal piedmont areas. Therefore, inferences based on these data regarding the temporal distribution of young fault activity should be used with caution.

In addition to the limitations imposed by map and photo scales, one other factor also significantly constrains the resolution of the present map. The photography used in this analysis, which was acquired under high sun-angle conditions, is not well suited for the discrimination and mapping of subtle topographic features. Consequently, reexamination of any of the fault systems shown on this map using larger scale and (or) lower sun-angle aerial photography would very likely reveal a substantial number of additional young fault segments.

PATTERNS OF LATEST TERTIARY AND QUATERNARY FAULTING

Several factors significantly influence the preservation of fault-related landforms and, therefore, the apparent distribution of young faults as indicated by the distribution of these landforms can be significantly biased. These factors include (1) composition, induration, and structural integrity of the rock or sediment type(s) underlying fault scarps; (2) geomorphic environment of the scarps or other fault-related landforms; (3) regional climatic conditions and paleoclimatic variations; and (4) magnitude and recurrence of fault movement (Wallace, 1977; Bucknam and Anderson, 1979; Nash, 1980, 1984; Hanks and others, 1984; Mayer, 1984; Pierce and Colman, 1986; Machette, 1986, 1988, 1989). Therefore, the distribution of young faults shown on this map provides, at best, only an approximate and somewhat biased picture of late Tertiary and Quaternary faulting within the quadrangle. Specifically, faults having a long history of recurrent movement, juxtaposing bedrock and alluvium, or cutting upper Cenozoic flows and (or) welded ash flows tend to be overrepresented whereas faults of pre-late Pleistocene age cutting unconsolidated surficial deposits and having other short histories of recurrent movement or long recurrence intervals tend to be underrepresented. Scarps developed on volcanic rocks may be preserved for periods of as long as 10 m.y. By comparison, scarps on the fluvially active parts of piedmont surfaces would likely be completely destroyed within a few thousand years at most, and even on inactive piedmont surfaces, fault scarps on unconsolidated alluvial fill are significantly rounded within 10,000 years (Wallace, 1977), and low-angle scarps are so significantly degraded by truncation on standard aerial photography within a few hundred thousand years (Wallace, 1977; Hanks and others, 1984; Machette, 1989).

Young faulting within the Lund 1° by 2° quadrangle is fairly typical of the central Great Basin. Fault orientations are predominantly north-northeasterly in areas along the west margin of Lake Valley, and are nearly bisected by the quadrangle from north to south whereas they are predominantly northerly in areas to the east of that range (table 1, fig. 1). This west-to-east change is part of a general change in young fault orientation across the central Great Basin from central Nevada to central Utah. Moreover, the density of young faults also changes from west to east across the quadrangle. Average young fault densities in the western part of the quadrangle are nearly twice as high as average densities in the eastern part (0.08 km/km² and 0.05 km/km², respectively; table 1).

Two major range-bounding fault zones, each showing evidence of late Pleistocene and (or) Holocene movement, form the most prominent young fault features in the quadrangle. These major fault zones define the west edge of the Quin Canyon Range-Grant Range-White Pine Mountains and the west edge of the Egan Range. Other areas with evidence of late Pleistocene and (or) Holocene movement are located on distal piedmonts and basin flats in southern Railroad Valley, central White River Valley, along the west margin of Lake Valley, and at the south end of Snake Valley (in the extreme northeast corner of the quadrangle). One of these areas, the central part of White River Valley, contains the largest cluster of young intrabasin faults in the west-central part of the Great Basin. The predominant north-northeasterly to northerly orientations are typical of most young intrabasin faults throughout the Great Basin.

REFERENCES CITED

Bucknam, R.C., and Anderson, E., 1979, Estimation of fault-scarp ages from a scarp-height-slope-angle relationship: *Geology*, v. 7, p. 11-14.

Bull, W.B., 1977, Tectonic geomorphology of the Mojave Desert, California: U.S. Geological Survey, Office of Earthquakes, Volcanoes, and Engineering, Contract Report 14-08-001-G-394, Menlo Park, California, 188 p.

Bull, W.B., and McFadden, L.D., 1977, Tectonic geomorphology north and south of the Garlock fault, California, in Doehring, D.C., ed., *Geomorphology in arid regions: Proceedings 8th Annual Geomorphology Symposium*, State University of New York, Binghamton, p. 113-137.

Christiansen, R.L., and McKee, E.H., 1978, Late Cenozoic volcanic and tectonic evolution of the Great Basin and Columbia intermontane basin: *Geological Society of America Memoir* 152, p. 283-311.

Eaton, G.P., 1982, The Basin and Range Province: Origin and tectonic significance: *Annual Reviews Earth and Planetary Science*, v. 10, p. 409-440.

Ertco Western, Inc., 1981, Faults and lineaments in the MX siting region, Nevada and Utah: unpublished report prepared for the Department of the U.S. Air Force, Norton A.F.B., Calif., 77 p., 11 maps, scale 1:250,000.

Hanks, T.C., Bucknam, R.C., Lajoie, K.R., and Wallace, R.E., 1984, Modification of wave-cut and faulting-controlled landforms: *Journal of Geophysical Research*, v. 89 (B7), p. 5771-5790.

Machette, M.N., 1986, History of Quaternary offset and paleoseismicity along the La Jencia fault, central Rio Grande rift, New Mexico: *Seismology Society of America Bulletin*, v. 76, p. 259-272.

———, 1988, Quaternary movement along the La Jencia fault, central New Mexico: U.S. Geological Survey Professional Paper 1460, 81 p.

———, 1989, Slope-morphometric dating, in Forman, S.L., ed., *Dating methods applicable to Quaternary geologic studies in the western United States*: Utah Geological and Mineral Survey Miscellaneous Publication 89-2, p. 30-42.

Mayer, L., 1984, Dating Quaternary fault scarps formed in alluvium using morphological parameters: *Quaternary Research*, v. 22, p. 300-313.

Nash, D.B., 1980, Morphological dating of degraded normal fault scarps: *Journal of Geology*, v. 88, p. 353-360.

———, 1984, Morphologic dating of fluvial terrace scarps and fault scarps near West Yellowstone, Montana: *Geological Society of America Bulletin*, v. 95, p. 1413-1424.

Pierce, K.L., and Colman, S.M., 1986, Effect of height and orientation (microclimate) on geomorphic degradation rates and processes, late-glacial terrace scarps in central Idaho: *Geological Society of America Bulletin*, v. 97, p. 869-885.

Stewart, J.H., 1983, Cenozoic structure and tectonics of the northern Basin and Range Province, California, Nevada and Utah, in *The role of heat in the development of energy and mineral resources in the northern Basin and Range Province: Geothermal Resources Council Special Report* 13, p. 25-40.

Thornbury, W.D., 1969, *Principles of Geomorphology*: John Wiley & Sons, Inc., New York, 594 p.

Wallace, R.E., 1977, Profiles and ages of young fault scarps, north-central Nevada: *Geological Society of America Bulletin*, v. 88, p. 1267-1281.

———, 1978, Geometry and rates of change of fault-generated range fronts, north-central Nevada: *U.S. Geological Survey Journal of Research*, v. 6, p. 637-650.

Zoback, M.L., Anderson, R.E., and Thompson, G.A., 1981, Cenozoic evolution of the state of stress and style of tectonism of the Basin and Range Province of the western United States, in Vine, F.T., and Smith, A.D., eds., *Extensional tectonics associated with convergent plate boundaries*: The Royal Society, London, p. 189-216.

RECONNAISSANCE PHOTOGEOLOGIC MAP OF YOUNG FAULTS IN THE LUND 1° BY 2° QUADRANGLE, NEVADA AND UTAH

By
John C. Dohrenwend, Bruce A. Schell, and Barry C. Moring
1991

Any use of trade, product, or firm names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Government.