

Base from U.S. Geological Survey, 1956
Revised 1981
Transverse Mercator projection

Table 1. Characteristics of young faults in the Wells 1° by 2° quadrangle
(See figure 1 for area boundaries)

Orientation	West		East	
	Length (km)	Percent	Length (km)	Percent
90.0 to 80°	---	---	2.8	1.1
79.9 to 70°	12.4	1.4	---	---
69.9 to 60°	10.8	1.2	6.6	2.7
59.9 to 50°	9.0	1.0	16.4	6.7
49.9 to 40°	20.3	2.3	4.8	2.0
39.9 to 30°	47.2	5.3	13.4	5.4
29.9 to 20°	98.0	11.1	26.0	10.6
19.9 to 10°	112.8	12.7	50.8	20.7
09.9 to 0.0°	251.4	28.4	35.8	14.6
359.9 to 350°	201.6	22.8	58.6	23.8
349.9 to 340°	76.8	8.7	19.6	8.0
339.9 to 330°	13.4	1.5	---	---
329.9 to 320°	18.4	2.1	3.4	1.4
319.9 to 310°	8.4	0.9	2.8	1.1
309.9 to 300°	5.6	0.6	---	---
299.9 to 290°	---	---	5.0	2.0
289.9 to 280°	---	---	---	---
279.9 to 270°	---	---	---	---
Total (km)	886.1	100.0	246.0	100.0
Area (km ²)	14360	---	4130	---
Density (km/km ²)	0.062	---	0.060	---
Number	207	---	53	---
Density (no./km ²)	0.014	---	0.013	---
Mean length (km)	4.3	---	4.6	---



SCALE 1:250 000
0 5 10 15 20 25 MILES
0 5 10 15 20 25 KILOMETERS

CONTOUR INTERVAL 200 FEET
WITH SUPPLEMENTARY CONTOURS AT 100 FOOT INTERVALS
NATIONAL GEODETIC VERTICAL DATUM OF 1929

Table 2. General photogeologic and geomorphic criteria used to estimate general ages of piedmont surfaces

Map unit	Age	Depth of dissection	Drainage net morphology	Interfluvial morphology	Geomorphic relations	Typical geomorphic environments	General field criteria
Unit Q3	Holocene (0 to 10 ka)	Shallow to none; generally <3 m	Predominantly radial from fan apex; channels typically poorly to moderately well defined	Typically poorly defined; bar and swale micro-topography common on most surfaces	Surfaces cut by late Pleistocene glacial moraines	Distal piedmont surfaces; channels and terraces in proximal areas; (proximal) surface along some highly active range fronts	Unweathered to slightly weathered; very weak to weak soil development
Unit Q2	Late Pleistocene (10 to 130 ka)	Shallow to moderate; typically 2-6 m	Predominantly radial from fan apex; however, some channels head on piedmont; well-defined channels	Typically well defined; broad and flat with abrupt margins	Surfaces overlain by pluvial shorelines and (or) late Pleistocene glacial moraines	Proximal to distal piedmont surfaces; some inset terraces	Weakly to moderately well developed soils; interlocking stone pavements
Unit Q1	Early to middle Pleistocene (0.13 to 1.5 Ma)	Moderate to deep; commonly >10 m	Predominantly subparallel; well-defined channels	Well defined; older interfluvial surfaces (ballenas) commonly narrow and irregular	Surfaces overlain by pluvial shorelines and (or) late Pleistocene glacial moraines	Generally confined to intermediate and proximal piedmont areas	Moderately to very well developed soils; interlocking stone pavements



Mapped in 1989-90 by J.C. Dohrenwend and M.A. McKittrick
Edited by C.L. Goring; prepared by Lori Moore
Manuscript approved for publication June 28, 1991




Figure 1. Rose diagrams summarizing orientation of young faulting. Number indicates mean fault trend. See table 1 for data.

INTRODUCTION

The State of Nevada occupies most of the central and western parts of the Great Basin, the largest tectonically active region within the Basin and Range province of North America. The topography of this region is typified by generally north-northeast to northeast-trending, subparallel mountain ranges separated by alluvial basins of similar plan form and orientation. This classic Basin and Range physiography is the product of at least two phases of middle to late Cenozoic extensional faulting (Zoback and others, 1981; Eaton, 1982; Stewart, 1983), and most of the basin and range of the region are at least partly bounded by late Cenozoic faults. The earlier phase of extension was marked by widespread shallow detachment faulting that began at approximately 35 Ma and locally continued into the time of the later phase (Eaton, 1982; Stewart, 1983). The later phase, which was dominated by high-angle, more deeply penetrating block faulting, may have begun locally at about 17 Ma and continues episodically to the present time (Christiansen and McKee, 1978; Eaton, 1982). This map, one of a series of 1° by 2° quadrangle maps showing young faults in Nevada, provides a generalized picture of the late Tertiary and Quaternary faulting that is associated with the latter extensional phase. These young faults are a primary determinant of the present configuration of ranges and basins within the quadrangle.

MAPPING PROCEDURE

Young faults are herein defined as those faults that have undergone latest Tertiary and (or) Quaternary movement. These faults are commonly marked by a variety of diagnostic structural landforms and other surficial phenomena that can be readily identified and mapped on aerial photographs. These features include (1) scarps on latest Tertiary and (or) Quaternary surficial deposits, volcanic strata, or geomorphic surfaces (either erosional or depositional); (2) prominent alignments of linear drainageways, ridges and swales, active springs and (or) spring deposits, and linear discontinuities of structure, rock type, and vegetation; and (3) abrupt, steeply sloping range fronts with basal scarps, faceted spurs, wineglass valleys, and elongate drainage basins with narrow valley floors (Thornbury, 1969; Bull, 1977; Bull and McFadden, 1977; Wallace, 1977, 1978).

National High Altitude Program (NHAP), 1:58,000 nominal-scale, color-infrared photographs were used for photogeologic analysis. This mapping was transferred directly to 1/2° by 1° topographic quadrangle maps that were enlarged to the scale of the photographs. These maps were then reduced, compiled at 1:250,000 scale, and digitized using a GTCO digitizing board connected to a Macintosh II microcomputer. The resulting vector file was converted to raster format (cell size = 200 m by 200 m) and analyzed to determine the approximate length and average orientation of each fault segment. These data are summarized in table 1 and figure 1.

General ages of surficial deposits and erosion surfaces cut by young faults were estimated using a variety of photogeologic and geomorphic criteria (table 2). These age estimates provide a general indication of the approximate timing of young faulting throughout the quadrangle. However, it should be emphasized that these data do not necessarily reflect the age of most recent surface rupture along any particular fault segment. Rather, they provide only very general (and commonly somewhat biased) age constraints on this surface faulting. Age estimates based on photogeologic analysis of surficial deposits and erosion surfaces are, at best, both tentative and imprecise. Moreover, the distribution of these deposits and surfaces is inherently biased by geomorphic process and environment. For example, in those areas of the Great Basin where range uplift rates are low, the remains of older geomorphic surfaces tend to be concentrated in proximal piedmont areas, whereas younger surficial deposits tend to accumulate on distal piedmonts and basin flats. Consequently, young faults located in intrabasin areas are more likely to offset younger surface deposits than are faults located along range fronts or in proximal piedmont areas. Therefore, inferences based on these data regarding the temporal distribution of young fault activity should be used with caution.

PATTERNS OF LATEST TERTIARY AND QUATERNARY FAULTING

Several factors significantly influence the preservation of fault-related landforms and, therefore, the apparent distribution of young faults as indicated by the distribution of these landforms can be significantly biased. These factors include (1) composition, induration, and structural integrity of the rock or sediment type(s) underlying fault scarps; (2) local geomorphic environment of the scarp or other fault-related landform; (3) regional climatic conditions and paleoclimatic variations; and (4) magnitude and recurrence of fault movement (Wallace, 1977; Bucknam and Anderson, 1979; Nash, 1980, 1984; Hanks and others, 1984; Mayer, 1984; Pierce and Colman, 1986; Machette, 1986, 1988, 1989). Therefore, the distribution of young faults shown on this map provides, at best, only an approximate and somewhat biased picture of late Tertiary and Quaternary faulting within the quadrangle. Specifically, faults having a long history of recurrent movement, juxtaposing bedrock and alluvium, or cutting upper Cenozoic lava flows and (or) welded ash-flow tuffs tend to be overrepresented whereas faults of pre-late Pleistocene age cutting unconsolidated surficial deposits and having either short histories of recurrent movement or long recurrence intervals tend to be underrepresented. Scarps developed on volcanic rocks may be preserved for periods of as long as 10 m.y. By comparison, scarps on the fluvially active parts of piedmont surfaces would likely be completely destroyed within a few thousand years at most, and even on inactive piedmont surfaces, fault scarps on unconsolidated alluvial fill are significantly rounded within 10,000 years (Wallace, 1977), and low relief would be sufficiently degraded to be unrecognizable on standard aerial photography within a few hundred thousand years (Wallace, 1977; Hanks and others, 1984; Machette, 1989).

Young faults in the Wells 1° by 2° quadrangle are less abundant and less well defined than in most other areas of Nevada. Young fault density is relatively low, averaging about 0.06 km² throughout most areas of the quadrangle (table 1). Major range-bounding faults are limited to three general areas: the east flank of the Independence Mountains on the extreme west edge of the quadrangle, the east flank of the Granite Range in the east-central part of the quadrangle, and a general north-south-trending alignment of three fault zones that nearly bisect the quadrangle just west of long 115° W. Only those major range-bounding fault zones along the Granite and East Humboldt Ranges show evidence of possible late Pleistocene (10-130 ka) faulting. Other areas of possible late Quaternary movement are present on the east flank of the Adobe Range, along the northwest margin of Goshute Valley and along the northeast and northwest margins of Pilot Creek valley. All of these areas lie in the south half of the quadrangle and most are located in the south-central to southeastern part. Evidence of possible Holocene (0-10 ka) faulting is restricted to the northwest flank of the East Humboldt Range.

REFERENCES CITED

Bucknam, R.C., and Anderson, E., 1979, Estimation of fault-scarp ages from a scarp-height-slope-angle relationship: *Geology*, v. 7, p. 11-14.
Bull, W.B., 1977, Tectonic geomorphology of the Mojave Desert, California: U.S. Geological Survey, Office of Earthquakes, Volcanoes, and Engineering, Contract Report 14-08-001-G-394, Menlo Park, Calif., 188 p.
Bull, W.B., and McFadden, L.D., 1977, Tectonic geomorphology north and south of the Garlock fault, California, in Doehring, D.C., ed., *Geomorphology in arid regions: Proceedings 8th Annual Geomorphology Symposium*, State University of New York, Binghamton, p. 115-137.
Christiansen, R.L., and McKee, E.H., 1978, Late Cenozoic volcanic and tectonic evolution of the Great Basin and Columbia intermontane basin: *Geological Society of America Memoir* 152, p. 283-311.
Eaton, G.P., 1982, The Basin and Range Province: Origin and tectonic significance: *Annual Reviews Earth and Planetary Sciences*, v. 10, p. 409-440.
Hanks, T.C., Bucknam, R.C., Lajoie, K.R., and Wallace, R.E., 1984, Modification of wave-cut and faulting-controlled landforms: *Journal of Geophysical Research*, v. 89 (B7), p. 5771-5790.
Machette, M. N., 1986, History of Quaternary offset and paleoseismicity along the La Jencia fault, central Rio Grande rift, New Mexico: *Seismology Society of America Bulletin*, v. 76, p. 259-272.
———, 1988, Quaternary movement along the La Jencia fault, central New Mexico: U.S. Geological Survey Professional Paper 1460, 81 p.
———, 1989, Slope-morphometric dating, in Forman, S.L., ed., *Dating methods applicable to Quaternary geologic studies in the western United States: Utah Geological and Mineral Survey Miscellaneous Publication* 89-7, p. 20-42.
Mayer, L., 1984, Dating Quaternary fault scarps formed in alluvium using morphological parameters: *Quaternary Research*, v. 22, p. 300-313.
Nash, D.B., 1980, Morphological dating of degraded normal fault scarps: *Journal of Geology*, v. 88, p. 353-360.
———, 1984, Morphological dating of fluvial terrace scarps and fault scarps near West Yellowstone, Montana: *Geological Society of America Bulletin*, v. 95, p. 1413-1424.
Pierce, K.L., and Colman, S.M., 1986, Effect of height and orientation (microclimate) on geomorphic degradation rates and processes along glacial terrace scarps in central Idaho: *Geological Society of America Bulletin*, v. 97, p. 869-885.
Stewart, J.H., 1983, Cenozoic structure and tectonics of the northern Basin and Range Province, California, Nevada and Utah, in *The role of heat in the development of energy and mineral resources in the northern Basin and Range Province: Geothermal Resources Council, Special Report* 13, p. 25-40.
Thornbury, W.D., 1969, *Principles of Geomorphology*: John Wiley & Sons, Inc., New York, 594 p.
Wallace, R.E., 1977, Profiles and ages of young fault scarps, north-central Nevada: *Geological Society of America Bulletin*, v. 88, p. 1267-1281.
———, 1978, Geometry and rates of change of fault-generated range fronts, north-central Nevada: U.S. Geological Survey *Journal of Research*, v. 6, p. 637-650.
Zoback, M.L., Anderson, R.E., and Thompson, G.A., 1981, Cenozoic evolution of the state of stress and style of tectonism of the Basin and Range Province of the western United States, in Vine, F.T., and Smith, A.D., eds., *Extensional tectonics associated with convergent plate boundaries: The Royal Society, London*, p. 189-216.

— Fault scarps on Quaternary surficial deposits or Quaternary erosional surfaces—
Scarps on Quaternary deposits or Quaternary erosion surfaces other than those displacements associated with known historic earthquakes. Hashmarks indicate downslope direction of scarp. Symbol indicates approximate age of offset surficial deposit or erosion surface:
Q3 Holocene (0-10 ka)
Q2-3 Latest Pleistocene and (or) Holocene (0-30 ka)
Q2 Late Pleistocene (10-130 ka)
Q1-2 Early to middle and (or) late Pleistocene (0.01-1.5 Ma)
Q1 Early to middle Pleistocene (0.13-1.5 Ma)
Photogeologic and geomorphic criteria used to estimate ages are summarized in table 2

— Fault-related lineaments on Quaternary surficial deposits or on Quaternary erosion surfaces—Alignments in Quaternary surficial deposits or on Quaternary erosion surfaces, of one or more of the following features: linear reaches of stream channels, linear stream valleys, shallow linear swales, springs, vegetation discontinuities. Commonly associated with fault scarps on Quaternary deposits and (or) erosional surfaces. Age designations as above

— Major range-front faults—Faults bounding tectonically active fronts of major mountain ranges. These range fronts are characterized by: fault juxtaposition of Quaternary alluvium against bedrock, fault scarps and lineaments on surficial deposits along or immediately adjacent to range fronts, absence of a general absence of piedmont, hilltop, hillside, transitions, steep bedrock slopes, faceted spurs, wineglass valleys, and subparallel systems of high-gradient, narrow, steep-sided canyons orthogonal to range front. Only mapped in areas along range front where fault scarps and (or) lineaments on Quaternary surficial deposits or on Quaternary erosion surfaces are absent

— Faults juxtaposing Quaternary alluvium against bedrock (other than major range-front faults)—Morphologically similar to major range-front faults except that associated fault systems are significantly less extensive and fault scarps are substantially lower, shorter, and less continuous. Solid lines indicate locations where fault scarps are abrupt and well-defined; long dashes indicate locations where scarps are less well defined.

— Faults forming scarps and (or) prominent topographic lineaments on Tertiary volcanic or sedimentary rocks—These scarps have morphologies ranging from partly rounded and moderately dissected to undisturbed. Topographic lineaments are composed of alignments of one or more of the following landforms: abrupt scarps, linear hillside ridges, benches and terraces, linear reaches of stream channels and small stream, ridge crest saddles and cols, linear depressions and small closed basins. Many of these faults form closely spaced groups in areas underlain by Tertiary ash-flow tuffs and lava flows

RECONNAISSANCE PHOTOGEOLOGIC MAP OF YOUNG FAULTS IN THE WELLS 1° BY 2° QUADRANGLE, NEVADA, UTAH, AND IDAHO

By
John C. Dohrenwend, Mary Anne McKittrick, and Barry C. Moring
1991

Any use of trade, product, or firm names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Government.

INTERNET—GEOLOGICAL SURVEY, RESTON, VA—1981
For sale by U.S. Geological Survey, Map Distribution, Box 2308, Federal Center, Denver, CO 80225