

UNITED STATES DEPARTMENT OF THE INTERIOR  
GEOLOGICAL SURVEY

MODERN CHRYSOMONAD CYSTS FROM  
ALTA MORRIS LAKE, ELDORADO COUNTY, CALIFORNIA

by

David P. Adam

and

Albert D. Mahood

OPEN-FILE REPORT  
80-822

This report is preliminary and has not  
been reviewed for conformity with  
U. S. Geological Survey editorial standards  
or stratigraphic nomenclature

## INTRODUCTION

This report is one of a series illustrating siliceous cysts from various localities. Although these cysts have been known for many years, they are so small that they cannot be observed in detail with optical microscopes. The recent development of the scanning electron microscope (SEM) has made possible much more detailed observations of the external features of these cysts, but as yet relatively few forms have been recorded in this way.

A major difficulty is taxonomic confusion. Many and perhaps all of these cysts are the resting stages of various algae of the phylum Chrysophyta; they will be referred to in the rest of this report as chrysoomonad cysts, although other groups in addition to the Chrysoomonadinae may be represented. Modern forms are found primarily in fresh water, and numerous authors have reported chrysoomonad cysts from Holocene sediments (for example, see Nygaard, 1956). Older fossils have been recovered mostly from marine deposits, and are known as archaeomonads; whether the two groups are as distinct as this terminology suggests is not clear.

For an introduction to the literature on chrysoomonad cysts and siliceous algal scales, the reader is referred to Adam and Mahood (1979b), a preliminary annotated bibliography on the subject.

captions. Sample preparation techniques are generally the same as those used for preparing diatom samples; details may be found in Mahood and Adam (1979b).

The purpose of these initial reports is to provide primary documentation of the occurrence of particular cyst forms at particular localities, and to provide a means by which the SEM photographs of the cysts may be placed in a permanent depository. Counts of the relative abundance of the various forms and interpretations of their significance have not yet been attempted, but must await a more complete understanding of the range of cyst morphologies.

We have illustrated all of the distinctive cyst forms found in the sample, using the best available photographs. In some instances we have included more than one photograph of a given form, but we have not included all of the photographs we have taken.

Negatives of the plates for this report are on deposit at the USGS Photo Library, and prints can be obtained (at your expense) by writing to:

U. S. Geological Survey Library  
Photo Library  
Stop 914  
Box 25406, Denver Federal Center  
Denver, Colorado 80225

The fossil archaeomonads have been described and named entirely on the basis of their cysts. This is not advisable with modern forms, because the cysts are the remains of one stage of the life cycle of algae that presumably already have legitimate taxonomic names. Proper cyst nomenclature therefore depends on establishing which cysts are produced by which algae. At the moment, we have only a very limited knowledge of the forms that exist, and almost no knowledge of the phylogenetical pedigrees of the various forms.

The present work is directed towards expanding our knowledge of the various cyst forms and their geographic and environmental distributions. Taxonomic problems are ignored, and the various cyst forms are simply given numbers, which have been assigned arbitrarily. These numbers are consistent throughout all reports in this series, and are being used to tabulate where the various forms occur. (A list of the previous reports in this series is given in Appendix A). The approach used has been that of "splitting", as opposed to "lumping"; it may well be desirable to lump together many of the forms described here when more is known about them.

The SEM photographs are the most important part of this paper, and no attempt has been made to reduce them to words. Supporting data has been placed in the

#### SITE DESCRIPTION

Alta Morris Lake is located at an elevation of 2485 m in the upper part of the Fallen Leaf Lake drainage, west of Lake Tahoe. The local bedrock is part of the Mt. Tallac roof pendant, and consists of metasedimentary rocks of Mesozoic age (Burnett, 1971). The sample was collected as a periphyton diatom sample by Dennis Marelli for A. Mahood during July 1972. The location of the lake, together with the locations of other sites where we have recovered cysts, is shown in Figure 1.



#### REFERENCES CITED

- Adam, David P., and Mahood, Albert M., 1979b, A preliminary working bibliography on siliceous algal cysts and scales. U. S. Geological Survey Open-File Report No. 79-1215, 34 p.
- Burnett, John, 1971, Geology of the Lake Tahoe Basin: California Geology, v. 24, no. 7, p. 119-127.
- Vygaard, Gunnar, 1956, Ancient and Recent flora of diatoms and Chrysophyceae in Lake Gribso, in Berg, Kaj, and Petersen, I. C., eds., Studies on the humic acid Lake Gribso: Folia Limnologia Scandinavica, No. 3, p. 32-94, 12 plates.

Alta Morris Lake, Plate A  
scale bar = 3 micrometers

- 1 - Type 143  
(with adhering debris)
- 2 - Type 212
- 3 - Type 212  
(aperture not visible)
- 4 - Type 213  
(with adhering debris)
- 5 - Type 214
- 6 - Type 215  
(with adhering debris)
- 7 - Type 216  
(with adhering debris; aperture is the lower  
hole with the collar around it)
- 3 - Type 171  
(with adhering debris; note small spines around  
collar)
- 9 - Type 136  
(with adhering debris)
- 10 - Type 136
- 11 - Type 298
- 12 - Type 60  
(with adhering debris)

Alta Morris Lake - Plate A



1



2



3



4




5




6




7



8



9



10



11




12

Alta Morris Lake, Plate 9  
scale bar = 3 micrometers

- 1 - Type 217  
(with adhering debris)
- 2 - Type 218  
(with adhering debris)
- 3 - Type 218  
(with adhering debris)
- 4 - Type 114  
(aperture at right)
- 5 - Type 220  
(with adhering debris; aperture barely visible at top right)
- 6 - Type 218?  
(with adhering debris; aperture not visible)
- 7 - Type 332  
(aperture not visible)
- 8 - Type 56  
(with adhering debris)
- 9 - Type 56  
(with adhering debris)
- 10 - Type 114?  
(with adhering debris; aperture not visible)
- 11 - Type 221  
(with adhering debris, including diatom at top)
- 12 - Type 221  
(enlarged view of (11))

Alta Morris Lake - Plate B




1




2



3



4



5



6




7




8




9



10



11




12


Alta Morris Lake, Plate C  
scale bar = 3 micrometers

- 1 - Type 98  
(with adhering debris)
- 2 - Type 98  
(with adhering debris)
- 3 - Type 98  
(with adhering debris)
- 4 - Type 222  
(aperture not visible)
- 5 - Type 160  
(with adhering debris)
- 6 - Type 223  
(aperture at top right (??))
- 7 - Type 333  
(with adhering debris)
- 8 - Type 224  
(with adhering debris)

Alta Morris Lake - Plate C



1



2



3



4




5



6



7










8

Alta Morris Lake, Plate D  
scale bar = 3 micrometers

- 1 - Type 225  
(with adhering debris)
- 2 - Type 226  
(with adhering debris)
- 3 - Type 280  
(with adhering debris)
- 4 - Type 227  
(note low reticulation on surface)
- 5 - Type 326  
(with adhering debris; aperture at top right (??))
- 6 - Type 228  
(with adhering debris; aperture not visible)
- 7 - Type 49  
(aperture not visible)
- 8 - Type 49  
(with adhering debris)

Alta Morris Lake - Plate D



APPENDIX A  
PREVIOUS REPORTS IN THIS SERIES

- Mahood, Albert D., and Adam, David P., 1979a, Late Pleistocene chrysomonad cysts from core 7, Clear Lake, Lake County, California: U. S. Geological Survey Open-file Report Number 79-971, 11 p., 4 plates. Defines types 1 through 44.
- Adam, David P., and Mahood, Albert D., 1979a, A preliminary annotated bibliography on siliceous algal cysts and scales: U. S. Geological Survey Open-file Report Number 79-1215, 34 p.
- Mahood, Albert D., and Adam, David P., 1979b, Techniques used for the cleaning, concentration, and observation of chrysomonad cysts from sediments: U. S. Geological Survey Open-file Report Number 79-1431, 5 p.
- Adam, David P., and Mahood, Albert D., 1979b, Chrysomonad cysts from Upper Echo Lake, Eldorado County, California: U. S. Geological Survey Open-file Report Number 79-1461, 21 p. + 12 plates.
- Adam, David P., and Mehringer, Peter J., Jr., 1980, Modern and Holocene chrysomonad cysts from Lost Trail Pass Bog, Montana: U. S. Geological Survey Open-file Report Number 80-797, 13 p. + 5 plates.
- Adam, David P., and Mahood, Albert D., 1980, Modern chrysomonad cysts from Fallen Leaf Lake, Eldorado County, California: U. S. Geological Survey Open-file Report Number 80-798, 9 p. + 2 plates.