

A plateau collapse model for the formation of the West Antarctic rift system / Transantarctic Mountains

Paul G. Fitzgerald,¹ Robert W. Bialas,² W. Roger Buck,² and Michael Studinger²

¹Department of Earth Sciences, Syracuse University, Syracuse, NY 13244, USA (pgfitzge@syr.edu)

²Lamont-Doherty Earth Observatory of Columbia University, Palisades, NY 10964, USA (rbialas@ldeo.columbia.edu, buck@ldeo.columbia.edu, mstudinger@ldeo.columbia.edu)

Summary A recent model (Bialas et al., 2007) proposes that the Transantarctic Mountains (TAM) are the remnant edge of a collapsed plateau. In this model, the West Antarctic rift system/TAM region was a plateau (the "West Antarctic Plateau") with thicker than normal crust before undergoing a topographic reversal. Plateau collapse was due to continental extension and concomitant denudation beginning in the Jurassic when widespread tholeiitic magmatism marked initial rifting between East and West Antarctica. This was followed by the major period of West Antarctic extension in the Cretaceous and then more localized extension in the western Ross Embayment in the Paleogene and Neogene. The West Antarctic plateau formed inboard of subduction zone that existed along the Pacific margin of Gondwana throughout the Paleozoic until the late Mesozoic. The remnant plateau edge, now supported by slightly thickened crust, represent the ancestral TAM that remained following plateau collapse.

Citation: Fitzgerald, P.G., R.W. Bialas, W.R. Buck and M. Studinger (2007), A plateau collapse model for the formation of the West Antarctic rift system/Transantarctic Mountains, in *Antarctica: A Keystone in a Changing World – Online Proceedings of the 10th ISAES X*, edited by A. K. Cooper and C. R. Raymond et al., USGS Open-File Report 2007-1047, Extended Abstract 087, 4 p.

Introduction

The Transantarctic Mountains (TAM) bisect the Antarctic continent, separating the two contrasting geological provinces of East and West Antarctica (Dalziel, 1992) and forming the western flank of the West Antarctic rift system (WARS) (Figure 1). The TAM stretch for >3000 km, reach elevations >4000 m and are capped by undeformed strata of the Devonian-Triassic Beacon Supergroup. Low-temperature thermochronology has been applied along the TAM to constrain the denudation history and by inference, the uplift and formation of the mountains. Episodes of denudation are recorded in the Jurassic (Fitzgerald and Baldwin, 2007), the early Cretaceous, late Cretaceous, the Eocene and Oligocene (e.g., Fitzgerald, 2002 and references therein). Denudation along the TAM in the Cretaceous is typically 1-2 km. The majority of denudation has occurred since the early Eocene, being constrained as ca. 4-9 km at the mountain front, decreasing inland. These periods of denudation have been correlated to regional tectonic events, except the onset of the major phase of denudation beginning in the early Eocene, which has always been problematic (Fitzgerald, 2002). Thermochronology is used to constrain the thermal history of a region, with the recorded cooling history related to denudation, which in the TAM is erosional not tectonic. However, erosion is largely associated with the formation of relief, a change in base-level, that can be created by either mountain uplift or subsidence due to extension.

The WARS is a broad region of extended continental lithosphere, ca. 750-1000 km wide, lying dominantly below sea-level. Sedimentary basins (Northern Basin, Victoria Land Basin, Central Trough and the Eastern Basin) are only delineated well in the Ross Sea and generally trend parallel to the TAM (e.g., Davey and Brancolini, 1995). Extension within the WARS was initiated in the Jurassic (ca. 180 Ma) during the initial stages of breakup of Gondwana, as marked by voluminous tholeiitic magmatism found along the TAM (e.g., Elliot and Fleming, 2004). The main phase of distributed extension within the WARS was during the Cretaceous (e.g., Lawver and Gahagan, 1994). Extension during the Cenozoic was more localized, being isolated to the basins adjacent to the TAM (e.g., Hamilton et al., 2001; Davey et al., 2006).

The TAM are the longest intracontinental rift flank mountains in the world and there are many models for their formation. These models include those that invoke mainly thermal processes (e.g., Smith and Drewry, 1984), a combination of mechanical and thermal processes related primarily to extension in the adjacent WARS (e.g., Fitzgerald et al., 1986; ten Brink et al., 1997) or the mechanical effects of extension of thick lithosphere (van der Beek et al., 1994; Busetti et al., 1999). One of the major failures of all these models is that they typically do not attempt to explain the multiple phases of denudation that are separated by periods of relative tectonic and thermal quiescence. Nor do they attempt to explain the disparity between wide rifting in the Cretaceous that may require hot, thin lithosphere (e.g., Buck, 1991) whereas rift-flank uplift of the TAM may require cold-thick lithosphere. Two new models or ideas do address these concerns. Huerta and Harry (2007) use dynamic finite element modeling to explore possible conditions (especially those of the pre-rift thermal state of the lithosphere) responsible for the two-stage evolution of the WARS. Their models, using constant regional extension rates, predict diffuse rifting in the WARS followed by focused rifting near the East-West Antarctic boundary without requiring a change in tectonic regime (e.g., plate motion change) provided upper mantle temperatures are initially $730 \pm 50^\circ\text{C}$. Bialas et al. (2007) propose a plateau collapse model and use numerical modeling to test the idea that the WARS/TAM was a high elevation plateau (West Antarctic plateau) with thicker than normal crust before the onset of continental extension. With extension the WARS underwent a topographic reversal, but leaving the TAM as the remnant edge of the former plateau. In essence, Bialas et al. (2007) are modeling

the Cretaceous portion of the extensional history of the WARS. Their results indicate that elevation of the remnant plateau edge is related to initial upper mantle temperatures, with an initial mocho temperature of 675-850°C required to retain the plateau edge, but still generate wide rifting in the middle of the former plateau. In general, the Huerta and Harry (2007) model and the Bialas et al. (2007) plateau collapse model agree well on the conditions required for the earlier diffuse extension, with later extension being focused on the edges. In this paper we briefly examine some geological evidence that supports the plateau collapse model.




Figure 1. Topographic and bathymetric map of part of the Transantarctic Mountains and West Antarctic rift system. The WARS is the region defined by the Ross Sea, Ross Ice Shelf, and the West Antarctic Ice Sheet. The Ross Embayment = Ross Sea + Ross Ice Shelf. The Ross Sea is characterized by elevated heat flow and a series of asymmetric basins separated by basement highs. The WARS tectonic history is usually taken to be the same as the Ross Sea. CT = Central Trough; EB = Eastern Basin; NB = Northern Basin; TR = Terror Rift; VLB = Victoria Land Basin. Image created using data from the BEDMAP compilation [2000].

Discussion

The TAM has been the location of multiple orogenic episodes, including Rodinia assembly followed by Rodinia break-up that led to the formation of a rifted passive margin along the paleo-Pacific margin of Australia and East Antarctica at least up until the late Neoproterozoic (e.g., Goodge et al., 2004b). The margin changed to an active continental arc ca. 530 Ma until ca. 480 Ma (Ross Orogeny in Antarctica, Delamarian Orogeny in Australia), resulting in deformation, calc-alkaline magmatism and crustal thickening. In Australia, rollback from ca. 508-460 Ma formed a back-arc basin (Foster et al., 2005), although this apparently did not occur in Antarctica. The along-strike Australian equivalent to the region that is now the extended portion of the Ross Embayment is the Lachlan Fold Belt (LFB). This fold belt records an extremely complex record of deposition, shortening, accretion and magmatism from the Cambrian into the Carboniferous (e.g., Foster and Gray, 2000). In Antarctica, the Robertson Bay Group of northern Victoria Land is equivalent to the western subprovince of the LFB and the Swanson Formation of Marie Byrd Land to the eastern

subprovince (Summerfield, 1991; Sutherland, 1999; Foster and Gray, 2000). Outboard of the Antarctic fold belt-equivalent and along the paleo-Pacific margin of Gondwana, in what is now Marie Byrd Land, a continental arc was active from at least 320 Ma to ca. 110 Ma (Mukasa and Dalziel, 2000). By analogy with the LFB, it was during the Late Ordovician to the Carboniferous that the West Antarctic Plateau, a region of thickened continental crust, was initially formed, but was maintained and likely enhanced as subduction continued along the paleo-Pacific margin of Antarctica until the late Mesozoic.

In the TAM following the Ross Orogeny there was ca. 15-20 km of erosion (Capponi et al., 1990) at denudation rates of up to ca. 0.5 km/my (Goodge et al., 2004a) to eventually form the Kukri erosion surface. Following a period of non-deposition, strata of the Beacon Supergroup, initially marine, glacial, then non-marine, were deposited from Devonian to Triassic (e.g., Barrett, 1991; Isbell, 1999). While paleocurrents in the lower units indicate that relief on the erosion surface controlled deposition patterns, at least in the central TAM (Isbell, 1999), as deposition continued paleocurrents indicate that Beacon basins were elongate parallel to the current TAM with high topography either side, and thus a topographic reversal has occurred since the end of the Triassic (Barrett, 1981).

The initial break-up of Gondwana is recorded in Antarctica by voluminous Jurassic Ferrar tholeiitic magmatism, preserved as thick sills and lava flows along the TAM and is also associated with rifting (and hence subsidence) within the WARS (e.g., Elliot, 1992). In the TAM, this voluminous magmatism in the Jurassic likely resulted in crustal thickening. At ~105 Ma, subduction of the Pacific-Phoenix spreading center along the Pacific margin off Marie Byrd switched the tectonic regime and ~20 m.y. of extensional tectonics followed in New Zealand (e.g., formation of the Paparoa core complex (Tulloch and Kimbrough, 1989)), in Marie Byrd Land (e.g., exhumation of the Fosdick Mountains metamorphic complex from mid-crustal levels (Richard et al., 1994)) and in the Ross Embayment (e.g., wide rifting (Cooper et al., 1991); detachment faulting at DSDP site 270 (Fitzgerald and Baldwin, 1997) and detachment faulting and formation of mylonites in the eastern Ross Sea at Cape Colbeck (Siddoway et al., 2004)). Wide rifting in the Ross Embayment likely ceased when New Zealand and the Campbell Plateau were rifted away from Antarctica ca. 83-79 Ma (Stock and Cande, 2002).

The Bialas et al. (2007) numerical model for plateau collapse does not generate a remnant plateau edge that has the present elevation (ca. 4 km) of much of the peaks of the TAM. Rather the elevation of the post-wide rifting (late Cretaceous) remnant plateau-edge (proto-TAM) was likely in the ca. 2 km range and has since been accentuated by the isostatic response along the front of the range due to erosion, especially wet-based glacial erosion after 34 Ma that has enhanced peak heights by up to 50% (~2 km) (Stern et al., 2005).

The strength of this plateau collapse model for the WARS/TAM is that is supported by regional geological information and it appears to solve a number of geological or geophysical discrepancies in previous models. The concept of plateau collapse is also testable in the geological record and provides a basis for further study. In addition, the plateau collapse model is not mutually exclusive with respect to any other model that seeks to explain the Cenozoic development of the TAM to high elevations by other means, for example the thermal effects of hot West Antarctic mantle lithosphere.

Acknowledgments. This study was supported by NSF grants OPP03-38281 to Buck and Studinger and OPP03-38009 to Fitzgerald.

References

- Barrett, P. J. (1981), History of the Ross Sea region during the deposition of the Beacon Supergroup 400-180 million years ago, *Journal of the Royal Society of New Zealand*, 11, 447-459.
- Barrett, P. J. (1991), The Devonian to Triassic Beacon Supergroup of the Transantarctic Mountains and correlatives in other parts of Antarctica, in *The Geology of Antarctica*, edited by R. J. Tingey, pp. 120-152, Clarendon Press, Oxford.
- Bialas, R. W., W. R. Buck, M. Studinger, and P. G. Fitzgerald (2007), Plateau collapse model for the Transantarctic Mountains/West Antarctic rift system: Insights from numerical experiments, *Geology*, 35.
- Buck, W. R. (1991), Modes of continental lithospheric extension, *Journal of Geophysical Research*, 96, 20161-20178.
- Buseti, M., G. Spadini, F. Van der Wateren, S. Cloetingh, and C. Zanolla (1999), Kinematic modelling of the West Antarctic rift system, Ross Sea, Antarctica, *Global and Planetary Change*, 23, 79-103.
- Capponi, G., B. Messiga, G. B. Piccardo, M. Scambelluri, G. Traverso, and R. Vannucci (1990), Metamorphic assemblages in layered amphibolites and micaschists from the Dessent Formation (Mountaineer Range, Antarctica), *Mem. Soc. Geol. Ital.*, 43, 87-95.
- Cooper, A. K., F. J. Davey, and K. Hinz (1991), Crustal extension and origin of sedimentary basins beneath the Ross Sea and Ross Ice Shelf, Antarctica, in *Geological evolution of Antarctica*, edited by T. M. R. A., et al., pp. 285-292, Cambridge University Press, New York.
- Dalziel, I. W. D. (1992), Antarctica: A tale of two supercontinents, *Annual Review of Earth and Planetary Sciences*, 20, 501-526.
- Davey, F. J., and G. Brancolini (1995), The Late Mesozoic and Cenozoic structural setting of the Ross Sea region, in *Geology and stratigraphy of the Antarctic margin*, edited, pp. 167-182, American Geophysical Union.
- Davey, F. J., S. C. Cande, and J. M. Stock (2006), Extension in the western Ross Sea region-links between Adare Basin and Victoria Land Basin, *Geophysical Research Letters*, 33, doi:10.1029/2006GL027383.
- Elliot, D. H. (1992), Jurassic magmatism and tectonism associated with Gondwanaland break-up: an Antarctic perspective, in *Magmatism and the causes of continental break-up*, edited by B. C. Storey, et al., pp. 165-184, Geological Society of London, London, United Kingdom.
- Elliot, D. H., and T. H. Fleming (2004), Occurrence and dispersal of magmas in the Jurassic Ferrar large igneous province, Antarctica, *Gondwana Research*, 7, 223-237.
- Fitzgerald, P. G. (2002), Tectonics and landscape evolution of the Antarctic plate since Gondwana breakup, with an emphasis on the West Antarctic rift system and the Transantarctic Mountains, in *Antarctica at the close of a Millennium. Proceedings of the 8th International Symposium on Antarctic Earth Science*, edited by J. A. Gamble, et al., pp. 453-469, Royal Society of New Zealand.

- Fitzgerald, P. G., and S. L. Baldwin (1997), Detachment fault model for the evolution of the Ross Embayment, in *The Antarctic Region: Geological Evolution and Processes*, edited by C. A. Ricci, pp. 555-564, Terra Antarctica Publication, Siena.
- Fitzgerald, P. G., and S. L. Baldwin (2007), Thermochronologic constraints on Jurassic rift flank denudation in the Thiel Mountains, Antarctica, in *Antarctica: A keystone in a Changing World*, edited by A. K. Cooper, et al., USGS Open-File Report 2007.
- Fitzgerald, P. G., M. Sandiford, P. J. Barrett, and A. J. W. Gleadow (1986), Asymmetric extension associated with uplift and subsidence in the Transantarctic Mountains and Ross Embayment, *Earth and Planetary Science Letters*, 81, 67-78.
- Foster, D. A., and D. R. Gray (2000), Evolution and structure of the Lachlan Fold Belt (Orogen) of eastern Australia, *Annual Reviews of Earth and Planetary Science*, 28, 47-80.
- Foster, D. A., D. R. Gray, and C. Spaggiari (2005), Timing of subduction and exhumation along the Cambrian East Gondwana margin, and the formation of Paleozoic backarc basins, *Geological Society of America Bulletin*, 117, 105-116.
- Goode, J. W., P. Myrow, D. Phillips, C. M. Fanning, and I. S. Williams (2004a), Siliclastic record of rapid denudation in response to convergent-margin orogenesis, Ross Orogen, Antarctica, in *Detrital thermochronology—Provenance analysis, exhumation, and landscape evolution of mountain belts*, edited by M. Bernet and C. Spiegel, pp. 105-126, Geological Society of America Special Paper 378, Boulder, Colorado.
- Goode, J. W., I. S. Williams, and P. Myrow (2004b), Provenance of Neoproterozoic and lower Paleozoic siliclastic rocks of the central Ross orogen, Antarctica: Detrital record of rift-, passive-, and active-margin sedimentation, *Geological Society of America Bulletin*, 116, 1253-1279.
- Hamilton, R. J., B. P. Luyendyk, C. C. Sorlien, and L. R. Bartek (2001), Cenozoic tectonics of the Cape Roberts Rift Basin and Transantarctic Mountains Front, Southwestern Ross Sea, Antarctica, *Tectonics*, 20, 325-342.
- Huerta, H., and D. L. Harry (2007), The transition from diffuse to focused extension: Modeled evolution of the West Antarctic rift system, *Earth and Planetary Science Letters*, 255, 133-147.
- Isbell, J. L. (1999), The Kukri Erosion Surface: a reassessment of its relationship to rocks of the beacon Supergroup in the central Transantarctic Mountains, Antarctica, *Antarctic Science*, 11, 228-238.
- Lawver, L. A., and L. M. Gahagan (1994), Constraints on the timing of extension in the Ross Sea region, *Terra Antarctica*, 1, 545-552.
- Mukasa, S. B., and I. W. D. Dalziel (2000), Marie Byrd Land, West Antarctica: Evolution of Gondwana's Pacific margin constrained by zircon U-Pb geochronology and feldspar common-Pb isotopic compositions, *Geological Society of America Bulletin*, 112, 611-627.
- Richard, S. M., C. H. Smith, D. L. Kimbrough, P. G. Fitzgerald, B. P. Luyendyk, and M. O. McWilliams (1994), Cooling history of northern Ford Ranges, Marie Byrd Land, West Antarctica, *Tectonics*, 13, 837-857.
- Siddoway, C. S., S. L. Baldwin, P. G. Fitzgerald, C. M. Fanning, and B. P. Luyendyk (2004), Ross Sea mylonites and the timing of intracontinental extension within the West Antarctic rift system, *Geology*, 32, 57-60.
- Smith, A. G., and D. J. Drewry (1984), Delayed phase change due to hot asthenosphere causes Transantarctic uplift? *Nature*, 309, 536-538.
- Stern, T. A., A. K. Baxter, and P. J. Barrett (2005), Isostatic rebound due to glacial erosion within the Transantarctic Mountains, *Geology*, 33, 221-224.
- Stock, J. M., and S. C. Cande (2002), Tectonic history of Antarctic seafloor in the Australia-New Zealand-South Pacific sector: Implications for Antarctic continental tectonics, in *Antarctica at the close of a Millennium. Proceedings of the 8th International Symposium on Antarctic Earth Science*, edited by J. A. Gamble, et al., pp. 251-259, Royal Society of New Zealand, Wellington, New Zealand.
- Summerfield, M. A. (1991), *Global Geomorphology: An Introduction to the Study of Landforms*, 537 pp., John Wiley, New York.
- Sutherland, R. (1999), Basement geology and tectonic development of the greater New Zealand region: An interpretation from regional magnetic data, *Tectonophysics*, 308, 341-362.
- ten Brink, U. S., R. I. Hackney, S. Bannister, T. A. Stern, and Y. Makovsky (1997), Uplift of the Transantarctic Mountains and the bedrock beneath the East Antarctic ice sheet, *Journal of Geophysical Research*, 102, 27603-27621.
- Tulloch, A. J., and D. L. Kimbrough (1989), The Paparoa metamorphic core complex, Westland-Nelson, New Zealand: Cretaceous extension associated with fragmentation of the Pacific Margin of Gondwana, *Tectonics*, 8, 1217-1234.
- van der Beek, P., S. Cloetingh, and P. Andriessen (1994), Mechanisms of extensional basin formation and vertical motions at rift flanks: Constraints from tectonic modeling and fission track thermochronology, *Earth and Planetary Science Letters*, 121, 417-433.