

(200)  
R29o



136654




no. 38

1949

(200)  
R290  
no. 38

U.S. Geological Survey.

Reports - Open file series.



OCT 4 - 1950



200)  
R290  
o. 38.

U.S. Geological Survey.  
Reports. Open file. No. 38.


UNITED STATES  
DEPARTMENT OF THE INTERIOR  
GEOLOGICAL SURVEY  
WASHINGTON

OPEN FILE RELEASE, September 1949

Geology of the Lippincott lead area,  
Inyo County, California

A preliminary report.

by James F. McAllister 1911-







(200)

R290

USGS LIBRARY - RESTON



3 1818 00083615 3

United States  
✓ Department of the Interior  
U.S. Geological Survey

*[Reports - Open file series]*

Geology of the Lippincott lead area, Inyo County, California

A preliminary report

---

by James F. McAllister

---

In cooperation with the  
State of California  
Division of Mines

1949



	Manuscript page
Introduction.....	3
Sedimentary and metamorphic rocks.....	5
Igneous rocks.....	8
Structure.....	8
Mineral deposits.....	9

## ILLUSTRATIONS

---

- Plate 1. Geologic and topographic map of the Lippincott lead area,  
Inyo County, California.
2. Map and projections of Main workings, Lippincott lead  
mine, Inyo County, California.
  3. Geologic map of Main workings, Lippincott lead mine,  
Inyo County, California.
- Figure 1. Index map showing location of the Lippincott lead area.

Geology of the Lippincott lead area, Inyo County, California  
is inaccessible and probably several hundred feet lower than the mine.  
Water has been hauled near by James F. McAllister  
miles from Scotty's Castle.

The geologic and topographic Introduction  
In recent years the only continuous mining activity in the Ubehebe  
mining district of Inyo County, California, has been at the Lippincott  
lead mine, which formerly was known as the Southern lead mine. The  
property is owned and operated by George Lippincott. It is 19 miles  
N. 59° E. from Keeler and 4 miles south of Ubehebe Peak (fig. 1; see map  
of Ballarat quadrangle, U. S. Geological Survey).  
The mine is somewhat inaccessible. It is 32 miles by an unsurfaced  
road through Racetrack Valley to the paved road that starts at Ubehebe  
Craters in Death Valley. On the paved road it is 85 miles farther to  
Death Valley Junction, or 125 miles to Lone Pine, which is on U. S. High-  
ways 6 and 395. A more direct route to Lone Pine is by a rough, narrow,  
and steep road from Racetrack Valley, through the south end of Saline  
Valley and up to the paved road between Darwin and Keeler. Over the shorter  
route it is advisable to use vehicles that have been well tested on rough  
mountain roads. At times torrential rains have made the more direct road  
impassable, whereas they have merely roughened the road to Death Valley.  
During 1948 and 1949 Mr. Lippincott has kept the roads open and scraped.  
The Racetrack playa, about 2½ miles north of the mine, has been used as  
a landing field. The mining area comprises Paleozoic dolomite and a little  
limestone, shale, and quartzite, which were intruded and granitized and  
surrounded by quartz monzonite. The principal mineral deposits are in the  
dolomite in a zone bordering the intrusive contact.



## Sedimentary and metamorphic rocks

The principal masses of sedimentary rocks are dolomites that range in color from nearly white or yellowish gray to dark gray. As a result of metamorphism of the dolomites near the contact with a stock of quartz monzonite, the colors are for the most part considerably lighter than the colors of the same unaltered formations. Metamorphism also has changed chert nodules and sandy and muddy impurities in the dolomites to such minerals as tremolite, diopside, antigorite, chrysotile, and—nearer the contact—garnet, epidote, idocrase, and scapolite. The dolomite grains have been recrystallized to a variety of sizes ranging from rather fine to moderately coarse. The largest carbonate grains, however, appearing in broad bands at the top and on the southwestern side of the highest ridge, are calcite, either white or colored brown by enclosed iron oxide. The metamorphic silicate minerals tend to be darker or weather darker than the dolomite, making conspicuous bands, which locally appear folded and twisted.

Certain smaller but nevertheless prominent masses of rocks consist of (1) quartzite grading into interstratified quartzite and dolomite; (2) small patches of shale and siltstone somewhat metamorphosed to schistose and hornfelsic rocks; and (3) some limestone.

The oldest unit of rock shown on the map (pl. 1) is a dark-gray dolomite marked da. It forms the eastern and southern flanks of the hill in the northeastern part of the area, and also the upper part of the southern slope of the hill in the northwestern corner. The dolomite contains some nodules of dark chert. In places this has been metamorphosed to tremolite radiating in white clots. The dolomite marked db, lying stratigraphically above the first dolomite, is very light gray and contrasts sharply with the first. The next layer of dolomite, dc, is also light gray but tends to weather in part yellowish or brownish, particularly near the base and near the top, where impurities have been metamorphosed to calc-silicate minerals. This rock unit forms an irregular band in the lower hills diagonally across the area. Stratigraphically above it, the next unit is dd, the very light gray dolomite that forms the prominent ridge above mine workings. A subdivision, ddq, in the bottom of unit dd, comprises white quartzite, quartzitic dolomite, and some interstratified dolomite. The lower boundary of the subdivision is sharp against medium-gray dolomite at the top of unit dc, but the upper boundary is gradational in that thinner and more widely spaced quartzite beds continue in the main part of dolomite dd. The lead deposits are in dd, especially in the lower part where thin beds of quartzite persist in the dolomite.



The rock units from da to dd are in continuous stratigraphic sequence. The remaining units are separated from dd either by faults or intrusive masses of quartz monzonite. Although these units are fragmentary in the mining area, it is known through comparison with the sedimentary sequence in nearby areas that the normal stratigraphic succession above dd is ls, qls, and sh, (pl. 1). Limestone ls, as exposed in patches in the southwestern part of the Lippincott area, is a medium-gray rock that has been bleached from its usual dark gray by metamorphism. Some parts are thinly bedded and contorted. A quartzite and impure limestone unit, qls, has been separated from unit ls by quartz monzonite, but still is associated with the metamorphosed silty shale sh. This association with dark shale on one side and limestone on the other serves to distinguish the quartzite of qls from the quartzitic member ddq at the base of dolomite dd.

The Paleozoic rocks in a few places are covered by patches of old gravel containing fragments as large as boulders, and consisting of a wide variety of rocks found in adjoining areas. Patches lower than the 3900- to 3950-foot contours have slumped down the hillsides or are remnants of re-worked material. The alluvium, al, is finer gravel left by recent intermittent torrents in canyon bottoms and on alluvial fans.

## Igneous rocks

The major intrusive rock is quartz monzonite, which forms a large stock south and west of the Lippincott area. Only the border was mapped to show the zone of contact with the sedimentary rocks. The border zone, especially in the southeastern part of the area, contains considerable aplite and pegmatite. The quartz monzonite consists of nearly equal quantities of orthoclase and plagioclase, less quartz, and a little hornblende. The volume percentage of minerals in a typical specimen of the quartz monzonite from just south of the area was determined by a micrometric analysis to be 41 percent orthoclase, 38 percent plagioclase, 16 percent quartz, 3 percent hornblende, and 2 percent accessory sphene and magnetite. The texture is coarse-grained, and somewhat porphyritic from larger crystals of orthoclase, which locally are parallel. A few small dikes of fine-grained gray mafic rock cut both quartz monzonite and dolomite dd. This mafic rock has been greatly altered and impregnated with fine-grained pyrite, which readily weathers staining the rock a characteristic brown.

## Structure

The major structure of the sedimentary rocks is an overturn of the openly folded upright sequence that occurs in the northern part of the area to the overturned sequence that occurs in the central and southern parts. The zone of overturning is marked by a fold that to the east becomes a fault. The short but conspicuous fold superficially resembles an overturned anticline plunging east, but as the younger beds are in the core, it is actually an inverted overturned syncline. Along the fault, which dips about  $70^{\circ}$  S., the south block has moved west relative to the north block. The horizontal component of movement, indicated by the relative locations of the vertical contact of dolomites da and db, is about 500 feet. Minor folds in the overturned sequence now are inverted anticlines and synclines.



A long fault is nearly parallel to the western margin of the area, and cuts across sedimentary rocks and quartz monzonite. The nature of the displacement is not shown in the map area, but northward beyond the area the west block has moved north relative to the east block. The fault intersects a broad zone of sheared rock which trends southeastward from the northwest corner of the area, and which is well exposed at the main junction of drainage. Displacement along the shear zone is not measurable, and may have been obscured by slippage along bedding planes. Minor faults, which in general trend north or northwest, and which perhaps are related to the shear zone, were the principal control of lead deposits.

#### Mineral deposits

Mineralization, presumably from the quartz monzonite, produced the following types of deposits; lead-bearing veins in the dolomite; irregular masses of copper- and iron-bearing minerals at the intrusive contact; brown-weathering siliceous replacement along fractures in dolomite; broad limonite-stained zones of coarse calcite in dolomite dd; black tourmaline veins in the more pegmatitic and aplitic facies of the quartz monzonite; a few quartz veins (some contain coarse barite) in the quartz monzonite; traces of scheelite in light-colored silicate rock well beyond the garnetiferous contact zones; and small poorly defined zones of talc replacing dolomite da. At present (1949) only the lead deposits are of economic interest.

The lead deposits are in siliceous veins and replacements along minor faults and breccia zones. All the deposits are in dolomite dd, and most are in the stratigraphically lower part that contains a little interbedded quartzite and sandy dolomite. The lead ore shoots are like pods and pipes along veins which pinch to mere stringers. Galena and cerussite are the lead ore minerals, in a little gangue of quartz and chalcedony; other associated minerals have not yet been studied. Both galena and cerussite occur from the surface to the deepest workings about 200 feet below the surface. Within this short vertical range the oxidation of the sulfide ore depended more on the local permeability of the enclosing material than on the nearness to the surface. Samples of the ore await analysis by the Geological Survey but the galena, according to Mr. Lippincott, carries considerable silver.

The largest ore body that has been mined was in the Main workings of the Lippincott mine (1, pl. 1). The pipelike ore shoot from its outcrop near the shaft plunged about  $70^{\circ}$  NW. for at least 200 feet (pl. 3). The diameter may have been as much as 10 feet. In the Addison workings of the Lippincott mine (4, pl. 1), ore was mined from a shoot about 125 feet long, inclined  $40^{\circ}$ . Ore was found also in the Confidence No. 1 and the Confidence No. 2 workings (3 and 2, pl. 1), and in 1949 a good pocket was being mined from the Taylor shaft of the Confidence No. 1.





Pocket contains 4 folded plates







USGS LIBRARY - RESTON



3 1818 00083615 3