



















770	< 0.1		68	NX Swivel barrel			Biotite-quartz-feldspar schist and gneiss. Biotite altered to chlorite, and feldspar partially to completely altered to clay. Chlorite slickensides along foliation and joints. Top 6 ft sheared.
	< 0.1		68				
780	< 0.1		68				
	0.1		68				
790	< 0.1		68				Granite, medium-grained, gneissic, and pegmatite. Granite similar to those above. Biotite altered to chlorite. Chlorite slickensides on joints.
	0.3		68				
800	< 0.2		68				
	< 0.1		68				
	< 0.1		68	NX Rigid barrel			Biotite-quartz-plagioclase gneiss and schist. Biotite altered to chlorite.
810	< 0.2		68				Biotite (chlorite) schist.
	0.6	>	69				Granite, medium-grained, gneissic. Massive with few joints. Biotite altered to chlorite.
820	< 0.2	>	69				Biotite-quartz-plagioclase schist and gneiss. Biotite altered to chlorite. Chlorite slickensides on joints.
830	< 0.1		69				Granite, medium- to coarse-grained gneissic, similar to granites above. Chlorite slickensides on joints. Top 1 ft sheared.
	< 0.2		70				
840	< 0.5	>	70				Biotite-plagioclase gneiss, medium-grained, only faintly foliated. Biotite altered to chlorite, and chlorite slickensides along joints.
	< 0.1		70				
850	< 0.7	>	70				Granite, medium-grained, gneissic. Consists of pink microcline, plagioclase feldspar, quartz, and biotite altered to chlorite. Schlieren of schist in lower part. Chlorite and calcite on joint surfaces.
860	< 0.5	>	70				
							Altered and sheared.
870	< 0.1		70		