

DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY

DESCRIPTION OF MAP UNITS

- Qac** ALLUVIUM AND COLLUVIUM (0-515 ft.)--stream, fan, and terrace gravels; locally caliche cemented.
- Tfer** RHYOLITES OF FORTYMILE CANYON.
- Tfco** Fluidal-flow rhyolite of Comb Peak (50+ ft.)--gray devitrified locally lithophysal flow-banded rhyolite. Only lower part exposed.
- Tfcp** Proclastic rock of Comb Peak (50-75 ft.)--white to pink vitric bedded tuff and massive tuff breccia that grade upward into flow breccia at base of overlying flow rhyolite.
- Tpf** PLATEAU CANYON FORMATION.
- Tuf of Crater Flat** (150+ ft.)--simple(?) cooling unit of pumiceous ash-flow tuff. Top is eroded. Light-gray partly welded devitrified tuff of interior of unit grades downward into white to pink non-welded glassy tuff at base. Phenocrysts (10-25 percent) are mainly quartz and alkali feldspar, with some plagioclase and rare biotite.
- Tpc** TIVA CANYON MEMBER (400+ ft.)--multiple-flow compound cooling unit of ash-flow tuff. Top is gray pumiceous vitric or vapor-phase crystallized, nonwelded to partly welded tuff less than 10 ft thick that grades down to densely welded red and black vitrophire about 5 ft thick. The vitrophire is underlain by red- or purple-brown outwashed devitrified welded tuff 50-100 ft thick that grades downward into a zone of purple-gray lithophysal welded tuff 100-200 ft thick. The lithophysal tuff grades downward into densely welded purple-gray devitrified tuff 75-150 ft thick that is characterized by closely spaced conchoidal fractures subparallel to foliation. Underlying the devitrified tuff is commonly a gray moderately welded crudely columnar-jointed unit 15-30 ft thick showing vapor-phase crystallization. Underlying this is 10-20 ft of brown or orange-brown partly welded vitric shards that grade down to gray or white nonwelded poorly bedded vitric tuff 15-30 ft thick at base of cooling unit. Phenocrysts, mainly alkali feldspar and quartz, base 2-5 percent pumice and about 10 percent phenocrysts. A lithologically distinctive bedded tuff, 5-10 ft thick and too thin to map, everywhere separates the Tiva Canyon and Yucca Mountain Members.
- Tpy** YUCCA MOUNTAIN MEMBER (0-175 ft.)--simple cooling unit of ash-flow tuff. Where less than 50 ft thick, entire cooling unit is non-welded to partly welded pink or gray vitric shard tuff. Where thicker, the center of the cooling unit is purple-brown welded devitrified tuff, and partly welded tuff in the upper part of the unit shows vapor-phase crystallization. The tuff contains 3-10 percent pumice lapilli, rare small lithic inclusions of angular red-brown aphanitic volcanic rock, and 1-3 percent phenocrysts of alkali feldspar with some oligoclase and rare biotite and quartz. The member is thickest in the northeastern part of quadrangle, and wedge out to the southeast is depositional.
- Tpp** Tuff of Pah Canyon (0-150 ft.)--multiple-flow cooling unit of ash-flow tuff. Periphery of cooling unit is pink or white nonwelded vitric tuff. In thick sections the center is pale orange-brown partly welded devitrified tuff. Contains 4-8 percent phenocrysts, mainly alkali feldspar and biotite with some oligoclase and rare quartz, 1-2 percent small (0.1-0.3 in.) but distinctive angular red-brown lithic inclusions of volcanic rock, about 5 percent pumice lapilli. Unit is thickest in northwest part of quadrangle; wedges out depositionally to southeast where it contains abundant large (4-10 in.) blocks of pumice.
- Tot** TOPPAH SPRING MEMBER (700+ ft.)--multiple-flow compound cooling unit of ash-flow tuff. At top is gray to brown poorly bedded nonwelded to partly welded pumiceous tuff 3-25 ft thick, commonly partly silicified or otherwise altered. The partly welded tuff is underlain by red and black densely welded vitrophire 3-10 ft thick. Below the vitrophire is red-brown devitrified welded tuff 75-150 ft thick characterized by distinctive gray-white outwashed pumice; underlying this tuff is purple-gray lithophysal welded tuff 100-250 ft thick with distinctive angular lithic inclusions of red-brown porphyritic volcanic rock near top, next lower zone of brown devitrified densely welded tuff 150-250 ft thick grades to black vitrophire 60-90 ft thick. The vitrophire grades downward, showing decreasing welding, to basal olive-brown vitric poorly bedded nonwelded tuff about 20 ft thick. At top, member contains about 15 percent pumice lapilli and 10-15 percent phenocrysts, mainly alkali feldspar and plagioclase with some biotite; these percentages decrease downward to about 5 percent pumice lapilli and 1-3 percent phenocrysts at base.
- Tpb** Bedded tuffs interlayered with ash-flow tuffs (0-40 ft.)--vitric white to buff ash fall, thin ash flow, and reworked tuffs. Ash-fall and ash-flow tuffs typically poorly to moderately bedded; reworked tuff well bedded, locally crossbedded. Contains 5-30 percent fine to coarse pumice, 1-10 percent phenocrysts, and less than 3 percent lithic fragments of dark aphanitic volcanic rock.
- Ticr** INDIAN TRAIL FORMATION.
- Ticr** Fluidal-flow rhyolite of Calico Hills (20+ ft.)--gray to red-brown, glassy to devitrified rhyolite. Flow layering is well developed and commonly has been locus of spherulitic crystallization. Phenocrysts (5-10 percent) are mainly feldspar and biotite. Only upper part exposed.
- Tico** Rhyolitic tuff breccia of Calico Hills (30+ ft.)--white to yellow tuff, containing large (4-15 in.) blocks of glassy flow rhyolite. Both tuff and rhyolite blocks are extensively zeolitized. Only upper part exposed.
- Tt** UNDIVIDED TUFFS (1,655 ft.)--white, buff, and pink vitric and zeolitic(?) tuff that is massive to well bedded. Bedding and local crossbedding best developed in upper part. Contains 10-30 percent pumice lapilli, 5-10 percent phenocrysts; abundant biotite in some beds. Only upper 50 ft exposed.

PREPARED IN COOPERATION WITH THE
U.S. ATOMIC ENERGY COMMISSION



EXPLANATION

Qac
Alluvium and colluvium

UNCONFORMITY

Tfer
Tfcp
Rhyolites of Forty-mile Canyon
Tfer, fluidal-flow rhyolite of Comb Peak;
Tfcp, proclastic rock of Comb Peak

UNCONFORMITY

Tpf
Tuf of Crater Flat
UNCONFORMITY

Tpc
Tiva Canyon Member

Tpy
Yucca Mountain Member

Tpp
Tuff of Pah Canyon

Tt
Topopah Spring Member

UNCONFORMITY

Ticr
Tico
Indian Trail Formation
Ticr, fluidal-flow rhyolite of Calico Hills;
Tico, tuff breccia of Calico Hills

Tt
Undivided tuffs

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY

UNCONFORMITY



UNCONFORMITY

This map is preliminary and has not been
edited or reviewed for conformity with
U.S. Geological Survey standards and
conventions.

GEOLOGIC MAP OF THE TOPPAH SPRING SW QUADRANGLE, NEVADA

By
R.W. Lipman and Edward J. McKay

Nevada (Topopah Spring SW quad). Geol. 1:24,000, 1964.
copr.



M(200)
R290
no. 64-104
c.1