

2000
R296
no. 67-120

~~DO NOT REMOVE THIS REPORT FROM FILE. 25~~

Geology and mineral deposits of the
San Cristóbal district, Villa Martin
Province, Potosi, Bolivia.

by

Herbert S. Jacobson and others

U. S. Geological Survey.

[Reports. Open file series]

(200)
R290
no. 67-120


GEOLOGY AND MINERAL DEPOSITS OF THE SAN CRISTOBAL DISTRICT,
VILLA MARTIN PROVINCE, POTOSI, BOLIVIA

By

Herbert S. Jacobson
U. S. Geological Survey

and

Carlos Murillo, Lorgio Ruiz, Oscar Tapia, Hugo Zapata,
Hugo Alarcon, Edgar Delgadillo, and Carlos Velasco,
Servicio Geologico de Bolivia



U. S. Geological Survey
OPEN FILE REPORT

This report is preliminary and has not been edited
for conformity with Geological Survey format.

Contents

	Page
Abstract	1
Introduction	2
History of the district	3
Previous investigations	5
Present investigations	5
Acknowledgments	6
Regional ^{geologic} setting	6
Geology	7
Sedimentary rocks	7
Potoco Formation	7
Quehua Formation	8
Dacite tuff.	9
Quaternary deposits.	9
Igneous rocks	10
Intrusive breccia.	10
Andesite porphyry.	10
Dacite porphyry.	10
Structure	11
Alteration.	12
Mineralization.	13
Geologic history	¹² 13

	Page
Mineral deposits	13
Mineralization.	14
Vein deposits	14
Disseminated and replacement deposits	15
Principal mines	15
Animas Mine.	15
Bertha Mine.	17
Colon Mine	18
Hedionda Mine.	18
Inca Mine.	19
Tesorera Mine.	19
Toldos Mine.	19
Trapiche Mine.	21
Geochemical surveys.	21
Sampling and analysis	21
Results and interpretation.	22
Economic potential	23
Recommendations.	23
References cited	25

Illustrations

	Page
Plate 1. Geology of the San Cristóbal district	7
2. Geology of the Toldos Mine.	19
Figure 1. Index Map of Bolivia.	2
2. Geologic Interpretation of the San Cristóbal district	7
3. Geology of the Animas Mine.	15
4. Electromagnetic map of the Animas Mine and vicinity	16
5. Geology of the Bertha Mine.	17
6. Geology of the Hedionda Mine.	18
7. Geochemical Soil Survey of the San Cristóbal district	22

GEOLOGY AND MINERAL DEPOSITS OF THE SAN CRISTÓBAL
DISTRICT, VILLA MARTIN PROVINCE, BOLIVIA

By

Herbert S. Jacobson, U. S. Geological Survey

and

Carlos Murillo, Lorgio Ruiz, Oscar Tapia, Hugo Zapata,

Hugo Alarcon, Edgar Delgadillo, and Carlos Velasco,

Servicio Geologico de Bolivia

Abstract

The San Cristóbal silver district near Uyuni, Bolivia, was studied in detail during the months of August to November 1966 jointly by members of the Servicio Geologico de Bolivia (GeoBol) and the U. S. Geological Survey, under the auspices of the Agency for International Development, U. S. Department of State.

The rocks of the district are all Tertiary or Quaternary. The oldest rocks are the red beds of the Potoco Formation, which are unconformably overlain by agglomerate, conglomerate, sandstone, and tuff of the flat-lying Quehua Formation. Both formations were intruded by dacite porphyry and andesite porphyry stocks and both are overlain by recent dacite porphyry lava flows. The intrusive and sedimentary rocks near the intrusive contacts are hydrothermally altered.

Mineral deposits of the district are associated with the intrusive rocks and are of four distinct types:

- 1) Sulfide veins containing lead, zinc, and silver associated with dacite porphyry intrusives (Colon and Tesorera Mines).

- 2) Oxide veins containing iron and silver associated with andesite porphyry intrusives (Toldos Mine).
- 3) Partial sulfide replacement of intrusive breccia containing silver, lead, and zinc (Animas Mine).
- 4) Disseminated lead-zinc-silver minerals in altered dacite porphyry intrusives and in adjacent altered sedimentary rocks (Hedionda Mine).

Disseminated silver minerals may possibly provide a basis for the development of open-pit silver mines. This possibility should be further investigated, beginning with geochemical surveys and hydrothermal alteration studies.

Introduction

The San Cristóbal District is located in southern Bolivia approximately 500 kilometers (airline) south of La Paz and 65 kilometers southwest ^{east} of Uyuni (fig. 1). The district is readily accessible by road from Uyuni, a trip of three hours by car, except during the rainy season (December-March). A detailed field investigation of the district, performed from August 26, 1966 to November 4, 1966, by the U. S. Geological Survey and the Servicio Geológico de Bolivia (GeoBol) was sponsored by the Agency for International Development Mission to Bolivia, U. S. Department of State.

This report presents the results of the field investigation but does not include the laboratory work which is in progress.

A detailed report summarizing the data obtained will be published in the near future by GeoBol.

(200)
R29

U. S. Geological Survey
OPEN FILE 602P




Figure 1.—Index map of Bolivia showing location of the San Cristobal district.

History of the district

Silver was discovered at the Hedionda Mine (Santa Barbara de Jayula) by the Spanish priest A. A. Barba over three centuries ago (Barba, 1637, book 1, chap. 2). Repeated attempts to mine the silver ended in failure because of the presence of carbon dioxide gas which caused the death of many miners. (The gas continues to issue from the wall rocks of the mine at the present time). Other silver mines in the district were exploited by the Spaniards in the 17th and 18th centuries, but no details are available. Considerable mining activity in this period is indicated by the numerous abandoned mine workings in the district.

In the past century the district has been the scene of intermittent mining activity. The Toldos Mine was operated from 1870 to 1921 by the Compania Minera de San Cristóbal producing ore containing 0.2 to 0.5 percent silver (Ahlfeld 1944, p. 15). Production from 1918 to 1921 averaged about 200 kilos of silver per month. The Hedionda Mine was re-discovered in 1896 by the Polish engineer J. Jackowski, who operated the mine from 1896 to 1901 and from 1927 to 1936, after having found a remedy for the carbon dioxide problem. This was simply the excavation of a canal in the floor of the mine workings which allowed the gas to flow out of the mine just like water. Also, mining was confined to work above the adits.

Partial smelter returns are an indication of the mine production
(J. Jackowski, unpub. data):

<u>Date</u>	<u>Kilos</u>	<u>Percent</u>	
		<u>Ag</u>	<u>Pb</u>
October 21, 1900	905	1.51	--
November 13, 1900	6195	0.76	17.59
September 6, 1901	4544	0.38	38.50
September 6, 1901	279	0.26	--
March 19, 1929	3300	0.70	43.0
April 4, 1929	600	0.53	42.1
May 10, 1929	336	0.53	38.2

No mines were active in the district from 1936 until the De Wett adit at the Hedionda Mine was opened in 1963 by P. Zubrzycki, who was given the right to operate the mine by the heirs of J. Jackowski in 1959. Recently this right has been transferred to the Lipez Mining Company which began operations in the district in 1966. This company now controls 8 properties totalling 627 hectares and is reopening some of the old mines in the district. There has been no recent production (O. B. Stoughton, manager, Lipez Mining Company, personal communication).

In addition to the Lipez Mining Company the Cooperativa Minera Litoral is now active in the district. This organization started mining lead-silver ore at the Animas Mine in December 1965. Current production is 2.3 tons of silver-bearing hand-sorted galena per month. The mine was started as a result of technical assistance to the Cooperativa Minera Litoral by one of the writers (Oscar Tapia), who examined five mineral prospects and selected the site of the Animas Mine.

Previous investigations

During the long history of the district many engineers and geologists have studied the district, but little information has been published. The principal known studies were made by: J. Jackowski, 1896 to 1901 and 1927 to 1936; F. Ahlfeld (Ahlfeld, 1944 and 1964); P. Zubrzycki, 1956 to 1966; O. Tapia, 1965-66; H. Torrico^{1/}, H. R. Cooke, 1966; and L. Ruiz, 1966. The work of Jackowski, Ahlfeld, Zubrzycki and Cooke emphasized the mineral deposits, whereas Tapia, Torrico, and Ruiz were mainly concerned with the regional geology. O. Tapia prepared the San Cristóbal quadrangle of the geologic map of Bolivia, which will be published by GeoBol at a scale of 1:100,000.

Present investigations

The aim of the present investigation was twofold:

1. To make a detailed study of the San Cristóbal district to assist in the development of the mineral deposits in the district.
2. To provide training for GeoBol geologists in modern geological, geochemical and geophysical field methods as a guide to future similar studies of other districts in Bolivia.

The methods employed in the investigation were:

1. Surface geological mapping with plane tables.
2. Underground geological mapping of the accessible mine workings.
3. Reconnaissance geochemical sampling and analysis to determine the limits of the mineral district.
4. Detailed geochemical ~~grid~~ sampling and analysis.
5. Geophysical magnetic and electromagnetic surveys.

^{1/} Torrico, H., 1966. Estudio Geologico de La Region de San Cristobal: Facultad de Ciencias Geologicas, U. S. M. A. Thesis, La Paz, Bolivia.

Acknowledgments

We are grateful for the support provided to the field party by Mr. Gaston Pereira, Director of GeoBol, and the GeoBol staff in La Paz. Many thanks are also due to the following GeoBol personnel who assisted us in the field: Salomon Rivas, Eduardo Mendoza, and Jose Torres. In addition, we are indebted to O. B. Stoughton (manager), R. Gamarra, and other staff members of the Lipez Mining Company for their helpful assistance in San Cristóbal. We also are thankful for the assistance received from members of the Cooperativa Mineral Litoral.

^{geologic} Regional setting

The San Cristóbal district lies within a northerly trending range of mountains surrounded by the high plateau (altiplano) of western Bolivia. The mountain range has a mature topography and maximum relief of about 800 meters. Elevations within the area studied range from 4080 to 4640 meters above mean sea level. The climate is dry, cold, and windy. Vegetation is sparse. The bedrock is relatively well exposed but in many places is covered by talus on the mountain slopes.

The rocks in the mountain range are all Tertiary and possibly Quaternary ⁴. The oldest rocks are Paleocene to Lower Miocene sandstones and shales of the 4500-meter-thick Potoco Formation.

These rocks are unconformably overlain by generally horizontal beds of agglomerate, conglomerate, sandstone, and tuff of the Quehua Formation (Pliocene) which has a thickness of more than 600 meters. This formation is covered by horizontal dacite porphyry lava flows (Pliocene?) ^{1/}, 5 to 30 meters thick, which cap the mountains.

1/ The age of the flows was considered Quaternary by Torrico (1966).

The rocks of the Quehua Formation are cut by dacite porphyry and andesite porphyry intrusive stocks (Pliocene or ^ePl~~is~~istocene?). The composition of dacite porphyry stocks and dacite porphyry lavas is similar and the stocks therefore probably were the feeders for the flows. The relative age of the dacite and andesite porphyry intrusives is unclear.

The valleys in the area are filled with unconsolidated Quaternary alluvium.

Geology

The geology of the San Cristóbal district is complex because a variety of rock types are present, many of which are hydrothermal alteration products. In the field several different rock types were mapped ^{as} ~~in~~ the same lithologic unit ^{owing} ~~due~~ to difficulties in identification. Laboratory studies of rock samples by the Servicio Geológico de Bolivia are now in progress. Rock names used in this report must therefore be regarded as provisional and subject to change.

Sedimentary rocks

Potoco Formation

The Potoco Formation is exposed over a large area (Plate 1, Fig. 2) in the southern half of the San Cristóbal district. A section ~~with an~~ ^{to be} estimated ~~thickness of~~ ^{thick} about 2000 meters is partly exposed, but exposures were not adequate for compilation of a detailed stratigraphic section.

PLEASE REPLACE IN POCKET
IN BACK OF BOUND VOLUME

EXPLANATION

SEDIMENTARY ROCKS

T1

Dacite tuff

T2

Quehua Formation
(Agglomerate, Conglomerate, tuff)

T3

Potoco Formation
Red shale, sandstone, and
conglomerate

IGNEOUS ROCKS

T4s T4l T4b T4l

Dacite porphyry

T4s, dacite porphyry stock;
T4l, dacite porphyry flow;
T4b, dacite porphyry breccia

T6

Intrusive breccia

Ranges from a breccia with mainly
sedimentary fragments to breccia
with mainly igneous fragments of
various compositions. Precise age
relation not determined

T5


Intrusive andesite porphyry

Hydrothermal alteration

U. S. Geological Survey
OPEN FILE MAP

This map is preliminary and has
not been edited or reviewed for
conformity with Geological Survey
standards or nomenclature.

Compiled by H. S. Jacobson, L. Ruiz,
O. Tapia, H. Zapata, H. Alarcon, C. Murillo,
C. Velasco and E. Mendoza



200 0 200 400 600 800 1000 METERS

METERS A
4100





Figure 2. -- Geologic Interpretation of the San Cristobal district, Plan Map and cross section.



(200)
R290
no. 67-120

The Potoco Formation consists mainly of steeply dipping red shale and siltstone. The shales are thin bedded and poorly indurated. Locally, beds of sandstone, quartzite, and conglomerate are present. The sandstone is a medium-grained red sandstone, and the quartzite is a yellow, medium coarse grained rock which was only observed in hydrothermally altered exposures. The quartzite beds are 0.5 to 6 meters thick. The conglomerate consists of angular, thin-bedded shale fragments and boulders from 2 to more than 50 centimeters in diameter, commonly 5-10 centimeters, in a red shale matrix. The conglomerate is probably the result of deformation during deposition of the shale. The age of the Potoco Formation, on the basis of regional studies (Torrico, 1966), is Paleocene to Early Miocene.

Quehua Formation

The Quehua Formation unconformably overlies the Potoco Formation and is horizontal except where it has been disturbed by intrusion. It is exposed in the center of the area mapped (Plate 1, Fig. 2), and the best exposures are just south of the village of San Cristóbal on the west bank of the creek.

The Quehua Formation is composed of a series of agglomerates, conglomerates, and tuff, plus minor amounts of sandstone. The uppermost bed of the Formation is an agglomerate rich in andesite porphyry fragments. These fragments are apparently derived from a source older than the andesite porphyry which intrudes the Quehua Formation.

Below the andesite agglomerate is a section of poorly sorted conglomerate and sandstone containing tuff. The conglomerates contain mostly sedimentary rock fragments, but some igneous fragments are also present. The base of the Quehua Formation in the district is a white tuff bed.

The age of the Quehua Formation is Pliocene because it overlies the Paleocene to Lower Miocene Potoco Formation, and because in adjacent areas it interfingers with an ignimbrite which is 7.5 million years old, according to a radiometric age determination.

Dacite tuff

In the southeastern margin of the area mapped (pl. 1 and fig. 2), a horizontal bed of dacite tuff overlies the Potoco Formation and is probably the youngest rock of the district. It is probably Pleistocene in age. The tuff consists of rock fragments (dacite?) up to one centimeter in diameter in a very fine grained white tuff matrix. Near the base of the tuff, red shale fragments from the underlying Potoco Formation are locally present.

Quaternary deposits

The Quaternary unconsolidated sediments (pl. 1) in the area include alluvium in the creek valleys, talus on some mountain slopes, and soil.

creek valleys, talus on some mountain slopes, and soil.

Igneous rocks

Intrusive breccia

An intrusive breccia is present in one large outcrop and four small outcrops in the area studied. This rock ranges from a breccia of largely sedimentary fragments to a breccia of largely igneous fragments of various compositions. Its precise age relation to other rocks is not yet determined. The Animas mine is in this rock.

Andesite porphyry

Andesite porphyry intrusive stocks crop out in the south-central and southeastern parts of the area mapped (pl. 1 and fig. 2). The rock consists of euhedral to subhedral plagioclase phenocrysts in a green, fine-grained matrix. Andesite porphyry breccia is present locally at the intrusive contacts.

The andesite is probably Pliocene or Pleistocene in age because it intrudes the Quehua Formation.

Dacite porphyry

Dacite porphyry is present in almost identical composition in three forms in the area studied: intrusive stocks, lava flows, and breccia pipes. All these rocks are probably derived from the same magma, the stocks being feeders for the flows. The stocks and the flows are thus contemporaneous, but the breccia pipes are younger because they cut the lava flows. Distribution of the dacite porphyry stocks is shown on plate 1 and figure 2; the flows cap the ridges at the higher elevations along the western, northern, and eastern borders of the district.

The dacite porphyry consists of quartz, plagioclase, and biotite phenocrysts in a fine-grained gray matrix. Quartz content is low and plagioclase content is high. Phenocrysts of other unidentified mafic minerals were also observed.

The stocks are younger than the Quehua Formation, as indicated by observed intrusive contacts. However, contacts between the dacite porphyry and the andesite porphyry intrusives were not found, and their relative ages are somewhat uncertain. The dacite porphyry flows overlie the andesite porphyry stocks, however, and if the dacite porphyry flows and stocks are presumed to be of the same age, then all the dacite porphyry is presumably younger than the andesite porphyry. The age of the dacite porphyry therefore probably is late Pliocene or possibly Pleistocene.

Structure

A regional N. 10° W.-trending anticlinal axis passes through the center of the San Cristóbal district, but is not readily apparent within the area mapped because it is obscured by the intrusive stocks. Only the beds of the Potoco Formation are folded in the anticline. The younger Quehua Formation and the dacite porphyry lava are horizontal and unconformably overlie the Potoco Formation. The beds of both the Potoco Formation and the Quehua Formation have been uplifted by the igneous intrusions. This is best illustrated at the Cuaresma Adit (fig. 6) where the beds dip at 20° in the portal and increase to a maximum dip of 50° near the intrusive contact. The dacite lavas on Mount Jayula (northeast corner of pl. 1) may also have been deformed by an underlying intrusion, though large-scale slumping may also explain the erratic bedding attitudes.

One major fault is inferred in the southwest corner of the area mapped (fig. 2). The fault has changed the strike of the Potoco Formation from N. 35° E. south of the fault to N. 50° E; north of the fault, but the direction of the movement is unknown.

In general there is little evidence for the existence of major faults in the district, but numerous faults with small displacement are present. The faults are best observed in the mine workings. Two fault systems were noted: one strikes N. 30°-40° E., and the other strikes N. 50°-70° W. The faulting was probably produced by regional east-westerly compressive forces. Some of the faults are mineralized.

Geologic history

The geologic history of the San Cristóbal district based on the geologic data obtained may be summarized as follows:

1. Deposition and subsequent folding of the sediments of the Potoco Formation.
2. Erosion of part of the Potoco Formation.
3. Deposition of the conglomerate, agglomerate, sandstone, and tuff of the Quehua Formation.
4. Intrusion of andesite porphyry stocks.
5. First period of hydrothermal activity resulting in rock alteration and deposition of iron oxide-siderite-barite veins containing silver minerals.
6. Intrusion of dacite porphyry stocks and deposition of dacite porphyry lava flows. The intrusives probably were feeders of the lava.
7. Intrusion of breccia pipes, faulting, and deposition of dacite tuff.
8. Second period of hydrothermal activity resulting in rock alteration and deposition of sulfide mineralization (galena-sphalerite-pyrite) with native silver and barite.
9. Erosion of the central valley in the district to create the present topography.

The exact dates of the above events are uncertain but all took place in Tertiary and Quaternary time.

Mineral deposits

Most of the Quehua Formation and part of the Potoco Formation as well as portions of the andesite and dacite porphyry stocks are hydrothermally altered. Clay alteration, silicification, and chloritization were observed megascopically and sericitization was noted microscopically.

Clay alteration commonly was observed in the shales of the Potoco Formation and the tuffs of the Quehua Formation. Intensity of alteration is variable, but locally a pure white clay rock was observed (kaolinite?). The dacite porphyry intrusives locally have clay alteration.

Silicification is common in the agglomerates of the Quehua Formation. In some localities strongly silicified agglomerate overlies tuff which has been largely altered to clay. Silicification is also locally present in the conglomerates of the Quehua Formation and in the dacite porphyry intrusives.

Chloritization was noted only in one area of dacite porphyry intrusive, and sericitization was noted locally in the andesite porphyry intrusives.

In addition, most surface exposures of both the andesite porphyry and dacite porphyry stocks contain coatings of iron oxides, particularly along joint planes. Some manganese oxide may also be present.

Mineralization

Sulfide or oxide mineralization is present in the district in all rock units. Disseminated and vein mineralization was observed in the dacite porphyry stocks and in adjoining sediments of the Quehua Formation. Sulfide minerals are also present as disseminations and partial replacements in the intrusive breccia at the Animas Mine. The sulfide minerals are galena, sphalerite, pyrite, and chalcopyrite. Silver is associated with these minerals as native silver or as argentiferous galena, and barite is the principal gangue mineral.

The oxide deposits consist of hematite veins in the andesite porphyry intrusives and in adjoining sedimentary rocks of the Potoco Formation. The veins contain siderite, barite, and quartz gangue, and silver as native silver and in tetrahedrite and stromeyerite (Ahlfeld, 1944, p. 13).

Vein deposits

Most of the deposits are veins that follow the principal structural trends; they bear approximately N. 35° E. and N. 60° W., and have steep dips. The veins are probably epithermal and are exposed in numerous small pits and adits plus some of the principal mines (Colon, Tesorera, Toldos) of the district. The majority of the veins contain sulfide minerals and barite gangue.

A second vein type is present both in an andesite porphyry intrusive (Toldos Mine) and in adjacent sediments ^{any rocks} (Inca Mine). The mineralization consists primarily of hematite, and barite and siderite gangue. The veins contain relatively small amounts of native silver and copper-silver sulfide minerals but no pyrite, galena, or sphalerite.

Disseminated and replacement deposits

Disseminated native silver, galena, sphalerite, and pyrite are present in hydrothermally altered portions of the dacite porphyry intrusives and in adjacent altered sediments ^{any rocks} (Hedionda Mine). The minerals are extremely fine grained and ^{in any case} often invisible to the naked eye.

A unique mineral deposit in the district is at the Animas Mine where a breccia pipe contains disseminated sulfide mineralization as well as local sulfide replacements of the breccia matrix. The sulfide minerals are argentiferous galena, sphalerite, and pyrite.

Principal mines

Animas Mine

Geology. -- ⁽¹⁹⁶⁶⁾ The Animas Mine is presently the only productive mine in the district. It is located in the southwest corner of the district (pl. 1 and fig. 2) in a mineralized breccia pipe about 40 meters wide and 300 meters long. The long axis of the pipe trends approximately N, 30° W. In the mine the breccia is cut off by a fault on the west and is in contact with fractured and altered siltstone and shale of the Potoco Formation (fig. 3) on the east. Surface exposures indicate that additional breccia should be present a short distance west of the fault in the mine.

(200)
R290
no. 67-120

U. S. Geological Survey
OPEN FILE MAP

This map is preliminary and has
not been edited or reviewed for
conformity with Geological Survey
standards or nomenclature.

EXPLANATION



Winze




Bedding




Fault, showing dip



Contact



Breccia with igneous
and sedimentary rock
fragments; Partly al-
tered and mineralized



Altered shale
and sandstone





Figure 3. -- Plan map of the Animas mine (main adit) Bolivia.

PLEASE REPLACE IN POCKET
IN BACK OF BOUND VOLUME

The breccia consists of angular to subrounded ~~rock~~ fragments of igneous and sedimentary rocks in a fine-grained matrix. The fragments are commonly 3 to 10 centimeters in diameter but occasionally are much larger. The percentage of sedimentary fragments within the breccia increases toward its eastern contact. The breccia is hydrothermally altered, but the alteration products have not been identified.

Mineralization at the surface consists of iron oxides. In the mine, sulfides are present mainly in the breccia matrix. Sulfide minerals are disseminated and locally they entirely replace the matrix in "pockets" commonly ¹⁻² meters in diameter. These pockets constitute the ore currently being mined and consist primarily of argentiferous galena with associated sphalerite and pyrite. The principal disseminated mineral observed was pyrite.

Geochemistry.--The wide-spaced geochemical soil sampling grid over the district (fig. ⁷~~8~~) shows no anomaly over the Animas Mine area.

Geophysics.--A turam electromagnetic survey was made over an area 400 meters by 300 meters above the mine along traverse lines spaced 20 meters apart in an attempt to detect strongly mineralized zones within the breccia. The electromagnetic survey indicated the presence of several weak electromagnetic anomalies (conductors). Conductors A, B, and C (fig. 4) represent a series of en echelon conductive zones parallel to the most prominent faults visible in the mine (fig. 3), and possibly ~~could~~ reflect other mineralized faults. The cause of the other conductors (fig. 4) is indeterminate.

(200)

R290

No. 67-120

U. S. Geological Survey

OPEN FILE MAP

This map is preliminary and has not been edited or reviewed for conformity with Geological Survey standards or nomenclature.




Fig. 4 Electromagnetic map of the Animas Mine


Conclusion.--The Animas Mine area exposes a mineralized breccia containing disseminated sulfide minerals as well as the local pockets of high-grade material now being mined. The amount of mineralized breccia above the level of present mine workings probably exceeds one million tons. Estimated possible reserves are several million tons. These reserves possibly could support an open-pit mine, if thorough testing of the deposit reveals a commercial grade of mineralization.

Bertha Mine

The Bertha Mine, one half kilometer south of the village of San Cristobal (pl. 1 and fig. 2), consists of a tunnel 195 meters in length (fig. 5) entirely within a horizontal bed of conglomerate of Figure 5 near here


the Quehua Formation. In the last 35 meters the tunnel follows a mineralized fault zone striking approximately N. 70° W. and dipping north at 45° to 64°. The fault zone ranges from 1 to 80 centimeters in width and contains lenses of sulfide mineralization as much as 30 centimeters thick. Mineralization consists of pyrite, sphalerite, galena, and an unidentified soft glassy yellow mineral. The conglomerate wallrock near the fault zone is partly silicified and contains disseminated pyrite. Inasmuch as the fault is well exposed in the face (fig. 5) and contacts with the dacite porphyry intrusive are nearby (pl. 1), the grade of mineralization may increase further to the west.

(200)
R290
no 67-
120



PLEASE REPLACE IN POCKET
IN BACK OF BOUND VOLUME

Figure 4.5 -- Plan map of the Bertha mine, Bolivia.



Colon Mine

The Colon Mine is a small mine located on the ridge 1.2 kilometers southwest of San Cristobal (pl. 1). Accessible mine workings total in length 25 meters. The mine has exposed two irregular lenticular veins. One vein strikes north-northwest, dips at a low angle to the northeast, and has a thickness of 10 to 50 centimeters. The other vein strikes northeast, dips 40° NW., and has a thickness of 15 to 20 centimeters. Both veins consist of massive barite with traces of galena. The wall-rock consists of altered Quehua Formation.

Hedionda Mine

The Hedionda Mine is the oldest known in the district (Barba, 1637). It is nearly one kilometer north of San Cristobal (pl. 1), and the main presently accessible mine openings are three adits, named Alkamari, Cuaresma, and DeWett. The largest adit is the Cuaresma (fig. 6). The adit exposes a series of east-striking and south-dipping

Figure 6 near here

sedimentary rocks in contact with a dacite porphyry intrusive. The dip of the sedimentary rocks increases as the contact is approached, and a 6- to 10-meter wide zone at the contact is brecciated. The sedimentary rocks consist of tuff, conglomerate, and agglomerate (Quehua Formation). They, as well as the dacite intrusive, are strongly hydrothermally altered and contain large amounts of white clay (kaolin?). All these rocks also contain disseminated sulfide minerals and native silver some of which is so finely disseminated that it is invisible to the naked eye. Several mineralized faults also are present in the mine (fig. 6). Mineralization consists of pyrite, galena, sphalerite, and native silver. The mine workings have recently been channel sampled in detail by the Lipez Mining Company, and preliminary results indicate the presence of disseminated silver, lead, and zinc. According to Mr. O. B. Stoughton, manager (oral commun., 1966), if results continue similar to those obtained to date, an orebody amenable to open-pit or related mining methods may be developed.




Figure 6. -- Plan map of the Cuaresma adit, Hedionda mine, Bolivia.

Mapped by H. Zapata, 1966

Inca Mine

The Inca Mine is located 1.5 kilometers southeast of San Cristóbal (pl. 1 and fig. 2) and may be regarded as an extension of the Toldos Mine. The mine is not readily accessible and was not mapped, but surface pits (pl. 1) show that the mine exploited several veins striking north to N. 30° W, dipping steeply northeast, and having an average thickness of 10 to 40 centimeters. The principal vein minerals are hematite and barite. Traces of other minerals present have not yet been identified, but silver is reportedly present. The wallrock is a sequence of hydrothermally altered sedimentary rocks (Potoco Formation).

Tesorera Mine

The Tesorera Mine, located 700 meters southwest of San Cristóbal, exploited a mineralized fault zone which strikes N. 55° E. and dips 85° SE. in the mine. At the surface (~~pl. 1~~) both walls of the fault are hydrothermally altered agglomerate, whereas in the mine the agglomerate forms the north wall of the fault and a dacite porphyry intrusive forms the south wall. Both the agglomerate and the dacite are hydrothermally altered. Most of the alteration ^{products are} is clay alteration (kaolin?), but silicification was also observed. The mineralized fault zone contains fault breccia approximately one half meter wide, ^{and} with pyrite and galena ~~mineralization~~. The dacite contains disseminated pyrite.

Toldos Mine

The Toldos Mine, located in the southeast corner of the district (pl. 1 and fig. 2), is the most extensively developed mine in the area. Most of the development is on the main level (pl. 2), but there are several levels above the main level and one below. The mine is now flooded below the main level.

Geology.--The mine entrance (pl. 2) exposes a series of altered shale and sandstone with local conglomerate of the Potoco Formation. Similar rocks crop out on the surface above the mine (pl. 1), but most of the mine workings (pl. 2) are in a green andesite porphyry intrusive. The intrusive is cut by a series of mineralized fault zones and veins striking N, 30-40° E, and dipping steeply northwest. The andesite porphyry wallrock is locally altered adjacent to the faults and veins. The wider veins and faults having a width of 0.5 to 1.0 meter were exploited for their silver content. Many veinlets and veins from one to 30 centimeters thick remain in the mine. The mineralization consists primarily of hematite, barite, and siderite. Quartz and magnetite are present locally, and the principal ore minerals reportedly are native silver and stromeyerite (Ahlfeld, 1964, p. 304).

Geophysics.--A magnetometer survey was conducted on the surface over the mine area along ^{northwest-trending} traverse lines 50 meters apart, ~~trending northwesterly,~~ perpendicular to the general strike of the veins and faults. A magnetometer traverse was also made in one section of the mine. The underground traverse detected magnetic anomalies of 300 to 800 gammas corresponding to some of the veins, whereas other veins had no associated magnetic anomaly. On the surface the magnetometer results outlined some weak linear northeast-trending magnetic anomalies with a magnetic relief of about 100 gammas. These anomalies are obscured by a magnetic gradient of 400 gammas over a distance of about 400 meters parallel to the strike of the anomalies. This gradient is probably related to the northwesterly contact between the Potoco Formation ~~sediments~~ and the andesite porphyry intrusive (pl. 2).

The two cross-cutting magnetic trends hamper magnetic interpretation and correlation with geology.

Conclusions.--The mineralized faults and veins in the mine are extensive but have been largely mined out above the main level. They no doubt continue below this level, where little mining was done in the past because of difficulties with water and carbon dioxide gas. Modern mining technology can overcome these difficulties if the silver content of the veins is sufficient to support mining. (The Lipez Mining Company is (1966) now sampling the veins and planning to unwater the flooded portion of the mine).

Trapiche Mine

The Trapiche Mine at the south end of San Cristóbal (pl. 1 and fig. 2) consists of 305 meters of drifts and crosscuts. It explored the area beneath some previously worked veins west of the mine entrance. No veins are cut by the mine workings, but several fault structures which are the probable roots of the veins above are exposed in the mine. This indicates that the mineralization in the San Cristóbal district is epithermal.

Geochemical surveys

Sampling and analysis

Two geochemical surveys were completed. The first was a reconnaissance soil sampling survey along the creeks. The second survey involved soil sampling on a 200-meter-square grid over an area exceeding 6 square kilometers. All samples were taken at a depth of 5 to 10 centimeters and were screened in the field to approximately -80 mesh with a nylon cloth screen.

The samples were all analyzed in a field laboratory for their citrate-extractable heavy metal (copper, lead, and zinc) content by standard techniques, dithizone being the test reagent (Ward and others, 1963, pp. 27-29).

Results and interpretation

The reconnaissance survey showed the limits of the mineralized district, and the grid sampling subsequently outlined the mineralized area in greater detail (fig. 7). The geochemical anomalies shown in fig. 6 extend about 1.5 kilometers south of the southern edge of the map. This extension anomaly is interpreted as representing drainage from the mineralized district and was therefore not outlined in detail.

The grid sampling indicated that the geochemical background in the district is less than 2 parts per million (ppm) combined citrate-extractable copper, lead, and zinc. The geochemical anomalies containing more than 20 ppm metal, or more than ten times the geochemical background, are therefore strong anomalies. Correlation of the anomalies (fig. 7) with the geology (pl. 1 and fig. 2) shows two significant features:

1. Anomalies B, C, D have an east-west trend which may be correlated with sulfide veins having a similar strike.
2. Anomalies F, G, H, ^{and} J1 are apparently related to the andesite porphyry intrusives.

The relationship of most of the other anomalies to the geology is not clear because of the wide sample spacing. Anomaly J2 is primarily due to contamination from mine dumps of the Toldos Mine.

(200)
R 290
no. 67-120

EXPLANATION


4 Soil sample site and citrate extractable metal content Cu + Pb + Zn in parts per million (ppm)

— 10 ppm contour

— 20 ppm contour

A - Geochemical anomaly referred to in text

200 100 0 200 400 600 800 1000 METERS



U. S. Geological Survey
OPEN FILE MAP
This map is preliminary and has not been edited or reviewed for conformity with Geological Survey standards or nomenclature.

Compiled by H. S. Jacobson, 1966

Note: Reconnaissance survey indicates that anomalous values continue 1-1/2 kilometers south of south edge of this map. (These anomalies are probably due to drainage)

Figure 7. -- Geochemical Soil Survey of the San Cristobal district.

U. S. GEOLOGICAL SURVEY
LIBRARY
SEP 11 1967
DENVER

PLEASE REPLACE IN POCKET
IN BACK OF BOUND VOLUME

Economic potential

The San Cristóbal district is mainly a silver district whose principal future economic potential lies in its possible resources of disseminated silver minerals, particularly in view of the current favorable silver market and of the possible utilization of low-cost open-pit mining methods. Disseminated native silver probably is present in several localities within the district in addition to the known deposit at the Hedionda Mine. The detection of such mineralization will require careful sampling and analysis of altered rocks throughout the district.

Underground mining of the veins has a relatively unfavorable economic potential because: (1) Most of the veins near the surface have already been mined, (2) the veins are epithermal and therefore are unlikely to have significant extensions in depth, and (3) underground mining in the district is hampered by the presence of water and carbon dioxide gas.

Recommendations

We recommend that mineral exploration of the district be continued in two stages:

- 1) A study of rock alteration combined with a geochemical survey

of all rock outcrops of the district except those of the dacite porphyry lavas (fig. 2). The survey would consist of sampling, at 25- to 50-meter intervals, of all the outcrops within the area indicated, followed by petrographic study of selected samples to identify the altered minerals and by geochemical analysis of all the samples. Geochemical analysis for silver, lead, and zinc is suggested. Analysis for silver can be made using the method described by H. H. Nakagawa and H. W. Lakin (Nakagawa and Lakin, 1965), and analysis for lead and zinc can be made with standard methods (Ward, and others, 1963).

- 2) Diamond drilling of favorable areas as determined by the above investigation and the geological data described in this report.

In addition we recommend that the following work be done at the principal mines:*

Animas Mine

1. Detailed channel sampling of the mine workings and analysis for silver, lead, and zinc.
2. Diamond drilling of two vertical holes 100 meters deep to test the grade and continuity of mineralization of the mineralized breccia. The holes should be collared in the breccia outcrops above the mine (pl. 1).
3. Pitting or mining to explore electromagnetic anomalies (conductors) A, B, and C.

Bertha Mine

Drifting ahead from the present face along the mineralized fault exposed in the face to explore for ore lenses in the fault zone.

Tesorera Mine

Detailed underground channel sampling to determine the grade of disseminated mineralization because the geology of the mine is similar to that of the Hedionda Mine.

* This does not include work already in progress being conducted by the Lipez Mining Company at the Hedionda and Toldos Mines.

References cited

- Ahlfeld, F., 1944, La Geologia del Mineral de San Cristobal de Lipez (Bolivia): Minería Boliviana: v. 1, no. 8, p. 9-15.
- Ahlfeld, F. and Schneider-Sherbina, A., 1964, Los Yacimientos Minerales y de Hidrocarburos de Bolivia: Dept. Nacional de Geol., Bull. 5, La Paz, Bolivia, p. 304-305.
- Barba, A. A., 1637, Arte de los Metales, Madrid.
- Nakagawa, H. J., and Lakin, H. W., 1965, A field method for the determination of silver in soils and rocks: In U. S. Geol. Survey Prof. Paper. 525-C, p. C172-C175.
- Ward, F. N., Lakin, H. W., Canney, F. C. and others, 1963. Analytical methods used in geochemical exploration by the U. S. Geological Survey: U. S. Geol. Survey Bull. 1152, 100 p.

POCKET CONTAINS
2 ITEMS.

Pocket contains 2 Items



3 1818 00082790 5