



U. S. Geological Survey.

[REPORTS-OPEN FILE SERIES], no. 1493: 1970.



(200)
R29c
no. 1493



(200)
R290
no. 1493



UNITED STATES
DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY
[Report in Open File Series]
Saudi Arabia Investigation Report
(IR) SA-3

GEOLOGICAL RECONNAISSANCE OF THE
WESTERN PART OF THE
WADI AR RIMAH QUADRANGLE,
SAUDI ARABIA

✓
cm
W.S.

by
James W. Mytton
U. S. Geological Survey

224611



U. S. Geological Survey
OPEN FILE REPORT
This report is preliminary and has
not been edited or reviewed for
conformity with Geological Survey
standards or nomenclature.

90
493

✓
U.S. GEOLOGICAL SURVEY
(WASHINGTON, D.C.)

20242
[Reports - Open file series]

For release NOVEMBER 6, 1970

The U.S. Geological Survey is releasing in open files the following reports. Copies are available for inspection in the Geological Survey Libraries, 1033 GSA Bldg., Washington, D. C. 20242; Bldg. 25, Federal Center, Denver, Colo. 80225; and 345 Middlefield Rd., Menlo Park, Calif. 94025:

1. Schmidt hammer test method for field determination of physical properties of zeolitized tuff, by John R. Ege, Danny R. Miller, and Walter Danilchik. 37 p., including 19 figs., 1 table. (*)
2. A geochemical evaluation of the Ash Sha'ib mineral prospect, Asir quadrangle, Kingdom of Saudi Arabia, by Glenn H. Allcott. 38 p., 6 pl., 3 figs., 6 tables.
3. Preliminary geophysical investigation of the Esh Qay'ib (Esh Sha'ib) mine area, Saudi Arabia, by Willard E. Davis and Hisham Kazzaz. 10 p., 6 figs.
4. Recommended geophysical investigations of mineral occurrences at Jabal Samran, Al Marasia, and Wutaileh, by D. H. Hase, V. J. Flanigan, and M. N. Akhrass. 5 p., 1 map, scale 1:500,000.
5. Qualitative analysis of airborne magnetometer data: The Arabian Shield, Kingdom of Saudi Arabia, by Donald H. Hase. 44 p., 1 pl.
6. Preliminary investigation of cement materials in the Taif area, Saudi Arabia, by Conrad Martin. 18 p., 1 fig., 1 table.
7. An appraisal of the Jabal Idsas magnetite mineralization based on diamond drill hole N40A, by Conrad Martin. 29 p., 4 figs., 1 table.
- ✓ 8. Geological reconnaissance of the western part of the Wadi Ar Rimah quadrangle, Saudi Arabia, by James W. Mytton. 3 p.
9. Reconnaissance ^{for} mineral deposits in the Precambrian rocks of the Wadi Ar Rimah quadrangle, Kingdom of Saudi Arabia, by James W. Mytton. 76 p., 1 pl., 3 figs., 3 tables.
10. Preliminary report on the Dawadami district, Saudi Arabia, by Paul K. Theobald, Jr. 4 p.

(*) Note: Item 1, above, is also available for inspection at 504 Custom House, San Francisco, Calif. 94111; 7638 Federal Bldg., Los Angeles, Calif. 90012; 1012 Federal Bldg., Denver, Colo. 80202; 8102 Federal Office Bldg., Salt Lake City, Utah; and in the Library, Mackay School of Mines, University of Nevada, Reno, Nev. 89507.

**GEOLOGICAL RECONNAISSANCE OF THE
WESTERN PART OF THE
WADI AR RIMAH QUADRANGLE,
SAUDI ARABIA**

by

James W. Mytton

PREFACE

In 1963, in response to a request from the Ministry of Petroleum and Mineral Resources, the Saudi Arabian Government and the U. S. Geological Survey, U. S. Department of the Interior, with the approval of the U. S. Department of State, undertook a joint and cooperative effort to map and evaluate the mineral potential of central and western Saudi Arabia. The results of this program are being released in USGS open files in the United States and are also available in the Library of the Ministry of Petroleum and Mineral Resources. Also on open file in that office is a large amount of material, in the form of unpublished manuscripts, maps, field notes, drill logs, annotated aerial photographs, etc., that has resulted from other previous geologic work by Saudi Arabian government agencies. The Government of Saudi Arabia makes this information available to interested persons, and has set up a liberal mining code which is included in "Mineral Resources of Saudi Arabia, a Guide for Investment and Development," published in 1965 as Bulletin 1 of the Ministry of Petroleum and Mineral Resources, Directorate General of Mineral Resources, Jiddah, Saudi Arabia.

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

GEOLOGICAL RECONNAISSANCE OF
THE WESTERN PART OF THE
WADI AR RIMAH QUADRANGLE,
SAUDI ARABIA

by

James W. Mytton

During the month of March the area bounded by 24°45' and 25°99'N. latitude and 42°00' and 42°30'E. longitude was mapped by Abdullah Ankary, Directorate General for Mineral Resources, who located and examined all the ancient mines and did geochemical sampling of the wadis. A brief visit was made to the area by J. W. Mytton during which time the major geological units were checked and reconnaissance trips were made as far north as Jabal ash Shabbah and as far south as Hadbat 'Akliyah.

In the area of Ankary's investigation, most of the attention was given to the regions around Jabals Ashumta and al Koom. Graywacke and associated conglomerate, siltstone and argillite are the dominant rocks at Jabal Ashumta and vicinity and the unit as a whole is similar to that north of Wadi Ar Rimah. At Jabal Ashumta, 24°51'N. x 42°19'E. where extensive workings are present, the graywacke has been converted to hornfels by the emplacement of gray biotite granite of the same character as that present in the northern part of the Wadi Ar Rimah quadrangle. Conglomerate is interbedded with the graywacke and is part of a cyclic sequence of graded bedding consisting of conglomerate beds about 4 inches in thickness overlain by coarse graywacke grading upwards into finer-sized wacke. Southeast of the Jabal, the graywacke and conglomerate give way to dirty siltstones. Northward, the wackes are associated with argillaceous shales and a major conglomerate unit which is quite extensive.

At Jabal Ablan, north of Jabal al Koom, the graywacke sequence includes basal greenish siltstone, graywacke and boulder conglomerate composed of diorite and monzonite derived from the complex of diorite and andesite which the sequence unconformably overlies. This basal unit is overlain by a succession of fine-grained siltstones, in turn overlain by a sequence of green argillites with intervening conglomerates containing pebbles and cobbles. The argillites grade upward into a mixture of graywacke and argillite along with relatively thick units of conglomerate.

The complex of diorite and andesite of Jabal al Koom and vicinity consists of a variety of rocks including fine-grained mafic diorite and andesite which are intruded by a series of monzonites, monzonite porphyries, latites, and adamellites. All these rocks are incorporated into the boulder conglomerate of Jabal Ablan and are therefore part of an epoch pre-dating the deposition of the graywacke sequence. Both the diorite-andesite complex and the graywacke unit are intruded by gray biotite granite. In the al Koom area a circular body of this granite contains xenoliths of diorite similar to that of the surrounding hills. The al Koom workings, 24°49'N.x 42°04'E., located to the southwest of this body, are in quartz veins in fine-grained mafic diorite. Some of the monzonites between al Koom and Jabal Ablan have slightly mineralized veins of quartz and calcite with traces of malachite and pyrite. Tungsten-bearing quartz was discovered by Abdulla Ankary in old mine workings, 24°50'N.x 42°07'E, located near the southeast end of the circular body. The composition of the material sampled as well as its mode of occurrence is discussed in a report by Ankary (unpub. data).

Numerous quartz veins were examined in the graywacke unit south of the village of Al Ba'jah close to its contact with the gray granite. All of these contained white to gray barren quartz and some silicified breccia occupying fractures parallel to the bedding. The diorite-andesite complex continues southward occurring as isolated segments in the granite terrain. Two mines were examined near the contact of the diorite and the gray granite namely Shobrom, 24°45'N.x 42°35'E., located northwest of Hadbat 'Akliyah

and Furdaykhinyah, 24°34'N.x 42°16'E. The workings at both localities are completely buried. At Shobrom the vein material observed on the dumps is fractured quartz containing specular hematite, pyrite, and a flake or two of gold. The workings are in epidotized granite which appears to have intruded diorite. Other than the Shobrom workings, no mines were found in the Hadbad 'Akliyah area. At Furdaykhinyah the vein material is fractured milky quartz which occurs in reddish felsites or rhyolites that have intruded mafic diorites and andesites. One piece of quartz was coated with malachite and contained pyrite in the fractures. Other quartz examined was barren with the exception of hematite. Grinding stones of granite and diabase were found at the site.

A very extensive linear body of bluish-white finely crystalline limestone or marble trending N. occurs just east of Wadi al Jarir in the south-west corner of the Wadi Ar Rimah quadrangle. It is in contact with dark siltstone and graywackes that have been intruded by sills of fine-grained diabase and andesite porphyry.

USGS LIBRARY-RESTON



3 1818 00077204 4