



A layer of windblown sand and silt containing sparse scattered ventifacts and generally mixed with underlying glacial deposits is present over much of the area but has not been mapped.




Alluvium

Stream deposits of light-grayish brown silt, sand and gravel along the Scantic River and smaller streams. Generally less than 20 feet thick: Qf; alluvial fan deposits; coarse grained fan-shaped poorly sorted and rudely stratified stream deposits. As much as 15 feet thick.




Stream terrace deposits

Yellowish-brown well-bedded sand, silt, and clay; locally gravelly. Highest terraces 50 feet above Scantic River; numerous small nonpaired terraces at lower levels on both sides of rivers and streams generally included in alluvium. Some deposits as much as 50 feet thick.




Outwash deposits

Proglacial kettled to unkettled outwash grading from coarse gravel to fine sand and silt downstream. Qsd₁ deposited when ice margin stood along line from Goat Rock to Somers. Qsd₂ deposited when ice margin stood along line trending WSW from Goat Rock. Qsd₃ deposited when ice margin was along an arcuate line trending SW from Mile Oak Corner into Long Meadow and Springfield. Qsd₄ deposited when ice margin stood along a line trending SW from Wilbraham to Springfield. Qsd₅ deposited when ice margin was north of quadrangle.




Kame terrace deposits

Coarse sand and gravel deposited between ice and valley walls by meltwater at higher altitudes than outwash deposits. Internal structure indicates collapse due to deposition on or against ice. Color ranges from reddish brown in western half of quadrangle to gray in east half of quadrangle. Includes several levels of kame terrace deposits. Locally as much as 100 feet thick but generally less than 50 feet thick.




Till

Compact to loose, non-sorted to poorly-sorted, nonstratified mixture of clay, silt, sand, pebbles, cobble, and boulder sized particles. Generally reddish-brown to pale reddish-brown in west half of quadrangle; gray to light gray in east half of area. Includes small lenses of stratified drift. Thickness varies but is generally less than 20 feet. May be more than 100 feet thick under crests of some drumlins. Deposited by ice moving southward in the lowlands and southeast in the highlands in east half of quadrangle. Till in highlands slightly older than that in lowlands in west half of quadrangle.




Bedrock

Individual outcrops shown by solid color; thinly veneered bedrock and scattered small outcrops shown by ruled pattern.



Artificial fill

Artificially emplaced earth, mainly sand and gravel, locally some till. Trash fills not mapped. Most fills less than 20 feet thick but locally as much as 40 feet thick.




Crest of ice-channel filling (eskers and crevasse fillings); angle of chevron points in direction of inferred flow.



Contact

Dashed where approximately located.



Glacial striations

Arrow shows inferred direction of movement of glacier; point of observation at tip of arrowhead



Strandline of inferred glacial lake




Abandoned quarry



Sand and gravel pit




Abandoned sand and gravel pit



Facewall of large sand and gravel pit




Spillway or large ice-marginal channel; arrow shows inferred direction of meltwater flow




Long axis of streamline molded form

Streamline hill (drumlin or drumlinoid) composed partly or entirely of till; smoothed and shaped by movement of glacier.




Swamp deposits

Grayish-brown to black peat, muck, sand, silt, and clay. Mineral matter accumulates by a combination of colluvial, alluvial, and eolian processes. Generally 5 to 10 feet thick but may be as much as 25 feet thick.




Stream terrace deposits

Yellowish-brown well-bedded sand, silt, and clay; locally gravelly. Highest terraces 50 feet above Scantic River; numerous small nonpaired terraces at lower levels on both sides of rivers and streams generally included in alluvium. Some deposits as much as 50 feet thick.




Outwash deposits

Proglacial kettled to unkettled outwash grading from coarse gravel to fine sand and silt downstream. Qsd₁ deposited when ice margin stood along line from Goat Rock to Somers. Qsd₂ deposited when ice margin stood along line trending WSW from Goat Rock. Qsd₃ deposited when ice margin was along an arcuate line trending SW from Mile Oak Corner into Long Meadow and Springfield. Qsd₄ deposited when ice margin stood along a line trending SW from Wilbraham to Springfield. Qsd₅ deposited when ice margin was north of quadrangle.




Kame terrace deposits

Coarse sand and gravel deposited between ice and valley walls by meltwater at higher altitudes than outwash deposits. Internal structure indicates collapse due to deposition on or against ice. Color ranges from reddish brown in western half of quadrangle to gray in east half of quadrangle. Includes several levels of kame terrace deposits. Locally as much as 100 feet thick but generally less than 50 feet thick.




Till

Compact to loose, non-sorted to poorly-sorted, nonstratified mixture of clay, silt, sand, pebbles, cobble, and boulder sized particles. Generally reddish-brown to pale reddish-brown in west half of quadrangle; gray to light gray in east half of area. Includes small lenses of stratified drift. Thickness varies but is generally less than 20 feet. May be more than 100 feet thick under crests of some drumlins. Deposited by ice moving southward in the lowlands and southeast in the highlands in east half of quadrangle. Till in highlands slightly older than that in lowlands in west half of quadrangle.




Bedrock

Individual outcrops shown by solid color; thinly veneered bedrock and scattered small outcrops shown by ruled pattern.



Artificial fill

Artificially emplaced earth, mainly sand and gravel, locally some till. Trash fills not mapped. Most fills less than 20 feet thick but locally as much as 40 feet thick.




Crest of ice-channel filling (eskers and crevasse fillings); angle of chevron points in direction of inferred flow.



Contact

Dashed where approximately located.



Glacial striations

Arrow shows inferred direction of movement of glacier; point of observation at tip of arrowhead



Strandline of inferred glacial lake




Abandoned quarry



Sand and gravel pit




Abandoned sand and gravel pit



Facewall of large sand and gravel pit




Spillway or large ice-marginal channel; arrow shows inferred direction of meltwater flow




Long axis of streamline molded form

Streamline hill (drumlin or drumlinoid) composed partly or entirely of till; smoothed and shaped by movement of glacier.




Swamp deposits

Grayish-brown to black peat, muck, sand, silt, and clay. Mineral matter accumulates by a combination of colluvial, alluvial, and eolian processes. Generally 5 to 10 feet thick but may be as much as 25 feet thick.




Stream terrace deposits

Yellowish-brown well-bedded sand, silt, and clay; locally gravelly. Highest terraces 50 feet above Scantic River; numerous small nonpaired terraces at lower levels on both sides of rivers and streams generally included in alluvium. Some deposits as much as 50 feet thick.




Outwash deposits

Proglacial kettled to unkettled outwash grading from coarse gravel to fine sand and silt downstream. Qsd₁ deposited when ice margin stood along line from Goat Rock to Somers. Qsd₂ deposited when ice margin stood along line trending WSW from Goat Rock. Qsd₃ deposited when ice margin was along an arcuate line trending SW from Mile Oak Corner into Long Meadow and Springfield. Qsd₄ deposited when ice margin stood along a line trending SW from Wilbraham to Springfield. Qsd₅ deposited when ice margin was north of quadrangle.




Kame terrace deposits

Coarse sand and gravel deposited between ice and valley walls by meltwater at higher altitudes than outwash deposits. Internal structure indicates collapse due to deposition on or against ice. Color ranges from reddish brown in western half of quadrangle to gray in east half of quadrangle. Includes several levels of kame terrace deposits. Locally as much as 100 feet thick but generally less than 50 feet thick.




Till

Compact to loose, non-sorted to poorly-sorted, nonstratified mixture of clay, silt, sand, pebbles, cobble, and boulder sized particles. Generally reddish-brown to pale reddish-brown in west half of quadrangle; gray to light gray in east half of area. Includes small lenses of stratified drift. Thickness varies but is generally less than 20 feet. May be more than 100 feet thick under crests of some drumlins. Deposited by ice moving southward in the lowlands and southeast in the highlands in east half of quadrangle. Till in highlands slightly older than that in lowlands in west half of quadrangle.




Bedrock

Individual outcrops shown by solid color; thinly veneered bedrock and scattered small outcrops shown by ruled pattern.



Artificial fill

Artificially emplaced earth, mainly sand and gravel, locally some till. Trash fills not mapped. Most fills less than 20 feet thick but locally as much as 40 feet thick.




Crest of ice-channel filling (eskers and crevasse fillings); angle of chevron points in direction of inferred flow.



Contact

Dashed where approximately located.



Glacial striations

Arrow shows inferred direction of movement of glacier; point of observation at tip of arrowhead



Strandline of inferred glacial lake




Abandoned quarry



Sand and gravel pit




Abandoned sand and gravel pit



Facewall of large sand and gravel pit




Spillway or large ice-marginal channel; arrow shows inferred direction of meltwater flow




Long axis of streamline molded form

Streamline hill (drumlin or drumlinoid) composed partly or entirely of till; smoothed and shaped by movement of glacier.




Swamp deposits

Grayish-brown to black peat, muck, sand, silt, and clay. Mineral matter accumulates by a combination of colluvial, alluvial, and eolian processes. Generally 5 to 10 feet thick but may be as much as 25 feet thick.




Stream terrace deposits

Yellowish-brown well-bedded sand, silt, and clay; locally gravelly. Highest terraces 50 feet above Scantic River; numerous small nonpaired terraces at lower levels on both sides of rivers and streams generally included in alluvium. Some deposits as much as 50 feet thick.




Outwash deposits

Proglacial kettled to unkettled outwash grading from coarse gravel to fine sand and silt downstream. Qsd₁ deposited when ice margin stood along line from Goat Rock to Somers. Qsd₂ deposited when ice margin stood along line trending WSW from Goat Rock. Qsd₃ deposited when ice margin was along an arcuate line trending SW from Mile Oak Corner into Long Meadow and Springfield. Qsd₄ deposited when ice margin stood along a line trending SW from Wilbraham to Springfield. Qsd₅ deposited when ice margin was north of quadrangle.




Kame terrace deposits

Coarse sand and gravel deposited between ice and valley walls by meltwater at higher altitudes than outwash deposits. Internal structure indicates collapse due to deposition on or against ice. Color ranges from reddish brown in western half of quadrangle to gray in east half of quadrangle. Includes several levels of kame terrace deposits. Locally as much as 100 feet thick but generally less than 50 feet thick.




Till

Compact to loose, non-sorted to poorly-sorted, nonstratified mixture of clay, silt, sand, pebbles, cobble, and boulder sized particles. Generally reddish-brown to pale reddish-brown in west half of quadrangle; gray to light gray in east half of area. Includes small lenses of stratified drift. Thickness varies but is generally less than 20 feet. May be more than 100 feet thick under crests of some drumlins. Deposited by ice moving southward in the lowlands and southeast in the highlands in east half of quadrangle. Till in highlands slightly older than that in lowlands in west half of quadrangle.




Bedrock

Individual outcrops shown by solid color; thinly veneered bedrock and scattered small outcrops shown by ruled pattern.



Artificial fill

Artificially emplaced earth, mainly sand and gravel, locally some till. Trash fills not mapped. Most fills less than 20 feet thick but locally as much as 40 feet thick.




Crest of ice-channel filling (eskers and crevasse fillings); angle of chevron points in direction of inferred flow.



Contact

Dashed where approximately located.



Glacial striations

Arrow shows inferred direction of movement of glacier; point of observation at tip of arrowhead



Strandline of inferred glacial lake




Abandoned quarry



Sand and gravel pit




Abandoned sand and gravel pit



Facewall of large sand and gravel pit




Spillway or large ice-marginal channel; arrow shows inferred direction of meltwater flow




Long axis of streamline molded form

Streamline hill (drumlin or drumlinoid) composed partly or entirely of till; smoothed and shaped by movement of glacier.




Swamp deposits

Grayish-brown to black peat, muck, sand, silt, and clay. Mineral matter accumulates by a combination of colluvial, alluvial, and eolian processes. Generally 5 to 10 feet thick but may be as much as 25 feet thick.




Stream terrace deposits

Yellowish-brown well-bedded sand, silt, and clay; locally gravelly. Highest terraces 50 feet above Scantic River; numerous small nonpaired terraces at lower levels on both sides of rivers and streams generally included in alluvium. Some deposits as much as 50 feet thick.




Outwash deposits

Proglacial kettled to unkettled outwash grading from coarse gravel to fine sand and silt downstream. Qsd₁ deposited when ice margin stood along line from Goat Rock to Somers. Qsd₂ deposited when ice margin stood along line trending WSW from Goat Rock. Qsd₃ deposited when ice margin was along an arcuate line trending SW from Mile Oak Corner into Long Meadow and Springfield. Qsd₄ deposited when ice margin stood along a line trending SW from Wilbraham to Springfield. Qsd₅ deposited when ice margin was north of quadrangle.




Kame terrace deposits

Coarse sand and gravel deposited between ice and valley walls by meltwater at higher altitudes than outwash deposits. Internal structure indicates collapse due to deposition on or against ice. Color ranges from reddish brown in western half of quadrangle to gray in east half of quadrangle. Includes several levels of kame terrace deposits. Locally as much as 100 feet thick but generally less than 50 feet thick.




Till

Compact to loose, non-sorted to poorly-sorted, nonstratified mixture of clay, silt, sand, pebbles, cobble, and boulder sized particles. Generally reddish-brown to pale reddish-brown in west half of quadrangle; gray to light gray in east half of area. Includes small lenses of stratified drift. Thickness varies but is generally less than 20 feet. May be more than 100 feet thick under crests of some drumlins. Deposited by ice moving southward in the lowlands and southeast in the highlands in east half of quadrangle. Till in highlands slightly older than that in lowlands in west half of quadrangle.




Bedrock

Individual outcrops shown by solid color; thinly veneered bedrock and scattered small outcrops shown by ruled pattern.



Artificial fill

Artificially emplaced earth, mainly sand and gravel, locally some till. Trash fills not mapped. Most fills less than 20 feet thick but locally as much as 40 feet thick.




Crest of ice-channel filling (eskers and crevasse fillings); angle of chevron points in direction of inferred flow.



Contact

Dashed where approximately located.



Glacial striations

Arrow shows inferred direction of movement of glacier; point of observation at tip of arrowhead



Strandline of inferred glacial lake



Abandoned quarry



Sand and gravel pit