


2458

Water resources studies in Alaska

WATER-QUALITY AND GEOHYDROLOGIC DATA AT TWO SANITARY LANDFILL SITES NEAR ANCHORAGE, ALASKA

BY CHESTER ZENONE AND D.E. DONALDSON



1974
DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY
OPEN-FILE REPORT

259900


DISCUSSION

Water-quality and geohydrologic data were collected at two sanitary landfill (solid-waste disposal) sites near Anchorage, Alaska. The study was undertaken by the U.S. Geological Survey in cooperation with the Greater Anchorage Area Borough and the U.S. Air Force. The purpose of the study was to evaluate the environmental effects of solid-waste disposal on the ground-water system and establish criteria for selecting future landfill sites. This report is a compilation of the data collected through August 1973.

The objectives of the Geological Survey's program are (1) to detect the possible presence of leachate components in ground water and to obtain detailed chemical analyses, (2) to determine the rate of attenuation in concentration of any leachate component with distance from the landfill area, and (3) to determine the direction and rate of movement of ground water containing the leachate.

For this study, 14 wells, designated by the letters BSL (Borough Sanitary Landfill) and ESL (Elmendorf Sanitary Landfill), were drilled within and near the two landfill areas. The initial sampling of ground water did detect presence of leachate in the ground water within and beneath the Anchorage Borough landfill site but not in the ground water beneath the Elmendorf Air Force Base site. Drillers' logs of the wells show the Anchorage Borough site to be a wet area--the refuse is deposited below the water table in some parts of the landfill. The Elmendorf Air Force Base site is in a dry area--the refuse is deposited above the water table. This is reflected by the water-level contours shown on the maps in this report. These contours are based on measurements in the BSL, ESL, and other nearby shallow wells. At the Anchorage Borough landfill site, the contours are terminated at International Airport Road, because the roadway is a barrier to shallow ground-water flow, and there are no reliable water-level data north of the road.

The initial collection of ground-water data at both landfill sites and the detection of leachate constituents in the ground water at the Anchorage Borough site have satisfied the first objective of this study. The other objectives are of long-term nature and are being pursued by a continuing program of water-quality and water-level monitoring.



EXPLANATION

MAPS

BSL-1 (or ESL-1)

Water-quality sampling and water-level observation wells drilled specifically for this study; upper number is altitude, in feet above mean sea level, of water level in the well.

5 (through 17)

Other water-quality sampling and (or) water-level observation points near sanitary landfill sites.

76
USGS A.H.

Water-level observation auger hole; drilled by U.S. Geological Survey; number is altitude, in feet above mean sea level, of water level in the well.

79

Water-level contours showing altitude of the water table (in feet above sea level). Three different contour intervals (1 foot, 2 feet, and 20 feet) are used on the maps in this report.

A--A'

Approximate lines of sections through the landfill areas.



SECTIONS

Measured water levels in wells

Perforated interval of well casings.

FEET X 0.3048=METERS

English-Metric conversion factor (for water-level figures on maps)




Water-quality data at Elmendorf Air Force Base Sanitary Landfill and nearby area, June 22-23, 1973. (Analyses by U.S. Geological Survey, Salt Lake City, Utah)



Sample no.	Location description	Depth to water below land surface (ft)	Water temperature	Specific conductance (microhm/cm at 25 °C)	pH units	Milligrams per liter																Micrograms per liter										
						Calcium (Ca) dissolved	Magnesium (Mg) dissolved	Sodium (Na) dissolved	Potassium (K) dissolved	Bicarbonate (HCO ₃) dissolved	Chloride (Cl) dissolved	Sulfate (SO ₄) dissolved	Nitrate (NO ₃) dissolved	Fluoride (F) dissolved	Phosphate (as P) dissolved	Carbon (C), organic dissolved	Silica (SiO ₂) dissolved	Phosphorus (P), inorganic dissolved	Phosphorus (P), organic dissolved	Nitrogen, dissolved (as N)	Ammonia (NH ₃), dissolved (as N)	Aluminum (Al) dissolved	Cadmium (Cd) dissolved	Cobalt (Co) dissolved	Chromium (Cr) dissolved	Iron (Fe) dissolved	Manganese (Mn) dissolved	Nickel (Ni) dissolved				
ESL-1*	Well in landfill	37.6 (11.5)	5.0	149	7.6	20	3.2	3.0	0.3	72	1.3	16	0.29	0.0	0.0	7.8	0.00	0.01	0.00	0.00	0	0	0	24	1	14	10	0	1400	0	30	
ESL-2*	Well in landfill	4.6 (1.4)	4.0	129	7.1	17	2.6	3.0	.4	49	1.0	19	.32	.0	.5	7.8	.00	.00	.00	.00	0	0	0	0	30	3	0	10	50	270	0	20
ESL-3*	Well 200 ft (61 m) south of landfill	13.0 (4.0)	3.5	132	7.1	18	2.9	2.4	.4	52	1.0	16	.32	.0	.5	7.8	.00	.00	.04	.00	0	0	0	0	25	1	1	10	50	200	0	0
ESL-4*	Well 320 ft (98 m) west of landfill	34.5 (10.5)	3.5	156	7.6	21	3.4	2.8	.4	59	1.4	23	.29	.1	.0	7.6	.00	.00	.00	.00	0	0	0	0	18	1	0	0	40	610	10	40
5	Well nr Ship C at Boniface Parkway	5.2 (1.6)**	3.5	137	7.0	19	2.9	2.7	.4	58	0.9	18	.25	.1	.0	7.6	.00	.00	.10	.01	0	0	0	0	27	0	0	220	110	650	10	20
6	Surface water - Ship C downstream from landfill	--	6.0	119	7.3	18	2.7	2.1	.4	54	.5	13	.11	.1	1.5	6.1	.00	.01	.10	.01	10	0	0	0	17	0	0	0	119	210	0	10
7	Gallery well nr Ship C downstream from landfill	9.0 (2.7)	5.0	189	7.0	27	4.9	3.0	.5	88	1.5	18	.25	.3	0.0	9.9	.01	.20	.00	.00	10	0	0	0	21	5	1	10	0	40	0	0
8	Surface water - Ship C upstream from landfill	--	8.0	121	8.0	17	2.6	2.1	.4	55	0.8	13	.15	.0	2.0	6.4	.01	.02	.13	.02	20	0	0	0	17	1	0	10	9	190	0	0
9	Well nr U.S. Air Force Hospital	26.9 (8.2)**	8.0	246	7.8	39	6.6	3.2	.7	134	1.3	15	.30	.1	0.0	12	.01	.01	.00	.00	10	0	0	0	21	1	1	10	9	920	0	0

* Elmendorf Sanitary Landfill well.

** Water level measured April 23, 1973.



Bulldozer spreading and compacting refuse in typical landfill operation. Refuse is covered with at least 6 inches of clean fill (material at lower right) at the end of each day's operation.




Water-quality data at Greater Anchorage Area Borough Sanitary Landfill and nearby area, June 8-9, 1973. (Analyses by U.S. Geological Survey, Salt Lake City, Utah)

Sample no.	Location description	Depth to water below land surface (ft)	Water temperature (°C)	Specific conductance (microhm/cm at 25°C)	pH units	Milligrams per liter												Micrograms per liter															
						Calcium (Ca) dissolved	Magnesium (Mg) dissolved	Sodium (Na) dissolved	Potassium (K) dissolved	Bicarbonate (HCO ₃) dissolved	Chloride (Cl) dissolved	Sulfate (SO ₄) dissolved	Nitrate (NO ₃) dissolved	Fluoride (F) dissolved	Carbon (C), organic dissolved	Silica (SiO ₂) dissolved	Phosphorus (P), inorganic dissolved	Phosphorus (P), organic dissolved	Nitrogen (N), inorganic dissolved	Nitrogen (N), organic dissolved	Ammonia (NH ₃) dissolved	Aluminum (Al) dissolved	Chromium (Cr) dissolved	Cadmium (Cd) dissolved	Copper (Cu) dissolved	Selenium (Se) dissolved	Zinc (Zn) dissolved	Barium (Ba) dissolved	Iron (Fe) total	Manganese (Mn) dissolved	Nickel (Ni) dissolved		
BSL-1*	In natural area approx. 150 ft (46 m) east of fill	4 (1.2)	1.0	252	7.3	39	6.1	2.6	0.6	139	3.8	13	0.00	0.2	3.0	31	0.06	0.06	0.02	0.01	110	0	0	0	0	1	2	0	30	5,600	5,500	490	470
BSL-2*	Within landfill area nr eastern edge	4 (1.2)**	3.0	514	6.3	75	16	27	2.0	270	9.8	4.2	.04	.1	130	41	.04	.04	.28	.04	0	0	0	3	11	3	20	1,900	43,000	2,400	2,100	18,000	
BSL-3*	Within landfill area	6 (1.8)	--	5350	6.7	--	--	--	--	3510	--	--	.00	--	2480	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	760,000	--	--
BSL-4*	Within landfill area	17 (5.2)	3.0	2440	6.2	440	64	76	6.6	1380	110	8.4	.00	.3	860	31	.07	.07	.00	5.9	0	10	0	10	7	16	80	76,000	94,000	7,800	6,700		
BSL-5*	Within landfill area	18 (5.5)	3.5	2660	5.7	160	100	140	4.5	1230	170	9.2	.00	.2	1030	34	.04	.18	1.4	2.1	0	20	0	10	12	0	260	34,000	--	14,000	--	--	
BSL-6*	In natural area approx. 80 ft (24 m) west of fill	10 (3.0)	8.5	1790	6.9	270	57	97	5.6	911	210	28	.00	.2	35	22	.02	.15	1.4	0.51	0	0	0	5	5	13	70	59,000	--	4,800	--	--	
BSL-7*	In natural area approx. 800 ft (240 m) west of BSL-6	12 (3.8)	2.5	80	7.1	10	1.7	2.1	0.5	30	1.4	11	.07	.1	1.5	15	.00	.11	2.7	.01	10	0	0	3	0	0	10	220	4,300	10	90		
BSL-8*	Within landfill area; nr northern edge at International Airport Road	13 (4.0)	5.0	676	6.3	77	17	36	1.8	314	68	3.4	.00	.1	23	22	.01	.09	0.65	.55	0	0	0	4	2	2	30	2,900	--	2,000	--	--	
BSL-9*	In undisturbed area approx. 440 ft (134 m) south of fill	11 (3.4)	2.0	363	7.1	46	16	4.9	1.1	230	2.7	4.0	.00	.1	6.0	26	.02	.03	.03	.06	10	0	0	1	10	3	20	33,000	--	470	460		
BSL-10*	Nr Northwood School 750 ft (229 m) north of fill	11 (3.4)	--	212	7.6	30	7.4	3.9	1.1	125	4.3	3.4	--	.1	5.5	1.5	.00	.08	.19	.19	--	0	0	4	7	0	10	30	--	280	--	--	
11	Bentzen Lake (at surface)	--	11.0	60	7.8	2.4	1.0	1.8	1.2	12	2.8	2.0	.01	.0	6.5	0.2	.02	.03	.56	.05	--	0	0	6	28	2	0	220	--	20	--	--	
12	Delaney Lake (at surface)	--	12.0	221	7.4	25	7.8	6.9	2.3	97	11	12	.01	.1	12	.6	.00	.04	.65	.06	--	0	1	9	4	3	10	120	--	70	--	--	
13	Conners Lake (at surface)	--	5.0	116	7.5	15	3.8	2.6	0.9	63	2.7	2.0	.01	.1	8.5	.4	.00	.04	.47	.05	--	0	0	13	8	2	10	220	--	30	--	--	
14	Sloppy area south of fill (at surface)	--	12.0	79	6.5	8.0	2.4	3.5	1.5	35	3.7	3.2	--	.2	38	.4	.01	.04	1.1	.12	--	0	0	12	5	8	20	450	--	40	--	--	
15	Central Alaska Utility well	unavailable	3.0	244	7.6	31	9.7	4.4	0.8	147	1.9	7.6	.00	.2	3.0	15	.04	.04	0.00	.02	10	0	0	6	0	2	10	170	340	100	100		
16	Well at Conners Lake	30 (9.2)	5.0	260	8.0	30	8.6	9.8	3.3	165	1.4	2.9	.00	.2	3.5	29	.29	.29	.09	.31	10	0	--	48	0	0	120	220	70	80	--	--	
17	Central Alaska Utility well	unavailable	3.0	201	7.7	33	7.9	5.8	1.5	158	1.6	2.5	.00	.2	2.5	27	.29	.31	.00	.49	10	0	0	6	1	0	10	260	400	90	100	--	--

* Borough Sanitary Landfill well.

** 10 ft (3.05 m) casing added after sampling date.



M(200)
R290
10,74-1131
C.1

