

INTRODUCTION AND ACKNOWLEDGEMENTS

This map is being released in preliminary form to make the basic geologic information available during the current field season. A more comprehensive report on this area, including detailed descriptions of the mapping units, fossil collections, and igneous rock age dates, is in the final stage of completion.

The mapping is based chiefly on data from ground stations and foot traverses, but includes some field observations from helicopters and interpretations from aerial photographs. Fieldwork was done during the periods August 11-19 and September 1-5, 1970, August 17-September 11, 1971, June 15-30, 1972, and August 20-September 1, 1973 by Chapman 1970-73, Yeend 1970-72, Brosgé 1971-73, and Reiser 1972-73. The Tanana A-1, A-2, and north edge of the Kantishna River D-1 and D-2 portion of this area is taken largely from the unpublished detailed mapping of David M. Hopkins and Bond Taber that was accomplished by extensive foot traverses in 1956, 1957, and 1959. Their significant contribution is gratefully acknowledged, and the present authors assume responsibility for modifications and generalizations that we have made to reduce the 1:63,360-scale compilation of Hopkins and Taber to the scale of this map.

Some supplementary data have been obtained from field notes of R. R. Coats and W. N. Laval in 1942 and R. M. Chapman in 1943 and 1952 on the Grant-Moran Domes area; from R. R. Coats and T. G. Payne in 1942 and R. M. Chapman in 1943 on the Morelock Creek area; from H. M. Eakin in 1911, J. B. Mertie, Jr. in 1922 and 1931, and R. L. Foster in 1967 on the area between the Yukon and Tanana Rivers; and from F. R. Weber in 1962 on part of the Kantishna River D-1 quadrangle area. The thin-section identification work of Glenn J. Mac Pherson and Michael L. Throckmorton is gratefully acknowledged.

DESCRIPTION OF MAP UNITS

UNCONSOLIDATED DEPOSITS

GLACIAL

- Q<sub>1</sub> Till, cirque glaciation
- Q<sub>2</sub> Till
- Q<sub>3</sub> Till
- Q<sub>4</sub> Till, undifferentiated

ALLUVIAL

- Qal Recent alluvium
- Qaf Alluvial fan deposits
- Qbg Low-level bench gravel of Minook Creek
- Qtr Terrace deposits
- Qgs Older alluvium
- Qrg High-level gravel

COLLUVIAL AND ALLUVIAL

- Qcal Colluvium and alluvium, undifferentiated

COLLUVIAL

- Qcs Recent slide and slump deposits
- Qls Older landslide deposits
- Qcl Colluvium, undifferentiated
- Qta Talus

EOLIAN

- Ql Loess
- Qss Sand and silt

EROSIONAL SURFACES

- Qat Alluviation terrace
- Qb Rock-defended terraces on Minook Creek

BEDROCK

SEDIMENTARY AND METAMORPHIC ROCKS


- Ts Sandstone, conglomerate, shale and lignite
- Kcs Sandstone, quartzite, conglomerate, siltstone and slaty shale
- Ps Slaty shale, siltstone, graywacke and conglomerate
- Psc Siltstone, slate, phyllite and argillite
- Pc Calcareous schistose siltstone and sandstone, and some phyllite
- Pw Quartz wacke
- Psr Quartzite-gneiss schist, with some quartzite schist, calcareous schist, marble and phyllite
- Dst Toluosa Limestone
- Pl Limestone, dolomite, basaltic greenstone and chloritic schist, with some argillite, phyllite, and quartzite schist
- Qc Chert and some slaty shale
- Plic Limestone, dolomite, basaltic greenstone, chert and chloritic schist
- Gal Maroon and green argillite and slate, with quartzite, grit and some phyllite
- Psq Quartzite schist, quartzite, phyllite and slate

SEDIMENTARY, METAMORPHIC AND IGNEOUS ROCKS

- Ks Hornfels and gneiss
- Kvs Rhyolitic lava and breccia, tuff, chert and shale
- RPv Extrusive and intrusive basaltic and diabasic rocks, tuff, chert, argillite, slate and rarely clastic limestone
- Bvs Foliated basaltic lava, tuff, slaty shale, phyllite and some limestone and chert

IGNEOUS ROCKS

- Qta Andesitic lava
- Tg Granite
- TRV Felsic volcanic rock
- Kg Granitic rocks
- Ka Gabbro and diorite
- Ksp Saponatite with some diabase, gabbro and mafic volcaniclastic rocks



EXPLANATION OF SYMBOLS

- Contact: solid where observed or approximately known; dashed and where inferred. May include some faults.
- Fault or zone of faulting approximately located or inferred; dotted where inferred beneath unconsolidated deposits. Arrows show directions of inferred lateral movement; U and D indicate up and down sides where vertical movement is inferred.
- Anticline or synform approximately located or inferred, showing direction of plunge.
- Syncline or synform approximately located or inferred, showing direction of plunge; dotted where inferred beneath unconsolidated deposits.
- Strike and dip of beds; may include some overturned beds.
- Strike of vertical beds.
- Strike and dip of beds interpreted from distant observations or aerial photographs.
- Strike and dip of foliation, inclined and vertical.
- Flowing spring, hot or warm.
- Fossil locality; F1 through F8 are pollen and/or plant material; F9 is foraminifera, and F10 is a rugose coral.

SELECTED REFERENCES

Brosgé, W. P., Lamphere, M. A., Reiser, R. N., and Chapman, R. M., 1969, Probable Pliocene age of the Rampart Group, central Alaska: U.S. Geol. Survey Bull. 1294-B, p. 31-318.

Chapman, R. M., Coats, R. R., and Payne, T. G., 1963, Placer tin deposits in central Alaska: U.S. Geol. Survey open-file report, 53 p.

Chapman, R. M., Weber, F. R., and Taber, Bond, 1971, Preliminary geologic map of the Livengood quadrangle, Alaska: U.S. Geol. Survey open-file report.

Chapman, R. M., and Yeend, W. E., 1972, Preliminary geologic map of the northeastern part of the Tanana quadrangle, Alaska: U.S. Geol. Survey Misc. Field Studies Map MF-342.

Cobb, E. H., 1972, Metallic mineral resources map of the Tanana quadrangle, Alaska: U.S. Geol. Survey Bull. 535, 36 p.

Eakin, H. M., 1913, A geologic reconnaissance of a part of the Rampart quadrangle, Alaska: U.S. Geol. Survey Bull. 535, 36 p.

1916, The Yukon-Koyukuk region, Alaska: U.S. Geol. Survey Bull. 531, 88 p.

Godsey, L. D., Shapiro, L. H., Van Vorster, J. D., and Weber, F. R., 1972, Correlation of epicenters with mapped faults, east-central Alaska, 1960-1971: U.S. Geol. Survey open-file report, 7 p.

MacNeil, F. S., Wolfe, J. A., Miller, D. J., and Hopkins, D. M., 1961, Correlation of Tertiary formations of Alaska: Am. Assoc. Petroleum Geologists Bull., v. 45, no. 11, p. 1801-1809.

Mertie, J. B., Jr., 1937, The Yukon-Tanana region, Alaska: U.S. Geol. Survey Bull. 872, 276 p.

Paige, R. A., 1959, Tertiary geology of the Cheyenne Creek area, Alaska: Seattle, Washington Univ. M.S. thesis, 66 p.

Fallon, W. M., Jr., and Miller, T. P., 1970, Preliminary geologic investigation in the Seward River region, Alaska: U.S. Geol. Survey Bull. 1312-J, p. 11-110.

1973, Bedrock geologic map of Bettles and southern part of Wiseman quadrangles, Alaska: U.S. Geol. Survey Misc. Field Studies Map MF-692.

Péwé, T. L., Wahrhaftig, Clyde, and Weber, F. R., 1966, Geologic map of the Fairbanks quadrangle, Alaska: U.S. Geol. Survey Misc. Geol. Inv. Map I-455.


U.S. Geological Survey, 1972, Geological Survey research 1972: U.S. Geol. Survey Prof. Paper 800-A, p. A31-A52.

1973a, Aeromagnetic survey, Tanana quadrangle, northeast Alaska: U.S. Geol. Survey open-file report, scale 1:250,000.

1973b, United States Geological Survey Alaska program, 1973: U.S. Geol. Survey Circ. 683, p. 17-18, 43.

Yeend, W. E., 1971, Glaciation of the Ray Mountains, central Alaska, in Geological Survey Research 1971: U.S. Geol. Survey Prof. Paper 750-D, p. D122-D126.

CORRELATION OF MAP UNITS



Base from U.S.G.S. 1:250,000 topo series: TANANA, 1956; KANTISHNA RIVER, 1952; 200 feet; ALASKA. Compiled by Menlo Park Base Map Unit. (1-73) (8-22)

SCALE 1:250,000

SCALE 1:250,000

PRELIMINARY GEOLOGIC MAP OF THE TANANA AND NORTHEAST PART OF THE KANTISHNA RIVER QUADRANGLES, ALASKA  
BY  
ROBERT M. CHAPMAN, WARREN E. YEEND, WILLIAM P. BROSGÉ, AND HILLARD N. REISER  
1975

This map is preliminary and has not been reviewed for conformity with U.S. Geological Survey standards and nomenclature.